

Mathematical Methods - III

Week 1

Jayampathy Ratnayake

Faculty of Science
Department of Mathematics
Room 209(B)
jratnaya@indiana.edu

August 18, 2017

This course has can be divided into two parts

This course has can be divided into two parts

- Numerical Methods

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing
 - How to solve complex mathematical problems using computers

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing
 - How to solve complex mathematical problems using computers
 - why?

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing
 - How to solve complex mathematical problems using computers
 - why?
 - analytical solutions are unknown or extremely difficult to find

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing
 - How to solve complex mathematical problems using computers
 - why?
 - analytical solutions are unknown or extremely difficult to find
 - even if the analytical solutions are known, it may be too difficult to carry out computations without computers

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing
 - How to solve complex mathematical problems using computers
 - why?
 - analytical solutions are unknown or extremely difficult to find
 - even if the analytical solutions are known, it may be too difficult to carry out computations without computers
 - numerical methods may produce sufficiently accurate results, while using less resources

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing
 - How to solve complex mathematical problems using computers
 - why?
 - analytical solutions are unknown or extremely difficult to find
 - even if the analytical solutions are known, it may be too difficult to carry out computations without computers
 - numerical methods may produce sufficiently accurate results, while using less resources
 - ...
- Algebra

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing
 - How to solve complex mathematical problems using computers
 - why?
 - analytical solutions are unknown or extremely difficult to find
 - even if the analytical solutions are known, it may be too difficult to carry out computations without computers
 - numerical methods may produce sufficiently accurate results, while using less resources
 - ...
- Algebra
 - Introduction to Groups and Rings (Abstract Algebra)

This course has can be divided into two parts

- Numerical Methods
 - Scientific computing
 - How to solve complex mathematical problems using computers
 - why?
 - analytical solutions are unknown or extremely difficult to find
 - even if the analytical solutions are known, it may be too difficult to carry out computations without computers
 - numerical methods may produce sufficiently accurate results, while using less resources
 - ...
- Algebra
 - Introduction to Groups and Rings (Abstract Algebra)
 - We will look at some interesting applications in computer science.

Part I:

- 1 Nonlinear Equations
- 2 Interpolation, Curve Fitting and Approximating Functions
- 3 Numerical Integration
- 4 Numerical Solutions to ODEs

Part I:

- 1 Nonlinear Equations
- 2 Interpolation, Curve Fitting and Approximating Functions
- 3 Numerical Integration
- 4 Numerical Solutions to ODEs

Part II:

- 1 Modulo Arithmetic
- 2 Euclidian Algorithm
- 3 Groups
- 4 Lagrange's Theorem
- 5 RSA Algorithm
- 6 Polynomial Rings

- We will start with "Numerical Methods"

- We will start with "Numerical Methods"
- There are many free online resources / references available

- We will start with "Numerical Methods"
- There are many free online resources / references available
E.g:

- We will start with "Numerical Methods"
- There are many free online resources / references available
E.g:
 - *Numerical methods that work*. Forman S. Acton.
 - *MIT Open Course Ware*
 - http://www.mathcs.emory.edu/~ale/NAbook_Aug_2008.pdf

- We will start with "Numerical Methods"
- There are many free online resources / references available
E.g:
 - *Numerical methods that work*. Forman S. Acton.
 - *MIT Open Course Ware*
 - http://www.mathcs.emory.edu/~ale/NAbook_Aug_2008.pdf
- Students are encouraged to use these resources as references for this course.

- Many numerical computations, particularly when using computers, generate sequences of approximations.

- Many numerical computations, particularly when using computers, generate sequences of approximations.
- To understand these methods, their behaviour and usage, we need to know how these sequence can be understood in terms of their:

- Many numerical computations, particularly when using computers, generate sequences of approximations.
- To understand these methods, their behaviour and usage, we need to know how these sequence can be understood in terms of their:
 - Convergence/ Divergence

- Many numerical computations, particularly when using computers, generate sequences of approximations.
- To understand these methods, their behaviour and usage, we need to know how these sequence can be understood in terms of their:
 - Convergence/ Divergence
 - Rate of convergence

- Many numerical computations, particularly when using computers, generate sequences of approximations.
- To understand these methods, their behaviour and usage, we need to know how these sequence can be understood in terms of their:
 - Convergence/ Divergence
 - Rate of convergence

Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

A numerical approach can be taken by

- dividing the interval in to some number of subintervals
- in each subinterval, approximate the area by an rectangle


Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

A numerical approach can be taken by

- dividing the interval in to some number of subintervals
- in each subinterval, approximate the area by an rectangle

Also, observe that our results gets better and better as the divisions gets finer and finer.



Example continued...

Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

Example continued...

Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

- Let n be the number of divisions.

Example continued...

Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

- Let n be the number of divisions.
- Choose the function value at the midpoint of each subinterval as the height of the rectangle.

Example continued...

Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

- Let n be the number of divisions.
- Choose the function value at the midpoint of each subinterval as the height of the rectangle.
- Compute the approximate areas given by the method for $n = 1, 2, 3$ and 4.

Example continued...

Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

- Let n be the number of divisions.
- Choose the function value at the midpoint of each subinterval as the height of the rectangle.
- Compute the approximate areas given by the method for $n = 1, 2, 3$ and 4.
- What properties do you observe in the sequence generated by this scheme?

Example continued...

Example

Approximate the area bounded by $f(x) = \frac{\sin(x)}{x}$, $x = 1$ and $x = 3$.

- Let n be the number of divisions.
- Choose the function value at the midpoint of each subinterval as the height of the rectangle.
- Compute the approximate areas given by the method for $n = 1, 2, 3$ and 4.
- What properties do you observe in the sequence generated by this scheme?

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- 1 Bounded above:
if there is some number K such that $a_n \leq K$ for all $n \in \mathbb{N}$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- ① Bounded above:
if there is some number K such that $a_n \leq K$ for all $n \in \mathbb{N}$.
- ② Bounded below:
if there is some number k such that $k \leq a_n$ for all $n \in \mathbb{N}$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- ① Bounded above:
if there is some number K such that $a_n \leq K$ for all $n \in \mathbb{N}$.
- ② Bounded below:
if there is some number k such that $k \leq a_n$ for all $n \in \mathbb{N}$.
- ③ Bounded:
if it is both bounded above and bounded below.
I.e. there are some numbers K and k such that $k \leq a_n \leq K$ for all $n \in \mathbb{N}$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- ① Bounded above:
if there is some number K such that $a_n \leq K$ for all $n \in \mathbb{N}$.
- ② Bounded below:
if there is some number k such that $k \leq a_n$ for all $n \in \mathbb{N}$.
- ③ Bounded:
if it is both bounded above and bounded below.
I.e. there are some numbers K and k such that $k \leq a_n \leq K$ for all $n \in \mathbb{N}$.
Equivalently, $(a_n)_{n \in \mathbb{N}}$ is bounded if there is some M such that $|a_n| \leq M$ for all $n \in \mathbb{N}$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- ① Bounded above:
if there is some number K such that $a_n \leq K$ for all $n \in \mathbb{N}$.
- ② Bounded below:
if there is some number k such that $k \leq a_n$ for all $n \in \mathbb{N}$.
- ③ Bounded:
if it is both bounded above and bounded below.
I.e. there are some numbers K and k such that $k \leq a_n \leq K$ for all $n \in \mathbb{N}$.
Equivalently, $(a_n)_{n \in \mathbb{N}}$ is bounded if there is some M such that $|a_n| \leq M$ for all $n \in \mathbb{N}$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- 1 Monotonically increasing:
if $a_n \leq a_{n+1}$ for all $n \in \mathbb{N}$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- ① Monotonically increasing:
if $a_n \leq a_{n+1}$ for all $n \in \mathbb{N}$.
- ② Monotonically decreasing:
if $a_n \geq a_{n+1}$ for all $n \in \mathbb{N}$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- ① Monotonically increasing:
if $a_n \leq a_{n+1}$ for all $n \in \mathbb{N}$.
- ② Monotonically decreasing:
if $a_n \geq a_{n+1}$ for all $n \in \mathbb{N}$.
- ③ Monotonic:
if it is either Monotonically increasing or decreasing.
- ④ Eventually Monotonically increasing:
if there is some $k \in \mathbb{N}$ such that $a_n \leq a_{n+1}$ for all $n \geq k$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- ① Monotonically increasing:
if $a_n \leq a_{n+1}$ for all $n \in \mathbb{N}$.
- ② Monotonically decreasing:
if $a_n \geq a_{n+1}$ for all $n \in \mathbb{N}$.
- ③ Monotonic:
if it is either Monotonically increasing or decreasing.
- ④ Eventually Monotonically increasing:
if there is some $k \in \mathbb{N}$ such that $a_n \leq a_{n+1}$ for all $n \geq k$.
- ⑤ Eventually Monotonically decreasing:
if there is some $k \in \mathbb{N}$ such that $a_n \geq a_{n+1}$ for all $n \geq k$.

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence of numbers. We say it is

Definition

- 1 Monotonically increasing:
if $a_n \leq a_{n+1}$ for all $n \in \mathbb{N}$.
- 2 Monotonically decreasing:
if $a_n \geq a_{n+1}$ for all $n \in \mathbb{N}$.
- 3 Monotonic:
if it is either Monotonically increasing or decreasing.
- 4 Eventually Monotonically increasing:
if there is some $k \in \mathbb{N}$ such that $a_n \leq a_{n+1}$ for all $n \geq k$.
- 5 Eventually Monotonically decreasing:
if there is some $k \in \mathbb{N}$ such that $a_n \geq a_{n+1}$ for all $n \geq k$.
- 6 Eventually Monotonic:
if it is either Eventually Monotonically increasing or decreasing.

We are primarily interested in the convergence properties of sequences generated by various numerical schemes.

We are primarily interested in the convergence properties of sequences generated by various numerical schemes.

Definition

We say $(a_n)_{n \in \mathbb{N}}$ converge to L and write $\lim_{n \rightarrow \infty} a_n = L$ if

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } |a_n - L| \leq \epsilon \text{ for all } n \geq N$$

Down sides of the above definition are

Down sides of the above definition are

- 1 Need to know L to prove the convergence.

- 1 Need to know L to prove the convergence.
Many numerical schemes generate sequences whose limit is unknown.

Down sides of the above definition are

- ① Need to know L to prove the convergence.
 Many numerical schemes generate sequences whose limit is unknown.
- ② Need to know the general formula for a_n .

Down sides of the above definition are

- ① Need to know L to prove the convergence.
Many numerical schemes generate sequences whose limit is unknown.
- ② Need to know the general formula for a_n .
For many numerical schemes general formula is not available or inconvenient to work with.

Down sides of the above definition are

- ① Need to know L to prove the convergence.
Many numerical schemes generate sequences whose limit is unknown.
- ② Need to know the general formula for a_n .
For many numerical schemes general formula is not available or inconvenient to work with.

We need alternative ways to get our hands on the convergence.

Theorem

if (a_n) is Monotonically increasing and Bounded above, then the sequence converge (to the supremum).

Theorem

if (a_n) is Monotonically increasing and Bounded above, then the sequence converge (to the supremum).

Theorem

if (a_n) is Monotonically decreasing and Bounded below, then the sequence converge (to the infimum).

Example

Consider the sequence given by

$$x_1 = 1, \quad x_{n+1} = 3 - \frac{1}{x_n}$$

- ① Show that the sequence is both bounded above and bounded below.
- ② Show, using induction, that the sequence is monotonically increasing.
- ③ Conclude that the sequence converge to some limit and find it.
- ④ Explain how you can use this sequence to approximate the value of $\sqrt{5}$.

Consider the sequence given by $x_1 = 1$ and $x_{n+1} = \frac{1}{2} \left(\frac{3}{x_n^2} + x_n \right)$.

- ① Is the sequence monotonic (i.e. monotonically decreasing or increasing)?
- ② Is the sequence bounded below and/or bounded above?
- ③ Show that the sequence converge to some limit.
- ④ What is the limit of this sequence?

Definition

The sequence $(a_n)_{n \in \mathbb{N}}$ is said to be Cauchy if

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } |a_n - a_m| \leq \epsilon \text{ for all } n, m \geq N$$

Definition

The sequence $(a_n)_{n \in \mathbb{N}}$ is said to be Cauchy if

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } |a_n - a_m| \leq \epsilon \text{ for all } n, m \geq N$$

We have the following fundamental result.

Definition

The sequence $(a_n)_{n \in \mathbb{N}}$ is said to be Cauchy if

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } |a_n - a_m| \leq \epsilon \text{ for all } n, m \geq N$$

We have the following fundamental result.

Theorem

$(a_n)_{n \in \mathbb{N}}$ converge to some limit if and only if $(a_n)_{n \in \mathbb{N}}$ is Cauchy.

Definition

The sequence $(a_n)_{n \in \mathbb{N}}$ is said to be Cauchy if

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } |a_n - a_m| \leq \epsilon \text{ for all } n, m \geq N$$

We have the following fundamental result.

Theorem

$(a_n)_{n \in \mathbb{N}}$ converge to some limit if and only if $(a_n)_{n \in \mathbb{N}}$ is Cauchy.

Fundamental importance of this result is, by showing a sequence is Cauchy one can show it converge without knowing the actual limit.

Consider a continuous function f and two points a and b such that $f(a)$ and $f(b)$ have opposite signs. A sequence (c_n) is constructed as follows.

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.
- Consider the midpoint between a and b ; $c = \frac{a+b}{2}$. Set $c_1 = c$.
- Not all three of $f(a)$, $f(b)$ and $f(c)$ have the same sign. Pick the midpoint and the point with the opposite sign. Take the mid-point of the chosen points to be the next term in the sequence (c_n)
- Repeat the process.

Example

Let $f(x) = -x^5 + 2x + 2 = 0$, $a = 0$ and $b = 2$.

- 1 Find the first few terms of the sequence (x_n) generated by this recipe.
- 2 Show that $|c_n - c_{n+1}| = \alpha^n$ for some α and explicitly mention the value of α .
- 3 Show that this sequence is Cauchy.
- 4 To which value does the sequence converge?

Given a series

$$x_1 + x_2 + \dots + x_i + \dots = \sum_{i=1}^{\infty} x_i$$

Definition

We say the sum is L and write $\sum_{i=1}^{\infty} x_i = L$ if the sequence of n -partial sums $s_n = x_1 + x_2 + \dots + x_n = \sum_{i=1}^n x_i$ converge to L .

- 1 This course will have weekly Homework.
- 2 HW1: Complete the first tutorial.