

Mathematical Methods - III

Week 2

Jayampathy Ratnayake

Faculty of Science
Department of Mathematics
Room 209(B)
jratnaya@indiana.edu

August 18, 2017

solutions to non linear-equations

In this lesson, we will ...

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.
- Use the Bisection method

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.
- Use the Bisection method to to solve non-liner equations.

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.
- Use the Bisection method to to solve non-liner equations.
- Learn the notion of Errors and Convergence of a numerical scheme.

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.
- Use the Bisection method to to solve non-liner equations.
- Learn the notion of Errors and Convergence of a numerical scheme.
- Analyze the errors

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.
- Use the Bisection method to to solve non-liner equations.
- Learn the notion of Errors and Convergence of a numerical scheme.
- Analyze the errors associated with the bisection method and

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.
- Use the Bisection method to to solve non-liner equations.
- Learn the notion of Errors and Convergence of a numerical scheme.
- Analyze the errors associated with the bisection method and order of convergence.

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations and Give Examples.
- Use the Bisection method to solve non-linear equations.
- Learn the notion of Errors and Convergence of a numerical scheme.
- Analyze the errors associated with the bisection method and order of convergence.
- Discuss other techniques such as

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.
- Use the Bisection method to to solve non-liner equations.
- Learn the notion of Errors and Convergence of a numerical scheme.
- Analyze the errors associated with the bisection method and order of convergence.
- Discuss other techniques such has
 - fixed point iteration and

solutions to non linear-equations

In this lesson, we will ...

- Define the non-linear equations.and Give Examples.
- Use the Bisection method to to solve non-liner equations.
- Learn the notion of Errors and Convergence of a numerical scheme.
- Analyze the errors associated with the bisection method and order of convergence.
- Discuss other techniques such has
 - fixed point iteration and
 - Newton-Raphson method.

Consider the following example from the previous lecture.

Consider the following example from the previous lecture.

Consider a continuous function f and two points a and b such that $f(a)$ and $f(b)$ have opposite signs. A sequence (c_n) is constructed as follows.

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.
- Consider the midpoint between a and b ; $c = \frac{a+b}{2}$. Set $c_1 = c$.
- Not all three of $f(a)$, $f(b)$ and $f(c)$ have the same sign. Pick the midpoint and the point with the opposite sign. Take the mid-point of the chosen points to be the next term in the sequence (c_n)
- Repeat the process.

Example

Let $f(x) = -x^5 + 2x + 2 = 0$, $a = 0$ and $b = 2$.

- 1 Find the first few terms of the sequence (x_n) generated by this recipe.
- 2 What does this sequence approximate?
- 3 How can we guarantee that this is finding what we expect to find?
- 4 Show that $|c_n - c_{n+1}| = \alpha^n$ for some α and explicitly mention the value of α .
- 5 Show that this sequence is Cauchy.
- 6 To which value does the sequence converge? Justify your answer.

Given a function $f : \mathbb{R} \rightarrow \mathbb{R}$,

Given a function $f : \mathbb{R} \rightarrow \mathbb{R}$,
we wish to find a solution to the problem $f(x) = 0$.

Given a function $f : \mathbb{R} \rightarrow \mathbb{R}$,
we wish to find a solution to the problem $f(x) = 0$.
I.e., an x^* such that $f(x^*) = 0$.

Given a function $f : \mathbb{R} \rightarrow \mathbb{R}$,
we wish to find a solution to the problem $f(x) = 0$.
I.e., an x^* such that $f(x^*) = 0$.
An x^* such that $f(x^*) = 0$ is called a **zero** or a

Given a function $f : \mathbb{R} \rightarrow \mathbb{R}$,
we wish to find a solution to the problem $f(x) = 0$.
I.e., an x^* such that $f(x^*) = 0$.
An x^* such that $f(x^*) = 0$ is called a **zero** or a **root** of f .

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method:

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method:

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method:

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.
- Consider the midpoint of a and b , $c = \frac{a + b}{2}$

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method:

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.
- Consider the midpoint of a and b , $c = \frac{a+b}{2}$
- Not all three of $f(a)$, $f(b)$ and $f(c)$ have the same sign.

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method:

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.
- Consider the midpoint of a and b , $c = \frac{a+b}{2}$
- Not all three of $f(a)$, $f(b)$ and $f(c)$ have the same sign.
- Pick the midpoint c and the endpoint with the opposite sign.

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method:

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.
- Consider the midpoint of a and b , $c = \frac{a + b}{2}$
- Not all three of $f(a)$, $f(b)$ and $f(c)$ have the same sign.
- Pick the midpoint c and the endpoint with the opposite sign.
- Repeat the process with the new interval whose endpoints are the ones that was chosen in the above step.

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method:

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.
- Consider the midpoint of a and b , $c = \frac{a + b}{2}$
- Not all three of $f(a)$, $f(b)$ and $f(c)$ have the same sign.
- Pick the midpoint c and the endpoint with the opposite sign.
- Repeat the process with the new interval whose endpoints are the ones that was chosen in the above step.
- In each step, take the mid-point of the chosen interval to be the approximated solution.

Bisection Method

Here, we discuss a simple, yet effective, way of finding roots of a **continuous** function f , based on some initial information.

Bisection Method:

- Begin with two points, a and b , such that $f(a)$ and $f(b)$ have opposite signs.
- Consider the midpoint of a and b , $c = \frac{a + b}{2}$
- Not all three of $f(a)$, $f(b)$ and $f(c)$ have the same sign.
- Pick the midpoint c and the endpoint with the opposite sign.
- Repeat the process with the new interval whose endpoints are the ones that was chosen in the above step.
- In each step, take the mid-point of the chosen interval to be the approximated solution.

Example

Use the bisection method to find a solution to the equation
 $f(x) = \sin(x) - x^3 + 2x + 2 = 0$.

Problem

Try applying bisection method to find a solution to $x - \frac{1}{x^2} + 2 = 0$. Take $[-2, 1]$ to be the initial interval.

Problem

Try applying bisection method to find a solution to $x - \frac{1}{x^2} + 2 = 0$. Take $[-2, 1]$ to be the initial interval.
What are your observations?

- When does Bisection method does not fail/fail?

- When does Bisection method does not fail/fail?
- Why does it work?

- When does Bisection method does not fail/fail?
- Why does it work?
- What are the errors associated?

Theory behind the Bisection Method

Theorem (Intermediate Value Theorem (IVT))

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **continuous** function.

Theory behind the Bisection Method

Theorem (Intermediate Value Theorem (IVT))

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **continuous** function.

For any α and β in $[a, b]$,

Theory behind the Bisection Method

Theorem (Intermediate Value Theorem (IVT))

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **continuous** function.

For any α and β in $[a, b]$, f achieves every value in between $f(\alpha)$ and $f(\beta)$.

Theory behind the Bisection Method

Theorem (Intermediate Value Theorem (IVT))

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **continuous** function.

For any α and β in $[a, b]$, f achieves every value in between $f(\alpha)$ and $f(\beta)$.

I.e. For any c between $f(\alpha)$ and $f(\beta)$, there is a γ between α and β such that $f(\gamma) = c$.

Theory behind the Bisection Method

Theorem (Intermediate Value Theorem (IVT))

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **continuous** function.

For any α and β in $[a, b]$, f achieves every value in between $f(\alpha)$ and $f(\beta)$.

I.e. For any c between $f(\alpha)$ and $f(\beta)$, there is a γ between α and β such that $f(\gamma) = c$.

Corollary

If $f(a)$ and $f(b)$ has opposite signs,

Theory behind the Bisection Method

Theorem (Intermediate Value Theorem (IVT))

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **continuous** function.

For any α and β in $[a, b]$, f achieves every value in between $f(\alpha)$ and $f(\beta)$.

I.e. For any c between $f(\alpha)$ and $f(\beta)$, there is a γ between α and β such that $f(\gamma) = c$.

Corollary

If $f(a)$ and $f(b)$ has opposite signs, then f has a zero in (a, b) .

Theory behind the Bisection Method

Theorem (Intermediate Value Theorem (IVT))

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **continuous** function.

For any α and β in $[a, b]$, f achieves every value in between $f(\alpha)$ and $f(\beta)$.

I.e. For any c between $f(\alpha)$ and $f(\beta)$, there is a γ between α and β such that $f(\gamma) = c$.

Corollary

If $f(a)$ and $f(b)$ has opposite signs, then f has a zero in (a, b) .

I.e. There is a $c \in (a, b)$ such that $f(c) = 0$.

Theory behind the Bisection Method

Theorem (Intermediate Value Theorem (IVT))

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **continuous** function.

For any α and β in $[a, b]$, f achieves every value in between $f(\alpha)$ and $f(\beta)$.

I.e. For any c between $f(\alpha)$ and $f(\beta)$, there is a γ between α and β such that $f(\gamma) = c$.

Corollary

If $f(a)$ and $f(b)$ has opposite signs, then f has a zero in (a, b) .

I.e. There is a $c \in (a, b)$ such that $f(c) = 0$.

Remark

Bisection method is guaranteed to work only when the function is continuous in the initial interval.

Example

- Show that the function $f(x) = \cos x - x$ has a zero in $(-\frac{\pi}{2}, \frac{\pi}{2})$.

Example

- Show that the function $f(x) = \cos x - x$ has a zero in $(-\frac{\pi}{2}, \frac{\pi}{2})$.
- Use the Bisection algorithm to estimate a root. Start from the above interval.

Example

- Show that the function $f(x) = \cos x - x$ has a zero in $(-\frac{\pi}{2}, \frac{\pi}{2})$.
- Use the Bisection algorithm to estimate a root. Start from the above interval.
 - What is the initial guess for the root? $c_0 =$

Example

- Show that the function $f(x) = \cos x - x$ has a zero in $(-\frac{\pi}{2}, \frac{\pi}{2})$.
- Use the Bisection algorithm to estimate a root. Start from the above interval.
 - What is the initial guess for the root? $c_0 =$
 - What is the approximated root after the first iterate? $c_1 =$

Example

- Show that the function $f(x) = \cos x - x$ has a zero in $(-\frac{\pi}{2}, \frac{\pi}{2})$.
- Use the Bisection algorithm to estimate a root. Start from the above interval.
 - What is the initial guess for the root? $c_0 =$
 - What is the approximated root after the first iterate? $c_1 =$
 - What is the approximated root after the second iterate? $c_2 =$

Example

- Show that the function $f(x) = \cos x - x$ has a zero in $(-\frac{\pi}{2}, \frac{\pi}{2})$.
- Use the Bisection algorithm to estimate a root. Start from the above interval.
 - What is the initial guess for the root? $c_0 =$
 - What is the approximated root after the first iterate? $c_1 =$
 - What is the approximated root after the second iterate? $c_2 =$
...
- What can you say about the error after n iterates?

Example

- Show that the function $f(x) = \cos x - x$ has a zero in $(-\frac{\pi}{2}, \frac{\pi}{2})$.
- Use the Bisection algorithm to estimate a root. Start from the above interval.
 - What is the initial guess for the root? $c_0 =$
 - What is the approximated root after the first iterate? $c_1 =$
 - What is the approximated root after the second iterate? $c_2 =$
...
- What can you say about the error after n iterates?
- Find the number of iterates required to achieve an accuracy of 10^{-k} .

Errors

- 1 Above numerical methods generates a sequence of solutions (indexed by n).

Errors

- 1 Above numerical methods generates a sequence of solutions (indexed by n).
- 2 We expect the accuracy of the solution to be greater (Error to become smaller and smaller) as we increase n .

Errors

- 1 Above numerical methods generates a sequence of solutions (indexed by n).
- 2 We expect the accuracy of the solution to be greater (Error to become smaller and smaller) as we increase n .
- 3 We can iterate n -times to guarantee that the solution has reached to a desired accuracy

Errors

- 1 Above numerical methods generates a sequence of solutions (indexed by n).
- 2 We expect the accuracy of the solution to be greater (Error to become smaller and smaller) as we increase n .
- 3 We can iterate n -times to guarantee that the solution has reached to a desired accuracy (when the numerical scheme converge).

Absolute error = $| \text{true value} - \text{estimated value} |$

Absolute error = $| \text{true value} - \text{estimated value} |$

$$\text{Relative error} = \frac{| \text{true value} - \text{approximation} |}{| \text{true value} |} \times 100$$

Absolute error = | true value - estimated value |

$$\text{Relative error} = \frac{|\text{true value} - \text{approximation}|}{|\text{true value}|} \times 100$$

Remark

A disadvantage of the above definitions is that one needs to know the true value to compute the error.

Sources of Errors in Numerical Calculations

Sources of Errors in Numerical Calculations

- Rounding Errors - due to limited number of significant figures in a computer.

Sources of Errors in Numerical Calculations

- Rounding Errors - due to limited number of significant figures in a computer.
- Truncation Errors - due to finite approximations in the numerical scheme

Sources of Errors in Numerical Calculations

- Rounding Errors - due to limited number of significant figures in a computer.
- Truncation Errors - due to finite approximations in the numerical scheme

Sources of Errors in Numerical Calculations

- Rounding Errors - due to limited number of significant figures in a computer.
- Truncation Errors - due to finite approximations in the numerical scheme
- ❶ It is important to know the error estimates of our numerical schemes.

Sources of Errors in Numerical Calculations

- Rounding Errors - due to limited number of significant figures in a computer.
- Truncation Errors - due to finite approximations in the numerical scheme
- ① It is important to know the error estimates of our numerical schemes.
- ② Errors will give us an idea of what we compute, compared to the exact solution.

Sources of Errors in Numerical Calculations

- Rounding Errors - due to limited number of significant figures in a computer.
- Truncation Errors - due to finite approximations in the numerical scheme
- ① It is important to know the error estimates of our numerical schemes.
- ② Errors will give us an idea of what we compute, compared to the exact solution.

Important Notions in Numerical Analysis

- Convergence.

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?
 - I.e., Does the Errors $e_n = |c_n - c^*|$ converge to zero?

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?
 - I.e., Does the Errors $e_n = |c_n - c^*|$ converge to zero?
- Order of Convergence?

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?
 - I.e., Does the Errors $e_n = |c_n - c^*|$ converge to zero?
- Order of Convergence?
 - How fast the numerical scheme is converging or how fast the error is decaying.

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?
 - I.e., Does the Errors $e_n = |c_n - c^*|$ converge to zero?
- Order of Convergence?
 - How fast the numerical scheme is converging or how fast the error is decaying.
 - What is it for the Bisection method?

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?
 - I.e., Does the Errors $e_n = |c_n - c^*|$ converge to zero?
- Order of Convergence?
 - How fast the numerical scheme is converging or how fast the error is decaying.
 - What is it for the Bisection method?

Other notions:

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?
 - I.e., Does the Errors $e_n = |c_n - c^*|$ converge to zero?
- Order of Convergence?
 - How fast the numerical scheme is converging or how fast the error is decaying.
 - What is it for the Bisection method?

Other notions:

- Consistency.

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?
 - I.e., Does the Errors $e_n = |c_n - c^*|$ converge to zero?
- Order of Convergence?
 - How fast the numerical scheme is converging or how fast the error is decaying.
 - What is it for the Bisection method?

Other notions:

- Consistency.
- Stability.

Important Notions in Numerical Analysis

- Convergence.
 - Does the sequence (c_n) converge to some limit?
 - I.e., Does the Errors $e_n = |c_n - c^*|$ converge to zero?
- Order of Convergence?
 - How fast the numerical scheme is converging or how fast the error is decaying.
 - What is it for the Bisection method?

Other notions:

- Consistency.
- Stability.

Let S_n be a sequence of approximations converging to L (obtained by a discretization with decreasing step sizes h_n).

Definition

We say that $e_n = |S_n - L|$ is $O(f(n))$, if there is a constant C such that

$$|S_n - L| \leq Cf(n) \text{ for all } n.$$

We write

$$\text{Error} = e_n = |S_n - L| = O(f(n)).$$

- We say S_n converge to L with order p , if $f(n) = n^{-p}$.
p=2 : Quadratic Convergence.
p=3 : Cubic Convergence.

- We say S_n converge to L with order p , if $f(n) = n^{-p}$.
 $p=2$: Quadratic Convergence.
 $p=3$: Cubic Convergence.
- Errors exponentially decay if $f(n) = 2^{-n}$.

- We say S_n converge to L with order p , if $f(n) = n^{-p}$.
 $p=2$: Quadratic Convergence.
 $p=3$: Cubic Convergence.
- Errors exponentially decay if $f(n) = 2^{-n}$.

- Does the bisection method always converge?

- Does the bisection method always converge?
- What is the order of convergence of the Bisection Method.

- Does the bisection method always converge?
- What is the order of convergence of the Bisection Method.
- What are the drawbacks of this method?

Error Analysis for the Bisection Method

Suppose we apply the Bisection Method to the function f starting with interval $[a, b]$.

- 1 Find the maximum possible error after n iterates.
- 2 Show that when f is continuous Bisection method always converge.
- 3 Find the order of convergence of the Bisection method.

Problem

Find a root of the function $f(x) = x^3 - x + 1$ with error less than 10^{-3} .

Solve the following problems and give error bounds.

- $f(x) = \ln x = 0$

Solve the following problems and give error bounds.

- $f(x) = \ln x = 0$
- $f(x) = \ln x + x = 0$

Solve the following problems and give error bounds.

- $f(x) = \ln x = 0$
- $f(x) = \ln x + x = 0$
- $f(x) = \cos x - x = 0$

Solve the following problems and give error bounds.

- $f(x) = \ln x = 0$
- $f(x) = \ln x + x = 0$
- $f(x) = \cos x - x = 0$
- $f(x) = \sqrt{x^3 - 5} = 0$

Solve the following problems and give error bounds.

- $f(x) = \ln x = 0$
- $f(x) = \ln x + x = 0$
- $f(x) = \cos x - x = 0$
- $f(x) = \sqrt{x^3 - 5} = 0$
- $f(x) = x^3 - x + 1 = 0$

Solving $f(x) = g(x)$ amounts to finding a **root** for the function $f(x) - g(x)$.

Solving $f(x) = g(x)$ amounts to finding a **root** for the function $f(x) - g(x)$.

Example

- Solve the equation $x = \cos x$.

Solving $f(x) = g(x)$ amounts to finding a **root** for the function $f(x) - g(x)$.

Example

- Solve the equation $x = \cos x$. This amounts to finding a root of the function $g(x) = \cos x - x$.

Solving $f(x) = g(x)$ amounts to finding a **root** for the function $f(x) - g(x)$.

Example

- Solve the equation $x = \cos x$. This amounts to finding a root of the function $g(x) = \cos x - x$.
- Solve $x^3 + 1 = x$.

Solving $f(x) = g(x)$ amounts to finding a **root** for the function $f(x) - g(x)$.

Example

- Solve the equation $x = \cos x$. This amounts to finding a root of the function $g(x) = \cos x - x$.
- Solve $x^3 + 1 = x$.
- Solve $x = (x - 1)^{\frac{1}{3}}$.
- How can you compare (2) and (3)?

More sophisticated methods than the Bisection Method can be developed around the idea of "Fixed point iterations".