

Crop Pathology Analysis: Cotton

Location: Nagpur, India | Date: Real-Time Field Telemetry

Primary Diagnosis: bacterial_blight

Confidence Score: 99.5%

AI Expert Summary: Active lesion propagation identified on Cotton tissue. Elevated humidity (91%) accelerates spore germination.

1. Climate Vitals & Soil Profile

Temperature: 32.1 C	Humidity: 91%	Wind: 12.4 km/h	Rain Risk: 0%
Soil Classification: Vertisol Clay		Soil pH: 6.8 Organic Matter: 1.5%	

2. N-P-K Fertilizer & Nutrient Prescription

Nitrogen (N): 45%	Phosphorus (P): 35%	Potassium (K): 40%
Formula: Potassium-Rich Recovery Blend NPK 00-52-34		
Dosage & Schedule: 3g/L Copper Oxychloride + Potassium Nitrate 4g/L (Apply immediately; repeat in 7 days after leaf drying)		

3. Eco-Friendly Organic Remedies & Companion Planting

Biopesticide: Fermented Sour Buttermilk (Khasti Chaas) Spray 10%
Recipe: Ferment 1L sour curd in a copper vessel for 10 days. Dilute with 10L water and spray.
Companion Plants: Garlic & Allium species (Natural anti-bacterial exudates)

4. Processed Visual Image Telemetry

