

Internet gaming disorder in children and adolescents: a systematic review

FRANK W PAULUS¹  | SUSANNE OHMANN² | ALEXANDER VON GONTARD¹ | CHRISTIAN POPOW²

1 Department of Child and Adolescent Psychiatry, Saarland University Hospital, Homburg, Germany. **2** Department of Child and Adolescent Psychiatry, Medical University of Vienna, Vienna, Austria.

Correspondence to Frank W Paulus at Department of Child and Adolescent Psychiatry, Saarland University Hospital, 66421 Homburg, Germany. E-mail: frank.paulus@uniklinikum-saarland.de

This article is commented on by Mueller on page 636 of this issue.

PUBLICATION DATA

Accepted for publication 2nd February 2018.
Published online 6th April 2018.

ABBREVIATIONS

IGD Internet gaming disorder
MMORPG Massively multiplayer online
role-playing game

AIM Internet gaming disorder (IGD) is a serious disorder leading to and maintaining pertinent personal and social impairment. IGD has to be considered in view of heterogeneous and incomplete concepts. We therefore reviewed the scientific literature on IGD to provide an overview focusing on definitions, symptoms, prevalence, and aetiology.

METHOD We systematically reviewed the databases ERIC, PsycARTICLES, PsycINFO, PSYINDEX, and PubMed for the period January 1991 to August 2016, and additionally identified secondary references.

RESULTS The proposed definition in the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition provides a good starting point for diagnosing IGD but entails some disadvantages. Developing IGD requires several interacting internal factors such as deficient self, mood and reward regulation, problems of decision-making, and external factors such as deficient family background and social skills. In addition, specific game-related factors may promote IGD. Summarizing aetiological knowledge, we suggest an integrated model of IGD elucidating the interplay of internal and external factors.

INTERPRETATION So far, the concept of IGD and the pathways leading to it are not entirely clear. In particular, long-term follow-up studies are missing. IGD should be understood as an endangering disorder with a complex psychosocial background.

Games are an integrative part of human behaviour and experience. During the past two decades, the availability and use of computer technology has dramatically increased and changed the world of leisure activities. Use of the Internet and computer game playing have become common activities for children and adolescents, in addition to social and traditional mass media. Recent data from the USA suggest that 8- to 10-year-olds are busy 8 hours per day, and adolescents more than 11 hours per day, with the recreational use of various electronic media (mobile phones, television and videos, computer use, music, print media, Web pages, social media, not including telephone conversations and text messages). This covers more time than they spend in school or with friends.^{1,2}

For most individuals, computer gaming is an enjoyable and stimulating activity.^{3–5} Persons with various intra- and interpersonal risk factors may, however, become attracted to using computer gaming as a strategy to overcome individual problems. Gaming and seeking for game-related pleasure may lead to neglecting ‘normal’ relationships, school or work-related duties, and even basic physical needs. Computer gaming may thus be conceptualized as a continuum from an enjoyable activity to pathological and even addictive use.^{6–8}

The Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)⁹ conceptualizes ‘Internet gaming disorder’ (IGD) in the chapter ‘Conditions for further study’ (p.795), suggesting that this proposal is not yet intended for clinical use but that research on this topic is encouraged. The essential feature of IGD is persistent and recurrent participation in computer gaming for typically 8 to 10 hours or more per day and at least 30 hours per week, typically in Internet-based group games (especially massively multiplayer online role-playing games [MMORPG]).

Since IGD has been defined only in 2013, previous literature uses various terms for describing this clinical entity, such as Internet or computer addiction (see ‘Definition’). While IGD is typical for adolescents and young adults, preschool and preadolescent children prefer non-Web-based games. Therefore, the American Psychiatric Association definition may not exactly fit for younger children and those playing offline.

Although IGD may be a serious disorder leading to significant impairment of personal and social functioning, the definition, symptoms, prevalence, and aetiology have to be seen in view of heterogeneous, incomplete, and changing concepts, and only a few available clinical studies of scientific quality. We therefore decided to review the scientific

literature on IGD of the past two and a half decades to provide an up-to-date overview on this disorder.

METHOD

This systematic review was conducted according to the structured Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines for systematic reviews and meta-analyses.¹⁰ We searched the literature databases ERIC, PsycARTICLES, PsycINFO, PSYINDEX, and PubMed for articles published in English between January 1991 and August 2016. We searched the databases using the following terms: pathological gaming [computer games OR video games OR online games OR Internet games] AND [abuse OR addiction OR compulsive OR dependence OR dependency OR disorder OR effects OR excessive OR habits OR misuse OR pathological OR problem OR problematic].

Applying the PRISMA guidelines for reviewing articles on IGD was beset by various problems: IGD is an expansive topic and includes subchapters such as definitions, symptoms, consequences, aetiology, and related and risk factors. IGD terminology is inconsistent (see ‘Definition’) and was only defined in 2013 in an amendment to the DSM-5.⁹ We therefore used the various search terms listed above to retrieve a maximum number of related articles.

The aim of our review was to provide an overview on the concept and background of IGD. We initially retrieved 3682 articles. Checking the titles and abstracts for relatedness of the content, we excluded: (1) duplicates and articles with content not relevant to IGD, for example articles on compulsive gambling or articles on the use of specific technologies such as mobile phones; (2) articles referring only to adults (we only selected articles investigating children and adolescents); (3) articles without reference to IGD. This reduced the number of relevant articles to 148 (4.0%). We additionally included 104 articles, books, and book chapters cited in the original literature or retrieved in specific searches on subtopics. This summed up to a total number of 252 references in total.

RESULTS

Definition

There are several terms and concepts for describing the pathological use of computer technology: IGD, Internet addiction, technology-based addiction, pathological technology use, pathological video gaming, pathological Internet use, computer addiction, screen addiction, addictions to video games and online role-playing games, game overuse, video game addiction, smartphone addiction (<http://www.helpguide.org/articles/addiction/smartphone-and-internet-addiction.htm>), cyber-relationship addiction, net compulsions, information overload, excessive, problem, dependent, pathological, and addicted gamers.

The concept of IGD has been questioned¹¹ because clinical data on symptoms of ‘true’ addiction are lacking, negative consequences may be overemphasized, and Internet

What this paper adds

- In representative samples of children and adolescents, on average, 2% are affected by Internet gaming disorder (IGD).
- The mean prevalences (overall, clinical samples included) reach 5.5%.
- Definitions are heterogeneous and the relationship with substance-related addictions is inconsistent.
- Many aetiological factors are related to the development and maintenance of IGD.
- This review presents an integrated model of IGD, delineating the interplay of these factors.

gaming could be related to inefficient strategies of dealing with life problems, poor time management, or other underlying mental health problems. Interestingly, there is no IGD Medical Subject Headings (MeSH) term available.

There is also a debate about what amount of time spent with gaming indicates addiction because increased screen time relates to IGD.^{12,13} Griffiths and Meredith¹⁴ distinguished between excessive and addicted gamers depending on the consequences of gaming because gaming may also have positive effects, enabling (virtual) social inclusion, increasing self-esteem and social acceptance, and structuring daily routine. Some authors related IGD symptoms to other disorders such as obsessive-compulsive disorder, attention-deficit-hyperactivity disorder, or depression.^{15–17}

For the first time, DSM-5 has clearly defined symptom criteria for IGD,⁹ requiring five or more of the following nine diagnostic criteria during a 12-month period: (1) pre-occupation with Internet games becomes the dominant activity in daily life; (2) withdrawal symptoms when removing the Internet (e.g. irritability, anxiety, or sadness, no physical signs of pharmacological withdrawal); (3) need to spend increasing amounts of time with Internet gaming (tolerance); (4) lacking control of Internet gaming; (5) loss of interest in previous hobbies and entertainment; (6) continued excessive use despite knowledge of psychosocial problems; (7) deception of family members, therapists, or others about the amount of Internet gaming; (8) escape from or relief of negative mood (e.g. feeling helpless, guilty, anxious); (9) loss of important aspects of life (e.g. significant relationships, jobs, or educational/career opportunities).

The nosological classification of IGD⁹ is inconsistent (the first of the nine proposed criteria includes ‘Internet games’ only, but does not definitely exclude ‘non-Internet based computerized games’ under the heading ‘Subtypes’, one page later) and still a matter of debate (e.g. Brunborg et al.,¹⁸ Müller et al.,¹⁹ Wölfling et al.)²⁰ DSM-5 excluded various differential diagnoses, excessive use of social media, online consumption of sexual content, and excessive online gambling, because these were considered to relate to different pathologies. The forthcoming 11th Revision of the International Statistical Classification of Diseases and Related Health Problems (ICD-11) will include online or offline ‘gaming disorder’ (6C51)²¹ as a distinct nosological entity after neglecting this disorder in previous editions.²²

In addition, the DSM-5 definition of IGD excluded offline and video gaming, probably because there is no sufficient evidence for this type of gaming disorder as a separate entity. It is not clear whether this restriction is justified because online and offline gaming share most features, with the exception of direct social reinforcement (team aspects) of online gaming. Furthermore, offline gaming is more predominant in younger children,²³ possibly because Internet access may more easily be restricted in this age group. With one exception,²⁴ there are no data available on IGD in preschool children.

The criteria for defining non-problematic, problematic, and addictive use of computer games are diverse and probably best characterized as a continuum that considers various dimensions such as severity of impairment, personality, cognitive distortions, psychopathology, and social functioning.^{25,26}

Considering problematic computer, online, or video gaming as a behavioural addiction has led to significant controversy. For classifying IGD, the recommendation is to differentiate between high engagement, addiction, gaming enthusiasm, and problems related to excessive gaming. High engagement with gaming, contrary to game addiction, does not lead to negative consequences.^{18,27–32}

Pathological computer gaming and behavioural and substance-related addictions share similarities.^{33,34} Pathological gamers share neurobiological correlates with behavioural and substance-related addictions.^{8,35–39} An important aspect in this debate relates to the definition of addiction,⁴⁰ which includes characteristic elements such as salience, tolerance, withdrawal, relapse, mood modification, and tolerating negative consequences as key components.^{40,41}

Symptoms

IGD-related symptoms include excessive screen time, preoccupation with gaming, salience, tolerance, loss of control, conflictuousness, mood imbalance, and relapse.^{8,42–44}

Conflictuousness means repeatedly having internal conflicts or conflicts with others in relation to the excessive gaming activity.⁴¹ Mood imbalance (e.g. feelings of guilt, depression) occurs as a consequence of losing control of gaming.⁴⁵ Salience describes a condition where gaming has become the most important activity in one's life; and relapse refers to the tendency to refer to earlier gaming patterns after periods of abstinence.^{46,47}

When gamers are consistently engaged or preoccupied with thoughts about gaming, or crave gaming during abstinence, they are about to lose control.⁴⁷ King et al.⁴⁶ described symptoms of withdrawal, loss of control, and conflictuousness as the core symptoms of pathological gaming, and considered other symptoms as peripheral (i.e. not essential for the diagnosis of IGD). Charlton and Danforth⁴¹ identified conflictuousness, symptoms of withdrawal, relapse/reinstatement, and behavioural salience as core symptoms; and cognitive salience, tolerance, and euphoria as peripheral symptoms. Addicted gamers,

primarily exhibiting core symptoms, spend more time with gaming than those exhibiting only peripheral symptoms.

Problematic use may lead to tolerance, an important symptom of addiction.⁸ Gunuc³³ indicated that tolerance and time spent with gaming may play an important role in developing addiction because screen time increases with addiction. Tolerance is not always a symptom of addictive behaviour, just as its absence does not exclude addiction. If a gamer does not increase his gaming time (e.g. because of restricted access), psychological but not behavioural tolerance may predominate.³³ If tolerance develops, symptoms of withdrawal are likely to increase.^{33,48}

Craving and withdrawal are important symptoms of pathological gaming.^{49,50} Different conceptualizations of withdrawal^{9,49} may explain why symptoms of craving and withdrawal were not detected in some studies. Kaptsis et al.⁴⁹ speculated that mild withdrawal symptoms may be overlooked because the environment of the gamers has adjusted to the addiction, or because symptoms had only occurred in specific addictive subtypes. According to the DSM-5 classification,⁹ withdrawal symptoms include sadness, irritability, and anxiety, whereas King et al.⁵⁰ found patients being bored and driven for mental stimulation. A recent article⁵¹ suggested that a better definition of withdrawal symptoms and their duration is required.

Craving is not listed as a symptom of IGD in the DSM-5. It might be considered an aspect of withdrawal.⁴⁹ Related cues may activate specific brain regions and induce craving sensations.^{36,52} Dong et al.⁵³ found craving and related changes in functional magnetic resonance imaging (MRI) in patients with IGD but not in subjects with recreational game use (RGU). Cue-induced craving is assumed to play an important role in maintaining addictive behaviour and producing relapse.⁵⁴

Although cue-induced craving was much stronger in Internet addicts, Niu et al.⁵² also described craving symptoms in non-addicted Internet users.

Online gaming is a highly heterogeneous condition and depends on a wide range of psychological factors. It is important to understand gamers' individual motives and psychological factors to plan effective interventions. Billieux et al.⁵⁵ described five subtypes of online gamer: two non-problematic and three problematic, as follows. (1) Regulated recreational gamers (non-problematic): 'low impulsivity traits, high self-esteem'. (2) Regulated social role-players (non-problematic): 'low levels of self-esteem, low impulsivity, and motivations related to social exchange and role-playing'. (3) Unregulated achievers (problematic): 'high scores on all impulsivity facets and primarily motivated by achievement in the game'. (4) Unregulated escapers (problematic): 'low achievement and high escapism, poor self-esteem'. (5) Hard-core gamers (highly problematic): 'highest adverse consequences resulting from gaming and high involvement in the game'.

Several studies^{56–59} investigated the validity of the DSM-5 criteria, showing that some of them were more suited than others: 'escape' and 'preoccupation' were judged as

less suited.^{56,57,59} 'Deception' in the study of Ko et al.⁵⁷ was less suited for identifying IGD, whereas other studies^{59,60} found the opposite. 'Giving up other activities' and 'tolerance',^{56,59} as well as 'withdrawal',⁵⁹ and 'loss of control',⁵⁶ were also debated. In addition, Ko et al.⁵⁷ suggested 'craving' as a candidate criterion. Another topic of controversy is the problem of a cut-off level because, so far, there is no criterion standard for defining IGD.^{56,57,59}

Consequences

Problematic gaming may lead to several negative psychosocial consequences and mental health problems affecting available time, work, education, family, partnership, friends, social life, psychosocial well-being, social competence, leisure activities, self-esteem, and loneliness.^{61–65} Professional and academic problems may include poor grades, academic failure, and financial problems.^{62,65} There is a negative relationship between academic performance and pathological gaming, affecting self-esteem and self-confidence.⁶⁵ A minority of gamers reported that gaming resulted in an overall negative effect on quality of life.⁶² Furthermore, Andreassen et al.⁶⁶ emphasized the relationship between psychiatric disorders such as anxiety and depression and gaming addiction. Messias et al.⁶⁷ found a higher risk of sadness, suicide ideation, and suicide planning in heavy gamers involved with screen times of more than 5 hours a day. Pathological media use may also reduce sleep duration and disrupt sleep patterns.^{62,68} IGD is also associated with various somatic health and medical consequences, such as auditory hallucinations, enuresis, encopresis, wrist, neck, and elbow pain, tendosynovitis ('nintendinitis'), obesity, skin blisters, calluses, sore tendons, hand-arm vibration syndrome, and peripheral neuropathy.^{63,69} In addition, psychosocial and medical consequences of IGD are similar in males and females.⁶⁴

To distinguish between causes and consequences, structural equation models have to be established on the basis of longitudinal study data. Only a few studies have applied such models, which mainly suffer from sample sizes that are too small.⁷⁰

Prevalence

Prevalence rates of IGD vary widely between studies from different countries, ranging from an estimated 0.6% in a Norwegian survey⁷¹ up to an extreme of 50% in Korea.⁷² This may be caused by differences in assessing instruments, study population,⁸ and diagnostic criteria of IGD. Prevalence rates are highest in Eastern Asian countries and male adolescents aged 12 years to 20 years.⁹ The median prevalence rate for computer and IGD as calculated from Table I was 5.5%. Selecting the 10 population-based studies listed in Table I, we obtained a median prevalence of 2.0%, which is close to the number of 1.6% of a recent European population-based study.¹⁹

A recent review on 13 longitudinal epidemiological studies⁷³ showed a moderate to high symptom stability over 2 years, and protective social factors.

Sex differences

IGD may be observed up to five times more often in male children: IGD was diagnosed in 11.9% of the males and in 2.9% of the females in a large US study.⁷⁴ Only two studies^{75,76} reported no sex-related difference in addicted gamers.

Age

The influence of age on IGD prevalence is not clear. Unfortunately there are no studies available for younger children, with one exception.²⁴ The highest prevalence of IGD was found in adolescents;⁷⁷ only one recent study identified more addicted adults than adolescents.³⁰

Aetiology

The aetiology of IGD is complex and not fully understood. IGD 'can last for years and is not solely a symptom of comorbid disorders'.¹³ There are internal and external factors related to developing and maintaining the disorder. In addition, genetic, learning, motivational, game-related, and neurobiological aspects must be considered. It is unclear whether psychiatric symptoms predispose for, or follow, IGD because longitudinal studies are scarce.⁷⁸

Internal factors

Among the internal factors, deficient self-regulatory and decision-making abilities as a consequence of dysexecutive problems, mood and reward system dysregulation, and avoidant behaviour (escapism, deficient coping with negative emotions), low self-esteem, poor self-efficacy, and neurobiological factors may be involved.

Dysexecutive problems, deficient self-regulation, and decision-making. Deficient inhibitory control, sensory-motor coordination, and executive control including self-control have been described in patients with IGD.^{37,79,80} Lower functional connectivity of executive control networks along with impaired executive functions were described in individuals with IGD.⁸⁰ Compared with healthy individuals, the anterior insula and the dorsolateral prefrontal cortex are less activated during decision-making tasks in patients with IGD.^{81,82} In addition, the volume of the striatum is increased in patients with IGD, which relates to impaired cognitive control.^{83,84} Decreased inhibition of impulsivity entails the disability to control compulsive gaming despite serious negative consequences.⁸⁵ Impaired functioning of the amygdala, responsible for emotional control, may be associated with increased emphasis of emotions and immediate rewards, leading to excessive gaming without paying attention to negative long-term consequences.⁸⁵

Dysregulation of mood and reward system, avoidant behaviour. Negative and depressed mood, stress, anxiety, and preference for immediate rewards^{86,87} are associated with pathological gaming.^{63,88} Kwon⁸⁹ hypothesized that IGD, such as drug abuse, may serve as a self-destructive strategy for escaping unpleasant real-life problems. Li et al.⁹⁰ suggested depression to mediate the relationship

Table 1: Prevalence of computer and Internet gaming disorder (IGD) across various countries

Country	Study	Investigation	Age group	Sex (M:F)	Method(s)	Prevalence (%)	Representative
Australia	Thomas and Martin ²⁰⁸	CGA	12–54y	42:58	YIAT	5.0	Y, students
Australia	Thomas and Martin ²⁰⁸	IA	12–54y	42:58	YDQ, DSM-IV-TR	4.6	Y, students
Austria	Batthyány et al. ¹²⁰	CGA	13–18y	54:46	CSVK-R	2.7	Y, students
China	Cao et al. ²³¹	IA	12–18y	50:50	YDQ	2.4	Y, students
China	Xu and Yuan ²³²	Game addiction	13–18y		DSM-IV	21.5	Y, students
China	Cao et al. ²³³	PIU	10–24y	51:49	YIAT	8.1	Y, students
Germany	Grüsser et al. ²³⁴	Excessive computer and VG playing	11–14y	54:46	CSVK, ICD-10, DSM-IV-TR	9.3	Y, students
Germany	Gruesser et al. ⁶	CGA	Average 21y 1mo	94:6	ICD-10	11.9	Y, students
Germany	Rehbein et al. ²⁹	VGD	Average 15y 4mo	51:49	VGDS	2.8	Y, students
Germany	Rehbein et al. ²⁹	VGD	Average 15y 4mo	51:49	KFN-CSAS	1.7	Y, students
Germany	Wölfling et al. ²³⁵	CGA	13–18y		ICD-10	7.5–8.4	Y, students
Germany	Müller et al. ²³⁶	PIU	8–17y	?	AICA-S	11.3	N, patients
Germany	Müller et al. ¹⁶⁵	IA	14–94y	48:52	AICA-S, CSV-S	2.1	Y
Germany	Dreier et al. ⁷⁵	IGD	12–18y		AICA-S	5.2	Y, students
Germany	Paulus et al. ²⁴	CGA	4y 5mo–8y 2mo	50:50	YC-CGD	1.9	Y, children
Europe	Müller et al. ¹⁹	IGD	14–17	47:53	AICA-S	1.6	Y
International	Porter et al. ¹¹⁴	Problem VG use	14–40y	93:07	VGUQ	8.0	N, students
International	Hussain et al. ¹⁴¹	Online game addiction	12–62y	77:23	DSM	3.6–44.5	N, gamers
Iran	Zamani et al. ²³⁷	CGA	Grade 2 students	53:47	QACG	17.1	Y, students
the Netherlands	van Rooij et al. ³⁴	Online VG addiction	13–16y	49:51	CIUS	1.5	Y, students
the Netherlands	Haagsma et al. ¹⁹⁶	PGB	14–81y	50:50	GAS	1.3	Y
the Netherlands	Lemmens et al. ²³⁸	IGD	13–40y	50:50	IGD scales (DSM-5)	5.5	Y
Norway	Bakken et al. ²³⁹	IA	16–74y	47:53	YDQ	1.0	Y
Norway	Arnesen ²⁴⁰	VGA	16–40y	44:56	DSM-IV-TR	0.6–4.0	Y
Norway	Mentzoni et al. ⁷¹	Problematic VG use	15–40y		GASA	0.6	Y
Norway	Wittek et al. ²⁴¹	VGD	16–74y	63:37	GASA	1.4	Y, gamers
Singapore	Choo et al. ²⁴²	Pathological VG	9–13y	73:27	DSM-IV	8.7	Y, students
Singapore	Gentile et al. ¹³	Pathological VG use	12–18y	73:27	DSM-IV-TR	9.0	Y, students
South Korea	Hur ⁷²	IAD			IALL and IAD	50.0	Y, students
South Korea	Kim et al. ²⁴³	IA	15–16y	35:65	YDQ	1.6	Y, students
South Korea	Lee and Han ²⁴⁴	Online game addiction	Grade 5 and 6 students		Own, based on YIAT	2.5	Y, students
South Korea	Park et al. ²⁴⁵	IA		70:30	IAS	10.7	Y, students
South Korea	Jeong and Kim ²⁴⁶	IA	15–16y	53:47	IAT	2.2	Y, students
Spain	Tejero Salguero and Morán ²⁴⁷	Problem VG playing	13–18y	54:46	PVP	9.9	Y, students
Spain	Lopez-Fernandez et al. ²⁴⁸	IGD	11–18y	53:47	PVP	7.7	Y
Taiwan	Ko et al. ²⁴⁹	IA	Average 13y 7mo	52:48	CIAS	7.5	Y, students
Taiwan	Wan and Chiou ²⁵⁰	Online game addiction	17–24y		OAST	34	N
UK	Fisher ²⁵¹	VGA	11–16y	48:52	DSM-IV	19.9	Y, students
UK	Griffiths and Hunt ²⁵²	Dependence on computer games	12–16y	58:42	DSM-III-R	19.9	N
UK	Lopez-Fernandez et al. ²⁴⁸	IGD	11–18y	67:33	PVP	14.6	Y
USA	Gentile ⁷⁴	Pathological VG use	8–18y	50:50	DSM-IV	8.5	Y

M, male; F, female; CGA, computer game addiction; YIAT, Young's Internet Addiction Test; Y, yes; IA, Internet addiction; YDQ, Youth Dependency Questionnaire; DSM-IV-TR, Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision; CSVK-R, Fragebogen zum Computerspielverhalten bei Kindern [Questionnaire for computer gaming behaviour in children]; PIU, problematic Internet use; VG, video game; ICD-10, International Statistical Classification of Diseases and Related Health Problems, 10th Revision; VGD, video game dependency; VGDS, Videogame Dependency Scale; KFN-CSAS, Lower Saxony Research Institute of Criminology: Computer Game Addiction Scale; AICA-S, Assessment for Internet and Computer Game Addiction; N, no; IA, Internet addiction; CSV-S, Scale for the Assessment of Pathological Computer-Gaming; YC-CGD, Young Children – Computer Gaming Disorder Questionnaire; VGUQ, Video Game Use Questionnaire; DSM, Diagnostic and Statistical Manual of Mental Disorders; QACG, Questionnaire of Addiction to Computer Games; CIUS, Compulsive Internet Use Scale; PGB, problematic gaming behaviour; GAS, Gaming Addiction Scale; VGA, video game addiction; GASA, Game Addiction Scale for Adolescents; IAD, Internet addiction disorder; IALL, Internet Addiction Impairment Index; IAS, Internet Addiction Scale; IAT, Internet Addictions Test; PVP, Problem Video Game Playing scale; CIAS, Chen Internet Addiction Scale; OAST, Online gaming Addiction Scale for Adolescents in Taiwan; DSM-III-R, Diagnostic and Statistical Manual of Mental Disorders, Third Edition, Revised.

between actual–ideal self-discrepancy and escapism. On the other hand, escapism and game immersion are mediating factors between depression and problematic gaming, promoting IGD (<https://bpistone.wordpress.com/2012/09/03/>).

Self-esteem and self-efficacy. Low self-esteem was also found to trigger pathological gaming behaviour.⁹¹ Excessive gamers are attracted to games because gaming stimulates the experience of power and autonomy, and strengthens self-esteem. Furthermore, pathological gamers tend to overvalue gaming rewards, activities, identities (avatars), or items. This stimulates preoccupation with gaming and loss of interest in less attractive real-life activities. Avatars (simulated identities in a game) may enhance the feelings of empowerment and strength, and facilitate escape from real-life problems.⁹¹

Neurobiology. Recent research describes neurobiological anomalies in pathological gamers such as altered grey matter volume, functional connectivity, and activation in specific brain regions (especially the ventromedial prefrontal cortex) and neurobiological correlates to disorder-specific behaviours and symptoms.⁹² It is not clear whether these alterations are causes and risk factors for IGD or consequences resulting from excessive gaming. The most commonly used methods (electroencephalography, positron emission tomography, functional MRI, structural MRI [voxel-based morphometry, diffusion-tensor imaging]) were mainly applied in adults.

Although pathological gaming is considered a behavioural addiction (e.g. Leeman and Potenza,³⁸ Griffiths et al.,⁶³ Ong et al.,⁹³ Vousooghi et al.),⁹⁴ multiple similarities have been described between addictive gaming and other addictions such as substance dependency or pathological gambling. For example, Fauth-Bühler and Mann⁹⁵ found loss of sensitivity, increased reactivity to related cues, increased impulsivity, and altered reward-based learning in addictive gamers and pathological gamblers compared with healthy individuals.

Impulsive behaviours are associated with deficits of frontostriatal networks, interhemispheric connectivity, and an altered structure of the insula.^{96,97} Dysfunction of the prefrontal cortex and disrupted frontostriatal connectivity indicate that the frontal lobe fails to regulate emotion-driven urges.^{96,98} Gaming-related stimuli induce increased urges and provoke enhanced activation of frontolimbic and mesocorticolimbic reward systems in pathological gamers.^{95,99} The same brain areas are also involved in cue-induced craving with other addictions.^{36,38,99,100} Moreover, the temporal homogeneity of the regional blood-oxygen-level dependent signal in functional MRI, which presumably indicates neural activity,¹⁰¹ was found to be increased in the posterior cingulate cortex in patients with IGD and with alcohol use disorder.¹⁰²

Substance-related and behavioural addictions affect the ‘reward circuit’.^{38,79,103} The mesolimbic dopamine pathway is particularly important and includes dopaminergic neurons from the ventral tegmental area to the nucleus

accumbens. Dopamine levels that are too low or too high are associated with impulsivity and risky decisions.¹⁰⁴ Excessive Internet and game use is associated with an overall reward deficiency that is related to reduced dopaminergic activity, similar to that in drug abuse.^{8,79} Videogame playing was found to induce dopamine releases similar to cocaine.⁷⁹ Some authors have suggested an association between IGD and a dysfunction of the dopaminergic system.⁷⁹ Vousooghi et al.⁹⁴ found a reduced number of D5 dopamine receptors in computer game addicts compared with healthy individuals.

Hoefl et al.¹⁰⁵ found sex differences when investigating activation and connectivity of brain regions associated with the mesocorticolimbic reward system.⁸ The higher risk of gaming addiction for males could be related to these neuronal differences.⁶⁵

In summary, neurobiological and functional studies provide strong evidence for structural and functional brain changes in pathological gamers.

Personal risk factors. Personal risk factors include low self-esteem,^{55,106–111} social isolation,^{108,109,111,112} decreased interpersonal relationships and few or no friends in real life,^{113,114} impaired social functioning,¹³ limited other leisure or recreational activities,^{19,59} low educational or academic achievement,^{18,19,115–118} and increased truancy, low commitment to school, and poor school or academic performance or unemployment.^{19,27,29,74,119–122}

Frölich et al.¹²² found no specific psychiatric disorders associated with computer game misuse or addiction. The authors speculated that their results supported the idea of IGD as a distinct psychiatric entity. Hart et al.,¹²³ in contrast to Kim et al.,¹²⁴ Rehbein et al.,²⁹ and Subramaniam et al.,¹²⁵ found no self-reported physical reactions or increased stress levels in adolescent students with IGD.

Social factors such as poverty, social exclusion, poor parental competence and supervision, and inconsistent parental behaviour play a role in developing IGD. Socio-cultural influences such as gender roles, family structure, cultural values, and personal beliefs should also be considered as related to IGD.¹²⁶ Only one study reported multiple comorbid conditions of IGD in younger children.¹²⁷

Individual factors such as sex, age, personality traits, and various psychological factors may be related to an increased risk of developing and maintaining IGD.^{43,55,128}

Sex. Males, independent of age, bear a higher risk^{8,19,43,74,107,129–131} of developing IGD, participating in MMORPGs,¹³² and playing more violent games (e.g. Lemmens et al.)¹³³ They play more often for excitement and for making friends¹⁰⁷ than females. Females engage in gaming for passing time, social networking, texting, online shopping, but not for fighting or re-enacting negative emotions.^{30,43,66,107,134–138} Wei et al.¹¹² reported that female gamers presented with shorter durations of online gaming and shorter screen times. Social phobia also increases the risk of IGD,¹¹² independent of sex.

Age. Age shows an inversely U-shaped relationship to IGD^{8,43,66,125,137,139} and correlates with the time spent

computer gaming.^{140,141} Older adolescents have the highest probability of becoming game addicted.¹⁴² This may be related to a greater technology affinity of 'digital natives'⁶⁶ and to the developmental tasks of this age period (personal goals vs life optimization).¹⁴³ Adolescents are especially vulnerable to the negative effects of IGD.¹³³

Personality traits. Personality traits are related to developing IGD,^{144,145} especially impulsivity,^{55,110,146–148} lower self-control/self-regulation ability,^{124,146,149–151} sensation-, stimulation-, and/or novelty-seeking,^{119,152,153} inclination to boredom,¹¹⁹ risky behaviour,¹⁵⁴ hostility/animosity,^{119,155,156} and enhanced levels of aggression.^{19,83,110,124,146,153,154,157,158}

Guesser et al.⁶ found only a weak association between IGD and aggressive behaviour. Because of the cross-sectional design of most studies, a causal link between aggressive behaviour and IGD at present cannot be proved.¹³³

The contribution of the 'Big Five' personality traits (neuroticism, extraversion, openness to experience, agreeableness, and conscientiousness)¹⁵⁹ to developing and maintaining IGD is not clear. Introversion, shyness, or low extraversion have been associated with IGD.^{116,160–168} Data about trait empathy are inconsistent, ranging from decreased empathy (e.g. Bartholow et al.)¹⁶⁹ to no differences except for soft skills deficits.^{150,170} Other personality traits related to IGD are decreased conscientiousness,^{128,160,164,165,167,168} decreased openness,¹⁶⁷ agreeableness, resourcefulness,^{128,150,161,171,172} irritability,^{110,146} and high neuroticism/low emotional stability.^{128,153,161,163,165,168,171–175} State and trait anxiety^{153,163} are characteristic of IGD. Narcissistic,^{124,176} avoidant, and schizoid interpersonal traits are also important for IGD.¹⁷⁷

It must be stated that personality traits may change during individual development.¹⁷⁸ Unfortunately, most studies investigating the relationship between personality traits and IGD did not control for this age dependency. Future studies should therefore consider such effects.

The preference of problematic game users for online relationships is related to loneliness, deficient soft skills,^{30,64} and social inhibition.¹¹⁴ Being single bears a higher risk for developing IGD.⁴³ Caplan¹⁷⁹ described that the relationship between loneliness and IGD could be mistaken as social anxiety.

Substance use disorder. Ong et al.⁹³ found various personality factors associated with adolescent substance use disorder and IGD. Substance use disorder was associated with a higher risk of childhood adversity and delinquency,¹³² and IGD more with traumatic childhood experiences.¹⁸⁰

Personal motives and preferences. Personal motives and preferences such as escapism,^{8,56,181,182} immersion, achievement, and affiliation are related to IGD.⁵⁵ Wan and Chiou¹⁸³ described IGD effects as more related to relief from dissatisfaction than to gaining satisfaction. Game overuse may be related to three motivational factors: emotional regulation against dysphoria, escaping from real-life problems, and developing personal relationships with other gamers.¹⁸⁴ The Motives for Online Gaming Questionnaire

cites seven main online gaming motives: escape, coping, fantasy, skill development, recreation, competition, and social.¹⁸⁵

Cognitive performance. The relationship between cognitive performance and IGD is not clear. Hyun et al.¹³¹ found no differences in cognitive performance between addicted and non-addicted gamers. Because of the cross-sectional design of most clinical studies, the direction of causality is not clear. Park et al.¹⁸⁶ found lower comprehension scores resulting in lower social intelligence, and a negative relationship between gaming activity and performance in attention tasks. The earlier IGD manifests and the longer it lasts, the lower the performance in attention tasks.

External factors

Several external factors promote and maintain IGD against better insight.¹⁸⁷ Possible explanations between external factors and IGD are provided by learning theory (classical and operant conditioning, and gratifications theory)¹⁸⁸ and social-cognitive theory (model learning).^{189,190} External factors include familial, social, and game-related factors.

Familial factors. Poor relationships with parents and peers are associated with IGD.⁴⁵ Poor parental control and negative role models, as well as a positive parental attitude towards adolescent substance use, predict both substance use and IGD.^{191,192} Particularly ignorant, oppressive, and hostile parents, single parent families, and a broken home context increase the screen time of adolescents.¹⁹³ In the Berlin Longitudinal Media Study, a controlled group study involving 1207 schoolchildren over a period of 4 years, 'family violence' and 'poor parental care' were identified as key risk factors of IGD.¹⁹⁴

Social factors. Team aspects are important motivational factors for IGD; like-minded players may interact in real life and in virtual environments. Communication and social interaction are essential ingredients of many games, especially MMORPGs. Haagsma et al.¹⁹⁵ reported that online social interaction strengthens meaningful feelings and self-regulation, and reduces negative feelings such as loneliness and boredom. Shy and isolated children find a safe and rewarding way of interacting with peers in the virtual world. The avatar skin may provide a desirable appearance. In particular, MMORPGs create and provide a social environment that is built solely on gaming skills and availability on screen.

The virtual social environment enables the gamer to interact with people having similar (gaming) interests. This rather reinforces gaming instead of criticizing it.⁹¹ Virtual relationships thus become increasingly more important for the gamer, and virtual rank and status increasingly satisfy otherwise deficient social needs and reinforce online presence.^{91,196} Moreover, the guild or clan may exert social pressure to attach gamers to the game and keep them active.⁴⁵ Gamers will receive social rewards for gaming skills in the real peer group as well. Pathological gamers tend to overestimate gaming rewards, activities, virtual identities, or hard-earned 'valuable' items (weapons, life

energy, treasures, etc.). Overestimating such features may be another reason for losing interest in real-life activities.⁹¹

Wan and Chiou¹⁸³ found no relationship between experiencing a 'flow state' and excessive gaming, whereas Hull et al.¹⁹⁷ highlighted the experience of a flow state as 'heightened levels of a sense of time being altered during play', and that pursuing this 'high' was a risk factor for IGD.

Game-related factors. Structural characteristics of specific game genres will promote a pathological gaming behaviour:¹⁹⁸ game 'immersion' and increased weekly and holiday screen time;^{13,118,164,199–201} strong gaming urges;^{201,202} non-age-appropriate shooter games or games with forbidden content (forbidden fruit hypothesis);^{19,203,204} (over-) involvement in social networks;¹⁹⁹ and becoming familiar with specific game characteristics (mental constructs developed during regular participation).^{205,206}

Internet and role-playing games possess more addictive potential than offline games because of their inherent social reinforcements.^{195,207,208} In particular, MMORPGs include addiction-promoting features that attach gamers to the game;^{45,209} for example, a never-ending design, the possibility of acquiring power, glory, and attractive items such as invincible armament, their hierarchical social organization (e.g. in guilds), and a regular membership fee,²¹⁰ rewards and punishments (earning scores, reaching levels), narrative elements (e.g. an interactive story), realistic graphical quality, and fast loading times.²⁰⁹ Another method to increase game involvement is monetization: the obligation to purchase items in 'free-to-play' (initially cost-free) games.⁷⁵ Kuss et al.¹⁹⁸ found specific motivation, especially 'escapism', and 'mechanics', a subcomponent of achievement, but not total screen time to predict MMORPG addiction. MMORPG addiction in college students was predicted by five critical factors: curiosity, role playing, belonging, obligation, and reward.²¹¹

Gaming equipment and permanent Internet access influenced adolescents' IGD rates: possessing a PlayStation but not their own personal computer, Game Boy, television, or a powerful Internet access in their own room promoted game addiction.^{74,122,199}

Additional aspects. Beranuy et al.⁴⁵ identified three factors involved in motivating gamers: entertainment, escapism, and virtual friendships. With increasing screen time, MMORPG gamers start to lose control and narrow their behavioural focus. Griffiths and Beranuy,²¹² investigating subjective deficiencies in MMORPG gamers, found lack of real friends, difficulties in relationships, and problems with physical appearance as key motivational factors for gaming.

Furthermore, the use of non-age-appropriate games or games with forbidden content ('forbidden fruit hypothesis'),^{19,203,204} involvement in social networks,¹⁹⁹ and Internet access at home²¹³ as well as psychopathological comorbidities such as attention-deficit-hyperactivity disorder, depression, social anxiety, autism, etc. and other additional risk factors will promote IGD.

Complex models

Considering the complex aetiology of IGD, various models of the disorder have been developed. These models include behavioural, psychopathological, and social aspects with various weights. Cognitive-behavioural dimensions^{144,214} include maladaptive cognitions and thoughts going along with negative self-value, self-doubts, and poor self-efficacy. These deficits will promote and maintain pathological Internet use. Defence mechanisms used by pathological gamers are 'stonewalling', minimizing, blaming (e.g. stress), excusing, rationalizing, and attacking.²¹⁴ Strong emotions such as feelings of excitement when winning or finishing a game, powerfulness, relief (escaping from real-world problems), and friendships with 'electronic friends' will promote excessive gaming behaviour. Some models were empirically evaluated.¹⁹⁶

Haagsma et al.¹⁹⁵ proposed and evaluated a cognitive behavioural model of IGD. The key features of this model relate to preferences of social interaction, deficits in mood and self-regulation, ruminations, and compulsive aspects. The first three factors explained 79% of the variance in their model. In another publication,²¹⁵ the authors examined the model of planned behaviour theory, and found clients' perceived lack of control over gaming as the main factor for predicting IGD.

Dong and Potenza¹⁰³ proposed a cognitive-behavioural model of IGD. Because of an increased reward sensitivity and decreased lack of considering losses,^{8,216,217} as well as deficient response inhibition^{218,219} and executive control,⁸⁰ the authors speculated that deficient cognitive control in association with enhanced reward seeking would lead to increased gaming. Gamers exhibit deficiencies in impulse control, attentional capabilities,^{220–222} overall cognitive functioning,^{8,223} and impaired risk perception,²²⁴ favouring risky decisions because they strive for immediate positive rewards and neglect negative long-term consequences for social or work domains in real life.²²⁵

The multi-disciplinary aetio-pathogenetic model of Dreier et al.²²⁶ and Müller et al.²²⁷ includes psychological (e.g. dysfunctional beliefs), social (e.g. familial dysfunctions, social insecurity), and neurobiological (e.g. neurotransmitter dysregulations, dysfunctions of the reward system) IGD-promoting factors. Game-related (e.g. success, competition, exploration), environmental (e.g. availability, accessibility, popularity), and personal characteristics (e.g. impulsivity, negative self-concept, neuroticism), motivational states (e.g. escapism, compensation, coping), and specific game features (e.g. reward, curiosity, obligation) help to maintain IGD within vicious circles despite negative consequences.

Another complex bio-psycho-social process model, the Interaction of Person-Affect-Cognition-Execution (I-PACE) model,²²⁸ integrates psychological and neurobiological aspects of IGD. In short, a person defined by their core characteristics (influenced by numerous biological, motivational, and social protective and risk-increasing factors) decides to use a certain application in response to




Figure 1: An integrated model of Internet gaming disorder (IGD). PFC, prefrontal cortex; EDF, executive dysfunction; ADHD, attention-deficit–hyperactivity disorder; OCD, obsessive–compulsive disorder.




Figure 2: The pathway of gaming fun, pathological gaming, and gaming addiction.

their actual situation and their affective and cognitive responses. Internet gaming disorder (IGD) then develops as a result of the described personal characteristics and

affective responses under aspects of gratification and the specific neurobiological conditions of diminished executive control, despite negative consequences in daily life.

DISCUSSION

Many IGD-affected adolescents will endanger their health, fail academically, and may become isolated and depressive. Reviewing the scientific literature, the evidence of a diagnostic entity of IGD and the pathways to the disorder are not entirely clear. In particular, long-term follow-up studies are still lacking. There is some evidence that IGD does not affect anyone but that multiple, often interacting personal, familial, and environmental risk factors and comorbid conditions will contribute towards developing and maintaining the disorder.

Because the paucity of available high-quality, long-term studies impeded us in analysing available studies according to the PRISMA guidelines,¹⁰ we can only provide a systematic review on IGD, summarizing the available evidence.

The DSM-5 tentative definition of IGD⁹ is a good starting point for collecting diagnostic evidence, tolerance, loss of control, giving up other interests, and excessive use despite clear-cut psychosocial and health-related problems, representing the most stringent criteria. There are, however, weaknesses of the DSM-5 definition; it excludes other differential diagnoses, and alternative media and content, and has the problem of defining 'hazardous' screen times.^{18–20}

Developing IGD requires several complex interacting internal and external factors.^{103,196} Among these, problems of self-regulation and decision-making, dysregulation of mood, and reward systems are based on neurobiological deficits that may also explain sex differences.⁶⁵ Among the external factors, familial deficits, including intrafamilial violence and poor parental care, and deficient social skills may increase the risk of developing IGD, especially of becoming addicted to MMORPGs¹⁹⁵ or action real-time strategy (shooter) games such as Multiplayer Online Battle Arena.²²⁹ Finally, several game-related personally and socially reinforcing factors, as well as steadily improving technical performance, will promote IGD.^{207,208,211}

The wide range in IGD prevalence rates relates to different definitions of IGD, regional differences, and different methods of assessment. We found a median IGD

prevalence of 5.5% (Table I) and a median of 2.0% for population-based studies that seems more accurate. Male adolescents usually manifest higher prevalences than females.²⁰⁶ Most studies cited in our review investigated adolescents. Only a few individual studies^{1,13,74,87,118} and one review article²³⁰ included data on children. To the best of our knowledge, there is only one study available investigating IGD in preschool children.²⁴

Predisposing comorbidities and health-related consequences are commonly observed in IGD.^{20,29,71,184} Causal associations cannot be analysed because IGD and comorbid disorders such as attention-deficit-hyperactivity disorder, depression, or social anxiety share common aetiologies, and may interact. Longitudinal studies that could resolve this question are lacking. Low parental competence, personality traits, and problems of self-esteem or social skills may contribute to developing and maintaining IGD.

Summarizing aetiological knowledge, we suggest an integrated model of IGD (Fig. 1) that delineates the interplay of external and internal factors. External and internal factors are interrelated with each other and with IGD. IGD may aggravate existing deficits and vice versa; poor social relations will motivate for gaming, and spending increasing time with gaming will aggravate poor relationships, thus further reinforcing IGD.

Gaming reinforces itself by multiple mechanisms, as explained in the section on 'External factors'. The pathway towards addiction is characterized by a change of emphasis from fun, through losing control, to obsession. Figure 2 summarizes this development.

In conclusion, we have reviewed the current literature for characterizing IGD as a complex, endangering disorder with an intricate psychosocial background, and many personal, neurobiological, familial, social, and game-inherent IGD-promoting factors.

ACKNOWLEDGEMENTS

The authors have stated that they had no interests that might be perceived as posing a conflict or bias.

REFERENCES

1. Rideout VJ, Foehr UG, Roberts DF. Generation M²: Media in the Lives of 8- to 18-Year-Olds. Menlo Park, CA: Henry J. Kaiser Family Foundation, 2010.
2. Council on Communications and Media. Children, adolescents, and the media. *Pediatrics* 2013; **132**: 958–61.
3. De Freitas S, Griffiths M. Online gaming as an educational tool in learning and training. *Br J Educ Technol* 2007; **38**: 535–7.
4. De Freitas S, Griffiths M. The convergence of gaming practices with other media forms: what potential for learning? A review of the literature. *Learn Media Technol* 2008; **33**: 11–20.
5. Loton D, Borkoles E, Lubman D, Polman R. Video game addiction, engagement and symptoms of stress, depression and anxiety: the mediating role of coping. *Int J Ment Health Addict* 2016; **14**: 565–78.
6. Grüsser SM, Thalemann R, Griffiths MD. Excessive computer game playing: evidence for addiction and aggression? *Cyberpsychol Behav* 2007; **10**: 290–2.
7. Larose R, Mastro D, Eastin MS. Understanding Internet usage: a social-cognitive approach to uses and gratifications. *Soc Sci Comput Rev* 2001; **19**: 395–413.
8. Kuss DJ, Griffiths MD. Internet gaming addiction: a systematic review of empirical research. *Int J Ment Health Addict* 2012; **10**: 278–96.
9. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th edn. Arlington, VA: American Psychiatric Publishing, 2013.
10. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 2009; **6**: e1000100.
11. Wood RTA. Problems with the concept of video game addiction: some case study examples. *Int J Ment Health Addict* 2008; **6**: 169–78.
12. Chan PA, Rabinowitz T. A cross-sectional analysis of video games and attention deficit hyperactivity disorder

- symptoms in adolescents. *Ann Gen Psychiatry* 2006; **5**: 16.
13. Gentile DA, Choo H, Liau A, et al. Pathological video game use among youths: a two-year longitudinal study. *Pediatrics* 2011; **127**: e319–29.
14. Griffiths MD, Meredith A. Videogame addiction and its treatment. *J Contemp Psychother* 2009; **39**: 247–53.
15. Vukosavljević-Gvozden T, Filipović S, Opacic G. The mediating role of symptoms of psychopathology between irrational beliefs and Internet gaming addiction. *J Ration Emot Cogn Behav Ther* 2015; **33**: 387–405.
16. Carli V, Durkee T, Wasserman D, et al. The association between pathological internet use and comorbid psychopathology: a systematic review. *Psychopathology* 2013; **46**: 1–13.
17. Weiss MD, Baer S, Allan BA, Saran K, Schibuk H. The screens culture: impact on ADHD. *Atten Defic Hyperact Disord* 2011; **3**: 327–34.
18. Brunborg GS, Mentzoni RA, Froyland LR. Is video gaming, or video game addiction, associated with depression, academic achievement, heavy episodic drinking, or conduct problems? *J Behav Addict* 2014; **3**: 27–32.
19. Müller KW, Janikian M, Dreier M, et al. Regular gaming behavior and internet gaming disorder in European adolescents: results from a cross-national representative survey of prevalence, predictors, and psychopathological correlates. *Eur Child Adolesc Psychiatry* 2015; **24**: 565–74.
20. Wölfling K, Beutel ME, Dreier M, Müller KW. Bipolar spectrum disorders in a clinical sample of patients with Internet addiction: hidden comorbidity or differential diagnosis? *J Behav Addict* 2015; **4**: 101–5.
21. World Health Organization. International Classification of Diseases 11th Revision ICD-11 Beta Draft. 2017. <https://icd.who.int/dev11/l-m/en> (accessed 21 March 2018).
22. Mann K, Fauth-Bühler M, Seifert N, Heinz A, Expertengruppe Verhaltenssuchte der DGPPN. [The concept of behavioral addiction and limits of the term addiction]. *Nervenarzt* 2013; **84**: 548–56. (In German).
23. Vandewater EA, Rideout VJ, Wartella EA, Huang X, Lee JH, Shim MS. Digital childhood: electronic media and technology use among infants, toddlers, and preschoolers. *Pediatrics* 2007; **119**: e1006–15.
24. Paulus FW, Sinzig J, Mayer H, Weber M, von Gontard A. Computer gaming disorder and ADHD in young children—a population-based study. *Int J Ment Health Addict* 2017 Nov 30; <https://doi.org/10.1007/s11469-017-9841-0>. [Epub ahead of print].
25. La Rose R, Lin CA, Eastin MS. Internet addiction, habits and deficient self-regulation. *Media Psychol* 2003; **5**: 225–53.
26. Lee MS, Ko YH, Song HS, et al. Characteristics of Internet use in relation to game genre in Korean adolescents. *Cyberpsychol Behav* 2007; **10**: 278–85.
27. Skoric MM, Teo LL, Neo RL. Children and video games: addiction, engagement, and scholastic achievement. *Cyberpsychol Behav* 2009; **12**: 567–72.
28. Charlton JP, Danforth IDW. Validating the distinction between computer addiction and engagement: online game playing and personality. *Behav Inf Technol* 2010; **29**: 601–13.
29. Rehbein F, Kleimann M, Mössle T. Prevalence and risk factors of video game dependency in adolescence: results of a German nationwide survey. *Cyberpsychol Behav Soc Netw* 2010; **13**: 269–77.
30. Ferguson CJ, Coulson M, Barnett J. A meta-analysis of pathological gaming prevalence and comorbidity with mental health, academic and social problems. *J Psychiatry Res* 2011; **45**: 1573–8.
31. Brunborg GS, Mentzoni RA, Melkevik OR, et al. Gaming addiction, gaming engagement, and psychological health complaints among Norwegian adolescents. *Media Psychol* 2013; **16**: 115–28.
32. Brunborg GS, Hanss D, Mentzoni RA, Pallesen S. Core and peripheral criteria of video game addiction in the Game Addiction Scale for Adolescents. *Cyberpsychol Behav Soc Netw* 2015; **18**: 280–5.
33. Gunuc S. Relationships and associations between video game and Internet addictions: is tolerance a symptom seen in all conditions. *Comput Hum Behav* 2015; **49**: 517–25.
34. van Rooij AJ, Schoenmakers TM, van de Eijnden RJJM, van de Mheen D. Compulsive Internet use: the role of online gaming and other Internet applications. *J Adolesc Health* 2010; **47**: 51–7.
35. Duven EC, Müller KW, Beutel ME, Wölfling K. Altered reward processing in pathological computer gamers – ERP-results from a semi-natural gaming-design. *Brain Behav* 2015; **5**: 13–23.
36. Han DH, Bolo N, Daniels MA, Arenella L, Lyoo IK, Renshaw PF. Brain activity and desire for Internet video game play. *Compr Psychiatry* 2011; **52**: 88–95.
37. Han DH, Kim SM, Bae S, Renshaw PF, Anderson JS. Brain connectivity and psychiatric comorbidity in adolescents with Internet gaming disorder. *Addict Biol* 2017; **22**: 802–12.
38. Leeman RF, Potenza MN. A targeted review of the neurobiology and genetics of behavioral addictions: an emerging area of research. *Can J Psychiatry* 2013; **58**: 260–73.
39. Wood RTA, Gupta R, Derevensky JL, Griffiths M. Video game playing and gambling in adolescents: common risk factors. *J Child Adolesc Subst Abuse* 2004; **14**: 77–100.
40. King DL, Delfabbro PH, Griffiths MD. Cognitive behavioral therapy for problematic video game players: conceptual considerations and practice issues. *J Cyber Ther Rehabil* 2010; **3**: 261–73.
41. Charlton JP, Danforth IDW. Distinguishing addiction and high engagement in the context of online game playing. *Comput Hum Behav* 2007; **23**: 1531–48.
42. Griffiths MD, van Rooij AJ, Kardefelt-Winther D, et al. Working towards an international consensus on criteria for assessing internet gaming disorder: a critical commentary on Petry et al. (2014). *Addiction* 2016; **111**: 167–75.
43. Kuss DJ, Griffiths MD, Karila L, Billieux J. Internet addiction: a systematic review of epidemiological research for the last decade. *Curr Pharm Des* 2014; **20**: 4026–52.
44. Sim T, Gentile DA, Bricolo F, Serpelloni G, Gulam-oydeen F. A conceptual review of research on the pathological use of computers, video games, and the Internet. *Int J Ment Health Addict* 2012; **10**: 748–69.
45. Beranuy M, Carbonell X, Griffiths MD. A qualitative analysis of online gaming addicts in treatment. *Int J Ment Health Addict* 2013; **11**: 149–61.
46. King DL, Delfabbro PH, Zwaans T, Kaptis D. Clinical features and axis I comorbidity of Australian adolescent pathological Internet and video game users. *Aust N Z J Psychiatry* 2013; **47**: 1058–67.
47. Hellman M, Schoenmakers TM, Nordstrom BR, van Holst RJ. Is there such a thing as online video game addiction? A cross-disciplinary review. *Addict Res Theory* 2013; **21**: 102–12.
48. Beard KW. In: Young KS, de Abreu CN, editors. Internet Addiction: A Handbook and Guide to Evaluation and Treatment. Hoboken, NY: Wiley, 2011: 173–90.
49. Kaptis D, King DL, Delfabbro PH, Gradisar M. Withdrawal symptoms in internet gaming disorder: a systematic review. *Clin Psychol Rev* 2016; **43**: 58–66.
50. King DL, Kaptis D, Delfabbro PH, Gradisar M. Craving for internet games? Withdrawal symptoms from an 84-h abstinence from Massively Multiplayer Online gaming. *Comput Hum Behav* 2016; **62**: 488–94.
51. Kuss DJ, Griffiths MD, Pontes HM. Chaos and confusion in DSM-5 diagnosis of Internet gaming disorder: issues, concerns, and recommendations for clarity in the field. *J Behav Addict* 2017; **6**: 103–9.
52. Niu GF, Sun XJ, Subrahmanyam K, Kong FC, Tian Y, Zhou ZK. Cue-induced craving for Internet among Internet addicts. *Addict Behav* 2016; **62**: 1–5.
53. Dong G, Wang L, Du X, Potenza MN. Gaming increases craving to gaming-related stimuli in individuals with Internet gaming disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging* 2017; **2**: 404–12.
54. Ashrafioun L, Rosenberg H. Methods of assessing craving to gamble: a narrative review. *Psychol Addict Behav* 2012; **26**: 536–49.
55. Billieux J, Thorens G, Khazaal Y, Zullino D, Achab S, der Linden MV. Problematic involvement in online games: a cluster analytic approach. *Comput Hum Behav* 2015; **43**: 242–50.
56. Kiraly O, Urbán R, Griffiths MD, et al. The mediating effect of gaming motivation between psychiatric symptoms and problematic online gaming: an online survey. *J Med Internet Res* 2015; **17**: e88.
57. Ko CH, Yen JY, Chen SH, Wang PW, Chen CS, Yen CF. Evaluation of the diagnostic criteria of Internet Gaming Disorder in the DSM-5 among young adults in Taiwan. *J Psychiatr Res* 2014; **53**: 103–10.
58. Pontes HM, Griffiths MD. Measuring DSM-5 internet gaming disorder: development and validation of a short psychometric scale. *Comput Hum Behav* 2015; **45**: 137–43.
59. Rehbein F, Kliem S, Baier D, Mössle T, Petry NM. Prevalence of internet gaming disorder in German adolescents: diagnostic contribution of the nine DSM-5 criteria in a state-wide representative sample. *Addiction* 2015; **110**: 842–51.

60. Király O, Slecka P, Pontes HM, Urbán R, Griffiths MD, Demetrovics Z. Validation of the Ten-Item Internet Gaming Disorder Test (IGDT-10) and evaluation of the nine DSM-5 Internet gaming disorder criteria. *Addict Behav* 2017; **64**: 253–60.
61. Sublette VA, Mullan B. Consequences of play: a systematic review of the effects of online gaming. *Int J Ment Health Addict* 2012; **10**: 3–23.
62. Chou C, Condrón L, Belland JC. A review of the research on Internet addiction. *Educ Psychol Rev* 2005; **17**: 363–8.
63. Griffiths MD, Kuss DJ, King DL. Video game addiction: past, present and future. *Curr Psychiatry Rev* 2012; **8**: 308–18.
64. Lemmens JS, Valkenburg PM, Peter J. Psychosocial causes and consequences of pathological gaming. *Comput Hum Behav* 2011; **27**: 144–52.
65. Tokar S, Baturay MH. Antecedents and consequences of game addiction. *Comput Hum Behav* 2016; **55**: 668–79.
66. Andreassen CS, Billieux J, Griffiths MD, et al. The relationship between addictive use of social media and video games and symptoms of psychiatric disorders: a large-scale cross-sectional study. *Psychol Addict Behav* 2016; **30**: 252–62.
67. Messias E, Castro J, Saini A, Usman M, Peeples D. Sadness, suicide, and their association with video game and Internet overuse among teens: results from the Youth Risk Behavior Survey 2007 and 2009. *Suicide Life Threat Behav* 2011; **41**: 307–15.
68. King DL, Delfabbro PH, Zwaans T, Kaptis D. Sleep interference effects of pathological electronic media use during adolescence. *Int J Ment Health Addict* 2014; **12**: 21–35.
69. Weinstein AM. Computer and video game addiction – a comparison between game users and non-game users. *Am J Drug Alcohol Abuse* 2010; **36**: 268–76.
70. Byun S, Ruffini C, Mills JE, et al. Internet addiction: metasynthesis of 1996–2006 quantitative research. *Cyberpsychol Behav* 2009; **12**: 203–7.
71. Mentzoni RA, Brunborg GS, Molde H, et al. Problematic video game use: estimated prevalence and associations with mental and physical health. *Cyberpsychol Behav Soc Netw* 2011; **14**: 591–6.
72. Hur MH. Demographic, habitual, and socioeconomic determinants of Internet addiction disorder: an empirical study of Korean teenagers. *Cyberpsychol Behav* 2006; **9**: 514–25.
73. Mihara S, Higuchi S. Cross-sectional and longitudinal epidemiological studies of Internet gaming disorder: a systematic review of the literature. *Psychiatry Clin Neurosci* 2017; **71**: 425–44.
74. Gentile D. Pathological video-game use among youth ages 8 to 18: a national study. *Psychol Sci* 2009; **20**: 594–602.
75. Dreier M, Wolfling K, Duvén E, Gíralt S, Beutel ME, Müller KW. Free-to-play: about addicted whales, at risk dolphins and healthy minnows. Monetization design and Internet gaming disorder. *Addict Behav* 2017; **64**: 328–33.
76. Festl R, Scharnow M, Quandt T. Problematic computer game use among adolescents, younger and older adults. *Addiction* 2013; **108**: 592–9.
77. Griffiths M, Wood RT. Risk factors in adolescence: the case of gambling, videogame playing, and the Internet. *J Gambl Stud* 2000; **16**: 199–225.
78. Griffiths MD, King DL, Demetrovics Z. DSM-5 internet gaming disorder needs a unified approach to assessment. *Neuropsychiatry* 2014; **4**: 1–4.
79. Weinstein A, Lejoyeux M. New developments on the neurobiological and pharmacogenetic mechanisms underlying Internet and videogame addiction. *Am J Addict* 2015; **24**: 117–25.
80. Dong G, Lin X, Potenza MN. Decreased functional connectivity in an executive control network is related to impaired executive function in Internet gaming disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 2015; **57**: 76–85.
81. Lee D, Lee J, Yoon KJ, Kee N, Jung YC. Impaired anterior insular activation during risky decision making in young adults with internet gaming disorder. *NeuroReport* 2016; **27**: 605–9.
82. Qi X, Du X, Yang Y, et al. Decreased modulation by the risk level on the brain activation during decision making in adolescents with internet gaming disorder. *Front Behav Neurosci* 2015; **9**: 296.
83. Cai C, Yuan K, Yin J, et al. Striatum morphometry is associated with cognitive control deficits and symptom severity in internet gaming disorder. *Brain Imaging Behav* 2016; **10**: 12–20.
84. Yuan K, Yu D, Cai C, et al. Frontostriatal circuits, resting state functional connectivity and cognitive control in internet gaming disorder. *Addict Biol* 2016; **22**: 813–22.
85. Lin X, Dong G, Wang Q, Du X. Abnormal gray matter and white matter volume in “Internet gaming addicts”. *Addict Behav* 2015; **40**: 137–43.
86. Yoo HJ, Cho SC, Ha J, et al. Attention deficit hyperactivity symptoms and internet addiction. *Psychiatry Clin Neurosci* 2004; **58**: 487–94.
87. Bioulac S, Arfi L, Bouvard MB. Attention/deficit hyperactivity disorder and video games: a comparative study of hyperactive and control children. *Eur Psychiatry* 2008; **23**: 134–41.
88. Ng BD, Wiemer-Hastings P. Addiction to the internet and online gaming. *Cyberpsychol Behav* 2005; **8**: 110–3.
89. Kwon JH. Toward the prevention of adolescent internet addiction. In: Young KS, de Abreu CN, editors. Internet Addiction: A Handbook and Guide to Evaluation and Treatment. Hoboken, NY: Wiley, 2011: 223–44.
90. Li D, Liao A, Khoo A. Examining the influence of actual-ideal self-discrepancies, depression, and escapism, on pathological gaming among Massively Multiplayer Online adolescent Gamers. *Cyberpsychol Behav Soc Netw* 2011; **14**: 535–9.
91. King DL, Delfabbro PH. Internet gaming disorder treatment: a review of definitions of diagnosis and treatment outcome. *J Clin Psychol* 2014; **70**: 942–55.
92. Yau YH, Potenza MN. Gambling disorder and other behavioral addictions: recognition and treatment. *Harv Rev Psychiatry* 2015; **23**: 134–46.
93. Ong RHS, Peh CX, Guo S. Differential risk factors associated with adolescent addictive disorders: a comparison between substance use disorders and Internet/gaming addiction. *Int J Ment Health Addict* 2016; **14**: 993–1002.
94. Vouseoghi N, Zeinolabedin S, Sadat-Shirazi M-S, Eghbali F, Zarrindast MR. mRNA expression of dopamine receptors in peripheral blood lymphocytes of computer game addicts. *J Neural Transm (Vienna)* 2015; **122**: 1391–8.
95. Fauth-Bühler M, Mann K. Neurobiological correlates of internet gaming disorder: similarities to pathological gambling. *Addict Behav* 2015; **64**: 349–56.
96. Chen CY, Yen JY, Wang PW, Liu GC, Yen CF, Ko CH. Altered functional connectivity of the insula and nucleus accumbens in Internet gaming disorder: a resting state fMRI study. *Eur Addict Res* 2016; **22**: 192–200.
97. Fineberg NA, Chamberlain SR, Goudriaan AE, et al. New developments in human neurocognition: clinical, genetic, and brain imaging correlates of impulsivity and compulsivity. *CNS Spectr* 2014; **19**: 69–89.
98. Meng Y, Deng W, Wang H, Guo W, Li T. The prefrontal dysfunction in individuals with Internet gaming disorder: a meta-analysis of functional magnetic resonance imaging studies. *Addict Biol* 2015; **20**: 799–808.
99. Ko CH, Liu GC, Yen JY, Yen CF, Chen CS, Lin WC. The brain activations for both cue-induced gaming urge and smoking craving among subjects comorbid with Internet gaming addiction and nicotine dependence. *J Psychiatr Res* 2013; **47**: 486–93.
100. Ko CH, Yen JY, Liua SC, Huang CF, Yen CF. The associations between aggressive behaviors and Internet addiction and online activities in adolescents. *J Adolesc Health* 2009; **44**: 598–605.
101. Liu H, Rainey C, Lauer KK, et al. Peripheral blood pressure changes induced by dobutamine do not alter BOLD signals in the human brain. *NeuroImage* 2006; **30**: 745–52.
102. Kim H, Kim YK, Gwak AR, et al. Resting-state regional homogeneity as a biological marker for patients with Internet gaming disorder: a comparison with patients with alcohol use disorder and healthy controls. *Prog Neuropsychopharmacol Biol Psychiatry* 2015; **60**: 104–11.
103. Dong G, Potenza MN. A cognitive-behavioral model of Internet gaming disorder: theoretical underpinnings and clinical implications. *J Psychiatr Res* 2014; **58**: 7–11.
104. Koob GF, Volkow ND. Neurocircuitry of addiction. *Neuropsychopharmacology* 2010; **35**: 217–38.
105. Hoeft F, Watson CL, Kesler SR, Bettinger KE, Reiss AL. Gender differences in the mesocorticolimbic system during computer game-play. *J Psychiatr Res* 2008; **42**: 253–8.
106. Armstrong L, Phillips JG, Saling LL. Potential determinants of heavier internet usage. *Int J Hum Comput Stud* 2000; **53**: 537–50.
107. Ko CH, Yen JY, Chen CC, Chen SH, Yen CF. Gender differences and related factors affecting online

- gaming addiction among Taiwanese adolescents. *J Nerv Ment Dis* 2005; **193**: 273–7.
108. Bernardi S, Pallanti S. Internet addiction: a descriptive clinical study focusing on comorbidities and dissociative symptoms. *Compr Psychiatry* 2009; **50**: 510–6.
109. Schmit S, Chauchard E, Chabrol H, Sejourne N. [Evaluation of the characteristics of addiction to online video games among adolescents and young adults]. *Encéphale* 2011; **37**: 217–23. (In French).
110. Walther B, Morgenstern M, Hanewinkel R. Co-occurrence of addictive behaviours: personality factors related to substance use, gambling and computer gaming. *Eur Addict Res* 2012; **18**: 167–74.
111. Chen YL, Chen SH, Gau SS. ADHD and autistic traits, family function, parenting style, and social adjustment for Internet addiction among children and adolescents in Taiwan: a longitudinal study. *Res Dev Disabil* 2015; **39**: 20–31.
112. Wei HT, Chen MH, Huang PC, Bai YM. The association between online gaming, social phobia, and depression: an internet survey. *BMC Psychiatry* 2012; **12**: 92.
113. Lo SK, Wang CC, Fang W. Physical interpersonal relationships and social anxiety among online game players. *Cyberpsychol Behav* 2005; **8**: 15–20.
114. Porter G, Starcevic V, Berle D, Fenech P. Recognizing problem video game use. *Aust N Z J Psychiatry* 2010; **44**: 120–8.
115. Smyth JM. Beyond self-selection in video game play: an experimental examination of the consequences of massively multiplayer online role-playing game play. *Cyberpsychol Behav* 2007; **10**: 717–21.
116. Peng W, Liu M. Online gaming dependency: a preliminary study in China. *Cyberpsychol Behav Soc Netw* 2010; **13**: 329–33.
117. Cho H, Kwon M, Choi JH, et al. Development of the Internet addiction scale based on the Internet Gaming Disorder criteria suggested in DSM-5. *Addict Behav* 2014; **39**: 1361–6.
118. Zorbaz SD, Ulaş O, Kızıldağ S. Relation between video game addiction and interfamily relationships on primary school students. *Educ Sci Theory Pract* 2015; **15**: 489–97.
119. Chiu SI, Lee JZ, Huang DH. Video game addiction in children and teenagers in Taiwan. *Cyberpsychol Behav* 2004; **7**: 571–81.
120. Bathyány D, Müller KW, Benker F, Wölfling K. [Computer game playing: clinical characteristics of dependence and abuse among adolescents]. *Wien Klin Wochenschr* 2009; **121**: 502–9. (In German).
121. Frangos CC, Frangos CC, Sotiropoulos I. Problematic Internet use among Greek university students: an ordinal logistic regression with risk factors of negative psychological beliefs, pornographic sites, and online games. *Cyberpsychol Behav Soc Netw* 2011; **14**: 51–8.
122. Frölich J, Lehmkuhl G, Orawa H, Bromba M, Wolf K, Görtz-Dorten A. Computer game misuse and addiction of adolescents in a clinically referred study sample. *Comput Hum Behav* 2016; **55**: 9–15.
123. Hart GM, Johnson B, Stamm B, et al. Effects of video games on adolescents and adults. *Cyberpsychol Behav* 2009; **12**: 63–5.
124. Kim EJ, Namkoong K, Ku T, Kim SJ. The relationship between online game addiction and aggression, self-control and narcissistic personality traits. *Eur Psychiatry* 2008; **23**: 212–8.
125. Subramaniam M, Chua BY, Abidin E, et al. Prevalence and correlates of Internet gaming problem among Internet users: results from an Internet survey. *Ann Acad Med Singapore* 2016; **45**: 174–83.
126. Kaess M, Durkee T, Brunner R, et al. Pathological Internet use among European adolescents: psychopathology and self-destructive behaviours. *Eur Child Adolesc Psychiatry* 2014; **23**: 1093–102.
127. Ha JH, Yoo HJ, Cho IH, Chin B, Shin D, Kim JH. Psychiatric comorbidity assessed in Korean children and adolescents who screen positive for Internet addiction. *J Clin Psychiatry* 2006; **67**: 821–6.
128. Kuss DJ, Shorter GW, van Rooij AJ, van de Mheen D, Griffiths MD. The Internet addiction components model and personality: establishing construct validity via a nomological network. *Comput Hum Behav* 2014; **39**: 312–21.
129. Rehbein F, Baier D. Family-, media-, and school-related risk factors of video game addiction. A 5-year longitudinal study. *J Media Psychol* 2013; **25**: 118–28.
130. van Rooij AJ, Kuss DJ, Griffiths MD, Shorter GW, Schoenmakers TM, Van de Mheen D. The (co-)occurrence of problematic video gaming, substance use, and psychosocial problems in adolescents. *J Behav Addict* 2014; **3**: 157–65.
131. Hyun GJ, Han DH, Lee YS, et al. Risk factors associated with online game addiction: a hierarchical model. *Comput Hum Behav* 2015; **48**: 706–13.
132. Kardefelt-Winther D. Problematising excessive online gaming and its psychological predictors. *Comput Hum Behav* 2014; **31**: 118–22.
133. Lemmens JS, Valkenburg PM, Peter J. The effects of pathological gaming on aggressive behavior. *J Youth Adolesc* 2011; **40**: 38–47.
134. Durkee T, Kaess M, Carli V, et al. Prevalence of pathological internet use among adolescents in Europe: demographic and social factors. *Addiction* 2012; **107**: 2210–22.
135. Andreassen CS, Griffiths MD, Gjertsen SR, Krossbakken E, Kvam S, Pallesen S. The relationships between behavioral addictions and the five-factor model of personality. *J Behav Addict* 2013; **2**: 90–9.
136. Chiu SI, Hong FY, Chiu SL. An analysis on the correlation and gender difference between college students' Internet addiction and mobile phone addiction in Taiwan. *ISRN Addict* 2013; **2013**: 360607.
137. van Deursen AJAM, Bolle CL, Hegner SM, Kommers PAM. Modeling habitual and addictive smartphone behavior: the role of smartphone usage types, emotional intelligence, social stress, self-regulation, age, and gender. *Comput Hum Behav* 2015; **45**: 411–20.
138. Vadlin S, Åslund C, Hellström C, Nilsson KW. Associations between problematic gaming and psychiatric symptoms among adolescents in two samples. *Addict Behav* 2016; **61**: 8–15.
139. Andreassen CS. Online social network site addiction: a comprehensive review. *Curr Addict Rep* 2015; **2**: 175–84.
140. Griffiths MD, Davies MN, Chappell D. Demographic factors and playing variables in online computer gaming. *Cyberpsychol Behav* 2004; **7**: 479–87.
141. Hussain Z, Griffiths MD, Baguley T. Online gaming addiction: classification, prediction and associated risk factors. *Addict Res Theory* 2012; **20**: 359–71.
142. Ahmadi K, Saghaei A. Psychosocial profile of Iranian adolescents' Internet addiction. *Cyberpsychol Behav Soc Netw* 2013; **16**: 543–8.
143. Freund AM, Baltes PB. Selection, optimization, and compensation as strategies of life management: correlations with subjective indicators of successful aging. *Psychol Aging* 1998; **13**: 531–43.
144. Davis RA. A cognitive-behavioral model of pathological Internet use. *Comput Hum Behav* 2001; **17**: 187–95.
145. Jiang Q, Leung L. Effects of individual differences, awareness-knowledge, and acceptance of Internet addiction as a health risk on willingness to change Internet habits. *Soc Sci Comput Rev* 2012; **30**: 170–83.
146. Oh WO. Factors influencing internet addiction tendency among middle school students in Gyeongbuk area. *Taejeon Kambo Hakboe Chi* 2003; **33**: 1135–44.
147. Lee HW, Choi JS, Shin YC, Lee JY, Jung HY, Kwon JS. Impulsivity in Internet addiction: a comparison with pathological gambling. *Cyberpsychol Behav Soc Netw* 2012; **15**: 373–7.
148. Ding WN, Sun JH, Sun YW, et al. Trait impulsivity and impaired prefrontal impulse inhibition function in adolescents with Internet gaming addiction revealed by a go/no-go fMRI study. *Behav Brain Funct* 2014; **10**: 20.
149. Seay AF, Kraut RE. Project Massive: Self-Regulation and Problematic Use of Online Gaming. San Jose, CA: CHI 2007 Proceedings: Games, 2007.
150. Collins E, Freeman J, Chamorro-Premuzic T. Personality traits associated with problematic and non-problematic massively multiplayer online role playing game use. *Pers Individ Diff* 2012; **52**: 133–8.
151. Son DT, Yasuoka J, Poudel KC, Otsuka K, Jimba M. Massively multiplayer online role-playing games (MMORPG): association between its addiction, self-control and mental disorders among young people in Vietnam. *Int J Soc Psychiatry* 2013; **59**: 570–7.
152. Ko CH, Hsiao S, Liu GC, Yen JY, Yang MJ, Yen CF. The characteristics of decision making, potential to take risks, and personality of college students with Internet addiction. *Psychiatry Res* 2010; **175**: 121–5.
153. Mehlroof M, Griffiths MD. Online gaming addiction: the role of sensation seeking, self-control, neuroticism, aggression, state anxiety, and trait anxiety. *Cyberpsychol Behav Soc Netw* 2010; **13**: 313–6.
154. Baer S, Bogusz E, Green DA. Stuck on screens: patterns of computer and gaming station use in youth seen in a psychiatric clinic. *J Can Acad Child Adolesc Psychiatry* 2011; **20**: 86–94.
155. Yen CF, Ko CH, Yen JY, Chang YP, Cheng CP. Multi-dimensional discriminative factors for Internet addiction among adolescents regarding gender and age. *Psychiatry Clin Neurosci* 2009; **63**: 357–64.
156. Yen JY, Yen CF, Chen CC, Chen SH, Ko CH. Family factors of internet addiction and substance use

- experience in Taiwanese adolescents. *Cyberpsychol Behav* 2007; **10**: 323–9.
157. Ko CH, Yen JY, Chen CS, Yen YC, Yen CF. Predictive values of psychiatric symptoms for internet addiction in adolescents: a 2-year prospective study. *Arch Pediatr Adolesc Med* 2009; **163**: 937–43.
158. Anderson CA, Shibuya A, Ihori N, et al. Violent video game effects on aggression, empathy, and prosocial behavior in Eastern and Western countries: a meta-analytic review. *Psychol Bull* 2010; **136**: 151–73.
159. Costa PT, McCrae RR. The NEO-PI-R Personality Inventory Manual. Odessa, FL: Psychological Assessment Resources, 1985.
160. Landers RN, Lounsbury JW. An investigation of Big Five and narrow personality traits in relation to Internet usage. *Comput Hum Behav* 2006; **22**: 283–93.
161. van der Aa N, Overbeek G, Engels RC, Scholte RH, Meerkerk GJ, van den Eijnden RJ. Daily and compulsive Internet use and well-being in adolescence: a diathesis-stress model based on Big Five personality traits. *J Youth Adolesc* 2009; **38**: 765–76.
162. Ayas T. The relationship between Internet and computer game addiction level and shyness among high school students. *Educ Sci Theory Pract* 2012; **12**: 632–6.
163. Cole SH, Hooley JM. Clinical and personality correlates of MMO gaming: anxiety and absorption in problematic internet use. *Soc Sci Comput Rev* 2013; **31**: 424–36.
164. Kuss DJ, van Rooij AJ, Shorter GW, Griffiths MD, van de M. Internet addiction in adolescents: prevalence and risk factors. *Comput Hum Behav* 2013; **29**: 1987–96.
165. Müller KW, Beutel ME, Egloff B, Wölfling K. Investigating risk factors for Internet gaming disorder: a comparison of patients with addictive gaming, pathological gamblers and healthy controls regarding the Big Five personality traits. *Eur Addict Res* 2014; **20**: 129–36.
166. Öztürk C, Bektas M, Ayar D, Özgüven Öztornacı B, Yağcı D. Association of personality traits and risk of Internet addiction in adolescents. *Asian Nurs Res (Korean Soc Nurs Sci)* 2015; **9**: 120–4.
167. Wang CW, Ho RT, Chan CL, Tse S. Exploring personality characteristics of Chinese adolescents with internet-related addictive behaviors: trait differences for gaming addiction and social networking addiction. *Addict Behav* 2015; **42**: 32–5.
168. Braun B, Stopfer JM, Müller KW, Beutel ME, Egloff B. Personality and video gaming: comparing regular gamers, nongamers, and gaming addicts and differentiating between game genres. *Comput Hum Behav* 2016; **55**: 406–12.
169. Bartholow BD, Sestir MA, Davis EB. Correlates and consequences of exposure to video game violence: hostile personality, empathy, and aggressive behavior. *Pers Soc Psychol Bull* 2005; **31**: 1573–86.
170. Collins E, Freeman J. Do problematic and non-problematic video game players differ in extraversion, trait empathy, social capital and prosocial tendencies? *Comput Hum Behav* 2013; **29**: 1933–40.
171. Peters CS, Malesky LA. Problematic usage among highly-engaged players of massively multiplayer online role playing games. *CyberPsychol Behav* 2008; **11**: 481–4.
172. Lehenbauer-Baum M, Klaps A, Kovacovsky Z, Witzmann K, Zahlbruckner R, Stelina BU. Addiction and engagement: an explorative study toward classification criteria for internet gaming disorder. *Cyberpsychol Behav Soc Netw* 2015; **18**: 343–9.
173. Yan W, Li Y, Sui N. The relationship between recent stressful life events, personality traits, perceived family functioning and internet addiction among college students. *Stress Health* 2013; **30**: 3–11.
174. Huh S, Bowman ND. Perception of and addiction to online games as a function of personality traits. *J Media Psychol* 2008; **13**: 1–31.
175. Wölfling K, Müller K, Beutel M. Diagnostic measures: scale for the assessment of internet and computer game addiction (AICA-S). In: Mücken D, Teske A, Rehbein F, te Wildt B, editors. Prevention, Diagnostics, and Therapy of Computer Game Addiction. Lengerich: Pabst Science, 2010: 212–5.
176. Stopfer JM, Braun B, Müller KW, Egloff B. Narcissus plays video games. *Pers Individ Diff* 2015; **87**: 212–8.
177. Allison SE, von Wahlde L, Shockley T, Gabbard GO. The development of the self in the era of the Internet and role-playing fantasy games. *Am J Psychiatry* 2006; **163**: 381–5.
178. Soto CJ, John OP, Gosling SD, Potter J. Age differences in personality traits from 10 to 65: Big Five domains and facets in a large cross-sectional sample. *J Pers Soc Psychol* 2011; **100**: 330–48.
179. Caplan SE. Relations among loneliness, social anxiety, and problematic Internet use. *Cyberpsychol Behav* 2007; **10**: 234–42.
180. Schimmenti A, Passanisi A, Gervasi AM, Manzella S, Famà FI. Insecure attachment attitudes in the onset of problematic internet use among late adolescents. *Child Psychiatry Hum Dev* 2014; **45**: 588–95.
181. Yee N. Motivations for play in online games. *Cyberpsychol Behav* 2006; **9**: 772–5.
182. Hellström C, Nilsson KW, Leppert J, Åslund C. Influences of motives to play and time spent gaming on the negative consequences of adolescent online computer gaming. *Comput Hum Behav* 2012; **28**: 1379–87.
183. Wan CS, Chiou WB. Psychological motives and online games addiction: a test of flow theory and humanistic needs theory for Taiwanese adolescents. *CyberPsychol Behav* 2006; **9**: 317–24.
184. Hilgard J, Engelhardt CR, Bartholow BD. Individual differences in motives, preferences, and pathology in video games: the gaming attitudes, motives, and experiences scales (GAMES). *Front Psychol* 2013; **4**: 608.
185. Demetrovics Z, Urbán R, Nagygyörgy K, et al. The development of the Problematic Online Gaming Questionnaire (POGQ). *PLoS ONE* 2012; **7**: e36417.
186. Park MH, Park EJ, Choi J, et al. Preliminary study of Internet addiction and cognitive function in adolescents based on IQ tests. *Psychiatry Res* 2011; **190**: 275–81.
187. Griffiths MD. Videogame addiction: fact or fiction? In: Willoughby T, Wood E, editors. Children's Learning in a Digital World. Oxford: Blackwell, 2008: 85–103.
188. Palmgreen P, Wenner LA, Rosengren K. Uses and gratifications research: the past ten years. In: Rosengren K, Wenner L, Palmgreen P, editors. Media Gratifications Research. Beverly Hills, CA: Sage, 1985: 11–37.
189. Bandura A. Social Foundations of Thought and Action: A Social-Cognitive Theory. Englewood Cliffs, NJ: Prentice Hall, 1986.
190. Bandura A. Self-efficacy: The Exercise of Control. New York, NY: Freeman, 1997.
191. Smith LJ, Gradisar M, King DL. Parental influences on adolescent video game play: a study of accessibility, rules, limit setting, monitoring, and cybersafety. *Cyberpsychol Behav Soc Netw* 2015; **18**: 273–9.
192. Yen JY, Ko CH, Yen CF, Wu HY, Yang MJ. The comorbid psychiatric symptoms of Internet addiction: attention deficit and hyperactivity disorder (ADHD), depression, social phobia, and hostility. *J Adolesc Health* 2007; **41**: 93–8.
193. Kwon JH, Chung CS, Lee J. The effects of escape from self and interpersonal relationship on the pathological use of Internet games. *Community Ment Health J* 2011; **47**: 113–21.
194. Mölle T, Rehbein F. Predictors of problematic video game usage in childhood and adolescence. *Sucht* 2013; **59**: 153–64.
195. Haagsma MC, Caplan SE, Peters O, Pieterse ME. A cognitive-behavioral model of problematic online gaming in adolescents aged 12–22 years. *Comput Hum Behav* 2013; **29**: 202–9.
196. Haagsma MC, King DL, Pieterse ME, Peters O. Assessing problematic video gaming using the theory of planned behavior: a longitudinal study of Dutch young people. *Int J Ment Health Addict* 2013; **11**: 172–85.
197. Hull DC, Williams GA, Griffiths MD. Video game characteristics, happiness and flow as predictors of addiction among video game players: a pilot study. *J Behav Addict* 2013; **2**: 145–52.
198. Kuss DJ, Louws J, Wiers RW. Online game addiction? Motives predict addictive play behavior in massively multiplayer online role-playing games. *Cyberpsychol Behav Soc Netw* 2012; **15**: 480–5.
199. Yilmaz FGK, Yilmaz R, Teker N, Keser H. Prediction of Internet addiction of university students based on various variables. *World J Educ Technol* 2014; **6**: 75–87.
200. Männikkö N, Billieux J, Kaariäinen M. Problematic digital gaming behavior and its relation to the psychological, social and physical health of Finnish adolescents and young adults. *J Behav Addict* 2015; **4**: 281–8.
201. Billieux J, Chanaal J, Khazaal Y, et al. Psychological predictors of problematic involvement in massively multiplayer online role-playing games: illustration in a sample of male cybercafé players. *Psychopathology* 2011; **44**: 165–71.
202. Rho MJ, Jeong JE, Chun JW, et al. Predictors and patterns of problematic Internet game use using a decision tree model. *J Behav Addict* 2016; **5**: 500–9.
203. Bijvank MN, Konijn EA, Bushman BJ, Roelofsma PH. Age and violent-content labels make video games forbidden fruits for youth. *Pediatrics* 2009; **123**: 870–6.
204. Doornwaard SM, van den Eijnden RJ, Baams L, Vanwesenbeeck I, ter Bogt TF. Lower psychological well-being and excessive sexual interest predict symptoms of compulsive use of sexually explicit Internet material

- among adolescent boys. *J Youth Adolesc* 2016; **45**: 73–84.
205. LaRose R. The problem of media habits. *Commun Theory* 2010; **22**: 194–222.
 206. Hartmann T, Jung Y, Vorderer P. What determines video game use? The impact of users' habits, addictive tendencies, and intentions to play. *J Media Psychol* 2012; **24**: 19–30.
 207. Eichenbaum A, Kattner F, Bradford D, Gentile DA, Green CS. Role-playing and real-time strategy games associated with greater probability of Internet gaming disorder. *Cyberpsychol Behav Soc Netw* 2015; **18**: 480–5.
 208. Thomas NJ, Martin FH. Video-arcade game, computer game and internet activities of Australian students: participation habits and prevalence of addiction. *Aust J Psychol* 2010; **62**: 59–66.
 209. King DL, Delfabbro PH, Griffiths MD, Gradisar M. Assessing clinical trials of internet addiction treatment: a systematic review and CONSORT evaluation. *Clin Psychol Rev* 2011; **31**: 1110–6.
 210. Elliott L, Ream G, McGinsky E, Dunlap E. The contribution of game genre and other use patterns to problem video game play among adult video gamers. *Int J Ment Health Addict* 2015; **18**: 480–5.
 211. Hsu SH, Wen MH, Wu MC. Exploring user experiences as predictors of MMORPG addiction. *Comput Educ* 2009; **53**: 990–9.
 212. Griffiths MD, Beranuy M. Adicción a los videojuegos: una breve revisión psicológica. *Rev Psychoter* 2009; **73**: 33–49.
 213. Ak S, Koruklu N, Yilmaz Y. A study on Turkish adolescent's Internet use: possible predictors of Internet addiction. *Cyberpsychol Behav Soc Netw* 2013; **16**: 205–9.
 214. King DL, Delfabbro PH, Griffiths MD. Clinical interventions for technology-based problems: excessive Internet and video game use. *J Cogn Psychoter* 2012; **26**: 43–56.
 215. Haagsma MC, Pieterse ME, Peters O. The prevalence of problematic video gamers in The Netherlands. *Cyberpsychol Behav Soc Netw* 2012; **15**: 162–8.
 216. Dong G, Lu Q, Zhou H, Zhao X. Precursor or sequel: pathological disorders in people with Internet addiction disorder. *PLoS ONE* 2011; **6**: e14703.
 217. Hahn T, Notebaert KH, Dressler T, Kowarsch L, Reif A, Fallbatter AJ. Linking online gaming and addictive behavior: converging evidence for a general reward deficiency in frequent online gamers. *Front Behav Neurosci* 2014; **8**: 385.
 218. Goudriaan AE, Oosterlaan J, de Beurs E, van den Brink W. Neurocognitive functions in pathological gambling: a comparison with alcohol dependence. Tourette syndrome and normal controls. *Addiction* 2006; **101**: 534–47.
 219. Dawkins L, Powell JH, Pickering A, Powell J, West R. Patterns of change in withdrawal symptoms, desire to smoke, reward motivation and response inhibition across 3 months of smoking abstinence. *Addiction* 2009; **104**: 850–8.
 220. Field M, Cox WM. Attentional bias in addictive behaviors: a review of its development, causes, and consequences. *Drug Alcohol Depend* 2008; **97**: 1–20.
 221. Boyer M, Dickerson M. Attentional bias and addictive behaviour: automaticity in a gambling-specific modified Stroop task. *Addiction* 2003; **98**: 61–70.
 222. Lusher J, Chandler C, Ball D. Alcohol dependence and the alcohol Stroop paradigm: evidence and issues. *Drug Alcohol Depend* 2004; **75**: 225–31.
 223. Wang L, Wu L, Lin X, et al. Dysfunctional default mode network and executive control network in people with Internet gaming disorder: independent component analysis under a probability discounting task. *Eur Psychiatry* 2016; **34**: 36–42.
 224. Lin X, Zhou H, Dong G, Du X. Impaired risk evaluation in people with Internet gaming disorder: fMRI evidence from a probability discounting task. *Prog Neuropsychopharmacol Biol Psychiatry* 2015; **56**: 142–8.
 225. Pawlikowski M, Brand M. Excessive Internet gaming and decision making: do excessive World of Warcraft players have problems in decision making under risky conditions? *Psychiatry Res* 2011; **188**: 428–33.
 226. Dreier M, Wölfling K, Müller KW. Psychological research and a sociological perspective on problematic and addictive computer game use in adolescence. *Int J Child Adolesc Health* 2013; **6**: 421–35.
 227. Müller KW, Dreier M, Wölfling K. Excessive and addictive use of the internet – prevalence, related contents, predictors, and psychological consequences. In: Reinecke L, Oliver MB, editors. The Routledge Handbook of Media Use and Well-Being. New York, NY: Routledge, Taylor and Francis, 2016: 223–36.
 228. Brand M, Young KS, Laier C, Wölfling K, Potenza MN. Integrating psychological and neurobiological considerations regarding the development and maintenance of specific Internet-use disorders: an interaction of person-affect-cognition-execution (I-PACE) model. *Neurosci Biobehav Rev* 2016; **71**: 252–66.
 229. Nuyens F, Deleuze J, Muraire P, Griffiths MD, Kuss DJ, Billieux J. Impulsivity in Multiplayer Online Battle Arena (MOBA) gamers: preliminary results on experimental and self-report measures. *J Behav Addict* 2016; **5**: 351–6.
 230. Kuss DJ, Griffiths MD. Online gaming addiction in children and adolescents: a review of empirical research. *J Behav Addict* 2012; **1**: 3–22.
 231. Cao F, Su L, Liu T, Gao X. The relationship between impulsivity and Internet addiction in a sample of Chinese adolescents. *Eur Psychiatry* 2007; **22**: 466–71.
 232. Xu Z, Yuan Y. The impact of motivation and prevention factors on game addiction. Special Interest Group on Human-Computer Interaction 2008 Proceedings 15.
 233. Cao H, Sun Y, Wan Y, Hao J, Tao F. Problematic Internet use in Chinese adolescents and its relation to psychosomatic symptoms and life satisfaction. *Ann Acad Med Singapore* 2010; **39**: 822–9.
 234. Grüsser SM, Thalemann R, Albrecht U, Thalemann CN. [Excessive Computernutzung im Kindesalter – Ergebnisse einer psychometrischen Erhebung]. *Wien Klin Wochenschr* 2005; **117**: 188–95. (In German).
 235. Wölfling K, Müller K, Beutel M. [Reliabilität und Validität der Skala zum Computerspielverhalten (CSV-S)]. *Psychoter Psychosom Med Psychol* 2010; **61**: 216–24. (In German).
 236. Müller KW, Ammerschläger M, Freisleder FJ, Beutel ME, Wölfling K. [Addictive Internet use as a comorbid disorder among clients of an adolescent psychiatry – prevalence and psychopathological symptoms]. *Z Kinder Jugendpsychiatr Psychoter* 2012; **40**: 331–7. (In German).
 237. Zamani E, Kheradmand A, Chesmi M, Hedayati N. Comparing the social skills of students addicted to computer games with normal students. *Addict Health* 2010; **2**: 59–69.
 238. Lemmens JS, Valkenburg PM, Gentile DA. The Internet gaming disorder scale. *Psychol Assess* 2015; **27**: 567–82.
 239. Bakken JJ, Wenzel HG, Götestam KG, Johansson A, Oren A. Internet addiction among Norwegian adults: a stratified probability sample study. *Scand J Psychol* 2017; **50**: 121–7.
 240. Arnesen AA. Video Game Addiction among Young Adults in Norway: Prevalence and Health. 2010. <http://bora.uib.no/bitstream/handle/1956/4114/69119310.pdf?sequence=1&isAllowed=y> (accessed 22 July 2017).
 241. Wittek CT, Finserås TR, Pallesen S, et al. Prevalence and predictors of video game addiction: a study based on a National representative sample of gamers. *Int J Ment Health Addict* 2016; **14**: 672–86.
 242. Choo H, Gentile DA, Sim T, Li D, Khoo A, Liau AK. Pathological video-gaming among Singaporean youth. *Ann Acad Med Singapore* 2010; **39**: 822–9.
 243. Kim K, Ryu E, Chon MY, et al. Internet addiction in Korean adolescents and its relation to depression and suicidal ideation: a questionnaire survey. *Int J Nurs Stud* 2006; **43**: 185–92.
 244. Lee C, Han S. Development of the scale for diagnostic online game addiction. 3rd WSEAS/IASME International Conference on Educational Technologies; Arachon, F2007.
 245. Park SK, Kim JY, Cho CB. Prevalence of Internet addiction and correlates with family factors among South Korean adolescents. *Adolescence* 2008; **43**: 895–909.
 246. Jeong EJ, Kim DH. Social activities, self-efficacy, game attitudes, and game addiction. *Cyberpsychol Behav Soc Netw* 2011; **14**: 213–21.
 247. Tejeiro Salguero RA, Morán RM. Measuring problem video game playing in adolescents. *Addiction* 2002; **97**: 1601–6.
 248. Lopez-Fernandez O, Honrubia-Serrano ML, Baguley T, Griffiths MD. Pathological video game playing in Spanish and British adolescents: towards the exploration of Internet Gaming Disorder symptomatology. *Comput Hum Behav* 2014; **41**: 304–12.
 249. Ko CH, Yen JY, Yen CF, Lin HC, Yang MJ. Factors predictive for incidence and remission of internet addiction in young adolescents: a prospective study. *CyberPsychol Behav* 2007; **10**: 545–51.
 250. Wan CS, Chiou WB. The motivations of adolescents who are addicted to online games: a cognitive perspective. *Adolescence* 2007; **42**: 179–97.
 251. Fisher S. Identifying video game addiction in children and adolescents. *Addict Behav* 1994; **19**: 545–53.
 252. Griffiths MD, Sutherland I. Adolescent gambling and drug use. *J Community Appl Soc Psychol* 1998; **8**: 423–7.

RESUMEN

TRASTORNO DEL JUEGO EN INTERNET EN NIÑOS Y ADOLESCENTES: UNA REVISIÓN SISTEMÁTICA

OBJETIVO El trastorno del juego por Internet (IGD, siglas en inglés) es un trastorno grave que provoca y mantiene un deterioro personal y social relevante. El IGD ha sido considerado en vista de conceptos heterogéneos e incompletos. Por lo tanto, nosotros revisamos la literatura científica sobre el IGD para proporcionar una visión general enfocada en sus definiciones, síntomas, prevalencia y etiología.

MÉTODO Se revisaron sistemáticamente las bases de datos ERIC, PsyARTICLES, PsycINFO, PSYINDEX y PubMed en el período desde enero del 1991 a agosto del 2016, y adicionalmente se identificaron referencias secundarias.

RESULTADOS La definición propuesta en el Manual Diagnóstico y Estadístico de los Trastornos Mentales, Quinta Edición proporciona un buen punto de partida para diagnosticar el IGD, pero conlleva algunas desventajas. El desarrollo del IGD requiere de varios factores internos que interactúan, tales como deficiencias en el yo la regulación del estado de ánimo y las recompensas, problemas en la toma de decisiones y factores externos, tales como antecedentes de familias deficientes y las habilidades sociales. Además, factores específicos relacionados con el juego pueden promover el IGD. Resumiendo, el conocimiento etiológico, sugerimos un modelo integrado del IGD dilucidando la interacción de factores internos y externos.

INTERPRETACIÓN Hasta ahora, el concepto del IGD y las vías que conducen a él no son del todo claras. En particular, faltan estudios de seguimiento a largo plazo. El IGD debe entenderse como un peligro de trastorno con un fondo psicosocial complejo.

RESUMO

DESORDEM DO JOGO DE INTERNET EM CRIANÇAS E ADOLESCENTES: UMA REVISÃO SISTEMÁTICA

OBJETIVO A desordem do jogo de internet (DJI) é uma desordem séria causando e mantendo dificuldades pessoais e sociais. A DJI tem sido considerada em vista de conceitos heterogêneos e incompletos. Portanto, revisamos a literatura científica sobre DJI para fornecer uma visão geral com foco em definições, sintomas, prevalência e etiologia.

MÉTODO Revisamos sistematicamente as bases de dados ERIC, PsyARTICLES, PsycINFO, PSYINDEX, e PubMed para o período de janeiro de 1991 a agosto de 2016, e adicionalmente identificamos referências secundárias.

RESULTADOS A definição proposta pelo Manual Diagnóstico e Estatístico de Desordens Mentais, quinta edição, fornece um bom ponto de partida para diagnosticar DJI, mas traz algumas desvantagens. Desenvolver DJI requer a interação de vários fatores internos como eu deficiente, regulação de humor e recompensa, problemas com tomada de decisões, e fatores externos como suporte familiar e habilidades sociais deficientes. Adicionalmente, fatores específicos relacionados aos jogos podem levar a DJI. Sintetizando o conhecimento etiológico, sugerimos um modelo integrado de DJI para elucidar a relação entre fatores internos e externos.

INTERPRETAÇÃO Até o momento, o conceito de DJI e as vias levando a ela não são inteiramente claros. Em particular, faltam estudos com acompanhamento de longo prazo. A DJI deve ser entendida como uma desordem perigosa, com complexas bases psicossociais.