

Chapter 1

Mathematical Preliminaries

Le interazioni matematiche sono complesse e non triviali (vedi un po' di articoli di introduzione a AQFT per ispirarti)

Tendenzialmente le teorie quantistiche di campi moderne sono di Quantizzazione.. Quindi richiedono di specificare bene la struttura del campo classico (vedi intro di Mangiaratti shardashivly)

Gli strumenti matematici per raccontare la teoria dei campi classici sono essenzialmente 3: Fibrati, S-T G-H, LDOP e GHOP.



IN questo paper non ci soffermeremo sulle strutture del framework puramente quantistico (* algebre e quant'altro). Per un primer vedere articolo di Dappiaggi o Libro Adv aqft.

Diamo per scontato un background di base in Geometria differenziale e derivate esterne (algebre di Grassman? global calculus? non so come chiamarlo!).

Potrei avere la tentazione a provare ad usare un po' di linguaggio basilare delle categorie... la mia fonte \hat{A} Joy of Cat.



Stile: Intro lapiadaria ai 3 argomenti (bundle cinematica di campo, Glob iper stage per descrivere dinamica di tipo propagativo, Operatori tipo onda). Poi smitragliata di definizioni come faceva Penati.

1.1 Fiber Bundles

 intro

1.1.1 Formal Definition

Although it would be possible to present the concept *bundle* in a more general way through the language of categories, for our argument will be sufficient to consider

only the case of *smooth bundles*.

Definition 1: Fiber Bundle

A *Fiber Bundle* consists in a 4-ple (E, M, Q, π) where:

- E, M, Q : smooth manifolds called respectively *Total Space*, *Base Space*, *Typical Fiber*.
- $\pi : E \rightarrow M$ continuous smooth function (called *Bundle Projection*)

Endowed with a *Local Trivialization*:

- $\forall x \in E \exists$ a couple (U, χ) (called *local trivialization*)
 - U : neighborhood of x
 - $\chi : \pi^{-1}(U) \rightarrow U \times Q$: diffeomorphism^{a b}

such that: $p_1 \circ \chi = \pi|_{\pi^{-1}(U)}$.


$$\begin{array}{ccc} \pi^{-1}(U) & \xrightarrow{\chi} & U \times Q \\ \pi \downarrow & \swarrow p_1 & \\ U & & \end{array}$$

i.e: the following graph commutes:

^a surjectivity $\Rightarrow \pi^{-1}(U) \neq \emptyset$.

^b cartesian product of topological space is a topological space with the direct product topology.

Figure 1.1: The complete fiber bundle Structure.



Notation fixing

It is customary to refer to a vector bundle specifying only its total space:

$$E = (E, \pi, M; Q)$$

In the following we adopt this convention whenever this does not lead to misunderstandings.

Observation 1

For all $p \in M$ we refer to the submanifold $E_p := \pi^{-1}(p) \subset E$ as *fiber over the point p* .
 Every fiber E_p is diffeomorphic to the typical fiber F through the local trivialization charts.

Notation fixing

We say that a smooth bundle E is (*globally*) *trivial* if $E \simeq M \times Q$ i.e there exists a trivialization of E which is defined everywhere. Note that definition 1 prescribes the existence of local trivializations only.

When a smooth fiber bundle $(E, \pi, M; Q)$ is considered, in addition to the typical functions of the bundle (π, χ_α) should be taken in account also the local charts $(U_\alpha, \phi_{\alpha_k})_{k=E, M, Q}$ provided by the atlases of E, M and Q :

Definition 2: Bundle atlas

Collection local chart which trivializes E . I.e. triples $(U_\alpha, \psi_\alpha, \chi_\alpha)$ where:

- U_α open set in M such that $\bigcup_\alpha U_\alpha \supseteq M$.
- χ_α is a local trivialization.
- (U_α, ψ_α) local chart on E constructed from charts on the base and fiber manifold:

$$\psi_\alpha^{(E)} = \psi_\alpha^{(M)} \times \psi_\alpha^{(Q)} \circ \chi_\alpha$$

Endowing the bundles manifolds with other additional structures, can be introduced important subclasses of smooth bundles:

Definition 3: Vector Bundle

Is a smooth bundle $E = (E, \pi, M; V)$ such that:

- The typical fiber manifold V is a finite dimensional vector space.
- All the trivialization χ_α are diffeomorphism such that:

$$\chi_\alpha|_{\pi^{-1}(p)} \in \mathbb{GL}(n, \mathbb{R})$$

1.1.2 Cross Sections

The notion of bundle is particularly interesting from the perspective of physics because provides the rigorous description of a Q -valued field over the space M :

Definition 4: Smooth Section

Function $\phi : M \rightarrow E$ such that:

- ϕ smooth.
- $\phi \cdot \pi = \mathbb{1}_M$

Notation fixing

We refer to:

- *Global section* $\Leftrightarrow \text{dom}(\phi) = B$
- *Local section* $\Leftrightarrow \text{dom}(\phi) \subset B^a$

We denote the set of all the smooth sections of the bundle E as:

$$\Gamma^\infty(E)$$

^aUsually the domain is an open set of B

Observation 2

$\Gamma^\infty(E)$ is an infinite dimensional manifolds.

If E is a vector bundle $\Gamma^\infty(E)$ is an infinite dimensional vector space, therefore we can introduce the decomposition on a base:

1.1.3 Mapping between Bundles**1.1.4 Tangent Bundles**

Tautological one-form and symplectic form.

Fomm: As a mathematical curiosity, we note that the cotangent bundle of any manifold is orientable. Indeed, it carries a symplectic structure and hence a volume element.

1.1.5 Jet Bundles

$Ob(A) \text{ hom}(A, B) \text{ Mor coddomran}$

1.2 Globally Hyperbolic Space-times

⚠ Mettere solo le definizioni che uso prese dagli articoli di review delle Fonti

Appunti che mi ero preso scrivendo il secondo capitolo:

This condition is strictly connected to the dynamic behaviour of the system.

⚠ Def di dominio di dipendenza footnote di definizione di spazio tempo
def cauchy surface Remark causal future past def globally hyperbolic Teorema sulle caratterizzazioni

Notation fixing

We denote the set of all the cauchy surfaces as $\mathcal{P}_C(M)$.

Glob iperbolico determina la fogliazione dello spazio tempo per superfici di cauchy
La superficie di cauchy Σ questa:

Definition 5: Cauchy surface

questo da la possibilit  della buona posizione dei problemi di cauchy.. fisicamente
 Σ la condizione minima per definire i dati iniziali dell'evoluzione dinamica. definisco
data...

Rapporto con la condizione sugli operatori...

No! La definizione di green hyperbolicity non garantisce invece l'esistenza
e unicit  del problema di cauchy associata

e non solo, anche l'esistenza degli operatori di green associati che sono
ingrediente fondamentale della costruzione di peierls

⚠ $M \hat{=}$ glob iper e $P \hat{=}$ green iper per tener conto del comportamento
propagativo definire sup cauchy definire s-t iperbolico (solo la caratteriz-
zazione di ammettere una sup di cauchy) definire op green iperbolico su spazio
tempo iperbolico (cio  ha delle green ope) Propr di buona definizione es-
istenza e unicit  della soluzione

Di particolare ricorrenza fisica sono gli operatori normally iperbolico espres-
sione in coordinate esempio K-g!

⚠ Far notare che minkowski e tanti esempi importanti sono GH

Observation 3

(che serve dopo) lo spazio R è banalmente iperbolico in quanto tutti i punti posso essere visti come superfici di cauchy.

(sono ripetizioni inutili per la tesi, sono informazioni che si ritrovano ovunque... sono informazioni adatta al knowledge base)

Recurring definitions in general Relativity (excluding the general smooth manifold prolegomena).

Definition 6: Space-Time

A quadruple (M, g, σ, τ) such that:

- (M, g) is a time-orientable n -dimensional manifold ($n > 2$)
- σ is a choice of orientation
- τ is a choice of time-orientation

Definition 7: Lorentzian Manifold

A pair (M, g) such that:

- M is a n -dimensional ($n \geq 2$), Hausdorff, second countable, connected, orientable smooth manifold.
- g is a Lorentzian metric.

Definition 8: Metric

A function on the bundle product of TM with itself:

$$g : TM \times_M TM \rightarrow \mathbb{R}$$

such that the restriction on each fiber

$$g_p : T_p M \times T_p M \rightarrow \mathbb{R}$$

is a non-degenerate bilinear form.

Notation fixing

- *Riemman* if the sign of g is positive definite, *Pseudo-Riemman* otherwise.
- *Lorentzian* if the signature is $(+, -, \dots, -)$ or equivalently $(-, +, \dots, +)$.

Observation 4: Causal Structure

If a smooth manifold is endowed with a Lorentzian manifold of signature $(+, -, \dots, -)$ then the tangent vectors at each point in the manifold can be classed into three different types.

Notation fixing

$\forall p \in M, \quad \forall X \in T_p M$, the vector is:

- *time-like* if $g(X, X) > 0$.
- *light-like* if $g(X, X) = 0$.
- *space-like* if $g(X, X) < 0$.

Observation 5: Local Time Orientability

$\forall p \in M$ the timelike tangent vectors in p can be divided into two equivalence classes taking

$$X \sim Y \text{ iff } g(X, Y) > 0 \quad \forall X, Y \in T_p^{\text{time-like}} M :$$

We can (arbitrarily) call one of these equivalence classes "future-directed" and call the other "past-directed". Physically this designation of the two classes of future- and past-directed timelike vectors corresponds to a choice of an arrow of time at the point. The future- and past-directed designations can be extended to null vectors at a point by continuity.

Definition 9: Time-orientation

A global tangent vector field $\mathfrak{t} \in \Gamma^\infty(TM)$ over the Lorentzian manifold M such that:

- $\text{supp}(\mathfrak{t}) = M$
- $\mathfrak{t}(p)$ is time-like $\forall p \in M$.

Observation 6

The fixing of a time-orientation is equivalent to a consistent smooth choice of a local time-direction.

Definition 10: Time-Orientable Lorentzian Manifold

A Lorentzian Manifold (M, g) such that exist at least one time-orientation $\mathfrak{t} \in$

$\Gamma^\infty(TM)$.

Notation fixing

Consider a piece-wise smooth curve $\gamma : \mathbb{R} \supset I \rightarrow M$ is called:

- *time-like* (resp. light-like, space-like) iff $\dot{\gamma}(p)$ is time-like (resp. light-like, space-like) $\forall p \in M$.
- *causal* iff $\dot{\gamma}(p)$ is nowhere spacelike.
- *future directed* (resp. past directed) iff is causal and $\dot{\gamma}(p)$ is future (resp. past) directed $\forall p \in M$.

Definition 11: Chronological $\begin{smallmatrix} \text{future} \\ \text{past} \end{smallmatrix}$ of a point

Are two subset related to the generic point $p \in M$:

$$\mathbf{I}_M^\pm(p) := \{q \in M \mid \exists \gamma \in C^\infty((0, 1), M) \text{ time-like } \begin{smallmatrix} \text{future} \\ \text{past} \end{smallmatrix} - \text{directed} : \gamma(0) = p, \gamma(1) = q\}$$

Definition 12: Causal $\begin{smallmatrix} \text{future} \\ \text{past} \end{smallmatrix}$ of a point

Are two subset related to the generic point $p \in M$:

$$\mathbf{J}_M^\pm(p) := \{q \in M \mid \exists \gamma \in C^\infty((0, 1), M) \text{ causal } \begin{smallmatrix} \text{future} \\ \text{past} \end{smallmatrix} - \text{directed} : \gamma(0) = p, \gamma(1) = q\}$$

Notation fixing

Former concept can be naturally extended to subset $A \subset M$:

- $\mathbf{I}_M^\pm(A) = \bigcup_{p \in A} \mathbf{I}_M^\pm(p)$
- $\mathbf{J}_M^\pm(A) = \bigcup_{p \in A} \mathbf{J}_M^\pm(p)$

Definition 13: Achronal Set

Subset $\Sigma \subset M$ such that every inextendible timelike curve intersect Σ at most once.

Definition 14: $\begin{smallmatrix} \text{future} \\ \text{past} \end{smallmatrix}$ Domain of dependence of an Achronal set

The two subset related to the generic achronal set $\Sigma \subset M$:

$$\mathbf{D}_M^\pm(\Sigma) := \{q \in M \mid \forall \gamma_{\text{future}}^{\text{past}} \text{ inextendible causal curve passing through } q : \gamma(I) \cap \Sigma \neq \emptyset\}$$

Notation fixing

$\mathbf{D}_M(\Sigma) := \mathbf{D}_M^+(\Sigma) \cup \mathbf{D}_M^-(\Sigma)$ is called *total domain of dependence*.


Definition 15: Cauchy Surface

Is a subset $\Sigma \subset M$ such that:

- closed
- achronal
- $\mathbf{D}_M(\Sigma) \equiv M$


1.3 Green Hyperbolic Operators

 Mettere solo le definizioni che uso prese dagli articoli di review delle Fonti

 Pensavo di utilizzare la definizione di Green hyperbolic data da Bar che si avvale del concetto di formally dual (che non richiede la presenza del pairing) invece di quella usata in Advances AQFT che richiede solo che ammetta almeno un G^\pm per poi dimostrare tramite teorema che se \tilde{A} anche autoaggiunto vale l'unicità. Si tratta solo di una piccola sfumatura.. Deve essere chiarito che in tutto ciò che faccio interessano che

$$\forall P \exists ! G^\pm$$

. Che poi questa condizione derivi da GH secondo bar o Gh secondo dap+selfadj. \tilde{A} una di quelle questioni propriamente matematiche che poco interessa ai fisici della commissione.

 Devo richiedere che il green operator sia unico? sia negli schemi di quantizzazione che nella definizione di peierls faccio largo uso dell'unicità. Per provare questa unicità si passa per la definizione di una forma bilineare che permette di parlare di aggiunto formale e quindi avvalersi del teorema.

Green-hyperbolic operators are not necessarily hyperbolic in any PDE-sense and

that they cannot be characterized in general by well-posedness of a Cauchy problem. [?] [?]

(sono ripetizioni inutili per la tesi, sono informazioni che si ritrovano ovunque... sono informazioni adatta al knowledge base) Basic Definition in L.P.D.O. on smooth vector sections.

Consider $F = F(M, \pi, V)$, $F' = F'(M, \pi', V')$ two linear vector bundle over M with different typical fiber

Definition 16: Linear Partial Differential operator (of order at most $s \in \mathbb{N}_0$)

Linear map $L : \Gamma(F) \rightarrow \Gamma(F')$ such that:

$\forall p \in M$ exists:

- (U, ϕ) local chart on M .
- (U, χ) local trivialization of F
- (U, χ') local trivialization of F'

for which:

$$L(\sigma|_U) = \sum_{|\alpha| \leq s} A_\alpha \partial^\alpha \sigma \quad \forall \sigma \in \Gamma(M)$$

Remark:

(multi-index notation)

A multi-index is a natural valued finite dimensional vector $\alpha = (\alpha_0, \dots, \alpha_{n-1}) \in \mathbb{N}_0^n$ with $n < \infty$.

On \mathbb{R}^n a general differential operator can be identified by a multi-index:

$$\partial^\alpha = \prod_{\mu=0}^{n-1} \partial_\mu^{\alpha_\mu}$$

(Until the Schwartz theorem holds, the order of derivation is irrelevant.)

The order of the multi-index is defined as:

$$|\alpha| := \sum_{\mu=0}^{n-1} \alpha_\mu$$

????????????????????

Hp:

Proposition 1.3.1 (Existence and uniqueness for the Cauchy Problem)

$M = (M, g, o, t)$ a globally hyperbolic space-time.

- $\Sigma \subset M$ a spacelike cauchy surface with future-pointing unit normal vector field \vec{n} .

Th:

Observation 7

"Green-hyperbolic operators are not necessarily hyperbolic in any PDE-sense and that they cannot be characterized in general by well-posedness of a Cauchy problem. " [?] [?]

However the existence and uniqueness can be proved for the large class of the *Normally-Hyperbolic Operators*.