Proof portfolio

Duncan Freeman 933-135-442

November 13, 2019

1 Multiplication and Sets

Prove the following inequality:

$$\{6x|x\in\mathbb{Z}\}=\{2x|x\in\mathbb{Z}\}\bigcap\{3x|x\in\mathbb{Z}\}$$

 $\forall c \in \mathbb{Z} (6c \in \{2x | x \in \mathbb{Z}\} \land 6c \in \{3x | x \in \mathbb{Z}\})$. This is because 6c = (2)(3)c, so any 6c can be summoned from 2c and 3c.

Because 6 is *only* divisible by 2 and 3, there are no numbers for which the intersection of $\{2x|x\in\mathbb{Z}\}$ and $\{3x|x\in\mathbb{Z}\}$ would not also be a multiple of 6.

Having shown both of the following:

$$\{6x|x\in\mathbb{Z}\}\subseteq\{2x|x\in\mathbb{Z}\}\bigcap\{3x|x\in\mathbb{Z}\}$$

$$\{2x|x\in\mathbb{Z}\}\bigcap\{3x|x\in\mathbb{Z}\}\subseteq\{6x|x\in\mathbb{Z}\}$$

The original equality must be true.

2 Even/Odd

Prove that, for all integers a, b, and c, if a+b+c is odd then either a, b, and c are odd, or exactly one of a, b, or c is odd.

E = Even, O = Odd. Proof by exhaustion:

a	+	b	+	$^{\mathrm{c}}$	=	
\mathbf{E}		\mathbf{E}		\mathbf{E}	=	E
\mathbf{E}		\mathbf{E}		Ο	=	O
\mathbf{E}		Ο		\mathbf{E}	=	O
\mathbf{E}		Ο		Ο	=	\mathbf{E}
O		\mathbf{E}		\mathbf{E}	=	O
O		\mathbf{E}		Ο	=	\mathbf{E}
O		Ο		\mathbf{E}	=	\mathbf{E}
O		Ο		Ο	=	O

3 Induction

Prove that for all $n \geq 1$,

$$\frac{1}{1\cdot 3} + \frac{1}{3\cdot 5} + \ldots + \frac{1}{(2n-1)(2n+1)} = \frac{n}{2n+1}$$

Inductive hypothesis:

$$\sum_{i=1}^{k} \frac{1}{(2i-1)(2i+1)} = \frac{k}{2k+1}$$

Initial case n = 1:

$$\frac{1}{(2(1)-1)(2(1)+1)} = \frac{1}{2(1)+1}$$

$$\frac{1}{(1)(3)} = \frac{1}{3}$$

Inductive step n = k + 1:

$$\sum_{i=1}^{k+1} \frac{1}{(2i-1)(2i+1)} = \frac{k+1}{2(k+1)+1}$$

$$\sum_{i=1}^{k} \frac{1}{(2i-1)(2i+1)} + \frac{1}{(2(k+1)-1)(2(k+1)+1)} = \frac{k+1}{2k+3}$$

Utilizing the I.H.:

$$\frac{k}{2k+1} + \frac{1}{(2k+1)(2k+3)} = \frac{k+1}{2k+3}$$

$$\frac{k(2k+3)+1}{(2k+3)(2k+1)} = \frac{(k+1)(2k+1)}{(2k+3)(2k+1)}$$

$$\frac{2k^2 + 3k + 1}{2k^2 + 3k + 1} = \frac{(k+1)(2k+1)}{(2k+3)(2k+1)}$$

True by induction.

4 Triangle

Suppose you have 4 colors to color the 3 edges of an equilateral triangle. How many ways can you color the triangle if two ways are considered the same if they differ by a rotation? Include a clear description of how you came to your number.

When choosing colors for the triangle edges, we have to avoid picking our first, second, and third colors such that a rotation makes the choices redundant. We must also consider that triangles completely composed of one color do not differ due to rotation.

Picking our first color is trivial, it can be any of the n colors in our set. The next, however, must avoid a collision with the first. Therefore (barring our n one-color triangles) we avoid picking the color we picked first. Finally, the third color must be picked so that it does not collide with either the first or second pick. The set of pickable colors gets smaller each time we color an edge. The formula for the number of uniquely colored edges follows our logic:

$$n + \sum_{a=1}^{n} \sum_{b=a}^{n} \sum_{c=a+1}^{n} 1$$

Calculating for n = 4 yields the number 24.

The python version may make a little more sense: