


AP Calculus AB Take-Home Final

Duncan Freeman

May 13, 2018

1



$$f(x) = \frac{1}{4} + \sin \pi x, \quad g(x) = 4^{-x}$$

Finding the area of R :

First, we find the first intersection of $f(x)$ and $g(x)$. We'll call the x value of the intersection k . We can then use a calculator approximate the integration of the difference between $g(x)$ and $f(x)$ from 0 until k to obtain the area R .

$$R = \int_0^k [g(x) - f(x)] dx \approx 0.064$$

Finding the area of S

Next, we can find the second intersection of $f(x)$ and $g(x)$ (which we will call j). We can then integrate similarly to get the area; Note that we reverse the order of $g(x)$ and $f(x)$ because $f(x)$ has a higher value of y .

$$S = \int_k^j [f(x) - g(x)] dx \approx 0.410$$

Revolving S :

To revolve S around the horizontal line $y = -1$, we have to adjust $f(x)$ and $g(x)$ by $+1$, and then shroud them in the circle area equation πr^2 , finally integrating the resulting difference between k and j to obtain volume.

$$S_{\text{vol}} = \pi \int_k^j [(f(x) + 1)^2 - (g(x) + 1)^2] dx \approx 4.56$$

Sorry there's no graphic for the revolve, it's *hard*.

2

$$f(0) = 2, \quad f'(0) = -4, \quad f''(0) = 3$$

2.1 Part A

$$g(x) = e^{ax} + f(x)$$

First, we need to know the derivatives of $g(x)$, so we evaluate them as such:

$$\frac{d}{dx}g(x) = g'(x) = ae^{ax} + f'(x)$$

$$\frac{d}{dx}g'(x) = g''(x) = e^{ax}a^2 + f''(x)$$

Next, we can evaluate the derivatives of $g(x)$ in line with the pre-defined derivatives of $f(x)$ like so:

$$g'(0) = ae^{a(0)} + f'(0) = ae^0 - 4 = a - 4$$

$$g''(0) = e^{a(0)}a^2 + f''(0) = a^2e^0 + 3 = a^2 + 3$$

2.2 Part B

$$h(x) = \cos(kx)f(x)$$

First, we derive $h(x)$:

$$\frac{d}{dx}h(x) = h'(x) = -k \sin(kx)f(x) + \cos(kx)f'(x)$$

This gave us the slope ($h'(0)$), but we still need to find the y value of $h(0)$ and the slope at $h'(0)$

$$y = h(0) = \cos[k(0)]f(0) = \cos(0)(2) = (1)(2) = 2$$

$$m = h'(0) = -k \sin(k0)f(x) + \cos(k0)f'(0) = 0 + \cos(1)(-4) = -4$$

We can finally find the tangent equation:

$$y - 2 = -4(x - 0)$$

3

$$R(t) = 2 + 5 \sin \frac{4\pi t}{25}, \quad S(t) = \frac{15t}{1+3t}$$

$$A(t) = \text{Total rate} = S(t) - R(t)$$

3.1 Part A

We can integrate the loss function to obtain the amount removed over 6 hours like so:

$$\int_0^6 R(t) \approx 6.723 \text{ yd}^3$$

3.2 Part B

We can simply integrate the total rate over time and add it to the initial condition:

$$Y(t) = \int_0^t A(x) dx + 2500$$


3.3 Part C

We evaluate $A(t)$ at the time point, because that's the total rate:

$$A(4.0) \approx 1.908 \text{ yd}^3$$

3.4 Part D

It's intuitive to look at the graph of $A(t)$:




At point k , $A(t)$ intersects the x axis. Up until then, the amount of total sand has been decreasing (because $A(t)$ is below the x axis), and at k it has been decreasing the longest. We can now integrate $A(t)$ from 0 to k to obtain the total amount of sand at that point:

$$Y(k) \approx 2492.37 \text{ yd}^3$$

4

4.1 Part A

$$\frac{dy}{dx} = (y - 1)^2 \cos(\pi x)$$



4.2 Part B

Despite being able to simply look at the graph, we can infer that the slope from the differential equation will be zero at $y = 1$ because the entire equation is multiplied by $(y - 1)^{-2}$.

4.3 Part C

$$f(1) = 0; \quad y = 0, \quad x = 1$$

$$\frac{dy}{dx} = (y - 1)^2 \cos(\pi x)$$

$$\frac{dy}{(y - 1)^2} = \cos(\pi x) dx$$

$$(y - 1)^{-2} dy = \cos(\pi x) dx$$

$$\int (y - 1)^{-2} dy = \int \cos(\pi x) dx$$

$$\frac{1}{1 - 0} = \frac{\sin(\pi 1)}{\pi} + C_3$$

$$1 = 0 + C_3$$

$$1 = C_3$$

$$\frac{1}{1 - y} = \frac{\sin(\pi x)}{\pi} + 1$$

5

5.1 Part A

$$\int_{-1}^{10} (\sqrt{x-1}) \, dx \approx$$

5.2 Part B

$$\pi \int_{-1}^{10} (x-1) \, dx \approx$$

5.3 Part C

$$y = \sqrt{x-1}$$

$$y^2 = x-1$$

$$y^2 + 1 = x$$

$$\pi \int_0^3 (y^2 + 1)^2 \, dy \approx$$

6

6.1 Part A

1, 3. This is because $f'(x)$ is zero (has a horizontal tangent) at these locations.

6.2 Part B

Min: $x = 4$, the graph had been negative until then, meaning that $f(x)$ decreased until this point.

Max: $x = -1$, the graph stays negative for more of the graph between $-1 \leq x \leq 5$ than it stays positive. This means that there is no higher value of $f(x)$ than the beginning.

6.3 Part C

$$g(x) = xf(x)$$

$$g'(x) = (1)f(x) + xf'(x)$$

$$m = g'(2) = (1)f(2) + xf'(2) = 6 + (2)(-1) = 6 - 2 = 4$$

$$y = f(2) = 6, \quad x = 2$$

$$y - 6 = 4(x - 2)$$

7

Not finished!

8

$$h(x) = f(g(x)) - 6$$

$$h'(x) = f'(g(x))g'(x)$$

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$	$h(x)$	$h'(x)$
1	6	4	2	5	3	2
2	9	2	3	1	4	-8
3	10	-4	4	2	-7	21
4	-1	3	6	7		

8.1 Part A

$h(r)$ is continuous, $h(2) = 4$, and $h(3) = -7$. By *IVT*, $\exists r \in [2, 3]$ such that $h(r) = -5$.

8.2 Part B

$h'(r)$ is continuous, $h(2) = -8$, and $h(3) = 21$. By *IVT*, $\exists r \in [2, 3]$ such that $h'(r) = -5$.

8.3 Part C

$$\int_a^b f(x) = F(b) - F(a)$$

$$w(x) = \int_1^{g(x)} f(t) dt = F(1) - F(g(x))$$

$$w'(x) = f(g(x))g'(x) - f(1)$$

$$w'(3) = f(g(3))g'(3) - f(1) = f(4)(2) - 6 = -2 - 6 = -8$$

8.4 Part D

$$y = g^{-1}(x), \quad x = 2$$

$$y = g^{-1}(2), \quad g(1) = 2, \quad g^{-1}(2) = 1$$

$$m = g^{-1'}(2), \quad g'(3) = 2, \quad g^{-1'}(2) = 3$$


$$y - 1 = 3(x - 2)$$

9

9.1 Part A

$$r - 30 = 2.0(t - 5)$$

$$r - 30 = 2.0(5.4 - 5), \quad r = 30.8$$



The approximation is above the actual value, because the line approximation will overshoot the decreasing (concave down) function.

9.2 Part B

$$r(5) = 30, \quad r'(5) = 2.0$$

$$\frac{dV}{dt} = \frac{d}{dt} \frac{4}{3} \pi r^3 = 4\pi r^2 r'$$

$$4\pi(30.0)^2(2.0) = 7200\pi \text{ ft}^3/\text{min}$$


9.3 Part C

$$\int_0^{12} r'(t) dt \approx \sum_{i=1}^6 r'(t)_i \Delta t_i \approx$$

$$(0.5)(1) + (0.6)(4) + (1.2)(2) + (2.0)(3) + (4.0)(2) = 19.3 \text{ ft}$$

$\int_0^{12} r'(t) dt$ represents the number of feet the balloon has expanded (in radius) past it's original condition, and it is **not** the absolute radius of the balloon.

9.4 Part D



The RHS will under approximate the graph because it's slope is always negative, meaning any nonzero area under the curve from the right will undershoot it travelling left