

Lecture 6

Quine-McCluskey Method

- A systematic simplification procedure to reduce a minterm expansion to a minimum sum of products.
- Use $XY + XY' = X$ to eliminate as many as literals as possible.
 - The resulting terms = prime implicants.
- Use a prime implicant chart to select a minimum set of prime implicants.

Determination of Prime Implicants

✓ Eliminate literals

Two terms can be combined if they differ in exactly one variable.

$$AB'CD' + AB'CD = AB'C$$

$$\begin{array}{ccccccc} \underline{1} & \underline{0} & \underline{1} & \underline{0} & + & \underline{1} & \underline{0} & \underline{1} & \underline{1} & = & \underline{1} & \underline{0} & \underline{1} \\ X & & Y & & & X & & Y' & & & X & & \end{array}$$

$$A'BC'D + A'BCD' \text{ (won't combine)}$$

$$0 \ 1 \ 0 \ 1 \ + \ 0 \ 1 \ 1 \ 0 \text{ (check \# of 1's)}$$

We need to compare and combine whenever possible.

Sorting to Reduce Comparisons

✓ Sort into groups according to the number of 1's.

$$F(a,b,c,d) = \sum m(0,1,2,5,6,7,8,9,10,14)$$

- No need for comparisons
 - (1) Terms in nonadjacent group
 - (2) Terms in the same group

Group	0	0	0000
-------	---	---	------

Group	1	1	0001
		2	0010
		8	1000

Group	2	5	0101
		6	0110
		9	1001
		10	1010

Group	3	7	0111
		14	1110

Comparison of adjacent groups

- Use $X + X = X$ repeatedly between adjacent groups
- Those combined are checked off.
- Combine terms that have the same dashes and differ one in the number of 1's. (for column II and column III)

$$f = \underset{(1, 5)}{a'c'd} + \underset{(5, 7)}{a'bd} + \underset{(6, 7)}{a'bc} + \underset{(0, 1, 8, 9)}{b'c'} + \underset{(0, 2, 8, 10)}{b'd'} + \underset{(2, 6, 10, 14)}{cd'}$$

$$f = a'bd + b'c' + cd'$$

Determination of Prime Implicants

	Column I			Column II			Column III		
group 0	0	0000	✓	0, 1	000–	✓	0, 1, 8, 9	–00–	
group 1	1	0001	✓	0, 2	00–0	✓	0, 2, 8, 10	–0–0	
	2	0010	✓	0, 8	–000	✓	0, 8, 1, 9	–00–	
	8	1000	✓	1, 5	0–01		0, 8, 2, 10	–0–0	
group 2	5	0101	✓	1, 9	–001	✓	2, 6, 10, 14	--10	
	6	0110	✓	2, 6	0–10	✓	2, 10, 6, 14	--10	
	9	1001	✓	2, 10	–010	✓			
	10	1010	✓	8, 9	100–	✓			
group 3	7	0111	✓	8, 10	10–0	✓			
	14	1110	✓	5, 7	01–1				
				6, 7	011–				
				6, 14	–110	✓			
				10, 14	1–10	✓			

Prime Implicants

- The terms that have not been checked off are called prime implicants.

$$\begin{aligned}
 f &= 0-01 + 01-1 + 011- + -00- \\
 &\quad + -0-0 + --10 \\
 &= \underline{a'c'd} + a'bd + \underline{a'bc} + b'c' + \\
 &\quad \underline{b'd'} + cd'
 \end{aligned}$$

- Each term has a minimum number of literals, but minimum SOP for f:

$$\begin{aligned}
 f &= a'bd + b'c' + cd' \\
 &\quad (a'bd, cd' \Rightarrow a'bc) \\
 &\quad (a'bd, b'c' \Rightarrow a'c'd) \\
 &\quad (b'c', cd' \Rightarrow b'd')
 \end{aligned}$$

Definition of Implicant

– Definition

- Given a function of F of n variables, a product term P is an implicant of F iff for every combination of values of the n variables for which $P = 1$, F is also equal to 1.

- Every minterm of F is an implicant of F .
- Any term formed by combining two or more minterms is an implicant.
- If F is written in SOP form, every product term is an implicant.

– Example: $f(a,b,c) = a'b'c' + ab'c' + ab'c + abc = b'c' + ac$

- If $a'b'c' = 1$, then $F = 1$, if $ac = 1$, then $F = 1$. $a'b'c'$ and ac are implicants.
- If $bc = 1$, (but $a = 0$), $F = 0$, so bc is not an implicant of F .

Definition of Prime Implicant

– Definition

- A prime implicant of a function F is a product term implicant which is no longer an implicant if any literal is deleted from it.
- Example: $f(a,b,c) = a'b'c' + ab'c' + ab'c + abc = b'c' + ac$
 - Implicant $a'b'c'$ is not a prime implicant. Why? If a' is deleted, $b'c'$ is still an implicant of F .
 - $b'c'$ and ac are prime implicants.
- Each prime implicant of a function has a minimum number of literals that no more literals can be eliminated from it or by combining it with other terms.

Quine McClusky Procedure

- QM procedure:
 - Find all product term implicants of a function
 - Combine non-prime implicants.
 - Remaining terms are prime implicants.
 - A minimum SOP expression consists of a sum of some (not necessarily all) of the prime implicants of that function.
 - We need to select a minimum set of prime implicants.
 - If an SOP expression contains a term which is not a prime implicant, the SOP cannot be minimum.

Prime Implicant Chart

- Chart layout
 - Top row lists minterms of the function
 - All prime implicants are listed on the left side.
 - Place x into the chart according to the minterms that form the corresponding prime implicant.
- Essential prime implicant
 - If a minterm is covered only by one prime implicant, that prime implicant is called essential prime implicant. (9 & 14).
 - » Essential prime implicant must be included in the minimum sum of the function.

		0	1	2	5	6	7	8	9	10	14
(0, 1, 8, 9)	$b'c'$	X	X					X	(X)		
(0, 2, 8, 10)	$b'd'$	X		X				X		X	
(2, 6, 10, 14)	cd'			X		X				X	(X)
(1, 5)	$a'c'd$		X		X						
(5, 7)	$a'bd$				X		X				
(6, 7)	$a'bc$					X	X				

Selection of Prime Implicants

- ✓ Cross out the row of the selected essential prime implicants
- ✓ The columns which correspond to the minterms covered by the selected prime implicants are also crossed out.
- ✓ Select a prime implicant that covers the remaining columns. This prime implicant is not essential.

		0	1	2	5	6	7	8	9	10	14
(0, 1, 8, 9)	$b'c'$	*	*					*	*		
(0, 2, 8, 10)	$b'd'$	*		*				*		*	
(2, 6, 10, 14)	cd'			*		*				*	*
(1, 5)	$a'c'd$		*		X						
(5, 7)	$a'bd$				X		X				
(6, 7)	$a'bc$					X	X				

A Cyclic Prime Implicant Chart

- Two or more X's in every column.
- $F = \Sigma m(0,1,2,5,6,7)$
 - $F = a'b' + bc' + ac$. (by try and error). No guarantee for this to be minimum.

0	000	All checked	0,1	00-
1	001	off	0,2	0-0
2	010		1,5	-01
5	101		2,6	-10
6	110		5,7	1-1
7	111		6,7	11-

			0	1	2	5	6	7
①	→	(0, 1)	*	*				
		(0, 2)	*		*			
		(1, 5)		*		*		
②	→	(2, 6)			*		*	
③	→	(5, 7)				*		*
		(6, 7)					*	*

Another Solution

- $F = a'c' + b'c + ab$
- Each minterm is covered by two different prime implicants.

			0	1	2	5	6	7
P_1	(0, 1)	$a'b'$	X	X				
P_2	(0, 2)	$a'c'$	X		X			
P_3	(1, 5)	$b'c$		X		X		
P_4	(2, 6)	bc'			X		X	
P_5	(5, 7)	ac				X		X
P_6	(6, 7)	ab					X	X

Petrick's Method

- A more systematic way to find all minimum solutions from a prime implicant chart.
- P is True when all the minterms in the chart have been covered. $P = f(P_1, P_2, \dots)$
- Label each row with P_i .
 - P_i is true when the prime implicant in row P_i is included in the solution.
 - For column 0, we must choose either P_1 or P_2 in order to cover **minterm 0**. Thus $(P_1 + P_2)$ must be true.

			0	1	2	5	6	7
P_1	(0, 1)	$a'b'$	X	X				
P_2	(0, 2)	$a'c'$	X		X			
P_3	(1, 5)	$b'c$		X		X		
P_4	(2, 6)	bc'			X		X	
P_5	(5, 7)	ac				X		X
P_6	(6, 7)	ab					X	X

Petrick's Method (cont.)

- For column 0, we must choose either P_1 or P_2 in order to cover minterm 0. Thus $(P_1 + P_2)$ must be true.
- To cover minterm 1, $P_1 + P_3$ must be true, and etc.
- $P = (P_1 + P_2)(P_1 + P_3)(P_2 + P_4)(P_3 + P_5)(P_4 + P_6)(P_5 + P_6) = 1$
- This means: We must choose row P_1 or P_2 , and row P_1 or P_3 , and row P_2 or P_4 , etc.

			0	1	2	5	6	7
P_1	(0, 1)	$a'b'$	*	*				
P_2	(0, 2)	$a'c'$	*		*			
P_3	(1, 5)	$b'c$		*		*		
P_4	(2, 6)	bc'			*		*	
P_5	(5, 7)	ac				*		*
P_6	(6, 7)	ab					*	*

Petrick's Method (cont.)

- Then we simplify

$$P = (P_1 + P_2)(P_1 + P_3)(P_2 + P_4)(P_3 + P_5)(P_4 + P_6)(P_5 + P_6)$$

$$= (P_1 + P_2P_3)(P_4 + P_2P_6)(P_5 + P_3P_6);$$

$$(X+Y)(X+Z) = X + YZ$$

$$= (P_1P_4 + P_1P_2P_6 + P_2P_3P_4 + P_2P_3P_6)(P_5 + P_3P_6)$$

$$= P_1P_4P_5 + P_1P_2P_5P_6 + P_2P_3P_4P_5 + P_1P_3P_4P_6 + P_2P_3P_6$$

In other words, for $P = 1$ (to cover all minterms), we must choose row P_1 and P_4 and P_5 or row P_1 and P_2 and P_5 and P_6 or etc

There are five to choose. We choose $P_1P_4P_5$ or $P_2P_3P_6$.

			0	1	2	5	6	7
P_1	(0, 1)	$a'b'$	*	*				
P_2	(0, 2)	$a'c'$	*		*			
P_3	(1, 5)	$b'c$		*		*		
P_4	(2, 6)	bc'			*		*	
P_5	(5, 7)	ac				*		*
P_6	(6, 7)	ab					*	*

Simplification of Incompletely Specified Functions

- An incompletely specified function

$$F(A, B, C, D) = \sum m(2, 3, 7, 9, 11, 13) + \sum d(1, 10, 15)$$

(the terms following d are don't cares)

The don't cares are treated like required minterms when finding the prime implicants:

1	0001	✓	(1, 3)	00-1	✓	(1, 3, 9, 11)	-0-1
2	0010	✓	(1, 9)	-001	✓	(2, 3, 10, 11)	-01-
3	0011	✓	(2, 3)	001-	✓	(3, 7, 11, 15)	--11
9	1001	✓	(2, 10)	-010	✓	(9, 11, 13, 15)	1--1
10	1010	✓	(3, 7)	0-11	✓		
7	0111	✓	(3, 11)	-011	✓		
11	1011	✓	(9, 11)	10-1	✓		
13	1101	✓	(9, 13)	1-01	✓		
15	1111	✓	(10, 11)	101-	✓		
			(7, 15)	-111	✓		
			(11, 15)	1-11	✓		
			(13, 15)	11-1	✓		

The don't care columns are omitted when forming the prime implicant chart:

	2	3	7	9	11	13
(1, 3, 9, 11)		*		*	*	
*(2, 3, 10, 11)	*	*			*	
*(3, 7, 11, 15)		*	*		*	
*(9, 11, 13, 15)				*	*	*

$$F = B'C + CD + AD$$

* indicates an essential prime implicant.

Simplification Using Map-Entered Variables

- Extend K-map for more variables.
 - When E appears in a square, if $E = 1$, then the corresponding minterm is present in the function G.
 - $G(A,B,C,D,E,F) = m_0 + m_2 + m_3 + \text{Em}_5 + \text{Em}_7 + \text{Fm}_9 + m_{11} + m_{15} + (\text{don't care terms})$

		AB			
		00	01	11	10
CD	00	1			
	01	X	E	X	F
	11	1	E	1	1
	10	1			X

G

(a)

		AB			
		00	01	11	10
CD	00	1			
	01	X		X	
	11	1		1	1
	10	1			X

$E = F = 0$

$MS_0 = A'B' + ACD$

(b)

		AB			
		00	01	11	10
CD	00	X			
	01	X	1	X	
	11	X	1	X	X
	10	X			X

$E = 1, F = 0$

$MS_1 = A'D$

(c)

		AB			
		00	01	11	10
CD	00	X			
	01	X		X	1
	11	X		X	X
	10	X			X

$E = 0, F = 1$

$MS_2 = AD$

(d)

Map-Entered Variable

- $F(A,B,C,D) = A'B'C + A'BC + A'BC'D + ABCD + (AB'C)$, (don't care)
 - Choose **D** as a map-entered variable.
 - When $D = 0$, $F = A'C$ (Fig. a)
 - When $D = 1$, $F = C + A'B$ (Fig. b)
 - » two 1's are changed to x's since they are covered in Fig. a.
- $F = A'C + D(C + A'B) = A'C + CD + A'BD$

		A	
		0	1
BC	00		
	01	1	x
	11	1	D
	10	D	

(a)

		A	
		0	1
BC	00		
	01	x	x
	11	x	1
	10	1	

(b)

		DA			
		00	01	11	10
BC	00				
	01	1	x	x	1
	11	1		1	1
	10				1

(c)

General View for Map-Entered Variable Method

- Given a map with variables P_1, P_2 etc, entered into some of the squares, the minimum SOP form of F is as follows:

- $F = MS_0 + P_1 MS_1 + P_2 MS_2 + \dots$

where

- MS_0 is minimum sum obtained by setting $P_1 = P_2 \dots = 0$
- MS_1 is minimum sum obtained by setting $P_1 = 1, P_j = 0 (j \neq 1)$, and replacing all 1's on the map with don't cares.

– Previously, $G = A'B' + ACD + EA'D + FAD$.