


Fundamentals of Image Processing

Lecture 02

Visual Perception, Sensors, and Image Acquisition

The Human Visual System



Retina

- A thin layer covering approximately 200° on the back of the eye
- Covered by two basic types of photoreceptors: **cones** and **rods**
- Fovea
 - Region of maximum visual acuity
 - Contains only cones (approximately 147,000 per linear millimeter, about 6 to 7 millions total)
- With the exception of the fovea, the retina is densely covered by rods

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Photoreceptors

- Cones
 - Responsible for the perception of **color**
 - There are three types of cones (different pigments and chemical compositions)
 - **L-Cones** (often improperly called **Red** cones) are more sensitive to long wavelengths
 - **M-Cones** (often improperly called **Green** cones) are more sensitive to medium wavelengths
 - **S-Cones** (often improperly called **Blue** cones) are more sensitive to the short wavelengths
- Rods
 - Highly sensitive, are responsible for vision under low-lighting conditions

Illustration courtesy of Webvision (<http://webvision.med.utah.edu/photo2.html#cones>)
Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Density of Photoreceptors






Image from the book *Principles of Digital Image Synthesis*, Vol. 1, page 10, by Andrew Glassner, Morgan Kaufmann Publishers, Inc.



Cones in retina



Rods in retina

Images courtesy of Christine Curcio

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Photoreceptors' Response

In fact, violet instead of blue, and yellow instead of red!!




Image by Dowling, 1987. Reproduced from Webvision (<http://webvision.med.utah.edu/photo2.html#cones>)

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS




Trichromatic Vision

- It comes from the existence of cones sensitive to three different bands of the visible spectrum
- As a result, color sensations can be produced as linear combinations of the primaries **red**, **green** and **blue** (additive color model)


Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS




Human Eye Optics




Human Eye Optics




Perfect Eye





Myopia (nearsightedness)



Myopia Correction





The Human Eye

- Capable of adapting itself to a large range of intensity values (of the order of 10^{10})
 - Brightness adaptation
 - Cannot operate in the entire range simultaneously
- Brightness perception does not depend only on intensity
 - Mach bands
 - Simultaneous contrast

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Mach Bands

- The human eye is very sensitive to discontinuities in light intensity
- It results from lateral inhibition of eye photoreceptors
- The more light a photoreceptor receives, the more it inhibits the response of the photoreceptors next to it

actual intensity

perceived intensity

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Simultaneous Contrast

- The perceived brightness in one region does not depend only on the light intensity in that region
- All three small squares below have exactly the same intensity

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Image Acquisition

- In principle, one can generate images using any form of energy that can be detected by a sensor

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Sensors for Image Capture

- The incident energy is transformed into a digital signal using an analog-to-digital converter
- The intensity of the resulting signal is proportional to the incident energy
- Different materials are used to build sensors for specific wavelengths


Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Image Capture Device

- Characteristics
 - Number of sensors
 - Number of sensor cells
 - Sensor geometry
 - Dimensions and shape
 - Spacing between sensor cells
 - Distribution of the sensor cells on the surface


Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Sampling and Quantization



Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Sampling



Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS


Digital Image

- Same scene captured at different resolutions (sampling rates)




Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Quantization




Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Quantization



Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Histogram



original image
The histogram does not say anything about the spatial distribution of the gray shades (colors) in the image
Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

The impact of quantization is not very evident


Digital Images Details versus number of shades



Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Histogram

- Image with relatively few high-frequency details




Original image

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Digital Images Details versus number of shades

- Image with relatively few high-frequency details



The impact of quantization is more evident!

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS

Image Spatial Details versus Number of Shades

- For images with lots of spatial details (high frequencies), a small number of gray shades is, in general, sufficient for the representation
- Images with smooth transitions among the gray shades, in general require a larger number of shades

Copyright Manuel Menezes de Oliveira Neto, Informática UFRGS