

ALGEBRA

RETROALIMENTAC HÓNO Nro 3









PROBLEMA 1 Halle la suma de los factores primos, luego de factorizar el polinomio:

$$(x^2-4x)^2-2(x^2-4x)-15$$

$$(x^{2} - 4x)^{2} - 2(x^{2} - 4x) - 15$$

$$x^{2} - 4x$$

$$x^{2} - 4x$$

$$-5 - 5(x^{2} - 4x)$$

$$-2(x^{2} - 4x)$$

$$(x^2 - 4x + 3)(x^2 - 4x - 5)$$
 $x - 1$
 $x - 3$
 $x - 5$

$$x - 1 +$$

$$x - 3$$

$$x + 1$$

$$(x-1)(x-3)(x+1)(x-5)$$

$$(x-1)(x-3) (x+1)(x-5)$$

$$\therefore \sum factores primos: 4x-8$$



PROBLEMA 2 Factorice: $m^4 - 2m^3 + 3m^2 - 6m$. Luego, Indique el número de factores primos lineales

Resolución

$$m^4 - 2m^3 + 3m^2 - 6m$$
 $m^3(m-2) + 3m(m-2)$
 $(m-2)(m^3 + 3m)$
 $(m-2)m(m^2 + 3)$
F.P lineal lineal cuadrático

∴ Hay 2 factores primos lineales



PROBLEMA 3 Calcule la mayor suma de coeficientes de une de los factores primos de:

Resolución

os de:
$$T = x^2 + 16y^2 - 36z^2 + 8xy$$

$$TCP = x^2 + 8xy + 16y^2 - 36z^2$$

$$T = (x+4y)^2 - 36z^2$$

$$\sqrt{(x+4y)^2} \sqrt{36z^2}$$

2 121		
x + 4y	6z	

$$T = (x + 4y + 6z)(x + 4y - 6z)$$


FACTOR	SUMA DE COEFICIENTES	RESULTADO
1er	1+4+6	11
2do	1+4-6	-1

∴ La mayor suma de coeficientes es 11



PROBLEMA 4 Pedro tiene 20 años y César tiene "6m" años. Halle la diferencia de edades, siendo "m" el número de factores primos de: $5x^4 - 18x^2 - 8$?

Resolución



$$(5x^2 + 2)(x^2 - 4)$$

$$\sqrt{x^2} \quad \sqrt{4}$$

$$x \quad 2$$

$$(5x^2+2)(x+2)(x-2)$$

Hay 3 factores primos

m = 3

	EDADES
Pedro	20
C é sar	18

: La diferencia es 2



PROBLEMA 5 Simplifique:
$$\frac{4\sqrt{8} + 5\sqrt{32}}{2\sqrt{50} - \sqrt{18}}$$

Resolución

$$\frac{4\sqrt{4}\sqrt{2}+5\sqrt{16}\sqrt{2}}{2\sqrt{25}\sqrt{2}-\sqrt{9}\sqrt{2}} = \frac{8+20}{10-3} = \frac{28}{7} = 4$$

∴ Rpta: 4



PROBLEMA 6 Carlitos tiene 6 canicas y Pedrito tiene "E" canicas, cuántas canicas tienen

juntos: si
$$E = \sqrt{9 + \sqrt{80}} + \sqrt{7 - \sqrt{48}} - \sqrt{8 - \sqrt{60}}$$

Resolución

$$E = \sqrt{9 + 2\sqrt{20}} + \sqrt{7 - 2\sqrt{12}} - \sqrt{8 - 2\sqrt{15}}$$

$$5 + 4 \quad 5x4 \quad 4 + 3 \quad 4x3 \quad 5 + 3 \quad 5x3$$

$$E = \sqrt{5} + \sqrt{4} + \sqrt{4} - \sqrt{3} - (\sqrt{5} - \sqrt{3})$$

$$E = 2 + 2 = 4$$

∴ Juntos tendrán 10 canicas

PROBLEMA 7



Reduzca:
$$K = \sqrt{5 + \sqrt{21}} - \sqrt{5 - \sqrt{21}}$$

$$\sqrt{x} - \sqrt{y} = \sqrt{x + y - 2\sqrt{xy}}$$
 ; $x > y$

$$K = \sqrt{\frac{5 + \sqrt{21}}{10} + (5 - \sqrt{21})} - 2\sqrt{\frac{5 + \sqrt{21}}{5 - \sqrt{21}}}$$

$$25 - 21$$

$$K = \sqrt{10 - 2\sqrt{4}} = \sqrt{10 - 4} = \sqrt{6}$$



Sean los complejos

Halle la parte imaginaria de

$$z_1 = -21 + 3i$$

$$z_1 = -21 + 3i$$
 $z_2 = -15 - 2i$

$$4z_2 + 3z_1$$

$$z_1 = -21 + 3i$$
 $z_2 = -15 - 2i$

$$z_2 = -15 - 2i$$

$$3z_1 = -63 + 9$$

$$3z_1 = -63 + 9i$$
 $4z_2 = -60 - 8i$

$$\therefore \operatorname{Im}(4z_2 + 3z_1) = 1$$

PROBLEMA 9

Hallar el valor de 'n' para que 'z' sea un complejo real puro. $z = \frac{n+3i}{n+3-5i}$

$$z = \frac{n+3i}{n+3-5i} \qquad ; n \in \mathbb{R}$$

Resolución

Recordar:

$$\mathbf{Si:} \ z = \frac{a+bi}{c+di}$$

es un complejo real puro

$$\rightarrow ad = bc$$

$$(-5)(n) = (3)(n + 3)$$

 $-5n = 3n + 9$
 $-8n = 9$

$$\therefore n = -\frac{9}{8}$$

PROBLEMA 10

En la igualdad (5+7i)x + (1-2i)y = 11+12iAdemás x, y son reales.

Determine xy

$$(5+7i)x + (1-2i)y = 11 + 12i$$

$$5x + 7xi + y - 2yi = 11 + 12i$$

$$(5x + y) + (7x - 2y)i = 11 + 12i$$

$$5x + y = 11 \times 2$$

$$\begin{cases} 10x + 2y = 22 \\ 7x - 2y = 12 \end{cases} + 17x = 34$$

$$x = 2$$

$$y = 1$$

$$\therefore xy = 2$$