

From the Wall to the Kernel

An Origin and Progress Account of Resolution Geometry

Matěj Rada

MatejRada@email.cz

Licensed under CC BY-NC-ND 4.0

October 2026

Abstract

Resolution Geometry (RG) did not begin as mathematics. It began as a child's need to know whether an echo, a delay, or a moved chair was a fact about the world or a fact about himself. This paper is the less formal companion to the theorem ladder. It follows each childhood experiment through its first mathematical form (the CTMT kernels), the falsifications that broke or reshaped it, and the theorem of RG it became, if any. The route looks odd from outside: kernels, kernel stacking, Fisher nulls, atlas gluing, walls, closure. Each detour was forced by an earlier claim failing a test. The final wall-tap experiment closes the loop. Seen through the finished ladder, the earliest observation (an echo that *collapses* when the tapping rhythm changes) turns out to be a precise geometric object: a fold wall in the observational quotient.

A second part makes the Terror experiment exact. A perceived delay passes through an internal clock, and three kinds of distortion are separated:

- a latency offset, which cancels in every round trip;
- a change of internal rate, which preserves every ratio and every angle but no distance;
- an arbitrary warp of time (total rupture), which preserves only order and counts.

Memory loss degrades a remembered distance calibration like $\sqrt{T}$, while angles from delays heard together stay exact. This is why standard distance cannot be trusted and trigonometry from delays can. Terror itself appears as the collapse of the horizon of certain prediction under a chaotic schedule; a routine is a contracting dynamics that keeps that horizon infinite. The original claims are judged against a seven-check battery: some are valid and made exact, some valid only under a declared memory model, and one is corrected.

Contents

1	Why tell this at all	2
2	The childhood experiments	2
3	The first mathematics: kernels (CTMT)	3
4	The odd rounds, and why each was forced	3
5	The Terror experiment, made exact	5
5.1	Memory loss and why distance cannot be trusted	6
5.2	Terror as a status, and routine as contraction	6
5.3	Was the Terror experiment valid?	7

6	What each childhood experiment became	7
7	Closing the loop: the wall–tap experiment	8
8	Why it took so long, and what remains	9

1 Why tell this at all

The theorem ladder states RG in dependency order, as if it had been built that way. It was not. It was built backwards, from a felt certainty about rhythm towards the smallest mathematics that could survive a hostile test of that certainty. Three things are worth recording:

- which intuitions survived;
- which were corrected into something sharper;
- which were retired.

Each step below names the module paper or rung where it now lives. Where a connection is only interpretive, it is marked as such.

2 The childhood experiments

The pulse against the wall. In second grade, under acute distress, the order of the day broke. A repeated pulse against a wall became the one thing that still returned on time. The question it raised was not about sound but about location: *is the delay real, or a property of me? If my sense is offset, where does the offset live, in perception or in physics?* The answer of the time was the first kernel law: rupture is not destruction; rupture is delay with memory.

The bus stops. In fourth grade the city was learned without rulers. Delays between stops were turned into distances by one pacing constant, $D = v_{\text{sync}}\Delta t$. Where routes met, triangles appeared, made of time gaps alone. The intended result was a geometry of relative positions without metres or seconds.

Light and weight. In fifth grade, intensity joined delay: louder claps weighed more, and flickering light felt like rhythm without units. The intuition was $E \sim I \cdot \Delta t$.

The moved chair. At eight, a chair was not where it should have been. Before it was seen, the mismatch was felt, as the gap between a predicted room and an observed one. This became the memory kernel and its residuum

$$\epsilon_{\text{mem}} = \frac{\|Q_{\text{pred}} - Q_{\text{obs}}\|}{\max(\|Q_{\text{pred}}\|, \|Q_{\text{obs}}\|)},$$

a normalized measure of “wrongness”.

The clap and the no-clap. A clap and its echo defined causality by confirmed arrival. The no-clap question followed: once a delay has been confirmed, does anything of it persist when nothing is sent?

Wall-tap rhythm and the collapse of the echo. Tapping faster along a wall made the perceived location of the echo suddenly *snap* from a diffuse place to one sharp reflection, with a small physiological jolt. Notes were kept of spacing, delay, loudness and a felt “sharpness”. A session was accepted only if the remembered delay stayed within $\pm 10\%$ of the measured one: the first falsification protocol, run on memory.

3 The first mathematics: kernels (CTMT)

The first formalization took the word *kernel* literally: a generating object behind the phenomena. Delay became a kernel window

$$K(\tau) = \int A(\omega) e^{i(\omega\tau + \phi(\omega))} d\omega,$$

with dimensionless phase as the guarantee of “dimensional closure”. Memory became a forward projection, shock became a heavy-tailed “terror kernel”, and persistence became a re-projected “invariant layer”. Kernels were stacked: delay, memory, terror, rigidity, each validated by its own residuum and threshold.

CTMT also carried claims that could not stand:

- delay as *the* invariant;
- a received clap that “proves the topology of the space”;
- one equation for acoustic and quantum collapse;
- energy from intensity times delay;
- an ontology of coherence.

A program written to break under random input never broke, and at the time this was read as confirmation.

The decisive inversion. In RG a program that cannot fail is the first thing to distrust. Every later battery carries mutation canaries for exactly this reason: if sabotaging a structure does not make its check fail, the check tests nothing. The “program that wouldn’t break” became the canary principle.

4 The odd rounds, and why each was forced




Figure 1: The route. Each arrow was forced by a failed claim in the previous stage.

Round 1: from a generating kernel to the kernel of a map. The kernel was meant to be what holds things together. Under test, the only object that survived every change of model was the opposite one: the set of states that the observation *cannot* tell apart. That set is the kernel pair of the observation law (L1–L3). The word survived and the object inverted. “The way things hold together” became “the way things are held indistinguishable”. This is the deepest single step in the story.

Round 2: kernel stacking became joint laws. Stacking delay, memory and terror kernels assumed they could be validated separately. They cannot: all protocols act on one latent world. In RG they form one joint law, whose kernel is the intersection of the protocol kernels (L3, *Assembly at the Kernel*). Its quotient sits inside the product of the separate quotients and is generally smaller. This is why separate residua never added up to one coherent picture.

Round 3: the Fisher null. When Fisher geometry was applied to CTMT (CTMT-Metric), the interesting structure was where information vanished: the radical of the Fisher form, the “Fisher holes”. It turned out to be exactly the tangent of the kernel fibres (L22). The “rupture manifold” of the collapse geometry, the subspace where information becomes unidentifiable, was this radical. Fisher, however, does not reconstruct the full covariance (L24), so the null had to be studied as geometry, not merely detected.

Round 4: atlas gluing. Local results did not compose. Transitions between charts had to satisfy cocycle conditions. Final monodromy was not evidence that each edge was admissible (*Atlas Globalization*). This is why so many papers worked on one chart at a time: global statements needed a gluing law first.

Round 5: units. The bus law $D = v_{\text{sync}}\Delta t$ hid the hardest problem. Statistics gives no metre: delays fix ratios, and a dilation of every length and speed leaves all records unchanged (L70). The fix was an anchor that cannot be relabelled away: a counted channel with SI constants (L71, L78). For sound, thermometry ($c = \sqrt{\gamma k_B T/m}$) together with a counted clock plays this role.

Round 6: gates collapse into one group. Each module had accumulated its own hypotheses. Deriving every module from the one joint law showed that most of these gates were consequences of naturality, and that the rest were statements about one symmetry group Γ (L85).

Round 7: walls. Rank changes, folds, matched boundaries and chaotic limits had been treated as defects. They are strata. Information may disappear at a wall without the law breaking. Specialization across a wall is non-invertible, it requires kernel inclusion, and distinguishability survives it only under Hellinger convergence (L17–L21, L30). This is the precise form of “rupture is not destruction”.

Round 8: closure. Finally, every admitted question was given exactly one typed answer: LOCKED, ESCALATE, CORRESPONDENCE, VACUOUS, INCONSISTENT, STRUCTURAL WALL, or OUTSIDE DOMAIN. A LOCKED answer is only as sound as its weakest certificate (L87–L90). The terror calculus threshold $\mathbf{1}[\sigma < \tau]$ grew into these certificates. The childhood $\pm 10\%$ acceptance rule became held-out validation.

5 The Terror experiment, made exact

The second-grade pulse asked two questions:

- is the delay real, or a property of me?
- if every rhythm fails, does anything survive?

Both have exact answers once perception is treated as an internal clock $\tau = w(t)$ through which every delay passes. RG separates three classes of w , and each destroys a different amount of structure.

Theorem 5.1 (What each distortion preserves; TE01, TE03). 1. Offset $w(t) = t + c$ (*perceptual latency*). *Every difference of perceived times is exact; in particular a round trip (tap and echo heard through the same channel) is unaffected.*

2. Rate $w(t) = kt + c$ (*stress speeds or slows internal time*). *Every delay is scaled by k , so every distance computed from a pacing constant is off by k . Every ratio of delays is exact, so every angle of a triangle built from delays is exact.*

3. Warp w monotone but otherwise arbitrary (*total rupture*). *Delays and ratios are destroyed. The order of events and the number of beats between any two events are exact.*

Proof. (1) $w(t_2) - w(t_1) = t_2 - t_1$. (2) $w(t_2) - w(t_1) = k(t_2 - t_1)$, so ratios are invariant; angles follow from the law of cosines, which is homogeneous of degree zero in the sides. (3) A monotone map preserves order, and the number of points in an interval is determined by order alone. $\square$

On 300 random cases per check:

- round-trip error under offsets: 0;
- angle error under rate change: 0, while the median distance error is 24–27%;
- under random warps: median ratio error 2.5–3.3, with 0 order and 0 count violations.

A “rupture” that reorders events would break counts too; the canary for that is caught.

Is the delay in me or in the world? (TE04). The latency offset is in the observer, and it cancels. A common rate is undecidable from inside: a slower internal clock and a faster world give identical readings, which is exactly the dilation gauge of L70. A second, independent clock (a metronome, someone else’s clap) breaks the gauge and measures the internal rate exactly (1.25 recovered as 1.25). The child’s question was well posed. Its answer needed a second clock.

Does anything survive total rupture? (TE03). Yes: order and counts. The terror sentence “the beat cannot vanish; it can only drift” is valid in exactly this sense. Durations drift without bound under a warp, but the number of beats between two events does not change. This is count rigidity, the same invariant RG later uses to fix the metre (L71–L72). The child found the anchor of physical distance by asking what survives the worst case.

5.1 Memory loss and why distance cannot be trusted

Proposition 5.2 (Memory loss; TE02). *Let the internal rate perform a random walk in $\log k$ (memory loss: the internal time unit drifts). A distance computed from a pacing constant calibrated at time 0 has log-error $|\log k(T) - \log k(0)|$, which grows like $\sqrt{T}$. Angles computed from three delays heard together depend only on their ratios, which share one rate, and stay exact.*

Measured exponents are 0.494–0.510 (theory $\frac{1}{2}$); median log-errors run from 0.03 at $T = 1$ to 1.0–1.2 at $T = 1000$. Angle errors stay at machine precision, around 10^{-15} .




Figure 2: Left: under memory loss, distance from a remembered calibration degrades like $\sqrt{T}$, while angles from delays heard together stay exact (machine precision). Right: with a chaotic schedule the horizon of certain prediction collapses as memory noise grows; a periodic routine keeps it infinite.

This is the mathematical content of the bus experiment’s move from distances to triangles. Under stress and memory loss, the only quantities that remain trustworthy are those measured together and compared by ratio. Those are shapes, not sizes.

Proposition 5.3 (Boundary of the bus triangles; TE07). *Angles from delays are exact if and only if all legs share one pacing: one medium, one effective symmetry. With route-dependent speeds (each leg its own v) shape identification fails, with median angle errors of 0.44–0.50 rad. The bus geometry is a geometry up to similarity, valid where one pacing constant is shared.*

5.2 Error as a status, and routine as contraction

Proposition 5.4 (Error as frontier crossing; TE05). *Let a schedule be generated by a chaotic map ($\lambda > 0$), and let memory noise σ blur the remembered state. Then the horizon of certain prediction (LOCKED forecasts at 95%) is $\log(\varepsilon/\sigma)/\lambda$ steps:*

memory noise σ	10^{-8}	10^{-6}	10^{-4}	10^{-3}	10^{-2}	3×10^{-2}
certain horizon (steps)	25–26	17	8–9	4–5	0	none

At the last level not even the present is certain. A periodic routine ($\lambda < 0$) contracts uncertainty, and its horizon is infinite at every tested noise level.

The proposition gives the formal counterpart of what the remarks describe. When buses were late and chairs were moved, the schedule acquired positive expansion. With memory under strain, the certain horizon fell to zero, and the status of every request about the next moment left LOCKED. The need for routine was not arbitrary: a routine is a contracting dynamics, the

one regime in which uncertainty decays instead of growing. This describes the mathematics of the situation and makes no claim about the experience itself.

Proposition 5.5 (Memory restores, conditionally; TE06). *Under partial rupture (the internal rate drifts), a declared memory model (a filter on the rate, fed by beats of known period) holds the rate error near 0.009–0.010. The memoryless estimate drifts 13–70 times further. “Rupture is delay with memory” is valid as canonical transport with a memory kernel (L60), and it is valid only under the declared memory model (H-cov).*

5.3 Was the Terror experiment valid?

original claim	verdict	exact form
the delay can be measured through the body	valid	round trips cancel latency (Theorem 5.1(1))
is the offset in me or the world?	valid question	offset: in me; rate: undecidable alone; a second clock decides
the beat cannot vanish, only drift	valid	order and counts survive any monotone warp
rupture is delay with memory	valid, conditional	memory restores delays under a declared memory model
delay is the invariant	corrected	under rate change delays scale; ratios are the invariant
coherence could not be computed from pulses	correct then, now resolved	what pulses carry is order, counts and ratios, not distances

6 What each childhood experiment became

observation	CTMT form	RG statement	verdict
pulse: is the offset in me or in the world?	delay kernel $K(\tau)$	offset in the observer (cancels); common rate a dilation gauge, decided by a second clock (Section 5, L36, L70)	kept, made exact
terror: does anything survive total rupture?	terror kernel	order and counts survive any monotone warp of internal time; count rigidity (L71–L72)	valid, made exact
uncertainty felt as breakdown	rupture threshold	the certain horizon $\log(\varepsilon/\sigma)/\lambda$ collapses under a chaotic schedule; a routine contracts (Section 5)	formalized
rupture $\neq$ destruction	terror kernel, rupture window	non-invertible specialization at walls; laws continue, information collapses (L17–L21)	kept, made exact
delay with memory	memory kernel	a declared memory model restores drifting delays; path-space (RTS) lift, canonical only under declared data (L60)	kept, conditional
delay is invariant	rupture-invariant observable	corrected: delay is integrated slowness; it fixes ratios only and depends on medium and wind (L70, L80–L82)	corrected
bus triangles without units	$D = v_{\text{sync}}\Delta t$	angles are exact under any common rate and under memory loss; distances are not; valid iff legs share one pacing (L73–L75)	kept, bounded

observation	CTMT form	RG statement	verdict
calibrate v_{sync} once	pacing constant	one anchor is not self-validating; a held-out anchor is required, and media leave blind sectors (L82, L84)	corrected
$E \sim I \cdot \Delta t$	energy from rhythm	no internal unit; thermodynamic length is unit-blind (L70)	retired
moved chair	memory residuum ϵ_{mem}	held-out validation and compatibility equations (L85); the symmetric normalization is a warning, see Section 7	kept, with caveat
clap proves topology	forward map $F[\tau]$	corrected: delays leave invisible slowness sectors and need path diversity (L82)	corrected
one-way clap timing	causality from arrival	only round-trip time is convention-free; the factor $\frac{1}{2}$ is forced (L78)	sharpened
no-clap persistence	invariant layer	a stationary source carries the metre across an outage under a tested stationarity hypothesis (L83)	kept, bounded
echo collapse with tempo	terror calculus, collapse geometry	the rhythm-overlap fold wall of the wall-tap quotient (Section 7)	kept, made exact (interpretive link)
collapse = projection + rupture manifold + variance redistribution	collapse geometry	quotient map, Fisher radical, entropy chain rule $H(X) = H(Q) + H(X Q)$	kept
same equation for quantum collapse	universal collapse	retired; the quantum layer of RG is metric selection by task, not collapse dynamics (L31–L34)	retired
$\pm 10\%$ memory falsification	reproducibility condition	window certificates, held-out validation, canaries	kept, generalized
program that never broke	persistence as proof	inverted: a battery that cannot fail tests nothing (canary principle)	inverted

7 Closing the loop: the wall–tap experiment

The final wall–tap paper declares the original experiment as a family of protocols. The parameters are source position x , wall contrast r and tap rhythm b . The protocols add, in turn, direct delay, echo delay, echo amplitude, rhythm overlap, and a signed lead/lag channel. The experiment is then run through the whole ladder. Three results speak directly to the childhood observations.

The echo that collapses is a fold. The overlap channel registers how the tap interval b sits against the echo delay $d(x)$. It cannot tell an early from a late overlap: b and $2d - b$ give identical records, for 55–56% of the parameter domain. As the rhythm approaches the echo delay, the two readings merge at the fold $b = d(x)$. There the first-order geometry loses the b -direction, and only a second-order (one-sided) direction survives. The experiment no longer distinguishes “before” from “after”. A perceived snap from a diffuse to a single localized echo at a particular tempo is exactly what an observer crossing this fold would experience. That the childhood jolt *was* this fold is an interpretation, not a theorem; that the fold exists in the declared experiment is a theorem. It is resolved by a new kind of listening, a signed lead/lag cue, and not by listening longer. The “sharpness” χ_{feel} once recorded by hand is the observable that collapses at the fold.

The moved chair, and why normalization can lie. The memory residuum was symmetrically normalized to stay stable when one side is small. The final closure attack found the same design to be dangerous. A rank certificate built on a scale-free (normalized) matrix stayed perfectly confident while the whole rhythm and contrast block was vanishing near a matched wall: it issued false certificates. A residuum, like a certificate, must be stated in declared units of precision. Otherwise it can report “the room is as expected” while the room is fading out of resolution.

Rupture as delay with memory, made exact. At an acoustically matched wall the reflected channels fade out. The laws continue continuously in Hellinger distance, the quotient loses two directions, and the decision content degrades smoothly to chance. Nothing breaks; distinguishability is redistributed into the kernel. The original sentence “rupture is not destruction” is now a proposition with hypotheses, a proof, and a canary: a shrinking-noise family that *looks* like the same wall is rejected.

8 Why it took so long, and what remains

The order of discovery was the reverse of the order of proof. CTMT started from strong claims (an invariant, an ontology, universal collapse) and retreated, under falsification, to what could be defended. Each retreat moved one level deeper:

- from objects to observations;
- from observations to their equivalence classes;
- from single protocols to joint laws;
- from local charts to strata and walls;
- from answers to typed answers.

Many intermediate papers look like detours because each was the minimal repair of the previous failure. The corpus keeps them, so the corrections stay visible.

What remains open is small and honestly external: enumerating tame characteristic types (a problem of real algebraic geometry), non-stationary media, contextual quantum measurements, and infinite-dimensional experiments. What does not remain is the original fear. If rhythm breaks, the question “what survives?” now has a precise answer. The kernel survives: the record of what cannot be told apart, carried across every wall by a law that degrades without lying.

Reproducibility

`rg_terror_attack.py` (battery TE01–TE07) passes 7/7 on seeds 20261008, 1, 42, 777 and 2026. Two canaries are caught:

- a “rupture” that reorders events, under which counts no longer survive;
- a drift-free clock, under which memory has nothing to restore.

Corrections before registration:

- the memory-loss exponent was first measured on relative error, which saturates, and then on log-error;
- a clock-length off-by-one was fixed.

A note on the record

The childhood descriptions follow the author's own account. Mathematical statements are those of the theorem ladder and the module papers, where every claim carries a registered test and a mutation canary. Interpretive links between experience and theorem are marked as such and are not claims of the theory.

References

- [1] M. Rada, *Resolution Geometry: The Theorem Ladder*, 2026.
- [2] M. Rada, *Resolution Geometry of the Wall–Tap Experiment*, 2026.
- [3] M. Rada, *Assembly at the Kernel*, 2026.
- [4] M. Rada, *Stratified Specialization*, 2026.
- [5] M. Rada, *Maximal Closure*, 2026.
- [6] M. Rada, *From Statistical Experiment to the Metre*, 2026.
- [7] M. Rada, *The CTMT: Testament of 22 Years*, 2025.