



# THE RAPID CHANGE OF DEVELOPING BRAINS



# DIFFUSION MRI IN DEVELOPING BRAINS


## ■ Research motivations

- Assessment of brain maturation and white matter development
- Macroscale connectivity
- Comparative analyses (pathologies, species)
- Longitudinal studies from *in utero*




## ■ Clinical motivations<sup>1</sup>

- Cytotoxic (intra-cellular) edema
- Vasogenic (extra-cellular) edema



Adapted from  
Song et al.,  
Frontiers in  
Neuroscience  
2017

Figure from Afif et al., Brain Structure and Function 2007




<sup>1</sup> Garel. MRI of the Fetal Brain. Springer (2004)

# DIFFUSION MRI IN DEVELOPING BRAINS



## ■ MAIN CHALLENGES




- Low signal-to-noise ratio/low (through-plane) spatial resolution
- Motion
- Diffusion MRI distortion artefacts
- Protocol restrictions (small and unique b-values, limited number of gradient directions)
- Rapidly developing anatomy
- Gradient directions mismatch




Low through-plane resolution



Intra-volume motion



Inter-volumes motion and signal drop



# Robust estimation of the microstructure of the early developing brain using deep learning

Hamza Kebiri<sup>1,2,3</sup>, Ali Gholipour<sup>3</sup>, Rizhong Lin<sup>2,4</sup>, Lana Vasung<sup>5</sup>, and Davood Karimi<sup>3,\*</sup>, Meritxell Bach Cuadra<sup>1,2,\*</sup>

<sup>1</sup> CIBM Center for Biomedical Imaging, Switzerland

<sup>2</sup> Department of Radiology, Lausanne University Hospital (CHUV) and University of Lausanne (UNIL), Lausanne, Switzerland

<sup>3</sup> Computational Radiology Laboratory, Department of Radiology, Boston Children's Hospital and Harvard Medical School, Boston, Massachusetts, USA

<sup>4</sup> Signal Processing Laboratory 5 (LTS5), École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland


<sup>5</sup> Department of Pediatrics, Boston Children's Hospital, and Harvard Medical School, Boston, Massachusetts, USA

**MICCAI 2023; ISMRM 2023; Medical Image Analysis (under revision) 2023**

[https://link.springer.com/chapter/10.1007/978-3-031-43990-2\\_28](https://link.springer.com/chapter/10.1007/978-3-031-43990-2_28)

# CONTEXT

- Microstructure best estimated with fiber orientation distribution function (FODs)




Adapted from Shi  
and Toga, Molecular  
Psychiatry, 2017



- Problem
  - High number of measurements and high/multiple b-values requirement for accurate estimation with widely used state-of-the-art methods<sup>1,2</sup>


**BUT:** Unrealistic for developing brains!



- Proposed solution
  - Use deep neural networks and available research dedicated datasets to learn a mapping between few measurements and target diffusion metrics (FOD, tensor, scalar map, etc.)

<sup>1</sup> Tournier et al., Neuroimage, 2007; <sup>2</sup> Jeurissen et al., Neuroimage, 2014

# PROPOSED MODEL



# NETWORKS & DATA



- Input: 6 or 12 b0-normalized diffusion measurements (in spherical harmonics basis)
- Output: 45 SH coefficients of FOD estimated with multi-shell multi-tissue constrained spherical deconvolution (MSMT-CSD)<sup>1</sup> using all 300 multi-shell dHCP<sup>2,3</sup> measurements
- Two networks:
  - (1) **Newborn network** trained on dHCP newborn subjects and evaluated **quantitatively** on dHCP newborn subjects and **qualitatively** on clinical newborns subjects from Boston Children's Hospital (BCH)
  - (2) **Pre-term network** trained on dHCP pre-term subjects and evaluated **qualitatively** on clinical fetal subjects from BCH



<sup>1</sup> Jeurissen et al., Neuroimage, 2014; <sup>2</sup> Hutter et al., MRM, 2018; <sup>3</sup> Tournier et al., NMR in Biomedicine, 2020

# NETWORKS & DATA



- *Newborns network* (trained and tested on dHCP newborns<sup>2,3</sup> and clinical newborns)
  - Training/validation: 145 subjects of post-menstrual age (PMA) [26-45] weeks
  - Testing: 320 dHCP subjects; 15 BCH subjects (PMA [38-48] weeks)
  - 6 input uniform<sup>1</sup> measurements
  - $b = 1000 \text{ s/mm}^2$
  
- *Pre-term network* (trained on pre-term newborns<sup>2,3</sup> and tested on fetuses)
  - Training/validation: 77 pre-terms of PMA [26, 38] weeks
  - Testing: 11 fetal subject ([24, 38] gestational weeks)
  - 12 input measurements
  - Pre-terms at  $b=400 \text{ s/mm}^2$  and fetuses at  $b=500 \text{ s/mm}^2$

<sup>1</sup> Skare et al., JMR, 2000

# MODEL EVALUATION



- Baseline models
  - Constrained Spherical Deconvolution (CSD)<sup>1</sup> with 148 measurements
  - Constant Solid Angle<sup>2</sup> with 300 measurements
  - Sparse Fascicle Model (SFM)<sup>3</sup> with 300 measurements
  - Multi-layer perceptron (MLP)<sup>4</sup>
  - CTtrack (CNN + Transformers)<sup>5</sup>
- Ground truth agreement ( $\Delta GS$ )
  - Spliting the ground truth into two disjoint gold standard (GS) subsets of 150 measurements each

<sup>1</sup> Tournier et al., Neuroimage, 2007; <sup>2</sup> Aganj et al., MRM, 2010; <sup>3</sup> Rokem et al., PloS one, 2015 ; <sup>4</sup> Karimi et al., Neuroimage, 2021; <sup>5</sup> Hosseini et al., Neuro. Informatics, 2022 43

# MODEL EVALUATION



- dHCP newborns evaluation
  - Agreement rate (AR) in number of peaks
  - Angular error
  - Apparent Fiber Density<sup>1</sup>

	1 peaks	2 peaks	3 peaks
1 peaks	a <sub>11</sub>	a <sub>12</sub>	a <sub>13</sub>
2 peaks	a <sub>21</sub>	a <sub>22</sub>	a <sub>23</sub>
3 peaks	a <sub>31</sub>	a <sub>32</sub>	a <sub>33</sub>

$$AR_{1\text{-peaks}} = a_{11} / (a_{11} + a_{12} + a_{13} + a_{21} + a_{31})$$

$$AR_{2\text{-peaks}} = a_{22} / (a_{22} + a_{21} + a_{23} + a_{12} + a_{32})$$


$$AR_{3\text{-peaks}} = a_{33} / (a_{33} + a_{32} + a_{31} + a_{23} + a_{13})$$

- Clinical newborns evaluation
  - Visual assessment by an expert neuroanatomist
- Fetal evaluation
  - Visual assessment by an expert neuroanatomist
  - Histological comparison with post-mortem brains




<sup>1</sup> Rafflet et al., Neuroimage, 2012

## QUALITATIVE RESULTS: DHCP NEWBORNS




# QUANTITATIVE RESULTS: DHCP NEWBORNS



Method	$b$ -values ( $s/mm^2$ )	$M$	Angular error		
			Single fibers	Two fibers	Three fibers
$DL_n$	{0, 1000}	7	10°( $\pm 0.2$ )	20°( $\pm 0.3$ )	30°( $\pm 0.1$ )
CSD	{0, 2600}	148	7°( $\pm 0.2$ )	16°( $\pm 0.3$ )	27°( $\pm 0.1$ )
CSA	{0, 400, 1000, 2600}	300	43°( $\pm 0.3$ )	37°( $\pm 0.1$ )	35°( $\pm 0.1$ )
SFM	{0, 400, 1000, 2600}	300	42°( $\pm 0.6$ )	37°( $\pm 2.0$ )	35°( $\pm 4.0$ )
$\Delta GS$	{0, 400, 1000, 2600}	150	6°( $\pm 0.1$ )	14°( $\pm 0.1$ )	25°( $\pm 0.1$ )

- Low agreement rate within the ground truth for multiple fibers
- Our method with 7 diffusion measurements ~ other methods with x21 and x43 more measurements
- Lowest error for our method in approximating the apparent fiber density (AFD)


# QUANTITATIVE RESULTS: DHCP NEWBORNS



Method	b-values ( $\text{s/mm}^2$ )	$M$	Agreement rate (Angular error)			AFD error
			Single fibers	Two fibers	Three fibers	
DL <sub>n</sub>	{0, 1000}	7	77.5% (10°)	22.2% (20°)	8.0% (30°)	0.178 ( $\pm 0.083$ )
MLP	{0, 1000}	7	74% (16°)	15.5% (28°)	7.7% (32°)	0.398 ( $\pm 0.104$ )
CTtrack	{0, 1000}	7	74.5% (16°)	16.8% (25°)	4.4% (32°)	0.263 ( $\pm 0.105$ )


- Low agreement rate within the ground truth for multiple fibers
- Our method with 7 diffusion measurements ~ other methods with x21 and x43 more measurements
- Lowest error for our method in approximating the apparent fiber density (AFD)
- DL also outperforming MLP & CTtrack

# CORRELATION TO QUALITY CONTROL METRICS



- No correlation to other QC metrics (Outlier-ratio, SNR) and scan age
- Correlation to motion as estimated by SHARD pipeline<sup>1</sup>

# UNCERTAINTY ESTIMATION



$$\sigma = \frac{1}{N_{WBS}} \sum_{i=1}^{N_{WBS}} \|FOD_i - \mu\|_2 \quad \text{where } \mu_c = \frac{1}{N_{WBS}} \sum_{i=1}^{N_{WBS}} FOD_{i,c} \quad \text{and } c \in \{1, 45\}$$


## ■ Wild bootstrap uncertainty

$$\sigma_{norm} = \frac{\sigma}{m_j} \quad \text{where } m_j = \frac{1}{45} \sum_{c=1}^{45} \|FOD_{j,c}\| \quad \text{and } j \in \{1, N_{WBS}\}$$


# QUALITATIVE RESULTS: CLINICAL NEWBORNS BCH



**Slice 1**




$\text{DL}_n$




CSD

**Slice 2**



$\text{DL}_n$



CSD

**$\text{DL}_n$  using 7 measurements**

**CSD using 35 measurements**

# FETAL EVALUATION

## Axial Slices

Frontal Crossroad Area C2


Genu of Corpus Callosum

Posterior Limb of Internal Capsule

Cortical Plate of Superior Temporal Gyrus

**DL<sub>f</sub>**

**CSD**




Regions of interests (ROIs) definition based on histology and immunohistochemistry as in previous works<sup>1</sup>

<sup>1</sup> Karimi et al., Neuroimage, 2021

# FETAL EVALUATION

Coronal Slices



Regions of interests (ROIs) definition based on histology and immunohistochemistry as in previous works<sup>1</sup>

<sup>1</sup> Karimi et al., Neuroimage, 2021

# FETAL EVALUATION: RESULTS



Fetal brain region	$DL_f$	Tied	CSD
Frontal crossroad region	10	0	1
Genu of corpus callosum	11	0	0
Cortex of Insula	2	5	4
Posterior limb of internal capsule	7	2	2
Cortex of superior temporal gyrus	6	2	3
Subplate of the precentral gyrus	4	0	7
Internal capsule	3	0	8
Cerebral peduncles	1	1	9
Intermediate zone, genuculocortical	4	3	4
Intermediate zones, callosal	10	0	1
Prefrontal subplate	6	4	1
Prefrontal cortical plate	8	2	1
<b>Count per ROI</b>	<b>7</b>	<b>1</b>	<b>3</b>
<b>Count per subject</b>	<b>9</b>	<b>0</b>	<b>2</b>

# GENERALIZABILITY & DOMAIN SHIFTS



EPFL



Features	Baby Connectome Project (BCP)
Number of Subjects	210
Age Span	1-5 years
Imaging Device	3T Siemens Magnetom Prisma
Resolution	$1.5 \times 1.5 \times 1.5 \text{ mm}^3$
B-values	{0, 500, 1000, 1500, 2000, 2500, 3000} s/mm <sup>2</sup>



- Assessing the performance of deep learning on a different dataset
- Generalization from one dataset to another

# DOMAIN SHIFT IMPACTS



## CROSS-AGE AND CROSS-SITE DOMAIN SHIFT IMPACTS ON DEEP LEARNING-BASED WHITE MATTER FIBER ESTIMATION IN NEWBORN AND BABY BRAINS

*Rizhong Lin* <sup>1,2,3</sup>

*Ali Gholipour* <sup>4</sup>  
*Hamza Kebiri* <sup>2,5,\*</sup>

*Jean-Philippe Thiran* <sup>1,2,5</sup>  
*Meritxell Bach Cuadra* <sup>5,2,\*</sup>

*Davood Karimi* <sup>4</sup>

<sup>1</sup> Signal Processing Laboratory 5 (LTS5), École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

<sup>2</sup> Department of Radiology, Lausanne University Hospital (CHUV) and University of Lausanne (UNIL), Lausanne, Switzerland

<sup>3</sup> College of Electronic and Information Engineering, Tongji University, Shanghai, China

<sup>4</sup> Computational Radiology Laboratory, Department of Radiology,  
Boston Children's Hospital and Harvard Medical School, Boston, MA, USA

<sup>5</sup> CIBM Center for Biomedical Imaging, Switzerland


Lin et al., ISBI 2024

<https://arxiv.org/abs/2312.14773>

# DISCUSSION



- Conclusion
  - Deep learning can achieve state-of-the-art results in ODF estimation with fewer measurements
  - Can have a big impact on scan time reduction
  - Low agreement for multiple-fiber voxels within the dataset
  - Deep learning can provide some level of generalization across centers/protocols and anatomies (i.e. fetuses)
  
- Limitations
  - Specific response functions learning
  - Generalization to pathological subjects
  - Tractography ?



# Through-Plane Super-Resolution With Autoencoders in Diffusion Magnetic Resonance Imaging of the Developing Human Brain


*Hamza Kebiri<sup>1,2\*</sup>, Erick J. Canales-Rodríguez<sup>3</sup>, Hélène Lajous<sup>1,2</sup>, Priscille de Dumast<sup>1,2</sup>,  
Gabriel Girard<sup>1,2,3</sup>, Yasser Alemán-Gómez<sup>1</sup>, Mériam Koob<sup>1</sup>, András Jakab<sup>4,5</sup> and  
Meritxell Bach Cuadra<sup>1,2,3</sup>*

<sup>1</sup> Department of Radiology, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland, <sup>2</sup> CIBM Center for Biomedical Imaging, Lausanne, Switzerland, <sup>3</sup> Signal Processing Laboratory 5 (LTS5), Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland, <sup>4</sup> Center for MR Research University Children's Hospital Zurich, Zurich, Switzerland, <sup>5</sup> Neuroscience Center Zurich, University of Zurich, Zurich, Switzerland

Kebiri et al., ISMRM 2022; *Frontiers in Neurology*, 2022


<https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2022.827816/>

# THROUGH-PLANE RESOLUTION




Adapted from Chai et al.,  
IEEE Access, 2020

- Interpolation
  - Typically performed in low-through-plane resolution series in dMRI of developing brains
  - Interpolation was shown to be beneficial in dMRI of adult brains<sup>1</sup>




1x1x4 mm<sup>3</sup>




1x1x1 mm<sup>3</sup>

# MODEL ARCHITECTURE



# MODEL ARCHITECTURE & INFERENCE



# DATA

## ■ PRE-TERM

- 31 subjects
- Gestational age: 29-37 weeks
- 25,920 slices of  $1.17 \times 1.17 \text{ mm}^2$
- Slice thickness:  $1.5 \text{ mm}$
- $b = 1000 \text{ s/mm}^2$




## ■ FETAL

- 6 subjects (3 with motion)
- Gestational age: 23-35 weeks
- In-plane:  $1 \times 1 \text{ mm}^2$
- Slice thickness:  $4-5 \text{ mm}$
- $b = 700 \text{ s/mm}^2$




# RAW SIGNAL MEAN SQUARED ERROR




- Autoencoder outperforming interpolations
- Higher gap for 

# DWI LATENT SPACE EXPLORATION


- Distance between all B1000 volumes & between their encodings



# PRE-TERM QUALITATIVE RESULTS: COLOR FA



# ANATOMICAL SIMILARITY PRE-TERM & FETAL SUBJECTS




Pre-term subject from dHCP data (Hutter et al., MRM, 2018; Tournier et al., NMR in Biomedicine, 2020)


# QUALITATIVE FETAL RESULTS: DTI MAPS

35 GW - still subject (color FA, FA)


Original slice i




Synthesized slice  
with autoencoder



Synthesized slice  
with autoencoder




Original slice i + 5 mm




23 GW - motion subject (MD, FA)


Original slice i




Synthesized slice  
with autoencoder



Synthesized slice  
with autoencoder



Original slice i + 4 mm



# DISCUSSION




## ■ Conclusions

- Autoencoders can be used for dMRI super-resolution and outperform conventional interpolations
- They can also be used in slice outlier recovery
- Training on b0 images can generalize to gradient diffusion volumes
- Generalizing from pre-term newborns to fetal might be possible

## ■ Limitations

- Shallow fetal evaluation
- Autencoders generate smooth outputs (similarly to interpolation methods)
- Potential network hallucinations
- Motion




# Conclusion

# DISCUSSION



- Conclusion
  - Overview on medical imaging tools with a focus on MRI
  - Introduction to machine learning and its applications to medical imaging
  - Two examples of supervised and unsupervised learning approaches on medical imaging tasks
- Future directions of ML in medical imaging (MRI)
  - Domain shift issues (training and testing on different machines/protocols/ages/ etc.)
  - AI hallucinations
  - Reproducibility/variability
  - Clinical translations
  - More (publicly available) data needed (synthetic data?)
  - Multi-modal large language models (LLMs)



# THANK YOU!



Željka Krsnik



Athena Taymourtash  
Georg Langs



Kelly Payette  
Andras Jakab




Davood Karimi  
Lana Vasung  
Camilo Calixto  
Clemente Velasco-Annis  
Hakim Ouaalam  
Sean Clancy  
Simon Warfield  
Ali Gholipour



Hélène Lajous  
Yasser Alemán Gómez  
Rizhong Lin  
Priscille de Dumast  
Francesco La Rosa  
Jaime Barranco  
German Barquero  
Thomas Sanchez  
Pedro Macias Gordaliza  
Nataliia Molchanova  
Tommaso Di Noto  
Cristina Martínez  
João Jorge  
Meritxell Bach Cuadra



Erick Canales Rodríguez  
Gabriel Girard  
Marco Pizzolato  
Thomas Yu  
Farnaz Delavari  
Jean-Philippe Thiran



Eleonora Fornari  
Sébastien Tourbier  
Jean-Baptiste Ledoux  
Andrea Moreno  
Jonas Richiardi  
Vincent Dunet  
Mériam Koob  
Patric Hagmann  
Reto Meuli



Thank you for your attention



C I B M . C H

 @HamzaKebiri 