

## Graded Problems

**Problem 1** (a) Suppose  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  is a smooth function. Prove that  $f_{xyzx} = f_{zxxy}$ , where these are fourth-order partial derivatives. You are free to use Theorem 4.3 for this problem, but not Theorem 4.5.

(b) Verify the result of (a) for the function  $f(x, y, z) = x^3y^2z$ .

**Problem 2** Suppose we have differentiable functions  $\mathbf{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  and  $\mathbf{g} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that

$$\begin{aligned}\mathbf{g}(1, 2) &= (3, 0), \\ D\mathbf{f}(x, y) &= \begin{pmatrix} xe^y & x^2y \\ 0 & yx^2 + 1 \end{pmatrix}, \\ D(\mathbf{f} \circ \mathbf{g})(1, 2) &= \begin{pmatrix} 1 & 4 \\ 2 & -1 \end{pmatrix}.\end{aligned}$$

Find  $D\mathbf{g}(1, 2)$ .

## Professional Problem

**Problem 3** (a) Consider differentiable functions  $g : \mathbb{R} \rightarrow \mathbb{R}$  and  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f(x, y) = 0$  when  $y = g(x)$ . Prove that if  $\partial f / \partial y \neq 0$ , then

$$\frac{dg}{dx} = -\frac{\partial f / \partial x}{\partial f / \partial y}.$$

(b) Consider the equation  $\sin(x) + \cos(y) = 0$ . Make a suitable choice of  $f$ , and use the result of (a) to compute  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ .

(c) Use implicit differentiation to verify your answer to (b).

## Completion Packet

**Problem 4** Consider the functions  $\mathbf{f}(x, y) = (x^2 + y, xy - \sin(xy))$  and  $\mathbf{g}(x, y) = (e^{xy}, x^2y)$ . Compute  $D(\mathbf{f} + \mathbf{g})$  in two ways:

Homework 9: Properties of Derivatives

- (a) By computing the derivative of  $\mathbf{f} + \mathbf{g}$
  - (b) By computing the derivatives of  $\mathbf{f}$  and  $\mathbf{g}$ , and using the sum rule.
- 

**Problem 5** Consider the functions  $f(x, y, z) = xy^2z^3$  and  $g(x, y, z) = xyz$ .

- (a) Verify that

$$D(fg)(\mathbf{a}) = g(\mathbf{a})Df(\mathbf{a}) + f(\mathbf{a})Dg(\mathbf{a}).$$

- (b) Verify that

$$D(f/g)(\mathbf{a}) = \frac{g(\mathbf{a})Df(\mathbf{a}) - f(\mathbf{a})Dg(\mathbf{a})}{g(\mathbf{a})^2}.$$

---

**Problem 6** Find all second-order partial derivatives of the function  $f(x, y) = \ln(xy)$ .

---

**Problem 7** Find all second-order partial derivatives of the function  $g(u, v) = e^{u^2+v^2}$ .

---

**Problem 8** Find all second-order partial derivatives of the function  $h(x, y, z) = xy^2z^3$ .

---

**Problem 9** Consider the functions  $y(s, t) = e^s + e^{st} + e^t$  and  $\mathbf{x}(t) = (t, t^2)$ . Compute  $D(\mathbf{x} \circ y)$  in two ways:

- (a) Determining a formula for the composition  $\mathbf{x} \circ y$ , then computing the total derivative.
  - (b) Computing total derivatives of  $\mathbf{x}$  and  $y$ , and using the chain rule.
- 

**Problem 10** Consider functions  $\mathbf{f} : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  and  $\mathbf{g} : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  such that  $\mathbf{f}(x, y, z) = (x^2y + e^z, e^x + y)$  and  $D\mathbf{g}(x, y) = \begin{pmatrix} y & x \\ 0 & 1 \\ 0 & xy \end{pmatrix}$ . For each of the given total derivatives, either explain why they do not exist, compute them, or explain what additional information we would need to compute them.

Homework 9: Properties of Derivatives

(a)  $D(\mathbf{f} \circ \mathbf{g})$

(b)  $D(\mathbf{g} \circ \mathbf{f})$

---

**Problem 11** Consider functions  $\mathbf{f} : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  and  $\mathbf{g} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $\mathbf{f}(x, y, z) = (\sin(xyz), \cos(xyz))$  and  $D\mathbf{g}(x, y) = \begin{pmatrix} e^{xy} & 0 \\ 0 & e^{xy} \end{pmatrix}$ . For each of the given total derivatives, either explain why they do not exist, compute them, or explain what additional information we would need to compute them.

(a)  $D(\mathbf{f} \circ \mathbf{g})$

(b)  $D(\mathbf{g} \circ \mathbf{f})$

---

**Problem 12** Consider the function  $\mathbf{g} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by  $\mathbf{g}(\rho, \theta, \phi) = (\rho \cos \theta \sin \phi, \rho \sin \theta \sin \phi, \rho \cos \phi)$ , which changes spherical coordinates to Cartesian coordinates. For any differentiable function  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ , compute  $\partial f / \partial \rho$ ,  $\partial f / \partial \theta$ , and  $\partial f / \partial \phi$  in terms of  $\partial f / \partial x$ ,  $\partial f / \partial y$ , and  $\partial f / \partial z$ .

---