

January

3 mon

2005

Vibrations

The to and fro motion of a body about its mean position is called oscillations or vibrations.



There are three parts of oscillations.

- 4 tue
- i) Free oscillation
 - ii) Damped "
 - iii) Forced. "

i) Free oscillation/vibration is defined as the body which vibrates with definite interval and with natural frequency.

Ex:- SHO. - Simple Harmonic Oscillation.

5 wed



Simple pendulum.

A particle is said to execute SHM when it vibrates periodically in such a way that at any instant, the restoring force acting on it is proportional to its displacement from mean position and oppositely directed.

December 2004

Sun	Mon	Tue	Wed	Thu	Fri	Sat
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

Notes

January

6 Thu

$$\therefore F_{\text{Restoring}} \propto -x \rightarrow \textcircled{1}$$

Where F is the restoring force acting on the oscillator when its displacement from equilibrium position is ' x '.

$$F_R = -kx. \rightarrow \textcircled{2}$$

7 Fri

Now according to Newton's 2nd law.

$$F = ma. \rightarrow \textcircled{3}$$

From eqn $\textcircled{2}$ and $\textcircled{3}$.

$$ma = -kx.$$

$$ma + kx = 0. \rightarrow \textcircled{4}$$

We know that, $a = \frac{d^2x}{dt^2}$

$$\text{Eqn } \textcircled{4} \Rightarrow m \frac{d^2x}{dt^2} + kx = 0.$$

Divide by 'm' to above eqn.

$$\frac{d^2x}{dt^2} + \left(\frac{k}{m}\right)x = 0.$$

February 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28					

Notes

January

10 min

2008

Let us substitute $\frac{k}{m} = \omega^2$ in above eqns.

$$\boxed{\frac{d^2x}{dt^2} + \omega^2 x = 0}$$

→ ⑤

$$\omega = \sqrt{\frac{k}{m}} \text{. Angular freq.}$$

Eqn ⑤ represents the differential of simple Harmonic oscillator/vibration.

11 min

Solution of differential equation can be given as

$$x = e^{\alpha t} \quad \text{where } x \text{ is the displacement or amplitude}$$

→ ⑥

Differentiating above 'x'

$$\frac{dx}{dt} = \alpha \cdot e^{\alpha t} \rightarrow ⑦ \quad \frac{d^2x}{dt^2} = \alpha^2 \cdot e^{\alpha t} \rightarrow ⑧$$

12 min

Substituting ⑦ and ⑧ in eqn ⑤.

$$\alpha^2 \cdot e^{\alpha t} + \omega^2 e^{\alpha t} = 0.$$

$$e^{\alpha t} (\alpha^2 + \omega^2) = 0.$$

$$e^{\alpha t} \neq 0; \therefore \alpha^2 + \omega^2 = 0.$$



Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	2	3	4	5	6	7	8	9	10	11	12
13	14	15	16	17	18	19	20	21	22	23	24
25	26	27	28	29	30	31					
31	1	2	3	4	5	6	7	8	9	10	11

January

2005

13 Thu

$$\alpha^2 = -\omega^2.$$

$$\therefore \alpha = \pm i\omega.$$

General solution of eqn ⑤ is given as

$$x = A e^{i\omega t} + B e^{-i\omega t} \rightarrow ⑥$$

where A and B are constants

14 Fri

$$\text{Now } e^{i\omega t} = \cos \omega t + i \sin \omega t.$$

$$e^{-i\omega t} = \cos \omega t - i \sin \omega t.$$

$$⑥ \Rightarrow x = A(\cos \omega t + i \sin \omega t) + B(\cos \omega t - i \sin \omega t).$$

$$\Rightarrow x = A \underline{\cos \omega t} + i \underline{A \sin \omega t} + B \underline{\cos \omega t} - B \underline{i \sin \omega t}.$$

15 Sat

$$x = \cos \omega t (A+B) + i \sin \omega t (A-B) \rightarrow ⑦$$

Replacing $(A+B)$ and $i(A-B)$ by ^{some} constants

$$A+B = R \sin \phi \quad \text{and} \quad i(A-B) = R \cos \phi.$$

$$⑦ \Rightarrow x = \cos \omega t \cdot R \sin \phi + \sin \omega t \cdot R \cos \phi.$$

$$x = R (\cos \omega t \cdot \sin \phi + \sin \omega t \cdot \cos \phi).$$

February 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
1	2	3	4	5	6	7
8	9	10	11	12		
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28					

January

17 mon

2004

$$x = R \sin(\omega t + \phi) \quad \rightarrow \textcircled{11}$$

Eqn $\textcircled{11}$ represents the final solⁿ of SHM.

where $R \rightarrow$ Amplitude of the oscillatory system.

$(\phi + \omega t) \rightarrow$ Phase of vibration.

$x = R \sin(\omega t + \phi) \rightarrow$ represents the displacements of SHM. (Amplitude)

18 tue

We now calculate the time period and frequency of SHM.

Time period is defined as the time required for one complete oscillation.

$$t = 2\pi/\omega \Rightarrow 2\pi/\sqrt{k/m} \quad \therefore \omega^2 = \frac{k}{m}$$

19 wed

$$\omega = \sqrt{k/m}$$

Frequency of SHM:

$$f = \frac{1}{T} = \frac{\omega}{2\pi} = \frac{\sqrt{k/m}}{2\pi}$$

Significance of ' ω ' in eqn $\textcircled{11}$.

December 2004

Sun	Mon	Tue	Wed	Thu	Fri	Sat
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

Notes

January

20 Thu

If 't' is increased by $2\pi/\omega$

$$\text{i.e. } t = t + \frac{2\pi}{\omega}.$$

$$\therefore (1) \Rightarrow x = R \sin(\omega(t + \frac{2\pi}{\omega}) + \phi) \\ = R \sin(\omega t + 2\pi + \phi).$$

21 Fri

$$x = R \sin(2\pi + (\omega t + \phi))$$

$$x = R \sin(\omega t + \phi)$$

$$\therefore \sin(2\pi + \theta) = \sin \theta.$$

which means displacement is same after time
 $t = 2\pi/\omega$.

22 Sat

So, the period $t = \frac{2\pi}{\omega}$ is the time period.

ii) Damped oscillation / vibration — For a free oscillation the energy remains constant and hence oscillator continues indefinitely.

February 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28					

Notes

January

24 mon

But in reality, the amplitude of the oscillatory system gradually decreases due to the damping forces like friction (air) and resistance of the media.

25 tue "The oscillator whose amplitude, in successive oscillations goes on decreasing due to the presence of resistive forces are called as damped oscillators and oscillations are called damped oscillations (DHO)."

The damping force always act in a opposite directions to the motion of oscillatory body and velocity dependent.

26 wed $F_{\text{damping}} \propto -V \Rightarrow F_{\text{damping}} = -\gamma V$.

$\therefore F_{\text{damping}} = -\gamma V$. \rightarrow Damping constant.
 \rightarrow ① $V \rightarrow$ Damping force velocity

Now $F_{\text{restoring}} = -kx$. \rightarrow ②

According to Newton's 2nd law of motion.

$$F = ma. \rightarrow ③$$

Dec 2004	Sun	Mon	Tue	Wed	Thu	Fri	Sat	Notes
	5	6	7	8	9	10	11	
	12	13	14	15	16	17	18	
	19	20	21	22	23	24	25	
	26	27	28	29	30	31		

January

27th

Total Force, $F = \text{Restoring} + \text{Frapping}$.

$$\therefore ma = -kx - \tau \frac{dx}{dt}.$$

$$m \frac{d^2x}{dt^2} + \tau \frac{dx}{dt} + kx = 0.$$

Divide by m in the above eqn.

$$\frac{d^2x}{dt^2} + \frac{\tau}{m} \frac{dx}{dt} + \frac{k}{m} x = 0.$$

Substitute $\tau/m = \alpha k$ and $k/m = \omega^2$ in above eqn.

$$\therefore \boxed{\frac{d^2x}{dt^2} + 2\alpha k \frac{dx}{dt} + \omega^2 x = 0.} \rightarrow ④$$

where " αk " is the damping coefficient and " ω " is the natural freq. of oscillating body.

Eqn ④ represents the differential eqn of Damped Harmonic Oscillation.

This is second order linear differential Homogeneous equation in x .

The general solution of the above eqn can be written as.

S.No	Mon	Tue	Wed	Thu	Fri	Sat
1	2	3	4	5	6	7
8	9	10	11	12		
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28					

January/February
31 mon

$$x = A \cdot e^{\alpha t} \quad x \text{ is the amplitude}$$

$$\frac{dx}{dt} = A \cdot \alpha \cdot e^{\alpha t} \Rightarrow \frac{d^2x}{dt^2} = A \cdot \alpha^2 e^{\alpha t}.$$

$$④ \Rightarrow A \cdot \alpha^2 \cdot e^{\alpha t} + 2k \cdot A \alpha e^{\alpha t} + w^2 A e^{\alpha t} = 0.$$

$$1 \text{ tue} \Rightarrow A e^{\alpha t} [\alpha^2 + 2k\alpha + w^2] = 0.$$

$$\text{Since } A e^{\alpha t} \neq 0 \text{ but } \alpha^2 + 2k\alpha + w^2 = 0$$

The two roots of the quad. eqn. are.

$$\alpha = \frac{-2k \pm \sqrt{4k^2 - 4 \cdot 1 \cdot w^2}}{2 \cdot 1} = \frac{-2k \pm 2\sqrt{k^2 - w^2}}{2}$$

$$\alpha = -k \pm \sqrt{k^2 - w^2} \rightarrow ⑤$$

2 wed $\therefore \alpha$ in eqn ⑤ has two values.

$$\alpha_1 = -k + \sqrt{k^2 - w^2} \quad \text{and} \quad \alpha_2 = -k - \sqrt{k^2 - w^2}$$

Therefore, general soln of differential eqn is.

$$x = A \cdot e^{\alpha_1 t} + B \cdot e^{\alpha_2 t}.$$

$$x = A \cdot e^{(-k + \sqrt{k^2 - w^2})t} + B \cdot e^{(-k - \sqrt{k^2 - w^2})t} \rightarrow ⑥$$

January 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
30	31					1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29

Notes

February

Eqn ⑥ represents the amplitude of D.H.O.

The actual form of eqn ⑥ depends on whether

- (a) $k^2 > \omega^2$ +/cany damping
- (b) $k^2 < \omega^2$ Low "
- (c) $k^2 = \omega^2$ Critical "

Case-I $k^2 > \omega^2$ (+/cany damping)

If the damping is high, then $\sqrt{k^2 - \omega^2} = \beta$ is a real quantity and positive.


Then eqn ⑥ \Rightarrow

$$x = e^{-kt} [A \cdot e^{(\sqrt{k^2 - \omega^2})t} + B \cdot e^{(-\sqrt{k^2 - \omega^2})t}] \rightarrow \text{F}$$

$$x = e^{-kt} [A \cdot e^{\beta t} + B \cdot e^{-\beta t}]$$

$$x = A \cdot e^{-(k-\beta)t} + B \cdot e^{-(k+\beta)t}$$

Since both the exponents are negative then the displacement 'x' decreases continuously with the time. That is if the particle when once displaced return to its equilibrium position quite slowly.



This is DEAD BEAT

March 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

Notes Non oscillatory or aperiodic motion.

Ex:- Pendulum in viscous medium.

February

7 mon

(b) Case - II $k^2 < \omega^2$ — Under damping / low damping.
Here $\sqrt{k^2 - \omega^2}$ is imaginary.

So. eqn ⑥ $\sqrt{k^2 - \omega^2} = \sqrt{-(\omega^2 - k^2)} = i\sqrt{\omega^2 - k^2} = iw$,
where $\sqrt{-1} = i$ and $w = \sqrt{\omega^2 - k^2}$.

∴ Eqn ⑥ $\Rightarrow x = e^{-kt} [A e^{i\omega t} + B e^{-i\omega t}]$

8 Tue

$$x = e^{-kt} [A(\cos \omega t + i \sin \omega t) + B(\cos \omega t - i \sin \omega t)]$$


$$x = e^{-kt} [(A+B) \cos \omega t + i(A-B) \sin \omega t]$$

$$\Rightarrow A+B = a_0 \sin \phi ; i(A-B) = a_0 \cos \phi .$$

∴ $x = e^{-kt} [a_0 \sin \phi \cos \omega t + a_0 \cos \phi \sin \omega t]$.

9 wed $x = a_0 e^{-kt} [\sin \phi \cos \omega t + \cos \phi \sin \omega t]$
 $x = a_0 e^{-kt} \sin(\omega t + \phi)$ $\rightarrow ⑧$.

$a_0 e^{-kt}$ — represents amplitude of oscillation and it decays exponentially with time due to e^{-kt} .
 e^{-kt} — Damping factor.



January 2005


Sun	Mon	Tue	Wed	Thu	Fri	Sat
30	31				1	
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29

Notes


⑥ Case III $k^2 = \omega^2$ (critical damping)
 Eqn ⑥ will be like $x = (A+B)e^{-kt}$

Therefore, the displacement approaches to zero asymptotically for given value of initial position.

Here, the critically damped oscillator approaches to the equilibrium position more rapidly than compared to heavy damped.



Curves of all the three cases:



Sun	Mon	Tue	Wed	Thu	Fri	Sat
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

February

14 mon

iii) Forced/Driven Oscillators (FHO):

When oscillator is subjected to external periodic force and it oscillates with the frequency of external periodic force, it is called as Forced Harmonic Oscillator (FHO).

15 tue

$$F = F_{\text{Restoring}} + F_{\text{Damping}} + F_{\text{ext-periodic force}}$$

$$= -kx - \gamma v + F_0 \sin pt.$$

$$ma = -kx - \gamma \frac{dx}{dt} + F_0 \sin pt.$$

$$m \frac{d^2x}{dt^2} + \gamma \frac{dx}{dt} + kx = F_0 \sin pt.$$

16 wed Divide the above eqn by m .

$$\frac{d^2x}{dt^2} + \frac{\gamma}{m} \frac{dx}{dt} + \left(\frac{k}{m}\right)x = \frac{F_0}{m} \sin pt.$$

$$\boxed{\frac{d^2x}{dt^2} + 2K \frac{dx}{dt} + \omega^2 x = f_0 \sin pt} \rightarrow ①$$

Eqn ① is the differential eqn of FHO.

$$\text{where } \frac{\gamma}{m} = 2K ; \frac{k}{m} = \omega^2 ; \frac{F_0}{m} = f_0.$$

January 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
30	31					1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29

Notes

20

February
17th

Let us assume the solution of eqn ① as.

$$x = A \sin(pt - \theta)$$

where p is the driving frequency.

$$\frac{dx}{dt} = Ap \cos(pt - \theta)$$

$$\frac{d^2x}{dt^2} = -Ap^2 \sin(pt - \theta).$$

Substituting above values in eqn ①.

$$-Ap^2 \sin(pt - \theta) + 2KAp \cos(pt - \theta) + \omega^2 A \sin(pt - \theta) \\ = f_0 \sin((pt - \theta) + \theta). \quad \rightarrow \sin(A+B)$$

$$\Rightarrow (\sin pt - \theta)(\omega^2 A - Ap^2) + 2KAp \cos(pt - \theta) = \\ f_0 \sin(pt - \theta) \cos \theta + f_0 \cos(pt - \theta) \sin \theta.$$

$$\Rightarrow A(\omega^2 - p^2) \sin(pt - \theta) + \cancel{2KAp \cos(pt - \theta)} = \\ f_0 \sin(pt - \theta) \cos \theta + f_0 \sin \theta \cos(pt - \theta). \quad \rightarrow ②$$

Now case-1 If $(pt - \theta) = 0$ then $\therefore \sin 0 = 0$
 eqn ② $\Rightarrow 2KAp = f_0 \sin \theta \rightarrow ③ \quad \cos \theta = 1$

case-2 If $(pt - \theta) = \pi/2$ then $\therefore \sin \pi/2 = 1$
 eqn ② $\Rightarrow (\omega^2 - p^2) A = f_0 \cos \theta \rightarrow ④ \quad \cos \pi/2 = 0.$

March 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

Notes

February

21 mon

20

Squaring and adding eqn ③ and ④ we get

$$4K^2 p^2 A^2 + A^2 (\omega^2 - p^2)^2 = f_0^2 (\sin^2 \theta + \cos^2 \theta)$$

$$A^2 [4K^2 p^2 + (\omega^2 - p^2)^2] = f_0^2$$

22 tue

$$A = \frac{f_0}{\sqrt{(\omega^2 - p^2)^2 + 4K^2 p^2}} \rightarrow ⑤$$

Now let us substitute eqn ⑤ in $x = A \sin(pt - \theta)$

$$x = \frac{f_0}{\sqrt{(\omega^2 - p^2)^2 + 4K^2 p^2}} \sin(pt - \theta) \rightarrow ⑥$$

23 wed

Eqn ⑥ is the General solution eqn of FHO.

② Case - I when $\omega > p$ and damping is low

i.e., Driving frequency is very less than the natural frequency.

$$\text{Eqn } ⑤ \Rightarrow A = \frac{f_0}{\omega^2} = \frac{F_0/m}{k/m} = \frac{F_0}{k}$$

January 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
30	31				1	
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29

Notes

February

2005

24 thu


In this case amplitude depends on 'k' force constant only but not on the mass, damping and frequency of driving force.

(b) Case - II When $\omega \ll p$ and damping is low.

then Eqn (5) $\Rightarrow A \approx \frac{f_0}{p^2}$

25 fri

\Rightarrow If 'p' increases then amplitude 'A' will decrease



26 sat

(c) Case - III When $\omega = p$ (Resonance case):

In this case the amplitude will be maximum.

Eqn (5) $\Rightarrow A = \frac{f_0}{\sqrt{4k^2p^2}} \Rightarrow A = \frac{f_0}{2kp}$

March 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

Notes

February/March

28 mon

but in order to get 'A' maximum then we should get denominator of eqn ⑤ be minimum.

Consider $V = (w^2 - p^2)^2 + 4k^2p^2$ should be minimum.

also $\frac{du}{dp}$ should be minimum.

1 tue

$$\therefore \frac{dV}{dp} = 2(w^2 - p^2)(-2p) + 8k^2p = 0.$$

$$\Rightarrow -4p(w^2 - p^2) + 8k^2p = 0.$$

$$8k^2p - 4p(w^2 - p^2) = 0.$$

$$4p(2k^2 + p^2 - w^2) = 0.$$

2 wed

In the above eqn, $4p \neq 0$ since driving frequency can't be zero.

$$\therefore p^2 - w^2 + 2k^2 = 0$$

$$p^2 = w^2 - 2k^2$$

$$p = \sqrt{w^2 - 2k^2} = p_r \rightarrow \text{Resonant frequency}$$

February 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28					

Notes

March

3 Tu

At this frequency ' ω_r ' the amplitude will be maximum.

Substitute ' ω_r ' in the above expression for amplitude

$$A = \frac{\omega_0}{\sqrt{(\omega^2 - \omega_r^2)^2 + 4k^2\omega^2}}$$

4 Mi

Replace $\omega \rightarrow \omega_r = \sqrt{\omega^2 - 2k^2}$

$$\therefore A = \frac{\omega_0}{\sqrt{(\omega^2 - (\omega^2 - 2k^2))^2 + 4k^2(\sqrt{\omega^2 - 2k^2})^2}}$$

5 Sat

$$A_{\max} = \frac{\omega_0}{\sqrt{(2k^2)^2 + 4k^2(\omega^2 - 2k^2)}}$$

$$A_{\max} = \frac{\omega_0}{2k\sqrt{\omega^2 - k^2}}$$

For a certain value of driving frequency the amplitude will be maximum which is called as amplitude maximum Resonance.

April 2005

Sun	Mon	Tue	Wed	Thu	Fri	Sat
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30

Notes

•