

Dongfang Angular Motion Law and Operator Equations

X. D. Dongfang

*Orient Research Base of Mathematics and Physics,
Wutong Mountain National Forest Park, Shenzhen, China*

Quantum mechanics based on Planck hypothesis and statistical interpretation of wave function has achieved great success in describing the discrete law of micro motion. However, the idea of quantum mechanics has not been successfully used to describe the discrete law of macro motion, and the causality implied in the Planck hypothesis and the application scope of the basic principles of quantum mechanics have not been clarified. Here, I first introduce the angular motion law and its application, which seems to be of no special significance as a supplement to the very perfect classical mechanics, but plays an irreplaceable role in testing whether the core mathematical program of quantum mechanics of the operator evolution wave equation meets the unitary principle. Then, the operator evolution wave equations corresponding to the angular motion law are discussed, and the necessity of generalized optimization of differential equations are illustrated by the form of ordinary differential equations. Finally, the real wave equation which is superior to the Schrödinger equation in physical meaning but not necessarily the ultimate answer is briefly introduced. The implicit conclusion is that Hamiltonian can not be the only inevitable choice of constructing wave equation in quantum mechanics, and there is no causal relationship between operator evolution wave equation and quantized energy in bound state system, which indicates that whether the essence of quantum mechanics can be completely revealed is the key step to unify macro and micro quantized theory.

Keywords: Centrifugal force equation; Angular motion law; Operator principle; Wave equations of multiwave functions; Optimal differential equation; Existence and uniqueness of solution.

PACS number(s): 02.30.Jr—Partial differential equations; 03.65.Ge—Solutions of wave equations: bound states; 03.65.Pm—Relativistic wave equations

1 Introduction

From Planck's hypothesis^[1,2] to Einstein's photon theory^[3], to Bohr's hydrogen atom theory^[4], to Schrödinger's equation^[5] and Dirac's equation^[6], quantum mechanics^[7-11] has achieved great success in explaining the laws of blackbody radiation^[12], photoelectric effect^[13] and hydrogen atom spectrum^[14], and has also constantly made new breakthroughs in the practice and applications^[15-19]. Many problems of quantum mechanics have been clearly concluded.

However, the essence of quantum mechanics has not been clarified. This fundamental factor has led to the neglect of some fundamental problems which play a decisive role in quantum mechanics. For example, can or how to use basic principles to prove Planck's quantized energy hypothesis^[20]? How to prove Bohr's^[21] or Sommerfeld's^[22,23] quantization conditions? How to derive Schrödinger wave equation^[24,25] according to the basic principle? What is the scope of application of the basic principles of quantum mechanics such as the representation of mechanical quantities by operators? Can the fine structure constants^[26] in the atomic system change? Is Bowen's statistical interpretation^[27] the

ultimate physical meaning of wave function? Is the concept of orbit compatible with the physical meaning of probabilistic wave? Is the quantum mechanics operation method of constructing wave equation by expressing mechanical quantities in the laws of mechanics as operators suitable for describing the discrete law of macroscopic motion? We need to find the answers to these questions and clarify the scope of application of the principles of quantum mechanics, so as to systematically describe the macro discrete law and micro discrete law in the same logical framework. Some basic principles and mechanical laws ignored by traditional theories will play an important role.

Here I introduce the Dongfang angular motion law and the concise method of solving Kepler's orbital equation of motion by using the angular motion law, and introduce the multi wave function wave equations of micro motion corresponding to the angular motion law, so as to reveal the rich and colorful new mathematical and physical problems hidden in quantum mechanics. The solution of all these problems is the premise to solve the basic difficulties of quantum mechanics, including revealing the essence of quantum mechanics, deciding the establishment of the basic principles of com quantum and

the development direction of com quantum theory, and will also promote the further development of the analytical theory of differential equations.

2 Dongfang angular motion law

It is one of the important basic principles of quantum mechanics that the mechanical quantities of classical mechanical laws are expressed by operators and used to action on wave functions. New laws of mechanics or different expressions of existing laws of mechanics may have different effects on the construction of the wave equation. So, in addition to the equation describing the relationship between momentum and energy, are there any other classical mechanical equations that can be used to establish wave equations? What about the solutions of wave equations established by new mechanical equations? In order to find the answers to these questions, a universal law of angular motion and its operator evolution wave equations are proposed.

Assuming that the object moves in a curve, the mass of the object is m , the size of the position vector is r , the direction angle is θ , the radial velocity of the motion is $\mathbf{v}_r = v_r \mathbf{e}_r$, and the transverse velocity is $\mathbf{v}_\theta = v_\theta \mathbf{e}_\theta$, then the radial force \mathbf{F}_r and the transverse force F_θ of the object satisfy the equations respectively,

$$\begin{aligned} F_r &= \frac{m}{r} \left(v_\theta \frac{dv_r}{d\theta} - v_\theta^2 \right) \\ F_\theta &= \frac{mv_\theta}{r} \left(\frac{dv_\theta}{d\theta} + v_r \right) \end{aligned} \quad (1)$$

Among them, the central force F_r is positive when it is repulsive force and negative when it is attractive; the transverse force F_θ is positive when it is dynamic force and negative when it is resistance force. Equation (1) is called the Dongfang angular motion law.

The centripetal force F_n of a body moving along a curve is usually expressed by the rate v and the radius ρ of curvature, $F_n = mv^2\rho^{-1} = p^2(m\rho)^{-1}$. The angular motion law is the generalized centripetal force equation. The common feature of the two methods is that they contain momentum squares and have no relation to energy, so they can not be used directly to construct Hamilton operators. This provides a basis for the comparison demonstration of the applicable scope of the basic operation methods of quantum mechanics.

Now let's prove the Dongfang angular motion law. The physical quantities are represented by complex numbers in exponential form^[28]. The polar coordinate system or complex plane xoy is established, as shown in Figure 1. To avoid confusion with the imaginary number unit i of quantum mechanics, $j = \sqrt{-1}$ is used here to represent the imaginary number of units. The complex exponential form of orbital equation $\mathbf{r} = \mathbf{r}(t)$ is $\mathbf{r} = r(t) e^{j\theta(t)}$, where θ is the polar angle and t is the time. The dynamic radial and transverse unit vectors

\mathbf{e}_r and \mathbf{e}_θ satisfy the right hand rule. The complex exponential form of the velocity vector is obtained by calculating the first derivative of \mathbf{r} with respect to time t ,

$$\mathbf{v} = \frac{d\mathbf{r}}{dt} = (v_r + jv_\theta) e^{j\theta} \quad (2)$$




Figure 1 The momentum, radial momentum and transverse momentum of a particle in curvilinear motion on the plane of polar coordinates

The real part and the imaginary part in parentheses correspond to the radial velocity $v_r = dr/dt$ and the transverse velocity $v_\theta = r d\theta/dt$, respectively. The derivative of transverse velocity with respect to time t is transformed into a derivative with respect to angle θ . One has $dt = r d\theta/v_\theta$, so $dv_r/dt = (v_\theta/r) dv_r/d\theta$ and $dv_\theta/dt = v_\theta dv_\theta/(r d\theta)$. From this, the derivative of velocity (2) with respect to time t is calculated, and the following complex exponential form of the acceleration vector is obtained,

$$\mathbf{a} = \frac{d\mathbf{v}}{dt} = \frac{1}{r} \left[\left(v_\theta \frac{dv_r}{d\theta} - v_\theta^2 \right) + j \left(v_\theta \frac{dv_\theta}{d\theta} + v_r v_\theta \right) \right] e^{j\theta} \quad (3)$$

The real part and the imaginary part in curly brackets are opposite to the radial acceleration a_r and the transverse acceleration a_θ , respectively. The transverse velocity formula $r d\theta/dt = v_\theta$ is used in the above expression.

The complex exponential representation of the central force is $\mathbf{F} = (F_r + jF_\theta) e^{j\theta}$. Where F_r takes positive value to represent repulsive force and the direction is the same as that of \mathbf{e}_r ; F_r takes negative value to represent gravitation and the direction is opposite to that of \mathbf{e}_r ; F_θ takes positive value to represent power and the direction is the same as that of \mathbf{e}_θ ; F_θ takes negative value to represent resistance and the direction is opposite to that of \mathbf{e}_θ . In the complex plane, the complex exponential representation of Newton's law of motion^[29] is as follows,

$$(F_r + jF_\theta) e^{j\theta} = \frac{m}{r} \left[\left(v_\theta \frac{dv_r}{d\theta} - v_\theta^2 \right) + j \left(v_\theta \frac{dv_\theta}{d\theta} + v_r v_\theta \right) \right] e^{j\theta} \quad (4)$$

Hence, from the definition of complex number equality, the equation of the angular motion law (1) is obtained.

The angular motion law is a generalized form of centripetal force equation describing curve motion. The characteristic of the angular motion law is that the derivative of time can be transformed into the derivative of angle. Linear motion has no angular motion, so the angular motion law is not applicable to linear motion.

3 Angular motion law in central force field

From classical mechanics to quantum mechanics, the problem of a central force is one of the standard subjects. Coulomb's law and the law of gravitation are both central forces, which control the law of atomic spectrum and the law of celestial motion respectively, and maintain the dynamic stability of the micro-world and the macro-world. The transverse force in the central force field is zero. According to the second formula of equation (1), we obtain the relation $dv_\theta/d\theta + v_r = 0$. So $v_r = -dv_\theta/d\theta$, $v_r^2 = v_r v_r = -v_r (dv_\theta/d\theta)$. From $v^2 = v_r^2 + v_\theta^2$, we get the formula $v_\theta^2 = v^2 + v_r dv_\theta/d\theta$, which is substituted for the first formula of (1) to obtain several expressions of the angular motion law of the central force field,

$$\begin{aligned} F_r &= \frac{m}{r} \left(-v_\theta \frac{d^2 v_\theta}{d\theta^2} - v_\theta^2 \right) \\ F_r &= \frac{m}{r} \left(v_\theta \frac{dv_r}{d\theta} - v_r \frac{dv_\theta}{d\theta} - v_r^2 - v_\theta^2 \right) \\ F_r &= \frac{m}{r} \left[\left(\frac{dv_\theta}{d\theta} \right)^2 - v_\theta \frac{d^2 v_\theta}{d\theta^2} - v_r^2 - v_\theta^2 \right] \end{aligned} \quad (5)$$

The radial velocity of the circular motion is $v_r = 0$, and the centripetal force is directed to the center of the cir-

cle, so F_r is negative. The centripetal force equation $F_r = mv_\theta^2/r$ is obtained by substituting this into the first formula (1) or (5) formula. It can be seen that the centripetal force equation is a special case of the angular motion law. Newton infers that the interaction force between planets and the sun follows the law of inverse square ratio by using the centripetal force equation of circular motion. By using the angular motion law or Newton's law of motion, it can be proved that the interaction force governing the motion of a conic curve has and only has the law of the inverse square ratio. The proof process is omitted here.

Dongfang's law of angular motion itself is not a breakthrough discovery. Perhaps because the mathematical process of proving the two-dimensional angular motion law is too ordinary, we even think that it should not be proposed as a new law or principle. However, if we try to write and prove the three-dimensional angular motion law, and apply the two-dimensional and three-dimensional angular motion laws to quantum mechanics to construct abundant wave equations, we will find that the difficulties of accurate solutions of new wave equations and the unknown physical meaning of new wave equations pose challenges to the operator principle of quantum mechanics, and we can realize that the proposal of Dongfang's angular motion law is not only necessary, but also of far-reaching significance.

According to Newton's law of motion and vector operation rules, we can derive the three-dimensional form of the angular motion law which contains many but not completely independent equations. The three-dimensional form of the angular motion law in spherical coordinate system of a central force field is a system of equations containing multiple equations,

$$\begin{aligned} v_\theta \frac{dv_\theta}{d\theta} + v_\theta v_r - v_\varphi^2 \cot \theta &= 0 \\ \frac{1}{\sin \theta} \frac{dv_\varphi}{d\varphi} + v_r + v_\theta \cot \theta &= 0, \quad \frac{dv_\varphi}{d\varphi} + v_r \sin \theta + v_\theta \cos \theta = 0 \\ v_\varphi \frac{dv_\theta}{d\varphi} - v_\theta \frac{dv_\varphi}{d\varphi} + v_r v_\varphi \frac{d\theta}{d\varphi} - (v_\varphi^2 + v_\theta^2) \cos \theta - v_r v_\theta \sin \theta &= 0 \\ F_r &= \frac{m}{r} \left(v_\theta \frac{dv_r}{d\theta} - v_\theta^2 - v_\varphi^2 \right), \quad F_r = \frac{m}{r} \left(\frac{v_\varphi}{\sin \theta} \frac{dv_r}{d\varphi} - v_\theta^2 - v_\varphi^2 \right) \\ F_r &= \frac{m}{r} \left(v_\theta \frac{dv_r}{d\theta} - v_r \frac{dv_\theta}{d\theta} - v_r^2 - v_\theta^2 \right) + m (v_r \cos \theta - v_\theta \sin \theta) v_\varphi \frac{d\varphi}{d\theta} \\ F_r &= \frac{m}{r \sin \theta} \left[v_\varphi \frac{dv_r}{d\varphi} - v_r \frac{dv_\varphi}{d\varphi} - (v_r^2 + v_\varphi^2) \sin \theta - v_r v_\theta \cos \theta - v_\varphi v_\theta \frac{d\theta}{d\varphi} \right] \end{aligned} \quad (6)$$

In the above equation, when F_r is the repulsive force, it takes a positive value, and when F_r is the attractive force, it takes a negative value. The momentum representation of the three-dimensional angular motion law is omitted here. Otherwise there are too many equations and the content appears redundant. The form of Newton's law of motion in the spherical coordinate sys-

tem can also be incorporated into the three-dimensional angular motion law. On the other hand, if you discuss the relativistic momentum form of the three-dimensional angular motion law, you will find the difficulties in relativistic logic. However, this is not the focus of studying the angular motion law and its operator evolution equation. The reader can discuss related issues separately.

It is very simple to solve Kepler orbital equation of motion of an object under the action of the inverse square ratio law $F_r = K/r^2$ with the angular motion law. Here F_r denotes the central force, K is a constant, $K > 0$ for repulsion, $K < 0$ for gravitation, and the transverse force is $F_\theta = 0$. According to the first formula of (5), there is

$$F_r = -\frac{m}{r} \left(v_\theta \frac{d^2 v_\theta}{d\theta^2} + v_\theta^2 \right) = -\frac{mrv_\theta}{r^2} \left(\frac{d^2 v_\theta}{d\theta^2} + v_\theta \right) \quad (7)$$

Where m denotes mass. $mr v_\theta = L$ is an angular momentum constant, from which an expression of transverse velocity $v_\theta = L/mr$ is obtained. Then the second order homogeneous linear differential equation with constant coefficients is obtained by using the central force $F_r = K/r^2$, and its simplified form is

$$\frac{d^2}{d\theta^2} \left(\frac{L}{mr} \right) + \left(\frac{L}{mr} \right) = -\frac{K}{L}$$

According to the general and special solutions of second order homogeneous linear differential equation with constant coefficients, the conic solution $L/mr = A \cos(\theta + \alpha) - K/L$ is obtained. Among them, A and α are undetermined coefficients, which are determined by the orientation of the selected coordinate axis and the initial value conditions. Take $\alpha = 0$, and remember that $r = r_0$ when $\theta = 0$, one obtains $A = L/mr_0 + K/L$. The above Kepler orbital equation in polar coordinates expressed by angular momentum is transformed into the following form,

$$r = \frac{L^2/mK}{(1 + L^2/mr_0 K) \cos \theta - 1} \quad (8)$$

Dongfang's angular motion law can also be used to solve the common physical problems related to satellite motion. Of course, in mechanics, because the development of classical mechanics has been very perfect, the angular motion law as a supplement to classical mechanics seems to have no special significance, although its physical meaning is more clear than the radial force formula of curvilinear motion because it describes radial force and transverse force at the same time, and it is more convenient to solve some complex problems of curvilinear motion. However, the angular motion law plays an irreplaceable role in testing the basic theoretical program of the operator evolution wave equation of quantum mechanics and thus the logical basis of quantum mechanics. This is because of the existence of a new law of mechanics expressed by momentum. If the mathematical art of wave equation evolution through mechanical law operator is an inevitable physical logic, then Hamiltonian is no longer the only choice for quantum mechanics to construct wave equation.

4 Operator evolution wave equations of angular motion law

The unitary principle is an important basic principle generally applicable to the logical self consistency test of natural science theory and social science theory^[30, 31]. I used the unitary principle to prove that the assumption of constant speed of light can not be universally established, and find out the morbid equation of quantum numbers implied in quantum mechanics. As long as a theory implies discordant logical contradictions, it can be tested by the unitary principle. According to the unitary principle, the exact solution of the operator evolution wave equation of angular motion law must be consistent with the exact solution of the operator evolution wave equation of Hamiltonian. Otherwise, the operator theory, which is the key mathematical art of quantum mechanics, will be severely challenged because it violates the unitary principle.

Now let's discuss the operator evolution wave equation of angular motion law. The momentum \mathbf{p} is decomposed into radial momentum $p_r = mv_r$ and transverse momentum $p_\theta = mv_\theta$. The momentum expressions of equation (1) and (5) of the angular motion law are as follows.

$$\begin{aligned} p_r + \frac{dp_\theta}{d\theta} &= 0 \\ F_r &= \frac{1}{mr} \left(p_\theta \frac{dp_r}{d\theta} - p_\theta^2 \right) \\ F_r &= -\frac{1}{mr} \left(p_\theta \frac{d^2 p_\theta}{d\theta^2} + p_\theta^2 \right) \\ F_r &= \frac{1}{mr} \left(p_\theta \frac{dp_r}{d\theta} - p_r \frac{dp_\theta}{d\theta} - p^2 \right) \end{aligned} \quad (9)$$

Where the third formula is derived by substituting the first formula into the second formula and eliminating p_r . The latter three formulas of this equation system are equivalent. It is one of the basic principles of quantum mechanics to replace mechanical quantities in mechanical laws by operators to construct the wave equation. Three-dimensional space usually constructs the rectangular coordinate form of the wave equation first, and then transforms it into spherical coordinate form. The mechanical quantity of the angular motion law is replaced by the operator to construct wave equation, which will encounter the problem of expression of the radial momentum operator which has been controversial. Historically, when solving Dirac equation of hydrogen atoms in spherical coordinates, the radial momentum operator was defined as $\hat{p}_r = -i\hbar(\partial/\partial r + 1/r)$. The result of the debate is to agree with this expression. However, the deduction of theoretical physics should follow the rules of the mathematical operation. Different mathematical forms of operators in different coordinates can be converted to each other, when they move from spherical coordinates to rectangular coordinates. The expres-

sion of radial momentum operators mentioned above will be mathematically difficult. The correct form of translating momentum operators in rectangular coordinates directly into radial and transverse momentum operators in plane polar coordinates is as follows,

$$\hat{p}_r = -i\hbar \frac{\partial}{\partial r}, \quad \hat{p}_\theta = -\frac{i\hbar}{r} \frac{\partial}{\partial \theta} \quad (10)$$

where \hbar is Planck constant and $\hbar = (2\pi)^{-1} h$. The above radial momentum operator and transverse momentum operator (10) are necessary mathematical inferences. If Dirac equation's rectangular coordinate system is directly transformed into a polar coordinate system, for the hydrogen atom, the expected solution of the standard theory can still be obtained by solving the equation with boundary conditions^[32,33]. To achieve this goal, it is not necessary to redefine the independent conservative quantity $\hbar\hat{k}$ in relativistic quantum mechanics. Constructing the expected solution must not destroy any mathematics rule. The redefinition of the radial momentum operator $\hbar = (2\pi)^{-1} h$ violates the basic mathematical operation rules and is unnecessary.

There is a more complex relationship between radial momentum and angular momentum in the momentum representation of the angular motion law. When mechanical quantities are replaced by operators, the problem of composite operators arises in the equivalence relations $p_\theta (dp_r/d\theta) = (dp_r/d\theta) p_\theta$, $p_\theta (d^2 p_\theta/d\theta^2) = (d^2 p_\theta/d\theta^2) p_\theta$ of mechanical quantity contained in equation (9). Because of the lack of corresponding math-

ematical basis, the standard answers to these questions are still unknown. According to the stochastic expression of the system of equations (9), the forms of compound operators in the system of equations are written by using the formula (10), respectively. How to calculate one-step $\hat{p}_\theta (d\hat{p}_r/d\theta)$, $\hat{p}_\theta (d^2 \hat{p}_\theta/d\theta^2)$ and $\hat{p}_r (d\hat{p}_\theta/d\theta)$? Is $\hat{p}_\theta (d\hat{p}_r/d\theta)$ and $(d\hat{p}_r/d\theta) \hat{p}_\theta$ commutative, $\hat{p}_\theta (d^2 \hat{p}_\theta/d\theta^2)$ and $(d^2 \hat{p}_\theta/d\theta^2) \hat{p}_\theta$ commutative, $\hat{p}_r (d\hat{p}_\theta/d\theta)$ and $(d\hat{p}_\theta/d\theta) \hat{p}_r$ commutative? If only a random expression of the system of equations (9) is taken, the forms of the related composite operators in the system of equations written by formula (10) are as follows,

$$\begin{aligned} \hat{p}_\theta^2 &= \left(-\frac{i\hbar}{r} \frac{\partial}{\partial \theta} \right) \left(-\frac{i\hbar}{r} \frac{\partial}{\partial \theta} \right) = -\hbar^2 \left(\frac{1}{r} \frac{\partial}{\partial \theta} \right) \left(\frac{1}{r} \frac{\partial}{\partial \theta} \right) \\ \hat{p}_\theta \frac{d\hat{p}_r}{d\theta} &= -\frac{i\hbar}{r} \frac{\partial}{\partial \theta} \left[\frac{d}{d\theta} \left(-i\hbar \frac{\partial}{\partial r} \right) \right] = -\frac{\hbar^2}{r} \frac{\partial}{\partial \theta} \left[\frac{d}{d\theta} \left(\frac{\partial}{\partial r} \right) \right] \\ \hat{p}_\theta \frac{d^2 \hat{p}_\theta}{d\theta^2} &= -\frac{i\hbar}{r} \frac{\partial}{\partial \theta} \frac{d^2}{d\theta^2} \left(-\frac{i\hbar}{r} \frac{\partial}{\partial \theta} \right) = -\frac{\hbar^2}{r} \frac{\partial}{\partial \theta} \frac{d^2}{d\theta^2} \left(\frac{1}{r} \frac{\partial}{\partial \theta} \right) \\ \hat{p}_\theta \frac{d\hat{p}_r}{d\theta} - \hat{p}_r \frac{d\hat{p}_\theta}{d\theta} &= \hbar^2 \left(\frac{\partial}{\partial r} \right) \frac{d}{d\theta} \left(\frac{1}{r} \frac{\partial}{\partial \theta} \right) - \frac{\hbar^2}{r} \frac{\partial}{\partial \theta} \frac{d}{d\theta} \left(\frac{\partial}{\partial r} \right) \end{aligned}$$

Since the basic method of quantum mechanics expressed by operators is generally applicable, the radial and transverse momentum in formula (9) should be replaced by (10) radial and transverse operators and used to action on wave functions to describe the motion of the same particle, which requires at least four wave functions ψ_1 , ψ_2 , ψ_3 and ψ_4 . The corresponding four wave equations constitute a wave equation system of real number,

$$\begin{aligned} \frac{\partial \psi_1}{\partial r} + \frac{d}{d\theta} \left(\frac{1}{r} \frac{\partial \psi_1}{\partial \theta} \right) &= 0 \\ \frac{\hbar^2}{mr^2} \left\{ \frac{\partial}{\partial \theta} \left(\frac{1}{r} \frac{\partial \psi_2}{\partial \theta} \right) - \frac{\partial}{\partial \theta} \left[\frac{d}{d\theta} \left(\frac{\partial \psi_2}{\partial r} \right) \right] \right\} - F_r \psi_2 &= 0 \\ \frac{\hbar^2}{mr^2} \left\{ \frac{\partial}{\partial \theta} \left[\frac{d^2}{d\theta^2} \left(\frac{1}{r} \frac{\partial \psi_3}{\partial \theta} \right) \right] + \frac{\partial}{\partial \theta} \left[\left(\frac{1}{r} \frac{\partial \psi_3}{\partial \theta} \right) \right] \right\} - F_r \psi_3 &= 0 \\ \frac{\hbar^2}{mr} \left\{ \nabla^2 \psi_4 + \frac{\partial}{\partial r} \left[\frac{d}{d\theta} \left(\frac{1}{r} \frac{\partial \psi_4}{\partial \theta} \right) \right] - \frac{1}{r} \frac{\partial}{\partial \theta} \left[\frac{d}{d\theta} \left(\frac{\partial \psi_4}{\partial r} \right) \right] \right\} - F_r \psi_4 &= 0 \end{aligned} \quad (11)$$

These equations are all linear equations, but unlike Schrödinger equation and Dirac equation, the equation does not contain time, and the central force replaces the energy parameter.

At present, it is difficult to find the exact solution of the wave equation system (11). This is not only because the exact solution method of each equation is not clear, but also because there is no answer to the question whether the wave functions in the equation system are consistent or not and how to explain the physical meaning of each wave function. Since the last two equations of the angular motion law equation (9) are derived from the first two equations, the last three equations of wave equation system (11) should be equivalent, but the forms of the last three wave equations are quite differen-

t. The more difficult problem is whether the operators are commutative or not because of the different writing order of mechanics quantities. In terms of mathematical form only, the third derivative of the last three equations of wave equation system (11) seems to be out of line with common sense. Whether the wave equation system means that the scope of application of momentum operators is actually very limited, so the momentum in the laws of mechanics can not always be replaced by operators to construct a wave equation. It is very interesting and important to study the answers to the questions.

The wave equation system (11) is derived by applying the operator operation principle of quantum mechanics to the angular motion law. The angular motion law describes the motion of curves, so the wave equa-

tion system (11) is applicable to the motion of curves. Schrödinger equation and Dirac equation do not seem to be confined to the angular motion conditions, but actually conceal more complex logic difficulties. For example, Schrödinger equation of linear harmonic oscillator has expected quantized energy solution, whereas Klein-Gordon equation^[34,35] or Dirac equation of relativistic linear harmonic oscillator has no quantized expected solution; the orbital differential equation of curvilinear motion under the action of the law of inverse square ratio has the exact solution of a conic curve, and the orbital differential equation of linear motion under the action of the law of inverse square ratio is a non-linear differential equation and can not give the exact solution of the orbital equation. These difficult mathematical problems of theoretical physics are more enlightening.

5 Ordinary differential form of wave equation

Wave equation system (11) contains composite differential operation, which reveals an important problem that has been neglected for a long time, that is, the classical concept of orbital equation $r = r(\theta)$ is not abandoned because of the principle of quantum mechanics. To determine the spatial orientation of particles, r and θ represent independent coordinate parameters. To study the law of motion of objects governed by interaction forces, the orbital equation $r = r(\theta)$ is a constraint equation, and r and θ are not independent parameter-

s. However, in the Schrödinger equation, Dirac equation and even other mathematical and physical equations, r and θ have been treated as independent parameters and achieved great success. Therefore, a wave equation expressed by a partial differential equation may actually be an ordinary differential equation. Then, is the exact solution of such an ordinary differential equation the same as that of the original partial differential equation? This is a new mathematical and physical problem worthy of further study.

Considering two-dimensional hydrogen atom, the Coulomb force between electrons and protons is $F_r = -e^2 (4\pi\epsilon_0 r^2)^{-1} = -\alpha\hbar c r^{-2}$. Among them, r is the distance between electron and proton, e is the quantity of elementary charge, ϵ_0 is the dielectric constant in vacuum, α is the fine structure constant, $\hbar = h(2\pi)^{-1}$ and h is the Planck constant, c is the speed of light in vacuum. The polar coordinate form of the orbital equation of an electron moving around a proton is,

$$r = \frac{\delta}{1 - \zeta \cos \theta} \quad (12)$$

where polar angle θ represents the direction of the position vector. In order to avoid confusion caused by using the same symbols for different physical quantities, δ is used to represent the focus parameter and ζ is used to represent the eccentricity. The Coulomb force between the electron and the proton is substituted for the wave equation group (11), and the wave equation group of the hydrogen atom is obtained,

$$\begin{aligned} \frac{\partial \psi_1}{\partial r} + \frac{d}{d\theta} \left(\frac{1}{r} \frac{\partial \psi_1}{\partial \theta} \right) &= 0 \\ \frac{\hbar^2}{m} \left\{ \frac{\partial}{\partial \theta} \left(\frac{1}{r} \frac{\partial \psi_2}{\partial \theta} \right) - \frac{\partial}{\partial \theta} \left[\frac{d}{d\theta} \left(\frac{\partial \psi_2}{\partial r} \right) \right] \right\} - \frac{e^2 \psi_2}{4\pi\epsilon_0} &= 0 \\ \frac{\hbar^2}{m} \left\{ \frac{\partial}{\partial \theta} \left[\frac{d^2}{d\theta^2} \left(\frac{1}{r} \frac{\partial \psi_3}{\partial \theta} \right) \right] + \frac{\partial}{\partial \theta} \left[\left(\frac{1}{r} \frac{\partial \psi_3}{\partial \theta} \right) \right] \right\} + \frac{e^2 \psi_3}{4\pi\epsilon_0} &= 0 \\ \frac{\hbar^2}{mr} \left\{ \nabla^2 \psi_4 + \frac{\partial}{\partial r} \left[\frac{d}{d\theta} \left(\frac{1}{r} \frac{\partial \psi_4}{\partial \theta} \right) \right] - \frac{1}{r} \frac{\partial}{\partial \theta} \left[\frac{d}{d\theta} \left(\frac{\partial \psi_4}{\partial r} \right) \right] \right\} + \frac{e^2 \psi_4}{4\pi\epsilon_0 r^2} &= 0 \end{aligned} \quad (13)$$

This system of equations has no energy parameters, so it is impossible to obtain quantized energy solutions like Schrödinger equation or the Dirac equation directly. However, since r and θ satisfy the constraint equation (10) and are no longer independent coordinate parameters, the energy parameters are actually expressed indirectly by the focal parameters δ and eccentricity ζ of the elliptic equation. This is only a qualitative mathematical conclusion, but the quantitative problem is much more complicated.

According to Dongfang unitary principle, if the principles of quantum mechanics and the statistical interpretation of wave functions are both correct, the solutions of each equation of the wave equation group (13) should be equivalent, otherwise, a reasonable choice must be

made among the non equivalent wave functions. This requires negating other wave function equations without scientific basis for trade-offs. Although the equation system (13) is a system of linear differential equations, it is still difficult to find the exact solutions of the equation system so as to obtain scientific conclusions. Are their exact solutions consistent? Are there any indirect quantized exact solutions representing energy? Whether there is an indirect expression of the exact solution of energy quantization? Does the exact solution of quantization conform to the prediction of quantum mechanics? These problems puzzle us.

The first equation in the wave equation system (11) or (13) is the simplest. It has no controversial question whether the operator is commutative or not. It can be

transformed into an ordinary differential equation by the constraint equation, i.e. orbital equation (12). Because r is a function of θ , the partial derivative of the equation is rewritten to ordinary differential. From the orbital equation (12), we get that,

$$\begin{aligned}\frac{dr}{d\theta} &= \frac{d}{d\theta} \left(\frac{\delta}{1 - \varsigma \cos \theta} \right) = -\frac{\varsigma \delta \sin \theta}{(1 - \varsigma \cos \theta)^2} \\ \frac{\partial \psi_1}{\partial r} &= \frac{d\psi_1}{d\theta} \frac{d\theta}{dr} = -\frac{(1 - \varsigma \cos \theta)^2}{\varsigma \delta \sin \theta} \frac{d\psi_1}{d\theta} \\ \frac{d}{d\theta} \left(\frac{1}{r} \frac{\partial \psi_1}{\partial \theta} \right) &= \frac{1 - \varsigma \cos \theta}{\delta} \frac{d^2 \psi_1}{d\theta^2} + \frac{\varsigma \sin \theta}{\delta} \frac{d\psi_1}{d\theta}\end{aligned}$$

By substituting these relations into the first equation of the wave equation system (11) or (13), the second order ordinary differential equation with respect to the parameter θ is obtained,

$$\frac{d^2 \psi_1}{d\theta^2} - \frac{(1 - \varsigma \cos \theta)^2 - \varsigma^2 \sin^2 \theta}{\varsigma \sin \theta (1 - \varsigma \cos \theta)} \frac{d\psi_1}{d\theta} = 0 \quad (14)$$

Let $\chi = \cos \theta$, then equation (14) is reduced to

$$\begin{aligned}\frac{d\psi_1}{d\theta} &= -\sin \theta \frac{d\psi_1}{d\chi} \\ \frac{d^2 \psi_1}{d\theta^2} &= (1 - \chi^2) \frac{d^2 \psi_1}{d\chi^2} - \chi \frac{d\psi_1}{d\chi}\end{aligned}$$

the equation (14) is transformed into

$$\frac{d^2 \psi_1}{d\chi^2} - \frac{3\varsigma^2 \chi^2 - 3\varsigma \chi + 1 - \varsigma^2}{\varsigma^2 \chi^3 - \varsigma \chi^2 - \varsigma^2 \chi + \varsigma} \frac{d\psi_1}{d\chi} = 0 \quad (15)$$

By using the orbital equation (12), the derivative of wave function to θ can also be transformed into the derivative to r , and then the first equation of wave equation system (11) can be simplified to the ordinary differential equation of the derivative to parameter r ,

$$\frac{d^2 \psi_1}{dr^2} - \frac{(1 - \varsigma^2) r^2 - 2\delta r + 2\delta^2}{(1 - \varsigma^2) r^3 - 2\delta r^2 + \delta^2 r} \frac{d\psi_1}{dr} = 0 \quad (16)$$

By using the above methods, the last three partial differential wave equations of the wave equation system (13) can be transformed into ordinary differential equations. The existence and uniqueness of the exact solutions of these equations must be studied first. Only by finding the exact solution of the equation and comparing the solutions of the wave equation corresponding to the equivalent mechanical equation of different forms, can we know the true physical meaning of the wave function, and then clarify the scope of application of the basic principles of quantum mechanics in which mechanical quantities are replaced by operators. In order to study the exact solutions of equations (15) and (16), we need to develop the analytical theory of differential equations. The relevant mathematical basis is the optimization

theorem of differential equations^[36]. We need to extend the optimization theorem of differential equations to the generalized optimization theorem to study the existence and uniqueness of exact solutions of equations (15) and (16). Of course, finding the exact solution of equation (15) or (16) by any other method will also lead to important new physical conclusions including but not limited to the equation itself.

6 Comments and conclusions

In this paper, the law of angular motion is proposed, and the operator evolution wave equations of the law of angular motion of the central force field are written by using the quantum mechanics calculation principle of constructing the wave equation by replacing mechanical quantities with operators. The simplest equations of the wave equations of the hydrogen atom are simplified into two kinds of ordinary differential equations. At the same time, the necessity of generalized optimization of the differential equations is pointed out.

However, does the exact solution of each wave equation exist? Or the exact solutions exist, but are they consistent? These problems challenge the basic and most important computational rule of operator evolution of the wave equation in quantum mechanics. For any theory, it is only a necessary condition that the principle^[37, 38], method and conclusion of the theory conform to the unitary principle. If it does not conform to the unitary principle, there must be major defects or even mistakes. The law of conservation of momentum, the law of conservation of energy, the law of conservation of angular momentum and the angular motion law are all the inevitable inferences of Newton's law of motion. Applying the angular motion law to quantum mechanics, the obtained wave equation set of the central force field implies the existence of many different wave functions. How to explain the physical meaning of these different wave functions is obviously an urgent topic. The statistical interpretation of wave functions may not be the only physical meaning of wave functions. Whether it belongs to functions that contain different physical meanings such as orbital density can not be affirmed or negated at present. From the point of view of statistical law, the concept of the electron cloud is very applicable, but the probability of a probability wave does not mean the negation of the concept of micro particle trajectory. The problem that quantized energy destroys the law of conservation of energy implied in the morbid equation of quantum numbers^[31] and the exact solutions of the operator evolution equations of angular motion law all show that quantum mechanics can not pass the logical test of the unitary principle, so it is actually a very imprecise theory. But the idea of quantum mechanics is meaningful, and quantum mechanics needs to be revised systematically. I have tested some important ba-

sic physical theory problems with the unitary principle, and found that the difficulty of fundamentally eliminating physical theory needs to correct or prove all basic physical assumptions that lack logical basis.

In quantum mechanics, it is considered that the energy and angular momentum of the interactive system are observable and can be expressed by a linear Hermitian operator. From the point of view of the com quantum phenomena of the macroscopic system, velocity is truly directly observable compared with angular momentum and energy. From the so-called plane wave function $\psi(\mathbf{r}, t) = A \exp[(i/\hbar)(m\mathbf{v} \cdot \mathbf{r} - Et)]$, we can write the velocity operator $\hat{v} = -(i\hbar/m)\nabla$ or the velocity square operator $\hat{v}^2 = -(\hbar^2/m^2)\nabla^2$. Therefore, a strict proof is needed to make a choice between the velocity operator and the momentum operator. Essentially, the steady wave equation means that the mathematical expression of the wave function independent of time is $\partial\psi/\partial t = 0$. But this accurate mathematical expression will lead to time-dependent Schrödinger equation $i\hbar\partial\psi/\partial t = -(\hbar^2/2m)\nabla^2\psi + U(r)\psi$ can not be transformed into the steady equation. The problem means different physical ideas, that is, the establishment of wave equations may not be limited to Hamiltonian operators. For example, considering the energy level transition of a quantum system, the absorbed or radiated energy of the system must follow the law of conservation of energy. Therefore, referring to the Maxwell equations, we can find a more reasonable reason than the Hamiltonian operator evolution method, but it may not be the true portrayal of the natural law, and write the following real number wave equation,

$$\nabla^2\psi + \frac{4\pi^2}{\sigma c^2}(E - U)\psi - \frac{1}{c^2}\frac{\partial^2\psi}{\partial t^2} = 0 \quad (17)$$

Among them, m denotes the mass of particle, E and U denote energy and potential energy respectively, and σ is a constant for com quantum theory. The real wave equation is one of the corollaries of com quantum theory, which shows that the virtual number is not an indispensable element of quantum mechanics. There is no doubt about how to explain the concept of steady state in real wave equation. Because $\partial\psi/\partial t = 0$ corresponds to the steady-state Schrödinger wave equation, and the solution of this steady-state equation is just that have been

given by quantum mechanics. Perhaps readers can write other forms of real wave equations and find the intrinsic relationship between different real wave equations.

It is generally believed that quantum mechanics is one of the perfect and precise scientific theories in mathematics. However, problems such as the scope of application of the basic principles of quantum mechanics and the nature of quantum mechanics have not been solved. The application of an operator operating method in quantum mechanics to the new laws of mechanics such as the angular motion law brings new mathematical and physical difficulties to be solved. Its scientific conclusion is of great significance to the design idea of the assumption that the quantum theory does not depend on the assumption that it is impossible to prove or is finally proved to be very limited. The assumption that accords with the natural law must have exact causality and can be proved and become a theorem. Otherwise, it will not be regarded as the basic principle in the com quantum theory. Even if some hypothetical form logic deduction can solve some difficult problems locally. The wave function of quantum mechanics is interpreted as a probability function. Why can't it be a new physical quantity that has not yet been discovered? What are the wave functions and wave equations of the com quantum theory, which can be used to describe both macroscopic and microscopic discrete laws? Let's find the answers to the questions together.

To sum up, Dongfang angular motion law and the energy momentum equation constitute two metrics for the unitary principle test of quantum mechanics. It is one of the most important basic principles of quantum mechanics to replace mechanical quantity with Hermite operators acting on wave functions to construct wave equations. Applying Dongfang angular motion law, there is enough derived wave equations in the same quantum system, and the existence of solutions of these equations seriously deviates from the expectations of quantum mechanics. Obviously, the operator principle of quantum mechanics does not follow the unitary principle, so it is not universally effective, which means that quantum mechanics has serious mathematical defects. This conclusion should have a far-reaching impact on physics and promote theoretical physics to enter a new period of vigorous development.

-
- [1] Planck, M. On the law of distribution of energy in the normal spectrum. *Annalen der physik* **4**, 1 (1901).
 - [2] Millikan, R. A. A direct photoelectric determination of Planck's "h". *Physical Review* **7**, 355 (1916).
 - [3] Einstein, A. Concerning an heuristic point of view toward the emission and transformation of light. *American Journal of Physics* **33**, 367 (1965).
 - [4] Bohr, N. On the Constitution of Atoms and Molecules. Part II: Systems containing only a Single Nucleus. *Philos. Mag.* **26**, 476 (1913).
 - [5] Schrödinger, E. An undulatory theory of the mechanics of atoms and molecules. *Physical review* **28**, 1049 (1926).
 - [6] Dirac, P. A. M. *The principles of quantum mechanics*. 70-72 (Clarendon Press Oxford, 1958).
 - [7] Jauch, J. M. *Foundation of quantum mechanics*. (Addison-Wesley, 1968).
 - [8] Schiff, L. *Quantum Mechanics*, McGraw-Hill Book Company. (New York, 1955).
 - [9] Landau, L. D. & Lifshitz, E. M. *Quantum mechanics: non-relativistic theory*. Vol. 3 (Elsevier, 2013).
 - [10] Bjorken, J. D. & Drell, S. D. *Relativistic quantum mechanics*. (McGraw-Hill, 1965).

- [11] Greiner, W. Relativistic quantum mechanics: wave equations. (1990).
- [12] Rybicki, G. B. & Lightman, A. P. *Radiative processes in astrophysics*. (John Wiley & Sons, 2008).
- [13] Serway, R. A. & Jewett, J. W. *Physics for scientists and engineers with modern physics*. (Cengage learning, 2018).
- [14] Rydberg, J. R. Recherches sur la constitution des spectres d'émission des éléments chimiques. *Kungliga vetenskapsakademiens handlingar* **23** (1890).
- [15] Bourennane, M. *et al.* Multiphoton entanglement and interferometry. *Fortschritte der Physik: Progress of Physics* **51**, 273-279 (2003).
- [16] Pan, J.-W. *et al.* Multiphoton entanglement and interferometry. *Reviews of Modern Physics* **84**, 777 (2012).
- [17] Wu, Z. *et al.* Realization of two-dimensional spin-orbit coupling for Bose-Einstein condensates. *Science* **354**, 83-88 (2016).
- [18] Liu, Y. *et al.* Device-independent quantum random-number generation. *Nature* **562**, 548 (2018).
- [19] Zeng, Y. *et al.* Entangling Two Individual Atoms of Different Isotopes via Rydberg Blockade. *Physical Review Letters* **119**, 160502 (2017).
- [20] Planck, M. Ueber das Gesetz der Energieverteilung im Normalspectrum. 553-563 (1901).
- [21] Hassoun, G. One and two dimensional hydrogen atoms. *American Journal of Physics* **49**, 143-146 (1981).
- [22] Wilson, W. LXXXIII. The quantum-theory of radiation and line spectra. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **29**, 795-802 (1915).
- [23] Sommerfeld, A. Zur quantentheorie der spektrallinien. *Annalen der Physik* **356**, 1-94 (1916).
- [24] Nelson, E. Derivation of the Schrödinger equation from Newtonian mechanics. *Physical Review* **150**, 1079 (1966).
- [25] Gottfried, K. Inferring the statistical interpretation of quantum mechanics from the classical limit. *Nature* **405**, 533-536 (2000).
- [26] Webb, J. K., Flambaum, V. V., Churchill, C. W., Drinkwater, M. J. & Barrow, J. D. Search for time variation of the fine structure constant. *Physical Review Letters* **82**, 884 (1999).
- [27] Pais, A. Max Born's statistical interpretation of quantum mechanics. *Science* **218**, 1193-1198 (1982).
- [28] Chen, R. Complex number analysis method for central force problem. *College Physics* **4**(1), 41-42 (1987).
- [29] Newton, I. *Newton's Principia : the mathematical principles of natural philosophy*. 83 (Daniel Adee, 1846).
- [30] Dongfang, X. D. [On the Relativity of the Speed of Light](#). *Mathematics & Nature* **1**, 202101 (2021).
- [31] Dongfang, X. D. [The Morbid Equation of Quantum Numbers](#). *Mathematics & Nature* **1**, 202102 (2021).
- [32] Chen, R. Correction to Solution of Dirac Equation. *arXiv preprint arXiv:0908.4320* (2009).
- [33] Chen, R. New exact solution of Dirac-Coulomb equation with exact boundary condition. *International Journal of Theoretical Physics* **47**, 881-890 (2008).
- [34] Sakurai, J. J. *Advanced quantum mechanics*. (Pearson Education India, 1967).
- [35] Baym, G. *Lectures on quantum mechanics*. 499 (CRC Press, 2018).
- [36] Chen, R. The Optimum Differential Equations. *Chinese Journal of Engineering Mathematics*, 82-86,91 (2000).
- [37] Dongfang, X. D. [Relativistic Equation Failure for LIGO Signals](#). *Mathematics & Nature* **1**, 202103 (2021).
- [38] Dongfang, X. D. [Dongfang Modified Equations of Molecular Dynamics](#). *Mathematics & Nature* **1**, 202104 (2021).