Rethinking Teaching Abstract Algebra

Colin Ford

Math/Physics Academy
@MathPhysics.Academy on social media
www.mathphysics.academy

Preface and Disclaimer

Preface and Disclaimer

- I personally struggled with learning abstract algebra, despite excellent help from friends and professors.

Preface and Disclaimer

- I personally struggled with learning abstract algebra, despite excellent help from friends and professors.
- Current approaches are often rather... abstract!

Preface and Disclaimer

- I personally struggled with learning abstract algebra, despite excellent help from friends and professors.
- Current approaches are often rather... abstract!
- Not necessarily a bad thing, but I imagine a different approach than what might be found in e.g. Dummit & Foote or Jacobson, etc.

Preface and Disclaimer

- I personally struggled with learning abstract algebra, despite excellent help from friends and professors.
- Current approaches are often rather... abstract!
- Not necessarily a bad thing, but I imagine a different approach than what might be found in e.g. Dummit & Foote or Jacobson, etc.
- The same friends have differing opinions, and somewhat disagree that my approach is "better."

Preface and Disclaimer

- I personally struggled with learning abstract algebra, despite excellent help from friends and professors.
- Current approaches are often rather... abstract!
- Not necessarily a bad thing, but I imagine a different approach than what might be found in e.g. Dummit & Foote or Jacobson, etc.
- The same friends have differing opinions, and somewhat disagree that my approach is "better."
- I am unaware whether or not others have attempted an approach like this before.

Prerequisites

Prerequisites

- Requires a semester or two of linear algebra

Prerequisites

- Requires a semester or two of linear algebra
- Approach tied to definition of vector space as given in Introduction to MPA video presentation

Prerequisites

- Requires a semester or two of linear algebra
- Approach tied to definition of vector space as given in Introduction to MPA video presentation
- Consequently linear algebra course would be reworked (boo, 8 axioms)

Prerequisites

- Requires a semester or two of linear algebra
- Approach tied to definition of vector space as given in Introduction to MPA video presentation
- Consequently linear algebra course would be reworked (boo, 8 axioms)
- A colleague offered the approach of introducing groups via a simplified study of Galois groups

Why?

Why?

- Groups contextualized, allows for what I see as a nice reveal of other algebraic structures

Why?

- Groups contextualized, allows for what I see as a nice reveal of other algebraic structures
- Unbeknownst to (most) students, the material in linear and abstract algebra courses highly interconnected

Why?

- Groups contextualized, allows for what I see as a nice reveal of other algebraic structures
- Unbeknownst to (most) students, the material in linear and abstract algebra courses highly interconnected
- Groups, rings, (algbebraic number) fields, vector spaces, modules, etc. are everywhere in the undergraduate curriculum

Why?

- Groups contextualized, allows for what I see as a nice reveal of other algebraic structures
- Unbeknownst to (most) students, the material in linear and abstract algebra courses highly interconnected
- Groups, rings, (algbebraic number) fields, vector spaces, modules, etc. are everywhere in the undergraduate curriculum
- Many students (at least in the US) do not see any of this material until late in their second year or their third year

Why?

- Groups contextualized, allows for what I see as a nice reveal of other algebraic structures
- Unbeknownst to (most) students, the material in linear and abstract algebra courses highly interconnected
- Groups, rings, (algbebraic number) fields, vector spaces, modules, etc. are everywhere in the undergraduate curriculum
- Many students (at least in the US) do not see any of this material until late in their second year or their third year
- This does not serve students well whatsoever, in my opinion

A rough layout of linear algebra

A rough layout of linear algebra

- Introduce students to groups and vector spaces similar to Introduction to MPA video presentation

A rough layout of linear algebra

- Introduce students to groups and vector spaces similar to Introduction to MPA video presentation
- Primarily leave a detailed study of groups for later

A rough layout of linear algebra

- Introduce students to groups and vector spaces similar to Introduction to MPA video presentation
- Primarily leave a detailed study of groups for later
- Perhaps mention Galois, Galois groups from a birds-eye point-of-view, but no more than "necessary"

A rough layout of linear algebra

- Introduce students to groups and vector spaces similar to Introduction to MPA video presentation
- Primarily leave a detailed study of groups for later
- Perhaps mention Galois, Galois groups from a birds-eye point-of-view, but no more than “necessary”
- Upon reaching a transition point, return to group and group-action definition of vector space

Perturbing the algebraic structure

Perturbing the algebraic structure

- A *vector space* is an abelian group $(V, +)$ along with a group action by $K = \mathbb{R}$ real or $K = \mathbb{C}$ complex numbers

Perturbing the algebraic structure

- A *vector space* is an abelian group $(V, +)$ along with a group action by $K = \mathbb{R}$ real or $K = \mathbb{C}$ complex numbers
- K is more than a group, it is a field. I.e. we have division, and our elements commute

Perturbing the algebraic structure

- A *vector space* is an abelian group $(V, +)$ along with a group action by $K = \mathbb{R}$ real or $K = \mathbb{C}$ complex numbers
- K is more than a group, it is a field. I.e. we have division, and our elements commute
- Proposal is to perturb this structure to reveal other algebraic structures

Introduce fields either before or concurrently

Introduce fields either before or concurrently

- A field K may be viewed simultaneously as an abelian group under addition and an abelian group under multiplication

Introduce fields either before or concurrently

- A field K may be viewed simultaneously as an abelian group under addition and an abelian group under multiplication
- Until now we swept delicate notions of multiplicative inverses and division under the rug

Some questions

Some questions

What if we have the same structure of a vector space, but

Some questions

What if we have the same structure of a vector space, but

- take away our ability to commute elements of K?

Some questions

What if we have the same structure of a vector space, but

- take away our ability to commute elements of K?
- take away our ability to divide by elements of K?

Some questions

What if we have the same structure of a vector space, but

- take away our ability to commute elements of K?
- take away our ability to divide by elements of K?

What becomes of K, and what becomes of V?

Hit the piñata

Hit the piñata

- K may still be viewed as an abelian group under addition $+$

Hit the piñata

- K may still be viewed as an abelian group under addition $+$
- Hit commutativity:

Hit the piñata

- K may still be viewed as an abelian group under addition $+$
- Hit commutativity: a *division ring* falls out!

Hit the piñata

- K may still be viewed as an abelian group under addition $+$
- Hit commutativity: a *division ring* falls out!
- Hit division:

Hit the piñata

- K may still be viewed as an abelian group under addition $+$
- Hit commutativity: a *division ring* falls out!
- Hit division: a *ring* falls out!

Define rings

Define rings

A *ring* is a set with two binary operations, addition $+$ and multiplication $\times$, satsifying:

Define rings

A *ring* is a set with two binary operations, addition $+$ and multiplication $\times$, satsifying:

- $(R, +)$ is an abelian group

Define rings

A *ring* is a set with two binary operations, addition $+$ and multiplication $\times$, satsifying:

- $(R, +)$ is an abelian group
- $\times$ is associative

  - $(a + b) \times c = (a \times c) + (b \times c)$

Define rings

A *ring* is a set with two binary operations, addition $+$ and multiplication $\times$, satsifying:

- $(R, +)$ is an abelian group
- $\times$ is associative
- the *distributive laws* hold in R: for all a, b, c in R
  - $(a + b) \times c = (a \times c) + (b \times c)$
  - $a \times (b + c) = (a \times b) + (a \times c)$

Define rings

A *ring* is a set with two binary operations, addition $+$ and multiplication $\times$, satsifying:

- $(R, +)$ is an abelian group
- $\times$ is associative
- the *distributive laws* hold in R: for all a, b, c in R
  - $(a + b) \times c = (a \times c) + (b \times c)$
  - $a \times (b + c) = (a \times b) + (a \times c)$
- If $\times$ is commutative, the ring is called *commutative*; if there is a multiplicative identity 1, we say R *has an identity* or *contains a* 1

Define rings

A *ring* is a set with two binary operations, addition $+$ and multiplication $\times$, satsifying:

- $(R, +)$ is an abelian group
- $\times$ is associative
- the *distributive laws* hold in R: for all a, b, c in R
  - $(a + b) \times c = (a \times c) + (b \times c)$
  - $a \times (b + c) = (a \times b) + (a \times c)$
- If $\times$ is commutative, the ring is called *commutative*; if there is a multiplicative identity 1, we say R *has an identity* or *contains a* 1

Rings are kind of weird! We should provide examples.

Examples of rings

Examples of rings

- the integers $\mathbb{Z}$ with familiar addition and multiplication

Examples of rings

- the integers $\mathbb{Z}$ with familiar addition and multiplication
- the quaternions $\mathbb{H}$ with component-wise addition and a distributive multiplication form a non-commutative ring with identity

Examples of rings

- the integers $\mathbb{Z}$ with familiar addition and multiplication
- the quaternions $\mathbb{H}$ with component-wise addition and a distributive multiplication form a non-commutative ring with identity
- polynomials with coefficients in a commutative ring R with identity form a commutative ring with identity, denoted $R[X]$, with component-wise addition and distributive multiplication

Examples of rings

- the integers $\mathbb{Z}$ with familiar addition and multiplication
- the quaternions $\mathbb{H}$ with component-wise addition and a distributive multiplication form a non-commutative ring with identity
- polynomials with coefficients in a commutative ring R with identity form a commutative ring with identity, denoted $R[X]$, with component-wise addition and distributive multiplication
- $n \times n$ matrices with entries from an arbitrary ring form a non-commutative ring $M_n(R)$ for $n \geq 2$. If R has identity, so does $M_n(R)$

What changes?

What if our vector space's group action was by a ring instead of a field?

What changes?

What if our vector space's group action was by a ring instead of a field?

- Do we lose linearity?

What changes?

What if our vector space's group action was by a ring instead of a field?

- Do we lose linearity?
- What kinds of things can be multiplied but not divided?

What changes?

What if our vector space's group action was by a ring instead of a field?

- Do we lose linearity?
- What kinds of things can be multiplied but not divided?
- What kinds of multiplicative things cannot be commuted?

What changes?

What if our vector space's group action was by a ring instead of a field?

- Do we lose linearity?
- What kinds of things can be multiplied but not divided?
- What kinds of multiplicative things cannot be commuted?

Any real world execution of this may be met with (pleasant) surprise.

Definition of a module

Definition of a module

Let R be an arbitrary ring. A *left R-module* is a set M together with

Definition of a module

Let R be an arbitrary ring. A *left R-module* is a set M together with

- a binary operation $+$ such that $(M, +)$ is an abelian group

Definition of a module

Let R be an arbitrary ring. A *left R-module* is a set M together with

- a binary operation $+$ such that $(M, +)$ is an abelian group
- an action of R on M satisfying, for all r, s in R and m, n in M:

Definition of a module

Let R be an arbitrary ring. A *left R-module* is a set M together with

- a binary operation $+$ such that $(M, +)$ is an abelian group
- an action of R on M satisfying, for all r, s in R and m, n in M:
    - $(r + s)m = rm + sm$

Definition of a module

Let R be an arbitrary ring. A *left R-module* is a set M together with

- a binary operation $+$ such that $(M, +)$ is an abelian group
- an action of R on M satisfying, for all r, s in R and m, n in M:
  - $(r+s)m = rm + sm$
  - $(rs)m = r(sm)$

Definition of a module

Let R be an arbitrary ring. A *left R-module* is a set M together with

- a binary operation $+$ such that $(M, +)$ is an abelian group
- an action of R on M satisfying, for all r, s in R and m, n in M:
  - $(r + s)m = rm + sm$
  - $(rs)m = r(sm)$
  - $r(m + n) = rm + rn$

Definition of a module

Let R be an arbitrary ring. A *left R-module* is a set M together with

- a binary operation $+$ such that $(M, +)$ is an abelian group
- an action of R on M satisfying, for all r, s in R and m, n in M:
  - $(r + s)m = rm + sm$
  - $(rs)m = r(sm)$
  - $r(m + n) = rm + rn$ if R contains a 1, then $1m = m$

Definition of a module

Let R be an arbitrary ring. A *left R-module* is a set M together with

- a binary operation $+$ such that $(M, +)$ is an abelian group
- an action of R on M satisfying, for all r, s in R and m, n in M:
  - $(r + s)m = rm + sm$
  - $(rs)m = r(sm)$
  - $r(m + n) = rm + rn$ if R contains a 1, then $1m = m$

And, as we more or less hinted at, if R is a field, M is a vector space.

Hand-wavy examples of modules

Hand-wavy examples of modules

- any vector space over a field F is trivially an F-module

Hand-wavy examples of modules

- any vector space over a field F is trivially an F-module
- if $R = \mathbb{Z}$, and A is *any* abelian group, A can be made into a $\mathbb{Z}$-module

Hand-wavy examples of modules

- any vector space over a field F is trivially an F-module
- if $R = \mathbb{Z}$, and A is *any* abelian group, A can be made into a $\mathbb{Z}$-module
- if M is a smooth manifold, the set of all real-valued smooth functions $C^\infty(M)$ is a ring, then the set of all smooth vector fields forms a $C^\infty(M)$-module

Hand-wavy examples of modules

- any vector space over a field F is trivially an F-module
- if $R = \mathbb{Z}$, and A is *any* abelian group, A can be made into a $\mathbb{Z}$-module
- if M is a smooth manifold, the set of all real-valued smooth functions $C^\infty(M)$ is a ring, then the set of all smooth vector fields forms a $C^\infty(M)$-module
- if $R = D$ a ring of differential operators, we can construct these things called D-modules

Hand-wavy examples of modules

- any vector space over a field F is trivially an F-module
- if $R = \mathbb{Z}$, and A is *any* abelian group, A can be made into a $\mathbb{Z}$-module
- if M is a smooth manifold, the set of all real-valued smooth functions $C^\infty(M)$ is a ring, then the set of all smooth vector fields forms a $C^\infty(M)$-module
- if $R = D$ a ring of differential operators, we can construct these things called D-modules

By now students have witnessed on some level a plethora of algebraic structures.

Thank you

Please consider subscribing or following Math/Physics Academy on

- YouTube @MathPhysics.Academy
- Instagram @mathphysics.academy
- TikTok @mathphysics.academy
- General email contact@mathphysics.academy
- For tutoring, email tutoring@mathphysics.academy, or visit us on Facebook