

# Second Year Mathematics Revision

## Calculus - Part 2

Kabir Agrawal, Ethan Brown

UNSW MathSoc

26 Apr 2019



# Table of Contents

- 1 Vector Calculus
  - Vector Fields and their properties
  - Scalar Integrals
  - Vector Integrals
  - Vector Integral Theorems
  - Orthogonal Curvilinear Co-Ordinates
- 2 Fourier Series
  - Preliminaries
  - Fourier Series
  - Function Convergence
- 3 Analysis
  - Types of Sets
  - Limits and Continuity



# Introduction to vector fields

## Definition 1

A **vector field** in 2 or 3-space is a function **F** that assigns a vector to a particular point  $(x, y)$  or  $(x, y, z)$ . The **gradient vector field** is the vector field defined by the function

$$\mathbf{F} = \nabla f = \left( \frac{\partial}{\partial x} f, \frac{\partial}{\partial y} f, \frac{\partial}{\partial z} f \right).$$

These vector fields are typically represented as

$$\mathbf{F}(x, y) = P(x, y)\hat{i} + Q(x, y)\hat{j} \text{ or}$$

$\mathbf{F}(x, y, z) = P(x, y, z)\hat{i} + Q(x, y, z)\hat{j} + R(x, y, z)\hat{k}$ . Here,  $\hat{i}, \hat{j}, \hat{k}$  are the 3 standard basis vectors in the associated space.



# Introduction to Vector fields

## Definition 2

A **conservative vector field** is a vector field  $\mathbf{F}$  which can be expressed as the gradient of a scalar function. That is, there is a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  such that  $\mathbf{F} = \nabla f$ . The corresponding function  $f$  is called the **potential** function.



# Properties of Vector Fields

## Definition 3

The **del/nabla** operator is the vector  $\left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}\right)$ .

The advantage of using this definition is that we now have the ability to use the vector operations as per what we already know, that is, scalar multiplication, dot product, and cross product. For example, taking  $\nabla$  and multiplying it by the scalar function  $f$ , we get the **gradient** vector.



# Divergence

Here, we let  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  to be a continuous, differentiable function in the appropriate vector space.

## Definition 4 (Divergence)

The **divergence** of a vector field describes the rate of change of how much fluid flows out of the neighbourhood of a point. That is, does water have a tendency to flow in or out of the point. The operation is defined as:

$$\operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F}$$

Here,  $n$  could take any value, and the definition will still make sense, but typically we shall only take the case  $n = 3$



# Curl

## Definition 5 (Curl)

The **curl** represents the tendency of water to spin about the point in the neighbourhood of the point. It is computed according to the expression:

$$\text{curl}\mathbf{F} = \nabla \times \mathbf{F}$$

## Restrictions for curl

The curl operator is only defined in  $\mathbb{R}^3$  because it is fundamentally a cross product.



# Examples

## Example 1

Consider the vector field defined by  $\mathbf{F}(x, y, z) = (x, y, z)$ . Evaluate the following:

- 1  $\nabla \cdot \mathbf{F}$
- 2  $\nabla \times \mathbf{F}$
- 3  $\nabla \times (\mathbf{a} \times \mathbf{F})$
- 4  $\nabla \frac{1}{\|\mathbf{F}\|^3}$





# Parameterising curves

The following are typical parameterisation of the curves:

- 1  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  corresponds to  $x = a \cos(t), y = b \sin(t), 0 \leq t \leq 2\pi$  in the anti-clockwise direction.
- 2  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  corresponds to  $x = a \cos(t), y = -b \sin(t), 0 \leq t \leq 2\pi$  in the clockwise direction
- 3 A line segment from  $\mathbf{x}$  to  $\mathbf{y}$  has parameterisation  $(1 - t)\mathbf{x} + t\mathbf{y}$
- 4  $y = f(x)$  corresponds to  $x = t, y = f(t)$
- 5  $x = g(y)$  corresponds to  $y = t, x = g(t)$



# Path Integrals

## Definition 6

A **path integral** is given by the formula:

$$\int_C f(x, y) ds = \int_{t=a}^{t=b} f(h(t), g(t)) \|\mathbf{r}'(t)\| dt$$

where the curve  $C$  has a parameteric vector equation  $\mathbf{r}(t)$



# Examples

## Example 2 [Q151 MATH2011]

- ①  $\int_C (x + y + z) ds$  where  $C$  is the curve parameterised by  $(\cos t, \sin t, t)$ ,  $t \in [0, 2\pi]$
- ②  $\int_C e^{\sqrt{z}} ds$  where  $C$  is the curve parameterised by  $(1, 2, t^2)$ ,  $t \in [0, 1]$



# Surface Integrals

Surfaces are functions in  $\mathbb{R}^3$  that are dependent on 2 variables. Thus in order to parameterise surfaces, we need to use 2 variables and express the 3rd variable in terms of the other 2.

## Definition 7

A **surface integral** is an integral which measures surface area. We denote such an integral by the formula

$$\iint_S f(x, y, z) ds$$

$f(x, y, z)$  represents some kind of density function, and if this density is equal to 1 over the surface  $S$ , then we get the surface area.

# Evaluating Surface Integrals I

To evaluate surface integrals, we need to parameterise them in one of the following ways:

- ①  $\mathbf{s}(x, y) = (x, y, g(x, y))$
- ②  $\mathbf{s}(u, v) = (x(u, v), y(u, v), z(u, v))$ .

In the first scenario, we express the surface as a function of the other 2 variables already existing in the expression [This is common in more elementary expressions for the surface].

In the second scenario, we construct 2 new variables to describe every component in the vector.



# Evaluating Surface Integrals II

In either case, we obtain the following definition:

## Theorem

$$\iint_S f(x, y, z) ds = \iint_{S'} f(x(u, v), y(u, v), z(u, v)) \|\mathbf{s}_u \times \mathbf{s}_v\| du dv$$

To see how this is applied to the parameterisation  $\mathbf{s} = (x, y, g(x, y))$ .

$$\mathbf{s}_x \times \mathbf{s}_y = (1, 0, g_x) \times (0, 1, g_y)$$

$$= \det \begin{bmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 0 & g_x \\ 0 & 1 & g_y \end{bmatrix}$$

$$= -\hat{i}g_x - \hat{j}g_y + \hat{k}$$



# Surface Integrals Examples

## Example 3 [MATH2111 Q11]

Find a parametric representations for the parts of the plane  
 $2x + 3y + z = 4$ :

- ①  $1 \leq x \leq 2, 2 \leq z \leq 4$
- ②  $0 \leq x + y + z \leq 7, 2 \leq x - y \leq 4$
- ③  $x \geq 0, x^2 + y^2 \leq 4$



# Surface Integrals Examples

## Example 4 [MATH2111 Q12]

Let  $\Omega$  denote the conical region  $\sqrt{x^2 + y^2} \leq z \leq 2$ .

- 1 Find a parametric representation  $\mathbf{x}(u, v)$  for the surface, the boundary of  $\Omega$ .
- 2 Use simple geometry to write down the outwards point unit normal vector at each point on  $S$ .
- 3 Verify that these vectors are parallel to the normal vectors obtained from the formula  $N(\mathbf{x}) = \frac{\partial \mathbf{x}}{\partial u} \times \frac{\partial \mathbf{x}}{\partial v}$
- 4 Find the surface area of the cone.





# Vector Integrals

Vector integrals involve the integration of vector fields across paths and surfaces. However, you can only ever integrate scalar functions. So the actual vector integrals are ever so slightly different in their formulas.



# Path integrals of vector fields

## Path Integral of a vector field

Consider a vector field

$\mathbf{F}(x, y, z) = P(x, y, z)\hat{i} + Q(x, y, z)\hat{j} + R(x, y, z)\hat{k}$ , and suppose a curve  $C$  has a parameterisation  $\mathbf{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ . Then we write:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{t=a}^{t=b} \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt = \int_C Pdx + Qdy + Rdz$$

In this case, orientation must be preserved, that is, path integrals of vector fields are dependent on what direction the path goes in.

Typically, the anti-clockwise direction is the default direction, unless otherwise specified.



# Conservative Vector Fields

## Theorem 1 (Fundamental Theorem of Line Integrals)

Suppose that  $C$  is a differentiable curve given by  $\mathbf{r}(t)$ ,  $a \leq t \leq b$ , and  $\mathbf{F} = \nabla f$  [That is,  $\mathbf{F}$  is conservative]. Then the following holds:

$$\int_C \nabla f \cdot d\mathbf{r} = f(\mathbf{r}(b)) - f(\mathbf{r}(a))$$



# Corollaries

## Corollaries

- ①  $\int_C \nabla f \cdot d\mathbf{r}$  is path independent
- ② If  $\mathbf{F}$  is a continuous vector field on an open connected region  $D$  and if  $\int_C \mathbf{F} \cdot d\mathbf{r}$  is path independent for any path in  $D$ , then  $\mathbf{F}$  is conservative.
- ③  $\int_C \mathbf{F} \cdot d\mathbf{r}$  is path independent, if and only if  $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$  for every closed path  $C$



# Examples of vector integrals along paths

## Example 5 [MATH2011 Q155, MATH2111 Q2]

Calculate  $\int_C \mathbf{F} \cdot d\mathbf{r}$  for the vector field

$\mathbf{F}(x, y, z) = (x^2 + y^2 + z^2)\hat{i} - z\hat{j} + (y + 1)\hat{k}$  and the curve  $C$  from  $(0, 0, -1)$  to  $(0, 0, 1)$  is

- 1 The straight line joining the 2 points
- 2 An arc of the circle  $y^2 + z^2 = 1$  in the plane  $x = 0$  oriented counterclockwise when viewed from the positive  $x$ -axis.

If  $\mathbf{F}$  conservative?



# Surface Integrals of vector fields

In a rather obvious sense, we can generalise the path integral into a surface integral, and we can arrive at the following conclusion:

## Definition

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot \hat{\mathbf{n}} dS = \iint_S \mathbf{F} \cdot \mathbf{n} dA$$

where  $A$  is image projected by  $S$  onto the  $xy$  plane.



# Examples

## Example 6 [MATH2111 Q16]

Evaluate  $\int_S \mathbf{F} \cdot d\mathbf{S}$  where  $\mathbf{F} = \sinh(x)\hat{i} + \cosh(y)\hat{k}$  and  $S$  is given by  $z = x + y^2$  for  $0 \leq y \leq x, 0 \leq x \leq 1$ .



# Examples

## Example 7 [MATH2011 Q171 a)]

Evaluate  $\iint_S \mathbf{F} \cdot \hat{n} dS$  where  $\mathbf{F} = y\hat{i} - x\hat{j} + z\hat{k}$  and  $z = 4 - x^2 - y^2, z \geq 0$  and  $\hat{n}$  pointing upwards.





# Integral Theorems

There are 3 major integral theorems:

## Theorems

- 1 Green's Theorem
- 2 Divergence Theorem
- 3 Stokes' Theorem



# Green's Theorem

## Green's Theorem

For a closed region  $D$  in  $\mathbb{R}^2$  we have the following equivalence:

$$\oint_{\partial D} P(x, y)dx + Q(x, y)dy = \iint_D \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dx dy$$

## Conditions

The relevant conditions required for this theorem to hold:

- 1  $D$  is a closed, non-self intersecting region. [Note: If this is the case, split the region into 2, but this won't be covered]
- 2  $P, Q$  are  $C^1(D)$ , meaning they are differentiable functions on  $D$ .

# Examples

## Example 8 [MATH2011 Q161, MATH2111 Q28]

Evaluate  $\oint_C (x^2 - 2xy)dx + (x^2 + 3)dy$  around the boundary  $C$  taken anti-clockwise around the region contained by  $y^2 = 8x$  and  $x = 2$  using Green's Theorem.



# Frame Title

## Example 9 [MATH2111 Q31]

Show that for any planar region  $\Omega$  that:

$$\text{area}(\Omega) = \frac{1}{2} \oint_{\partial\Omega} (x dy - y dx)$$



# Divergence Theorem

## Divergence Theorem

For a  $C^1(R)$  vector field  $\mathbf{F}$  on a closed and bounded solid  $V$ . Then the following holds:

$$\iiint_V \nabla \cdot \mathbf{F} dV = \iint_{\partial V} \mathbf{F} \cdot d\mathbf{r}$$



# Examples for Divergence Theorem I

## Example 10 [MATH2111 Past Paper]

Let  $\mathbf{F} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a vector defined by  $\mathbf{F} = xy\hat{i} + z^3\hat{j} + y^2\hat{k}$  and  $W$  the box defined by  $[0, 1] \times [0, 2] \times [0, 4]$  and the orientation defined outward. Use divergence theorem to evaluate:

$$\iint_{\partial W} \mathbf{F} \cdot d\mathbf{S}$$



# Examples for Divergence Theorem II

## Example 11 [MATH2011 Q180]

Evaluate  $\iint_S \mathbf{F} \cdot \hat{\mathbf{n}} dS$  for  $\mathbf{F} = (x, -y, z^2 - 1)$  and  $S$  is the boundary surface of the solid  $\Omega$  bounded by the planes  $z = 0, z = 1, x^2 + y^2 = 1$



# Stokes' Theorem

## Stokes' Theorem

Consider a closed and bounded surface  $S$  and a  $C^1(S)$  vector field  $\mathbf{F}$ . Then we have:

$$\iint_S \nabla \times \mathbf{F} \cdot d\mathbf{s} = \int_{\partial S} \mathbf{F} \cdot d\mathbf{r}$$





# Examples for Stokes' Theorem

## Example 12 [MATH2011 Q175]

Verify Stokes' Theorem for the vector field

$\mathbf{F} = (2x - y, -yx^2, -y^2z)$  and  $S$  is the top half-surface of the sphere  $x^2 + y^2 + z^2 = 1, z \geq 0$  and  $C$  is the bounding curve of  $S$ .



# Examples for Stokes' Theorem

## Example 13 [MATH2011 Q178]

Calculate  $\int_C \mathbf{F} \cdot d\mathbf{s}$  where  $\mathbf{F} = (4z + x^2, -2x + 3y^5, 2x^2 + 5 \sin(z))$  and  $C$  is the curve of intersection of the surfaces  $x^2 + y^2 = 1, z = y + 1$



# Orthogonal Curvilinear Co-ordinates

Orthogonal curvilinear co-ordinates are a form of substitution that enables conversion into a more "convenient" form for the co-ordinate system. In effect, it is a parameterisation.

## Definition

We say that  $(\xi_1, \xi_2, \xi_3)$  are orthogonal co-ordinates if the following hold true:

$$\mathbf{b}_{\xi_i} \cdot \mathbf{b}_{\xi_j} = 0$$

where  $\mathbf{b}_{\xi_i} = \frac{d}{d\xi_i} \mathbf{x}$



# Example

## Example 14 [MATH2111 Q10]

Toroidal coordinates  $(w, \theta, \psi)$  are defined by  
 $x = (a + w \cos(\psi)) \cos(\theta)$ ,  $y = (a + w \cos(\psi)) \sin(\theta)$ ,  $z = w \sin(\psi)$ ,  
with  $a$  being a constant such that  $0 < w < a$ .

- ① Verify that  $(w, \theta, \psi)$  is a right-handed, orthogonal coordinate system. Find the scale factors and orthonormal basis vectors.
- ② Describe the curve  $C_1$  where  $\theta = \frac{\pi}{2}$ ,  $w = b$  if  $b$  is a constant such that  $0 < b < a$ .
- ③ Describe the curve  $C_2$  where  $\theta = \psi$ ,  $w = b$ ,  $0 < b < a$  and  $b$  is a constant.
- ④ Calculate the length of the curves  $C_1, C_2$  for  $a = 1, b = \frac{1}{2}$

# Table of Contents

- 1 Vector Calculus
  - Vector Fields and their properties
  - Scalar Integrals
  - Vector Integrals
  - Vector Integral Theorems
  - Orthogonal Curvilinear Co-Ordinates
- 2 Fourier Series
  - Preliminaries
  - Fourier Series
  - Function Convergence
- 3 Analysis
  - Types of Sets
  - Limits and Continuity



# Periodic Functions

Fourier Series deals with periodic functions, so we define a few terms.

## Definition 1

A function  $f : D \rightarrow Y$  is  **$L$ -periodic** if, for all  $x \in D$ ,

$$f(x + L) = f(x).$$

Further,  $f$  has **frequency** or **period**  $L$ , and the smallest  $K$  such that  $f$  is  $K$ -periodic is called the **fundamental frequency** of  $f$ .



# Real Trigonometric Polynomials

To define Fourier series, we first define trigonometric polynomials.

## Definition 2

A **real trigonometric polynomial** of degree  $n$  and period  $2L$  is a function of the form

$$f(x) = \frac{a_0}{2} + \sum_{k=1}^n \left( a_k \cos \frac{k\pi x}{L} + b_k \sin \frac{k\pi x}{L} \right),$$

where  $a_n \neq 0$  or  $b_n \neq 0$ , and each  $a_i, b_i \in \mathbb{R}$ .

Dividing  $a_0$  by 2 will simplify things later.



# Complex Trigonometric Polynomials (MATH2111)

Since  $e^{ik\pi x/L} = \cos \frac{k\pi x}{L} + i \sin \frac{k\pi x}{L}$ , we can extend real trigonometric polynomials as follows.

## Definition 3

A **complex trigonometric polynomial** of degree  $n$  and period  $2L$  is a function of the form

$$f(x) = \sum_{k=-n}^n c_k e^{ik\pi x/L} = c_0 + \sum_{k=1}^n \left( c_k e^{ik\pi x/L} + c_{-k} e^{-ik\pi x/L} \right),$$

where  $c_n \neq 0$  or  $c_{-n} \neq 0$ , and each  $c_i \in \mathbb{C}$ .

If each  $c_i = \overline{c_{-i}}$ , then each term is real, and the function is a real trigonometric polynomial.





# Fourier Series

Now we define Fourier Series.

## Definition 4

Suppose  $f : \mathbb{R} \rightarrow \mathbb{R}$  is bounded, piecewise continuous, and  $2L$ -periodic. Then the  $n^{\text{th}}$  **real Fourier polynomial** of  $f$  is

$$(S_n f)(x) = \frac{a_0[f]}{2} + \sum_{k=1}^n \left( a_k[f] \cos \frac{k\pi x}{L} + b_k[f] \sin \frac{k\pi x}{L} \right),$$

where

$$a_0[f] = \frac{1}{L} \int_{-L}^L f(x) dx,$$

and for  $1 \leq k \leq n$ ,

$$a_k[f] = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{k\pi x}{L} dx, \quad b_k[f] = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{k\pi x}{L} dx.$$

# Fourier Series (MATH2111)

## Definition 5

Suppose  $f : \mathbb{R} \rightarrow \mathbb{C}$  is bounded, piecewise continuous, and  $2L$ -periodic. Then the  $n^{\text{th}}$  **complex Fourier polynomial** of  $f$  is

$$\begin{aligned}(S_n f)(x) &= \sum_{k=-n}^n c_k[f] e^{ik\pi x/L} \\ &= c_0[f] + \sum_{k=1}^n \left( c_k[f] e^{ik\pi x/L} + c_{-k}[f] e^{-ik\pi x/L} \right),\end{aligned}$$

where

$$c_k[f] = \frac{1}{2L} \int_{-L}^L f(x) e^{-ik\pi x/L} dx$$

for  $-n \leq k \leq n$ .

# Fourier Series

A couple of results help us when dealing with even or odd functions.

## Theorem 1

If  $f$  is an even function, then

$$a_0[f] = \frac{2}{L} \int_0^L f(x) dx, \quad a_k[f] = \frac{2}{L} \int_0^L f(x) \cos \frac{k\pi x}{L} dx,$$

and  $b_k[f] = 0$ .

If  $f$  is an odd function, then

$$b_k[f] = \frac{2}{L} \int_0^L f(x) \sin \frac{k\pi x}{L} dx,$$

and  $a_0[f] = a_k[f] = 0$ .

# Fourier Series

A **Fourier series** is the limit of the sequence  $\{S_n f\}_{n=1}^{\infty}$  denoted  $Sf$ .

## Example 1

Find the real Fourier series of  $f$  where  $f(x) = x$  for  $-1 < x \leq 1$ , and requiring that  $f(x+2) = f(x)$  for all  $x \in \mathbb{R}$ .

So  $f$  is 2-periodic, and odd. Thus  $a_0[f] = a_k[f] = 0$ . Then


$$b_k[f] = 2 \int_0^1 x \sin k\pi x dx = 2 \frac{\sin \pi k - \pi k \cos \pi k}{\pi^2 k^2}.$$

Now,  $\sin \pi k = 0$  and  $\cos \pi k = (-1)^k$  as  $k$  is an integer, so

$$(Sf)(x) = \frac{2}{\pi} \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \sin k\pi x.$$



# Gibbs Phenomenon



# Fourier Series

At jump discontinuities, the Fourier series approaches the average of the function value either side of the discontinuity.

## Theorem 2

Suppose  $f$  has a jump discontinuity at  $a$ . That is, the limits

$$f(a^+) = \lim_{x \rightarrow a^+} f(x), \quad f(a^-) = \lim_{x \rightarrow a^-} f(x)$$

both exist, but  $f(a^+) \neq f(a^-)$ . Then

$$\lim_{n \rightarrow \infty} (S_n f)(a) = \frac{f(a^+) + f(a^-)}{2}.$$


For those in MATH2111, this is conditioned on the one-sided derivatives existing at  $a$ . For the functions you deal with, this will almost always be the case.



# Fourier Series

## Example 2

For the previous example, draw the graph of  $(Sf)(x)$  for  $-2 \leq x \leq 1$



Wherever the function is continuous, we can simply draw  $f$ . At each jump discontinuity, the Fourier series approaches the average of the function, in this case 0.



# Function Convergence

We can say the Fourier series “converges” to the function, but to do so, we need to define what we mean by convergence.

## Definition 6

A sequence of functions  $\{f_i\}_{i=1}^{\infty}$  **converges pointwise** to  $f$  on an interval  $I$  if, for every  $x \in I$ , we have

$$f(x) = \lim_{n \rightarrow \infty} f_n(x).$$

## Definition 7

A sequence of functions  $\{f_i\}_{i=1}^{\infty}$  **converges uniformly** to  $f$  on an interval  $I$  if, for every  $\epsilon > 0$ , there exists an  $N > 0$  such that

$$|f_n(x) - f(x)| < \epsilon$$

for all  $n \geq N$  and  $x \in I$ .





# Function Convergence

Uniform convergence implies pointwise convergence, however the converse is not true.

## Example 3

Prove that  $f_n(x) = x^n$  converges pointwise to  $f(x) = 0$  on  $[0, 1)$ , but not uniformly.

If  $x \in [0, 1)$ , then  $\lim_{n \rightarrow \infty} x^n = 0$ , so  $f_n$  converges pointwise.

Now take  $\epsilon = \frac{1}{2}$ . Then if the sequence converges uniformly we have

$$|f_n(x) - f(x)| = |x^n| = x^n < \frac{1}{2}$$

for sufficiently large  $n$ , say  $n \geq N > 0$ , and all  $x \in [0, 1)$ . So,

$$x < \frac{1}{\sqrt[N]{2}} < 1.$$

Take  $x \in (1/\sqrt[N]{2}, 1)$  to derive a contradiction.



# Function Convergence

## Theorem 3

If a sequence of continuous functions  $\{f_i\}_{i=1}^{\infty}$  converges uniformly to  $f$  on an interval  $I$ , then  $f$  is continuous on  $I$ .

This means we can disprove uniform convergence by showing that the sequence converges pointwise to a discontinuous limit.



# Series Convergence

Just as with sequences, we can define convergence for series of functions similarly.

## Definition 8

Let  $\{f_i\}_{i=1}^{\infty}$  be a sequence of functions. The infinite series

$$\sum_{k=1}^{\infty} f_k(x)$$

**converges pointwise** to  $S$  on an interval  $I$  if the partial sums converge pointwise to  $S$  on  $I$ . That is, for every  $x \in I$ ,

$$S(x) = \lim_{n \rightarrow \infty} \sum_{k=1}^n f_k(x).$$

# Series Convergence

## Definition 9

Let  $\{f_i\}_{i=1}^{\infty}$  be a sequence of functions. The infinite series

$$\sum_{k=1}^{\infty} f_k(x)$$

**converges uniformly** to  $S$  on an interval  $I$  if the partial sums converge uniformly to  $S$  on  $I$ . That is, for every  $\epsilon > 0$ , there exists an  $N > 0$  such that

$$|S_n(x) - S(x)| < \epsilon$$

for all  $n \geq N$  and  $x \in I$ , and where  $S_n$  is the  $n^{\text{th}}$  partial sum.

# Series Convergence

## Theorem 4 (Weierstrass M-test)

If the sequence of functions  $\{f_i\}_{i=1}^{\infty}$  satisfies

$$|f_n(x)| \leq M_n$$

for all  $x \in I$  and  $n \geq 1$ , and if

$$\sum_{n=1}^{\infty} M_n$$

converges, then

$$\sum_{n=1}^{\infty} f_n(x)$$

converges uniformly on  $I$ .

# Series Convergence

## Example 4 (Question 10 from MATH2111 2018)

Prove that

$$\sum_{n=1}^{\infty} \frac{\cos nx}{n^2 + x^2}$$

converges uniformly on  $\mathbb{R}$ .

$$\left| \frac{\cos nx}{n^2 + x^2} \right| \leq n^{-2}.$$

So, since

$$\sum_{n=1}^{\infty} n^{-2} = \frac{\pi^2}{6},$$

by Weierstrass M-test, the series converges uniformly on  $\mathbb{R}$ .



# Convergence of Fourier Series

With these definitions, we can say the following.

## Theorem 5

Suppose  $c_k$  are the complex, and  $a_k, b_k$  the real Fourier coefficients of  $f$ . Then if

$$\sum_{k=-\infty}^{\infty} |c_k|, \text{ or } \sum_{k=1}^{\infty} (|a_k| + |b_k|)$$

converge, then the corresponding Fourier series converges uniformly to  $f$ . Further,  $f$  is continuous.

So, if the function we are trying to represent as a Fourier series is discontinuous, then the series does not converge uniformly.



# Convergence of Fourier Series

## Example 5

Let  $f$  be defined such that  $f(x) = x$  for  $0 \leq x \leq 1$ ,  $f$  is even, and  $f(x+2) = f(x)$  for all  $x \in \mathbb{R}$ . You are given that

$$(Sf)(x) = \frac{1}{2} - \frac{4}{\pi^2} \sum_{k \text{ odd}} \frac{\cos k\pi x}{k^2}.$$

Does  $Sf$  converge uniformly on  $\mathbb{R}$ ? What does this say about  $f$ ?

So

$$\sum_{k=1}^{\infty} \left( \left| \frac{-4}{k^2 \pi^2} \right| + |0| \right) = \frac{4}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^2}$$

converges by p-test. Thus,  $Sf$  converges uniformly to  $f$ . Since the partial sums are continuous, uniform convergence implies that  $f$  is continuous.





# Table of Contents

- 1 Vector Calculus
  - Vector Fields and their properties
  - Scalar Integrals
  - Vector Integrals
  - Vector Integral Theorems
  - Orthogonal Curvilinear Co-Ordinates
- 2 Fourier Series
  - Preliminaries
  - Fourier Series
  - Function Convergence
- 3 Analysis
  - Types of Sets
  - Limits and Continuity



# Balls

To generalise results about open and closed intervals, we first define a ball in  $\mathbb{R}^n$ .

## Definition 10

A **ball** around  $\mathbf{a} \in \mathbb{R}^n$  of radius  $\epsilon > 0$  is

$$B(\mathbf{a}, \epsilon) = \{\mathbf{x} \in \mathbb{R}^n : d(\mathbf{a}, \mathbf{x}) < \epsilon\}.$$

Here,  $d(\mathbf{a}, \mathbf{x})$  is the Euclidean distance between  $\mathbf{a}$  and  $\mathbf{x}$ .

In  $\mathbb{R}^2$  and  $\mathbb{R}^3$  this is a disk and sphere, **without** its boundary.



# Types of Points

## Definition 11

Consider  $\Omega \subseteq \mathbb{R}^n$ . Then  $\mathbf{a} \in \mathbb{R}^n$  is

- An **interior point** of  $\Omega$  if there is a ball around  $\mathbf{a}$  contained entirely in  $\Omega$ .
- A **boundary point** of  $\Omega$  if every ball around  $\mathbf{a}$  contains both points in  $\Omega$  and not in  $\Omega$ .
- A **limit point** or **accumulation point** of  $\Omega$  if there is a sequence  $\{\mathbf{x}_n\}_{n=1}^{\infty} \subseteq \Omega$  with limit  $\mathbf{a}$ , and  $\mathbf{x}_i \neq \mathbf{a}$  for all  $\mathbf{x}_i$ .

Every interior point is also a limit point, but the converse is not necessarily true.



# Types of Points

## Example 6

Classify the points  $(0, 0)$ , and  $(1, 1)$  of the set

$$\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < 1\} \cup \{(1, 1)\}.$$

There is a ball around  $(0, 0)$  that is contained entirely in  $\Omega$ , so it is an interior point. There is a sequence contained in  $\Omega$  with limit  $(0, 0)$ , so it is a limit point. It is not a boundary point.

No ball around  $(1, 1)$  is contained entirely in  $\Omega$ , so it's not an interior point. There is also no sequence in  $\Omega$  with limit  $(1, 1)$  that isn't eventually constant, so it's not a limit point. It is a boundary point, as any ball around  $(1, 1)$  contains an element in  $\Omega$  (namely  $(1, 1)$ ), and not in  $\Omega$  (namely  $(1, 1 + \epsilon)$  for sufficiently small  $\epsilon$ ).



# Types of Sets

Now, we use these point definitions to define types of sets.

## Definition 12

A set  $\Omega \subseteq \mathbb{R}^n$  is

- **Open** if every element  $\mathbf{a} \in \Omega$  is an interior point.
- **Closed** if its complement  $\mathbb{R}^n - \Omega$  is open.

In  $\mathbb{R}^n$ ,  $\mathbb{R}^n$  and  $\emptyset$  are the only sets which are both open **and** closed.

## Theorem 6

The following definitions are equivalent:

- $\Omega$  is closed.
- $\Omega$  contains all of its boundary points.
- $\Omega$  contains all of its limit points.

# Types of Sets

## Example 7

Prove that  $\Omega = (a, b)$  is an open set in  $\mathbb{R}$  where  $a < b$ .

Suppose  $x \in \Omega$ . Then let  $\epsilon = \min \{x - a, b - x\}$ . Now, if  $c \in (x - \epsilon, x + \epsilon)$ , then

$$c > x - \epsilon \geq x - (x - a) = a.$$

Similarly,

$$c < x + \epsilon \leq x + (b - x) = b,$$

so  $c \in \Omega$ . That is,  $B(x, \epsilon) = (x - \epsilon, x + \epsilon) \subseteq (a, b)$ . Then every element  $x \in \Omega$  has a ball around it contained entirely in  $\Omega$ , and thus is an interior point. Thus,  $\Omega$  is open.



# Types of Sets

## Example 8

Is the set  $\Omega = B(\mathbf{0}, 1) \cup \{(x, y) \in \mathbb{R}^2 : x \geq 0\}$  open? Is it closed?

$\Omega$  is not open, as  $(0, 2) \in \Omega$  but  $(-\epsilon, 2) \notin \Omega$  for any  $\epsilon > 0$ , so any ball around  $(0, 2)$  does not lie entirely in  $\Omega$ . That is, not every point in  $\Omega$  is an interior point.

Any ball around  $(-1, 0)$  contains  $(-1, 0) \notin \Omega$  and  $(\epsilon - 1, 0) \in \Omega$  for some  $\epsilon > 0$ . Thus  $(-1, 0)$  is a boundary point not contained in  $\Omega$ , and so  $\Omega$  is not closed.



# Properties of Sets

## Definition 13

Suppose  $\Omega \subseteq \mathbb{R}^n$ . Then

- The **interior** of  $\Omega$  is the set of all interior points of  $\Omega$ .
- The **boundary** of  $\Omega$  is the set of all boundary points of  $\Omega$  denoted  $\partial\Omega$ .
- The **closure** of  $\Omega$  is  $\overline{\Omega} = \Omega \cup \partial\Omega$ .

The interior is the largest open subset of  $\Omega$ , and the closure is the smallest closed set containing  $\Omega$ .





# Limits in $\mathbb{R}^n$

A precise definition of limits in  $\mathbb{R}^n$  is as follows (similar to  $\epsilon - \delta$  in first year).

## Definition 14

Suppose  $\mathbf{f} : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $\mathbf{a} \in \Omega$ . We say

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} \mathbf{f}(\mathbf{x}) = \mathbf{b}$$

if for all  $\epsilon > 0$ , there exists  $\delta > 0$  such that if  $\mathbf{x} \in \Omega$  and  $0 < d(\mathbf{x}, \mathbf{a}) < \delta$ , then  $d(\mathbf{f}(\mathbf{x}), \mathbf{b}) < \epsilon$ . Alternatively,

$$\mathbf{x} \in \Omega \cap B(\mathbf{a}, \delta) - \{\mathbf{a}\} \implies \mathbf{f}(\mathbf{x}) \in B(\mathbf{b}, \epsilon).$$

# Limits in $\mathbb{R}^n$

## Example 9 (Question 33ii from MATH2111 2019)

Show that  $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2 + x^2 y^2}{x^2 + y^2} = 1$  using the definition of a limit.

Note that  $|f(x, y) - 1|$  is:

$$\left| \frac{x^2 + y^2 + x^2 y^2}{x^2 + y^2} - 1 \right| = \left| \frac{x^2 y^2}{x^2 + y^2} \right| \leq \frac{x^4 + 2x^2 y^2 + y^4}{x^2 + y^2} = x^2 + y^2.$$

So for some  $\epsilon > 0$  choose  $\delta = \sqrt{\epsilon}$ . Then if  $\sqrt{x^2 + y^2} < \delta$ , we have  $|f(x, y) - 1| \leq x^2 + y^2 < \delta^2 = \epsilon$ , as required.



# Limits in $\mathbb{R}^n$

Dealing with limits of vector functions can be difficult, so instead it suffices to consider each component's limit individually.

## Theorem 7

Suppose  $\mathbf{f} : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$  and denote the components of  $\mathbf{f}$   $f_1, f_2, \dots, f_m$ . Let  $\mathbf{a} \in \Omega$ . If

$$L_1 = \lim_{\mathbf{x} \rightarrow \mathbf{a}} f_1(\mathbf{x}), \quad L_2 = \lim_{\mathbf{x} \rightarrow \mathbf{a}} f_2(\mathbf{x}), \quad \dots, \quad L_m = \lim_{\mathbf{x} \rightarrow \mathbf{a}} f_m(\mathbf{x})$$

all exist, then

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} \mathbf{f}(\mathbf{x}) = (L_1, L_2, \dots, L_m)^T.$$

This can be used in conjunction with other methods on scalar functions to evaluate limits of vector functions.



# Pinching Principle

Just as single variable limits can be evaluated using the pinching principle, so can multivariable limits.

## Theorem 8 (Pinching Principle)

Suppose  $f, g, h : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $\mathbf{a} \in \Omega$ , and

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} f(\mathbf{x}) = \lim_{\mathbf{x} \rightarrow \mathbf{a}} h(\mathbf{x}) = L.$$

If  $f(\mathbf{x}) \leq g(\mathbf{x}) \leq h(\mathbf{x})$  for all  $\mathbf{x} \in B(\mathbf{a}, \epsilon)$  where  $\epsilon > 0$ . That is, all  $\mathbf{x}$  in a neighbourhood of  $\mathbf{a}$ . Then

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} g(\mathbf{x}) = L.$$

This can also be called the pinching theorem.



# Pinching Principle

## Example 10

Find  $\lim_{(x,y) \rightarrow (0,0)} \left( \frac{x^2 + y^2 + x^2 y^2}{x^2 + y^2}, \frac{xy(x+y)}{x^2 - xy + y^2} \right)^T$ .

So considering each component separately, we have

$$1 \leq \frac{x^2 + y^2 + x^2 y^2}{x^2 + y^2} \leq 1 + \frac{x^4 + 2x^2 y^2 + y^4}{x^2 + y^2} = 1 + x^2 + y^2,$$
$$0 \leq \left| \frac{xy(x+y)}{x^2 - xy + y^2} \right| \leq |x + y|.$$

The last inequality follows from expanding  $(|x| - |y|)^2 \geq 0$  and noting that  $x^2 + y^2 - xy \geq x^2 + y^2 - |xy|$ .

So, by pinching theorem, the limit is  $(1, 0)^T$ .



# Continuity in $\mathbb{R}^n$

Now that we have defined limits, we can define continuity.

## Definition 15

$\mathbf{f} : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$  is **continuous** at  $\mathbf{a} \in \Omega$  if

- $\mathbf{a}$  is a limit point of  $\Omega$ ,  $\lim_{\mathbf{x} \rightarrow \mathbf{a}} \mathbf{f}(\mathbf{x})$  exists and equals  $\mathbf{f}(\mathbf{a})$ , or
- $\mathbf{a}$  is not a limit point of  $\Omega$ .

## Theorem 9

Suppose  $\mathbf{f} : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ . Then the following are equivalent.

- $\mathbf{f}$  is continuous at  $\mathbf{a} \in \Omega$ .
- For all  $\epsilon > 0$ , there exists some  $\delta > 0$  such that if  $\mathbf{x} \in \Omega \cap B(\mathbf{a}, \delta) - \{\mathbf{a}\}$  then  $\mathbf{f}(\mathbf{x}) \in B(\mathbf{f}(\mathbf{a}), \epsilon)$ .
- For every  $\{\mathbf{x}_i\}_{i=1}^{\infty} \subseteq \Omega$  with limit  $\mathbf{a}$ ,  $\{\mathbf{f}(\mathbf{x}_i)\}_{i=1}^{\infty}$  has limit  $\mathbf{f}(\mathbf{a})$ .
- $\mathbf{f}(\mathbf{a})$  is in the interior of  $\mathbf{f}(\Omega)$  implies  $\mathbf{a}$  is in the interior of  $\Omega$ .

# Continuity in $\mathbb{R}^n$

## Example 11

Consider  $\mathbf{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  where  $\mathbf{f}(\mathbf{0}) = \mathbf{0}$  and otherwise

$$\mathbf{f}(x, y) = \left( \frac{x^3}{x^2 + y^2}, \frac{y^3}{x^2 + y^2} \right)^T.$$

Prove  $\mathbf{f}$  is continuous on  $\mathbb{R}^2$ .

Since  $\mathbb{R}^2$  is open, every  $\mathbf{a} = (a, b) \in \mathbb{R}^2$  is a limit point. When  $\mathbf{a} \neq \mathbf{0}$ ,  $\lim_{\mathbf{x} \rightarrow \mathbf{a}} \mathbf{f}(\mathbf{x})$  exists and is equal to  $\mathbf{f}(\mathbf{a})$ . When  $\mathbf{a} = \mathbf{0}$  we have

$$\lim_{\mathbf{x} \rightarrow \mathbf{0}} \frac{x^3}{x^2 + y^2} = \lim_{\mathbf{x} \rightarrow \mathbf{0}} \frac{y^3}{x^2 + y^2} = 0.$$

So,  $\lim_{\mathbf{x} \rightarrow \mathbf{0}} \mathbf{f}(\mathbf{x}) = \mathbf{0} = \mathbf{f}(\mathbf{0})$ , and thus  $\mathbf{f}$  is continuous on  $\mathbb{R}^2$ .



# Continuity in $\mathbb{R}^n$

Just like limits, we can determine continuity of a vector function by its components.

## Theorem 10

Suppose  $\mathbf{f} : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ .  $\mathbf{f}$  is continuous at  $\mathbf{a} \in \Omega$  **if and only if** all components of  $\mathbf{f}$  are continuous at  $\mathbf{a}$ .

Continuous functions preserve openness of sets in their preimage.

## Theorem 11 (Preimage Theorems)

Suppose  $\mathbf{f} : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ . Then the following are equivalent.

- $\mathbf{f}$  is continuous on  $\Omega$ .
- $U \subseteq \mathbb{R}^m$  is open implies  $\mathbf{f}^{-1}(U)$  is open.
- $U \subseteq \mathbb{R}^m$  is closed implies  $\mathbf{f}^{-1}(U)$  is closed.

Here,  $\mathbf{f}^{-1}(U) = \{\mathbf{x} \in \Omega : \mathbf{f}(\mathbf{x}) \in U\}$ .



# More Types of Sets

Now we define a few more types of sets.

## Definition 16

A set  $\Omega \subseteq \mathbb{R}^n$  is

- **Bounded** if there is some finite  $\epsilon > 0$  such that  $\Omega \subseteq B(\mathbf{0}, \epsilon)$ .
- **Path connected** if, for all  $\mathbf{x}, \mathbf{y} \in \Omega$ , there is a continuous function  $\phi : [0, 1] \rightarrow \mathbb{R}^n$  such that  $\phi(0) = \mathbf{x}$ ,  $\phi(1) = \mathbf{y}$ , and  $\phi([0, 1]) \subseteq \Omega$ . That is, there is a continuous path between all  $\mathbf{x}$  and  $\mathbf{y}$  contained entirely within  $\Omega$ .
- **Compact** if  $\Omega$  is closed and bounded.
- **Simply connected** if any closed curve in  $\Omega$  can be continuously deformed to a point without leaving  $\Omega$ .

# More Types of Sets

## Example 12

Let  $\Omega = B(\mathbf{0}, 2) - B(\mathbf{0}, 1)$ . Is  $\Omega$  path connected? Is it compact?

Let  $\mathbf{x}_1 = (r_1 \cos \theta_1, r_1 \sin \theta_1)^T$ ,  $\mathbf{x}_2 = (r_2 \cos \theta_2, r_2 \sin \theta_2)^T \in \Omega$ .

Consider

$$\phi(t) = (r(t) \cos \theta(t), r(t) \sin \theta(t))^T,$$

where  $r(t) = r_1 + (r_2 - r_1)t$  and  $\theta(t) = \theta_1 + (\theta_2 - \theta_1)t$ . Then  $r(t)$  lies between  $r_1$  and  $r_2$  for  $t \in [0, 1]$  and thus  $\phi([0, 1]) \subseteq \Omega$ . Further,  $\phi(0) = \mathbf{x}_1$  and  $\phi(1) = \mathbf{x}_2$ . Since the components of  $\phi$  are continuous,  $\phi$  is continuous, and so  $\Omega$  is path connected. Clearly  $\Omega \subseteq B(\mathbf{0}, 2)$ , so  $\Omega$  is bounded. However, it is not closed, so it isn't compact.



# Monotone Convergence Theorem

In  $\mathbb{R}$ , a sequence that is bounded and monotone has a limit in  $\mathbb{R}$ . We can say a similar thing for  $\mathbb{R}^n$ .

## Theorem 12 (Bolzano-Weierstrass Theorem)

Suppose  $\Omega \subseteq \mathbb{R}^n$ . Then the following are equivalent.

- $\Omega$  is closed and bounded (compact).
- Every sequence in  $\Omega$  has a subsequence that converges to an element of  $\Omega$ .

Since there's no real concept of monotone, we say that there simply is a subsequence that converges in  $\Omega$ .



# Bolzano-Weierstrass Theorem

## Example 13

Let  $\Omega = \{(x, y) \in \mathbb{R}^2 : -1 \leq x, y \leq 1\}$ . Does the sequence  $\{(\cos \frac{i\pi}{2}, \sin \frac{i\pi}{2})\}_{i=1}^{\infty}$  converge in  $\Omega$ ? Does it have a subsequence that converges in  $\Omega$ ?

The sequence does not converge in  $\Omega$ . It cycles through four points. However,  $\Omega$  is bounded and closed, so any sequence in  $\Omega$  has a subsequence that does converge. One such subsequence is the constant sequence  $\{(1, 0)\}_{i=1}^{\infty}$ .



# Image Theorems

Compactness and path connectedness are preserved by continuous functions.

## Theorem 13 (Image Theorems)

Suppose  $\mathbf{f} : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a continuous function. Then

- If  $K \subseteq \Omega$  is compact, then  $\mathbf{f}(K)$  is compact.
- If  $B \subseteq \Omega$  is path connected, then  $\mathbf{f}(B)$  is path connected.



# Image Theorems

## Example 14 (Question 65 from MATH2111 2019)

Let

$$S_1 = \{(x, y) \in \mathbb{R}^2 : 0 \leq x, y \leq 1\},$$

$$S_2 = \{(x, y) \in \mathbb{R}^2 : x, y \geq 0\}.$$

Is there a continuous function that maps  $S_1$  onto  $S_2$ ?  $S_2$  onto  $S_1$ ?

$S_1$  is compact, however  $S_2$  is not, so there is no continuous map from  $S_1$  onto  $S_2$ .

$S_2$  is path connected, but not compact, and  $S_1$  is path connected, so there may be a continuous function from  $S_2$  onto  $S_1$ .

$$\mathbf{f}(x, y) = (|\sin x|, |\sin y|)^T$$

is such a function.

