



MATH1231/1241 MathSoc Calculus Revision Session 2019 T1 Solutions

August 8, 2019

These answers were written and typed by Abdellah Islam. Please be ethical with this resource. The questions here were discussed in MathSoc's 2019 Term 2 MATH1231/1241 Final Exam Revision Session and several were taken or adapted from UNSW past exam papers and homework sheets, and all copyright of the original questions belongs to UNSW's School of Mathematics and Statistics. It is for the use of MathSoc members, so do not repost it on other forums or groups without asking for permission. If you appreciate this resource, please consider supporting us by coming to our events! Also, happy studying :)

We cannot guarantee that our answers are correct - please notify us of any errors or typos at unswmathsoc@gmail.com, or on our Facebook page. There are sometimes multiple methods of solving the same question. Remember that in the real class test, you will be expected to explain your steps and working out.

Example 1: 1231 2015 Q1.v

Prove that

$$S = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \in \mathbb{R}^3 : x_1 - 2x_2 + 4x_3 = 0 \right\}$$

is a subspace of \mathbb{R}^3 .

Clearly, $S \subseteq \mathbb{R}^3$ where \mathbb{R}^3 is a known vector space. Since $0 - 2(0) + 4(0) = 0$, $\mathbf{0} \in S$. So S contains a zero element.

Now suppose that $\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \in S$ and $\mathbf{y} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \in S$. Then

$$x_1 - 2x_2 + 4x_3 = 0, \quad (1)$$

$$y_1 - 2y_2 + 4y_3 = 0. \quad (2)$$

(1) + (2) gives us $(x_1 + y_1) - 2(x_2 + y_2) + 4(x_3 + y_3) = 0$, i.e. $\begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{pmatrix} \in S$. Hence $(\mathbf{x} + \mathbf{y}) \in S$

and so S is closed under vector addition.

If $\lambda \in \mathbb{R}$ and $\mathbf{x} \in S$ then $\lambda \times (1)$ gives us $\lambda x_1 - 2\lambda x_2 + 4\lambda x_3 = 0$, i.e. $\begin{pmatrix} \lambda x_1 \\ \lambda x_2 \\ \lambda x_3 \end{pmatrix} \in S$. Hence $\lambda \mathbf{x} \in S$

so S is closed under scalar multiplication.

Since S is a subset of \mathbb{R}^3 and contains a zero element, is closed under vector addition, and is closed under scalar multiplication, then by the Subspace Theorem S is a subspace of \mathbb{R}^3 .

Example 2: 1231 2013 Q1.i

Let

$$S = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \in \mathbb{R}^3 : x_1^3 + x_2^3 + x_3^3 = 0 \right\}.$$

a) Prove that S is closed under scalar multiplication.

b) Show that S is **not** a subspace of \mathbb{R}^3 .

Suppose that $\mathbf{x} \in S$ and $\lambda \in \mathbb{R}$. Then we have

$$x_1^3 + x_2^3 + x_3^3 = 0.$$

Multiplying by λ^3 ,

$$(\lambda x_1)^3 + (\lambda x_2)^3 + (\lambda x_3)^3 = 0.$$

Hence $\lambda \mathbf{x} \in S$, i.e. S is closed under scalar multiplication.

Note that $\mathbf{0}$ is an element of S , so to prove that S is not a subspace we will show that S is not closed under vector addition. Take $\mathbf{x} = (1, -1, 0)^T$ and $\mathbf{y} = (-2, 0, 2)^T$. Clearly $\mathbf{x}, \mathbf{y} \in S$, but

$$\mathbf{x} + \mathbf{y} = \begin{pmatrix} -1 \\ -1 \\ 2 \end{pmatrix} \notin S.$$

Hence S is not closed under vector addition. By the Subspace Theorem, S is not a subspace of \mathbb{R}^3 .

Example 3: 1231 2015 Q1.vi

Consider the vectors in \mathbb{R}^3 ,

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 2 \\ -3 \\ 3 \end{pmatrix}, \mathbf{b} = \begin{pmatrix} -1 \\ 6 \\ 3 \end{pmatrix}.$$

Prove that $\mathbf{b} \in \text{span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$.

We want examine the nature of solutions $(x_1, x_2, x_3)^T$ to

$$x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + x_3\mathbf{v}_3 = \mathbf{b}.$$

$\mathbf{b} \in \text{span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ if there is at least one solution. Notice that our equation can be written in the form $A\mathbf{x} = \mathbf{b}$:

$$\begin{pmatrix} 1 & 1 & 2 \\ -1 & 2 & -3 \\ 2 & 5 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -1 \\ 6 \\ 3 \end{pmatrix}.$$

Applying Gaussian elimination, we have

$$\left(\begin{array}{ccc|c} 1 & 1 & 2 & -1 \\ -1 & 2 & -3 & 6 \\ 2 & 5 & 3 & 3 \end{array} \right) \sim \left(\begin{array}{ccc|c} 1 & 1 & 2 & -1 \\ 0 & 3 & -1 & 5 \\ 0 & 0 & 0 & 0 \end{array} \right).$$

Clearly there are infinitely many solutions, so $\mathbf{b} \in \text{span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$.

Example 4

Let \mathbb{P}_2 be the vector space of all real polynomials of degree at most 2. Find three polynomials f_1, f_2, f_3 in \mathbb{P}_2 such that $f_i(0) = 1$ for $i = 1, 2, 3$ and $\{f_1, f_2, f_3\}$ is linearly independent.

We want three polynomials such that each has a different span. The easiest way to do this is

to consider the set of functions

$$\begin{aligned}f_1 &= a_1, \\f_2 &= a_2 + b_2x, \\f_3 &= a_3 + b_3x + c_3x^2.\end{aligned}$$

Since $f_i(0) = 1$ then $a_i = 1$. The other coefficients are arbitrary constants, so set all other constants to 1:

$$\begin{aligned}f_1 &= 1, \\f_2 &= 1 + x, \\f_3 &= 1 + x + x^2.\end{aligned}$$

Example 5: 1241 2016 Q3.iii

The field $\mathbb{F} = GF(4)$ has elements $\{0, 1, \alpha, \beta\}$ with addition and multiplication defined by the following tables. For the vectors

+	0	1	α	β	\times	0	1	α	β
0	0	1	α	β	0	0	0	0	0
1	1	0	β	α	1	0	1	α	β
α	α	β	0	1	α	0	α	β	1
β	β	α	1	0	β	0	β	1	α

$$\mathbf{b}_1 = \begin{pmatrix} 1 \\ \alpha \\ \beta \end{pmatrix}, \mathbf{b}_2 = \begin{pmatrix} \beta \\ 1 \\ 1 \end{pmatrix}, \mathbf{b}_3 = \begin{pmatrix} 1 \\ 0 \\ \alpha \end{pmatrix},$$

- a) show that $\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ is a basis for \mathbb{F}^3 ;
b) explain why $\{\mathbf{b}_1, \mathbf{b}_1 + \mathbf{b}_2, \mathbf{b}_2 + \mathbf{b}_3, \mathbf{b}_3\}$ is a spanning set but not a basis for \mathbb{F}^3 .

First we prove two important results. Suppose $\mathbf{x} \in \mathbb{F}^3$. Since $a + a = 0 \ \forall a \in \mathbb{F}$, then

$$\mathbf{x} + \mathbf{x} = \mathbf{0} \quad \forall \mathbf{x} \in \mathbb{F}^3. \quad (*)$$

Also, since $a + 0 = a \ \forall a \in \mathbb{F}$, then

$$\mathbf{x} + \mathbf{0} = \mathbf{x} \quad \forall \mathbf{x} \in \mathbb{F}^3. \quad (**)$$

Now, note that $\dim \mathbb{F}^3 = 3$. If we can show that $\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ is a linearly independent set then

we can apply the Dimension Theorem. For $x, y, z \in \mathbb{F}$, if

$$x\mathbf{b}_1 + y\mathbf{b}_2 + z\mathbf{b}_3 = \mathbf{0}$$

then we can represent this in the form $A\mathbf{x} = \mathbf{b}$:

$$\begin{pmatrix} 1 & \beta & 1 \\ \alpha & 1 & 0 \\ \beta & 1 & \alpha \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Carefully applying Gaussian elimination:

$$\begin{aligned} \left(\begin{array}{ccc|c} 1 & \beta & 1 & 0 \\ \alpha & 1 & 0 & 0 \\ \beta & 1 & \alpha & 0 \end{array} \right) &\xrightarrow{R1=\alpha R1} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & 0 \\ \alpha & 1 & 0 & 0 \\ \beta & 1 & \alpha & 0 \end{array} \right) \xrightarrow{R2=R2+R1} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & 0 \\ 0 & 0 & \alpha & 0 \\ \beta & 1 & \alpha & 0 \end{array} \right) \\ &\xrightarrow{R3=\beta R3} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & 0 \\ 0 & 0 & \alpha & 0 \\ \alpha & \beta & 1 & 0 \end{array} \right) \xrightarrow{R2 \leftrightarrow R3} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & 0 \\ \alpha & \beta & 1 & 0 \\ 0 & 0 & \alpha & 0 \end{array} \right) \\ &\xrightarrow{R2=R2+R1} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & 0 \\ 0 & \alpha & \beta & 0 \\ 0 & 0 & \alpha & 0 \end{array} \right). \end{aligned}$$

Clearly, our solution $(x, y, z)^T = (0, 0, 0)^T$. So the only solution is the trivial solution, hence $\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ is a linearly independent set. Therefore $\dim(\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}) = \dim(\mathbb{F}^3) = 3$, and so by the Dimension Theorem $\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ is a basis for \mathbb{F}^3 .

Now consider the second set of vectors in \mathbb{F}^3 .

$$\begin{aligned} (\mathbf{b}_1 + \mathbf{b}_2) + (\mathbf{b}_2 + \mathbf{b}_3) + (\mathbf{b}_3) &= \mathbf{b}_1 + (\mathbf{b}_2 + \mathbf{b}_2) + (\mathbf{b}_3 + \mathbf{b}_3) && \text{(associative law)} \\ &= \mathbf{b}_1 + \mathbf{0} + \mathbf{0} && \text{(Using (*))} \\ &= \mathbf{b}_1. && \text{(Using (**))} \end{aligned}$$

Since we have written \mathbf{b}_1 as a linear combination of $\mathbf{b}_1 + \mathbf{b}_2$, $\mathbf{b}_2 + \mathbf{b}_3$ and \mathbf{b}_3 , then

$$\{\mathbf{b}_1, \mathbf{b}_1 + \mathbf{b}_2, \mathbf{b}_2 + \mathbf{b}_3, \mathbf{b}_3\}$$

is a linearly dependent set. Hence the set cannot be a basis for \mathbb{F}^3 . However since

$$\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3 \in \text{span}(\{\mathbf{b}_1, \mathbf{b}_1 + \mathbf{b}_2, \mathbf{b}_2 + \mathbf{b}_3, \mathbf{b}_3\})$$

and $\{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$ is a spanning set for \mathbb{F}^3 , then $\{\mathbf{b}_1, \mathbf{b}_1 + \mathbf{b}_2, \mathbf{b}_2 + \mathbf{b}_3, \mathbf{b}_3\}$ is a spanning set for \mathbb{F}^3 .

Example 6: 1241 2016 Q3.iii

Consider the field $\mathbb{F} = GF(4)$, as defined in the previous example. Let $\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3$ be the vectors from the previous example. Set

$$\mathbf{v} = \begin{pmatrix} \alpha \\ 0 \\ 0 \end{pmatrix}.$$

c) Find the coordinate vector of \mathbf{v} with respect to the ordered basis $B = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3\}$.

The coordinate vector $\mathbf{x} = (x, y, z)^T$ of \mathbf{v} will satisfy the equation

$$x\mathbf{b}_1 + y\mathbf{b}_2 + z\mathbf{b}_3 = \mathbf{v}.$$

Writing this in the form $A\mathbf{x} = \mathbf{b}$, we have

$$\begin{pmatrix} 1 & \beta & 1 \\ \alpha & 1 & 0 \\ \beta & 1 & \alpha \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \alpha \\ 0 \\ 0 \end{pmatrix}.$$

Carefully applying Gaussian elimination:

$$\begin{aligned} \left(\begin{array}{ccc|c} 1 & \beta & 1 & \alpha \\ \alpha & 1 & 0 & 0 \\ \beta & 1 & \alpha & 0 \end{array} \right) &\xrightarrow{R1=\alpha R1} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & \beta \\ \alpha & 1 & 0 & 0 \\ \beta & 1 & \alpha & 0 \end{array} \right) \xrightarrow{R2=R2+R1} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & \beta \\ 0 & 0 & \alpha & \beta \\ \beta & 1 & \alpha & 0 \end{array} \right) \\ &\xrightarrow{R3=\beta R3} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & \beta \\ 0 & 0 & \alpha & \beta \\ \alpha & \beta & 1 & 0 \end{array} \right) \xrightarrow{R2 \leftrightarrow R3} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & \beta \\ \alpha & \beta & 1 & 0 \\ 0 & 0 & \alpha & \beta \end{array} \right) \\ &\xrightarrow{R2=R2+R1} \left(\begin{array}{ccc|c} \alpha & 1 & \alpha & \beta \\ 0 & \alpha & \beta & \beta \\ 0 & 0 & \alpha & \beta \end{array} \right). \end{aligned}$$

From $R3$ we have $z = \alpha$ since $\alpha \times \alpha = \beta$. In $R2$ we have

$$\begin{aligned} \alpha y + \beta \times \alpha &= \beta \\ \alpha y + 1 &= \beta && (\beta \times \alpha = 1) \\ \alpha y + 1 + 1 &= \beta + 1 && (\text{Adding 1 to both sides}) \\ \alpha y &= \alpha && (1 + 1 = 0 \text{ and } \beta + 1 = \alpha) \\ y &= 1. && (\text{Since } \alpha \times 1 = \alpha) \end{aligned}$$

In $R1$ we have

$$\begin{array}{ll}
 \alpha x + 1 \times 1 + \alpha \times \alpha = \beta & \\
 \alpha x + 1 + \beta = \beta & (1 \times 1 = 1 \text{ and } \alpha \times \alpha = \beta) \\
 \alpha x + (1 + \beta) + (1 + \beta) = \beta + 1 + \beta & (\text{Adding } (1 + \beta) \text{ to both sides}) \\
 \alpha x = \alpha + \beta & ((1 + \beta) + (1 + \beta) = 0 \text{ and } \beta + 1 = \alpha) \\
 \alpha x = 1 & (\alpha + \beta = 1) \\
 x = \beta & (\text{Since } \alpha \times \beta = 1)
 \end{array}$$

Hence the coordinate vector of \mathbf{v} with respect to the basis B is

$$\mathbf{x} = \begin{pmatrix} \beta \\ 1 \\ \alpha \end{pmatrix}.$$

Example 7: 1241 2014 S2 Q3.i

Prove that the function $T : \mathbb{P}(\mathbb{R}) \rightarrow \mathbb{R}^2$ defined by

$$T(p) = \begin{pmatrix} p(0) \\ p(1) \end{pmatrix}, \text{ for all polynomials } p \in \mathbb{P}(\mathbb{R}),$$

is a linear transformation.

For the map T to be linear, we need to show that T preserves addition and scalar multiplication. First consider addition. For any $p, q \in \mathbb{P}(\mathbb{R})$,

$$\begin{aligned}
 T(p + q) &= \begin{pmatrix} (p + q)(0) \\ (p + q)(1) \end{pmatrix} = \begin{pmatrix} p(0) + q(0) \\ p(1) + q(1) \end{pmatrix} = \begin{pmatrix} p(0) \\ p(1) \end{pmatrix} + \begin{pmatrix} q(0) \\ q(1) \end{pmatrix} \\
 &= T(p) + T(q).
 \end{aligned}$$

So T preserves addition. Now consider scalar multiplication. For any $p \in \mathbb{P}(\mathbb{R})$ and $\lambda \in \mathbb{R}$,

$$\begin{aligned}
 T(\lambda p) &= \begin{pmatrix} (\lambda p)(0) \\ (\lambda p)(1) \end{pmatrix} = \begin{pmatrix} \lambda p(0) \\ \lambda p(1) \end{pmatrix} = \lambda \begin{pmatrix} p(0) \\ p(1) \end{pmatrix} \\
 &= \lambda T(p).
 \end{aligned}$$

Hence T preserves scalar multiplication. Therefore since T preserves addition and scalar multiplication, then T is linear.

Example 8: 1241 2016 Q3.ii

Let V and W be vector spaces, let $T : V \rightarrow W$ be a linear transformation, and let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ be vectors in V .

- a) Prove that if $T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_m)$ are linearly independent, then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ are linearly independent.
- b) Suppose that $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ are linearly independent. Is $T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_m)$ linearly independent?

Suppose that $T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_m)$ are linearly independent, and assume $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ are linearly dependent. Then for some $i \in \{1, 2, \dots, m\}$ and constants λ_j , we have

$$\mathbf{v}_i = \sum_{j \neq i} \lambda_j \mathbf{v}_j.$$

So then

$$\begin{aligned} T(\mathbf{v}_i) &= T\left(\sum_{j \neq i} \lambda_j \mathbf{v}_j\right) \\ &= \sum_{j \neq i} T(\lambda_j \mathbf{v}_j) && \text{(Since } T \text{ preserves addition)} \\ &= \sum_{j \neq i} \lambda_j T(\mathbf{v}_j). && \text{(Since } T \text{ preserves scalar multiplication)} \end{aligned}$$

Hence we have a contradiction, since $T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_m)$ are linearly independent. Hence our assumption is incorrect, i.e. $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ are not linearly dependent. So we have proven, by contradiction, that $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ are linearly independent.

However, linear independence of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ does not imply linear independence of $T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_m)$. Consider, for example, the linear map $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\mathbf{x}) = \mathbf{0} \quad \forall \mathbf{x} \in \mathbb{R}^2.$$

The standard basis vectors $\mathbf{e}_1, \mathbf{e}_2$ are linearly independent, however $xT(\mathbf{e}_1) + yT(\mathbf{e}_2) = \mathbf{0}$ for any choice of $x, y \in \mathbb{R}$. So $T(\mathbf{e}_1), T(\mathbf{e}_2)$ are not linearly independent. Interestingly enough, part b would be true if T were injective.

Example 9: 1231 2013 Q2,iv

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear map which rotates a vector \mathbf{x} about the origin through $\frac{\pi}{6}$ anti-clockwise and doubles its length.

a) Show that $T(\mathbf{e}_1) = \begin{pmatrix} \sqrt{3} \\ 1 \end{pmatrix}$, where $\mathbf{e}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

b) Find the matrix A such that $T(\mathbf{x}) = A\mathbf{x}$ for all $\mathbf{x} \in \mathbb{R}^2$.

Since T rotates a vector $(x, y)^T$ anticlockwise by $\frac{\pi}{6}$, we know that for $x > 0$ and $y \geq 0$,

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} L \cos \left(\frac{\pi}{6} + \tan^{-1} \left(\frac{y}{x} \right) \right) \\ L \sin \left(\frac{\pi}{6} + \tan^{-1} \left(\frac{y}{x} \right) \right) \end{pmatrix}.$$

Since T also doubles the length of a vector $(x, y)^T$, then $L = 2\sqrt{x^2 + y^2}$. Hence

$$T \begin{pmatrix} x \\ y \end{pmatrix} = 2\sqrt{x^2 + y^2} \begin{pmatrix} \cos \left(\frac{\pi}{6} + \tan^{-1} \left(\frac{y}{x} \right) \right) \\ \sin \left(\frac{\pi}{6} + \tan^{-1} \left(\frac{y}{x} \right) \right) \end{pmatrix}.$$

So

$$\begin{aligned} T \begin{pmatrix} 1 \\ 0 \end{pmatrix} &= 2\sqrt{1+0} \begin{pmatrix} \cos \left(\frac{\pi}{6} + \tan^{-1} 0 \right) \\ \sin \left(\frac{\pi}{6} + \tan^{-1} 0 \right) \end{pmatrix} = 2 \begin{pmatrix} \sqrt{3}/2 \\ 1/2 \end{pmatrix} \\ &= \begin{pmatrix} \sqrt{3} \\ 1 \end{pmatrix}. \end{aligned}$$

Also,

$$\begin{aligned} T \begin{pmatrix} 0 \\ 1 \end{pmatrix} &= 2\sqrt{0+1} \begin{pmatrix} \cos \left(\frac{\pi}{6} + \frac{\pi}{2} \right) \\ \sin \left(\frac{\pi}{6} + \frac{\pi}{2} \right) \end{pmatrix} = 2 \begin{pmatrix} -1/2 \\ \sqrt{3}/2 \end{pmatrix} \\ &= \begin{pmatrix} -1 \\ \sqrt{3} \end{pmatrix}. \end{aligned}$$

Using the Matrix Representation Theorem, $T(\mathbf{x}) = A\mathbf{x}$ where

$$A = \begin{pmatrix} \sqrt{3} & -1 \\ 1 & \sqrt{3} \end{pmatrix}.$$

Example 10: 1231 2018 Q1.iv

Consider the matrix $M = \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix}$.

a) Find a basis for $\ker(M)$.

b) Find a basis for $\text{im}(M^T)$.

c) Give a geometric description of $\ker(M)$ and $\text{im}(M)$ as subspaces of \mathbb{R}^2 .

If $\mathbf{x} \in \ker(M)$ then $M\mathbf{x} = \mathbf{0}$. Hence

$$\begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$
$$\begin{pmatrix} x + y \\ 2x + 2y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

So $x + y = 0$, i.e. $y = -x$. So

$$\mathbf{x} = x \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

Therefore

$$\ker(M) = \text{span} \left\{ \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\},$$

so a basis for $\ker(M)$ is

$$\left\{ \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\}.$$

Now, consider $\mathbf{y} \in \text{im}(M^T)$. Then

$$\mathbf{y} = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x + 2y \\ x + 2y \end{pmatrix} = (x + 2y) \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

Therefore

$$\text{im}(M^T) = \text{span} \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\},$$

so a basis for $\text{im}(M^T)$ is

$$\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}.$$

The kernel of M is the line in \mathbb{R}^2 , in the direction $(1, -1)^T$. The image of M is the line in \mathbb{R}^2 , in the direction $(1, 2)^T$.

Example 11: 1241 2015 Q3.ii

Consider the mapping $T : \mathbb{P}_2 \rightarrow \mathbb{P}_3$ defined by

$$T(p)(x) = (x^2 + 1)p'(x) - 2xp(x).$$

Assuming T is linear, find the rank and nullity of T .

Let $p(x) = ax^2 + bx + c$. Then $p'(x) = 2ax + b$, and so

$$\begin{aligned} T(p)(x) &= (x^2 + 1)(2ax + b) - 2x(ax^2 + bx + c) \\ &= -bx^2 + 2(a - c)x + b. \end{aligned}$$

Hence

$$\begin{aligned} T(p) &= \begin{pmatrix} -b \\ 2a - 2c \\ b \end{pmatrix} = a \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix} + b \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} + c \begin{pmatrix} 0 \\ -2 \\ 0 \end{pmatrix} \\ &= 2(a - c) \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + b \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}. \end{aligned}$$

So if $q \in \text{im}(T)$ then

$$q = \text{span} \left\{ \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right\}.$$

So $\text{rank}(T) = 2$. Since $\dim(\mathbb{P}_2) = 3$ (standard basis is $\{1, x, x^2\}$), then by the Rank-Nullity Theorem, $\text{nullity}(T) = 1$.