



# MATH1131/41 Calculus

## Lab Test 2 Sample Solutions

July 20, 2019

These answers were written and typed by Supriya Segal, and edited by Abdellah Islam. Please be ethical with this resource. The questions here were taken from the MATH1131/41 Test 2 Calculus Question bank, and all copyright of the original questions belongs to UNSW's School of Mathematics and Statistics. It is for the use of MathSoc members, so do not repost it on other forums or groups without asking for permission. If you appreciate this resource, please consider supporting us by coming to our events! Also, happy studying :)

We cannot guarantee that our answers are correct - please notify us of any errors or typos at [unswmathsoc@gmail.com](mailto:unswmathsoc@gmail.com), or on our [Facebook page](#). There are sometimes multiple methods of solving the same question.

---

### Question 1

*Find, to 10 significant figures, the unique turning point  $x_0$  of*

$$f(x) = 5 \sin\left(\frac{1}{2}x^2\right) - \sin\left(\frac{5}{2}x\right)^2$$

*in the interval  $[1, 2]$ . Find, to 10 significant figures, the value of the second derivative  $f''(x)$  at the turning point, that is  $f''(x_0)$ .*

**Solution:** The answers are 1.580743250 and  $-9.652901450$ .

In Maple, assign the variable **f** to the function. Now assign a variable **p** to the derivative of the function

```
p := diff(f,x).
```

To solve for **p** on the interval  $[1, 2]$ , use the command

```
r := fsolve(p, x = 1..2).
```

This will output 1.580743250, which is the answer to the first part of the question.

Now differentiate **p** to get  $f''(x)$  and assign it to a variable using the command

```
q := diff(p,x).
```

To find  $f''(x_0)$ , enter the command

```
evalf(subs(x=r,q)).
```

This will output  $-9.652901450$ , which is the answer to the second part of the question.

## Question 2

The Maple expression for the constant  $\pi$  is \_\_\_\_\_. The Maple expression for  $\infty$  is \_\_\_\_\_. Evaluate, to 10 significant figures,

$$\int_1^{\infty} \frac{e^{-x} \cos\left(\frac{x^2}{3}\right)}{3+x} dx.$$

**Solution:** The answers are Pi, infinity and 0.03239018036

To find the integral using Maple, assign the variable a to the fraction to be integrated, then enter the Maple commands

```
int(a,x=1..infinity)
```

then

```
evalf(%)
```

to get the answer to 10 significant figures.


## Question 3

Select the option below which is the plot of the polar curve  $r = \sin(\theta) - 3 \cos(4\theta)$  for  $0 \leq \theta \leq 2\pi$ .

**Solution:** In Maple, activate `with(plots)` then enter the equation of the curve and assign it to a variable `r`. Then enter the Maple command

```
polarplot([r], theta = 0..2*Pi)
```

which will plot the polar curve.



## Question 4

Find the largest interval of the form  $[a, b]$  containing  $-4$  on which the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by the rule

$$f(x) = 2x^3 + 9x^2 - 60x - 7$$

has an inverse.

**Solution:** The answer is  $[-5, 2]$ .

A function is invertible if it is monotonically increasing or monotonically decreasing and continuous over an interval.

We can differentiate  $f(x)$ , either using Maple or by hand, resulting in

$$f'(x) = 6x^2 + 18x - 60.$$

To find the stationary points, set  $f'(x) = 0$ , which results in  $x = 2, -5$ . By testing points on either side of these  $x$  values, we can see that  $x = -5$  is a max point and  $x = 2$  is a min point. Since the curve is continuous between these two  $x$  values with a negative first derivative, we can deduce that it is monotonically decreasing over  $[-5, 2]$ .

## Question 5

Find the slope of the line tangent to

$$-y^6 + x^3 + 2x^2y = 1$$

at  $(1, 0)$  and enter it as a fraction or integer.

**Solution:** The answer is  $-\frac{3}{2}$ .

Differentiate the curve implicitly, either by hand or by setting the curve to the variable `eqn` then using the Maple command

`implicitdiff(eqn,y,x).`

This gives the output  $\frac{-3x^2 - 4xy}{-6y^5 + 2x^2}$ . Then substitute  $(1, 0)$  into this equation to find the answer.

## Question 6

Consider the function  $f : \mathbb{R} \rightarrow \mathbb{R}$ , defined by the rule  $f(x) = |x^3 + 5x^2|$ .

(a) Find all critical points of  $f$  on the interval  $[-5, -1]$  as exact values.

(b) Complete the following sentence. The function  $f$  is guaranteed to have \_\_\_ [choices: a max but no min / either a max or a min but not both / both a max and a min / stationary / min but no max] value on  $[-5, -1]$  because it is \_\_\_ [choices: integrable / bijective / continuous / differentiable / surjective / bounded / injective] on the interval  $[-5, -1]$  which is \_\_\_ [choices: invertible / both open and closed / continuous / real / closed / open] and bounded.

(c) The maximum value of  $f$  on  $[-5, -1]$  is \_\_\_\_\_. The minimum value of  $f$  on  $[-5, -1]$  is \_\_\_\_\_.

(d) At which type of critical point or points does the maximum occur? Tick all that apply. [Choices: An end point of  $[-5, -1]$ ; A stationary point of  $f$  on  $(-5, -1)$ ; A point on  $(-5, -1)$  where  $f$  is not differentiable].

(e) At which type of critical point or points does the minimum occur? Tick all that apply. [Choices: An end point of  $[-5, -1]$ ; A stationary point of  $f$  on  $(-5, -1)$ ; A point on  $(-5, -1)$  where  $f$  is not differentiable].

**Solution:**


(a) The answer is  $\{-5, -10/3, -1\}$ .

In Maple, assign the variable `f` to the function,

`f := abs(x^3 + 5*x^2).`

Then plot the graph on the interval  $[-6, 0]$  by using

```
plot(f, x=-6..0).
```



We can see from the graph that there is a point where  $f$  is not differentiable at  $x = -5$ , and there is a maximum point near  $x = -3$ .

To find this max, differentiate  $f$  and find where the first derivative is equal to 0, by using the command

```
g = diff(f, x)
```

which outputs  $(3x^2 + 10x)|1, x^3 + 5x^2|$ . To find the stationary point,

```
solve((3x^2 + 10x)*abs(1, x^3 + 5x^2) = 0)
```

which outputs  $-\frac{10}{3}$ .

(b) The answers are both a max and a min, continuous, closed. (By the Extreme Value Theorem.)

(c) The answers are  $\frac{500}{27}$ , 0.

We can see that the minimum is 0 from the graph we plotted for part (a). The maximum can be obtained by substituting  $-\frac{10}{3}$  into  $f$ .

(d) The answer is: A stationary point of  $f$  on  $(-5, 1)$ .

We can see this from the graph we plotted in part (a).

(e) The answers are: An end point, a point where it is not differentiable.