

# MATH1231/1241 Lab Test 1

## Algebra Solutions to Samples

October 7, 2019

These solutions were written and typed by Jeffrey Yang, and edited by Ethan Brown. Please be ethical with this resource. It is for the use of MathSoc members - do not repost it on other forums or groups without asking for permission. If you appreciate our resources, please consider supporting us by coming to our events! Also, happy studying :)

We cannot guarantee that our answers are correct - please notify us of any errors or typos at [unswmathsoc@gmail.com](mailto:unswmathsoc@gmail.com), or on our [Facebook page](#). There are sometimes multiple methods of solving the same question.

---

### Question 1

*Select the correct definition for  $\text{span}\{S\}$ , the set of vectors  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  in a vector space  $V$  with a scalar field  $\mathbb{F}$  from the choices below:*

1.  $\text{span}\{S\} = \{\mathbf{x} \in V : \mathbf{x} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n \text{ for } \lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}\}$
2.  $\text{span}\{S\} = \{\mathbf{x} \in V : \text{if } \mathbf{x} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n \text{ for } \lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}, \text{ then } \lambda_1 = \lambda = \dots = \lambda_n = 0\}$
3.  $\text{span}\{S\} = \{\mathbf{x} \in V : \mathbf{x} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n\}$
4.  $\text{span}\{S\} = \{\mathbf{x} \in V : \mathbf{x} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n \text{ for } \lambda_1, \lambda_2, \dots, \lambda_n \in V\}$
5.  $\text{span}\{S\} = \{\mathbf{x} \in V : \text{if } \mathbf{x} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n \text{ for } \lambda_1, \lambda_2, \dots, \lambda_n \in V, \text{ then } \lambda_1 = \lambda = \dots = \lambda_n = 0\}$

Recall that the span of a set refers to the set of all possible **linear combinations** of its elements. More specifically, each element in the span is formed by multiplying each vector in the set by a scalar (that is, an element from the scalar field  $\mathbb{F}$ ) and then adding the results. As such, the correct definition here is (1.). We require each  $\lambda_i$  to be a scalar and thus, have to be taken from the scalar field  $\mathbb{F}$  (and NOT the vector space  $V$ ). Moreover, they do not (and should not) all equal to 0.

## Question 2

Select all of the statements below that are true.

1. There exists a set of 4 vectors in  $\mathbb{R}^3$  that is linearly independent.
2. All sets of 4 vectors in  $\mathbb{R}^3$  must span  $\mathbb{R}^3$ .
3. There exists a set of 4 vectors in  $\mathbb{R}^3$  that is linearly independent.
4. All sets of 3 vectors in  $\mathbb{R}^4$  are linearly independent.
5. All sets of 4 vectors in  $\mathbb{R}^3$  are linearly dependent.
6. All sets of 3 vectors in  $\mathbb{R}^4$  do not span  $\mathbb{R}^4$ .
7. A set can have at most  $n$  mutually orthogonal non-zero vectors in  $\mathbb{R}^n$ .

In general,  $\mathbb{R}^n$  can have at most  $n$  linearly independent vectors and any spanning set of  $\mathbb{R}^n$  must have  $n$  **linearly independent** vectors. Expanding on the latter, it means that any spanning set of  $\mathbb{R}^n$  must have *at least*  $n$  vectors (however, simply having  $n$  or more vectors in a set does not imply that it spans  $\mathbb{R}^n$ ). Hence, (5.), (6.) and (7.) are true.

## Question 3

Select the correct completion of the definition of **linear independence** from the choices below.

The vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  in a vector space  $V$  over a scalar field  $\mathbb{F}$  are **linearly independent** means

1. if there are scalars  $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}$  such that  $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n = \mathbf{0}$  then  $\mathbf{v}_1 = \mathbf{v}_2 = \dots = \mathbf{v}_n = \mathbf{0}$
2. there are scalars  $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}$  such that  $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n = \mathbf{0}$  and  $\lambda_1 = \lambda_2 = \dots = \lambda_n = 0$
3. if there are scalars  $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}$  such that  $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n = \mathbf{0}$  then  $\lambda_1 = \lambda_2 = \dots = \lambda_n = 0$
4. there are scalars  $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}$  such that  $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n = \mathbf{0}$

The correct answer is (3.). It is important to note that  $\lambda_1 = \lambda_2 = \dots = \lambda_n = \mathbf{0}$  has to be the ONLY solution to the equation  $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n = \mathbf{0}$  for the vectors to be linearly

independent. The significance of this result is that no vector in  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  can be written as a linear combination of other vectors in  $S$ . If  $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n = \mathbf{0}$  had a non-trivial solution, we could just move any vector  $\lambda_i \mathbf{v}_i$  to the other side (assuming  $\lambda_i \neq 0$ ) and divide both sides by  $\lambda_i$  to write  $\mathbf{v}_i$  as a linear combination of the other vectors in  $S$  – this would imply that  $S$  is not linearly independent.

## Question 4

Select from the choices below the correct definition from a set  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  in a vector space  $V$  over a scalar field  $\mathbf{F}$  to **span**  $V$ .

1. If there are scalars  $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}$  such that  $\lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n = \mathbf{0}$  then  $\lambda_1 = \lambda_2 = \dots = \lambda_n = 0$ .
2. For any vector  $\mathbf{x} \in V$ , if there are scalars  $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}$  such that  $\mathbf{x} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n$  then  $\lambda_1 = \lambda_2 = \dots = \lambda_n = 0$ .
3. For every vector  $\mathbf{x} \in V$ , there are scalars  $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}$  such that  $\mathbf{x} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n$ .
4. For every vector  $\mathbf{x} \in V$ , there are **unique** scalars  $\lambda_1, \lambda_2, \dots, \lambda_n \in \mathbb{F}$  such that  $\mathbf{x} = \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n$ .

For  $S$  to span  $V$ , we need to be able to express any vector in  $V$  as a linear combination of the vectors in  $S$ . Note that the linear combination does not have to be unique - this is especially true when the number of elements in  $S$  is greater than the dimension of  $V$  (if each linear combination was unique, then  $S$  would be a **basis** for  $V$ ). As such, the correct answer is (3.). The condition that  $\lambda_1 = \lambda_2 = \dots = \lambda_n = 0$  pertains to linear independence and is irrelevant here.

## Question 5

Let  $\mathbb{R}^3$  have the usual componentwise vector space operations.

Let  $S$  be the subset of  $\mathbb{R}^3$  defined by

$$S = \left\{ \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \in \mathbb{R}^3 : x_3^3 - x_1 x_2^2 = 0 \right\}$$

Note that  $S$  contains the zero vector  $\mathbf{0}$ .

1. EITHER state  $S$  is closed under vector addition by writing the word *CLOSED* in the answer box below OR show  $S$  is NOT closed under vector addition by specifying two different non-zero vectors  $\mathbf{x}, \mathbf{y}$  in  $S$  with integer coefficients which are not scalar multiples of each other such that  $\mathbf{x} + \mathbf{y}$  is **not** in  $S$ .

Enter your answer in the box below using Maple notation, e.g. enter  $\langle 1, 2, 3 \rangle, \langle 7, 8, 9 \rangle$

for  $\mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \mathbf{y} = \begin{pmatrix} 4 \\ 5 \\ 6 \end{pmatrix}$ .

2. EITHER state  $S$  is closed under scalar multiplication by entering the word *CLOSED* in the answer box below OR show  $S$  is not closed under scalar multiplication by specifying (in this order) a non-zero vector  $\mathbf{x}$  in  $S$  with integer coefficients and an integer scalar  $\lambda$  such that the vector  $\lambda\mathbf{x}$  is **not** in  $S$ .

Enter your answer in the box below using Maple notation, e.g. enter  $\langle 1, 2, 3 \rangle, 4$  for

$\mathbf{x} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \lambda = 4$ .

1. We have  $\begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \in S$  but  $\begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 0 \end{pmatrix} \notin S$ . Thus, we would enter  $\langle 1, 1, -1 \rangle, \langle 1, 1, 1 \rangle$ . As a general tip, when there is a coefficient being squared in the equation ( $x_2$  in this case), consider two vectors which are the same EXCEPT with opposite signs for the squared coefficient.

2. CLOSED. As a general tip, ensure that the degree of each term is the same in the equation - here, both terms are of degree three.

### Question 13

Let

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} 2 \\ 3 \\ 5 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 3 \\ 2 \\ 0 \end{pmatrix}$$

Find  $A$ , an augmented matrix (with no bar), for the problem of when deciding when

$$\mathbf{x} \in \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$$

Enter your matrix in the box below.

Now use this matrix to determine what condition(s), if any, are required on  $x, y, z$  so that

$$\mathbf{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \in \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$$

A condition will be an equation in  $x, y, z$ . Use  $*$  for multiplication (i.e. enter expression in Maple format). For example, a typical answer would be  $2*x-y+3*z=0$

If there is more than one condition give answer as a set of equations separated by commas and enclosed in braces ,, e.g.  $2*x-y+3*x=0, 4*y-z=0$

If there are NO conditions, give the answer as: **None**

We consider the matrix whose columns are formed by  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  and  $\mathbf{x}$ . That is,

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & x \\ 2 & 3 & 2 & y \\ 4 & 5 & 0 & z \end{array} \right)$$

Reducing this to row echelon form, we have

$$\left( \begin{array}{ccc|c} 1 & 2 & 3 & x \\ 0 & 1 & 4 & 2x - y \\ 0 & 0 & 0 & 2x - 3y + z \end{array} \right)$$

This will only have a solution if  $2x - 3y + z = 0$ .

## Question 17

1. Let  $p_1(t) = 1 + 3t + 5t^2$ ,  $p_2(t) = 2 + 3t + t^2$ ,  $p_3(t) = 5 + 8t + 4t^2$  and  $q(t) = 1 + t + t^2$ .  
Give an augmented matrix which represents the linear system

$$\mu_1 p_1 + \mu_2 p_2 + \mu_3 p_3 = q$$

with real scalars  $\mu_1, \mu_2, \mu_3$ .

2. Suppose four polynomials  $p_1, p_2, p_3, q$  in  $\mathbb{P}_2$  have augmented matrix  $A$  corresponding to the equation above has echelon form

$$U = \left( \begin{array}{ccc|c} 1 & 2 & -3 & 4 \\ 0 & 1 & 7 & 6 \\ 0 & 0 & 0 & -3 \end{array} \right)$$

Are the three polynomials  $p_1, p_2, p_3$  **linearly independent**? (Yes/No)

Give the correct reasoning from

1.  $U$  has an all zero row in its first 4 columns
2. The first 3 columns of  $U$  are leading columns
3. There are non-leading columns in the first 3 columns of  $U$

If they are **linearly dependent**, then give a linear combination of  $p_1$  and  $p_2$  which equals  $p_3$ .

If they are **linearly independent**, then give a linear combination of  $p_1, p_2$  and  $p_3$  which equals  $q$ .

1. The corresponding augmented matrix  $A$  is

$$A = \left( \begin{array}{ccc|c} 1 & 2 & 5 & 1 \\ 3 & 3 & 8 & 1 \\ 5 & 1 & 4 & 1 \end{array} \right)$$

Note that the columns of the matrix are formed by the coefficients of each polynomial.

2. The polynomials  $p_1, p_2, p_3$  are NOT linearly independent since there are non-leading columns in the first 3 columns of  $U$  (i.e. (3.)). (2.) would imply linear independence and (1.) would imply linear dependence. From  $U$ , we deduce that  $-17p_1 + 7p_2 = p_3$  – ignore the 4th column and imagine that there was a bar between the 2nd and 3rd column.

## Question 23

Consider the following sets

$$A = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \in \mathbb{R}^3 : -9x - 8y + 5z = 0 \text{ where } -5 < x < 5 \text{ and } -4 < y < 4 \right\}$$

$$B = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} \in \mathbb{R}^3 : \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \lambda \begin{pmatrix} -2 \\ 0 \\ 4 \end{pmatrix} + \mu \begin{pmatrix} 0 \\ -8 \\ 8 \end{pmatrix} \text{ and } 6x + 9y - 2z = 0 \right\}$$

$$C = \{p \in \mathbb{P}_3 : p(x) = x^3 + ax^2 + bx + c \text{ where } a, b, c \text{ are real numbers}\}$$

Select all of the following which are true.

1.  $A$  is a subspace of  $\mathbb{R}^3$
2.  $B$  is a subspace of  $\mathbb{R}^3$
3.  $C$  is a subspace of  $\mathbb{P}^3$

Only (2.) is true. Neither  $A$  nor  $C$  are closed under scalar multiplication.