



MATH2621 Revision Seminar

Solutions

November 25, 2019

These solutions were written and typed by Kabir Agrawal and Ethan Brown. Please be ethical with this resource. It is for the use of MathSoc members - do not repost it on other forums or groups without asking for permission. If you appreciate our resources, please consider supporting us by coming to our events! Also, happy studying :)

We cannot guarantee that our answers are correct - please notify us of any errors or typos at unswmathsoc@gmail.com, or on our [Facebook page](#). There are sometimes multiple methods of solving the same question.

Example 1

Set	Open/Closed	Bounded	Compact	Connected	Simply C.	Region	Domain
S_1	Open	Yes	No	Yes	Yes	Yes	Yes
S_2	Closed	Yes	Yes	Yes	Yes	Yes	No
S_3	Closed	Yes	Yes	Yes	Yes	No	No
S_4	Open	Yes	No	Yes	No	Yes	Yes
S_5	Open	No	No	Yes	No	Yes	Yes

Note: Simply C. = Simply Connected. Important observations:

1. To justify S_1 is open, you can generally just quote the fact that a disc that doesn't contain it's boundary is automatically open. The more formal proof entails the following:

Let $\epsilon = 1 - |z|$, where $z = x + iy \in S_1$. Let $y \in B(z, \epsilon)$. Then $|y - z| < \epsilon = 1 - |z|$. Hence, $|y - z| + |z| < 1$. By triangle inequality, $|y - z| + |z| \geq |y - z + z| = |y|$ and

thus $|y| < 1$. Since the choice of $y \in B(z, \epsilon)$ was arbitrary, and for every $z \in S_1$, $B(z, \epsilon) \subseteq S_1$, S_1 must therefore be open.

2. S_3 is connected. We can use the path $c = \{z \in S_3 : z = tp + (1 - t)p, t \in [0, 1]\} = \{p\}$, and since such a set is always contained in S_3 , S_3 must be connected. Additionally, the point p is an boundary point of S_3 because every epsilon ball about p contains p but also contains elements outside p , and so any other point that is not p is an exterior point, and S_3 has no interior. This is why S_3 also is NOT a region.
3. There are sets that are neither open nor closed, and sets that are both open and closed. An example of a set that is neither open nor closed is:

$$S = \{z \in \mathbb{C} : \operatorname{Re}(z) \in (0, \infty), \operatorname{Im}(z) \in [0, \infty)\}$$

Since it contains some of it's boundary points (on the positive imaginary axis) but not all (for example, those on the positive real axis). The only sets in \mathbb{C} that are open and closed are \emptyset, \mathbb{C} . \emptyset is open because it doesn't contain any of its boundary points, and thus \mathbb{C} is closed for that reason. And \mathbb{C} is open because it can be written as a union of open balls in \mathbb{C} , and for this reason, \emptyset is closed.

Example 2

1. Note that it suffices to work out what happens to the boundaries, the region will be that which is inside the boundary. Consider the boundary line $z = c + iy$ (where $c \in [0, 4]$), then the image of that will be $(1 + i) \cdot (c + iy) + 2 = i(y + c) + 2 - y + c$. Letting $u = 2 - y + c, v = y + c$, we obtain the new equation to be $u + v = 2c + 2$, and since we restrict $y \in [-0.5, 5]$, we obtain the image:
2. The line passing through $z = 1, z = 3 + 4i$ is given by $z = t(1) + (1 - t)(3 + 4i) = (1 - 2t) + (4 - 4t)i$. Thus, the image will be $(1 + i)z + 2 = (1 + i)((1 - 2t) + i(4 - 4t)) + 2 = (1 - 2t - (4 - 4t)) + i(1 - 2t + 4 - 4t) = (-3 + 2t) + i(5 - 6t)$. Letting $u = -3 + 2t, v = 5 - 6t$, we get $3u + v = -4$ as the image, with no restriction because t is unrestricted.

Example 3

By Extended Triangle inequality, we have $|z^4 - 1| \geq ||z|^4 - 1| = |R^4 - 1| = R^4 - 1 \geq 15$ since $R \geq 2$ and thus $R^4 - 1 > 0$. Since both sides of the inequality are greater than 0, we may

reciprocate both sides to yield:

$$\left| \frac{1}{z^4 - 1} \right| \leq \frac{1}{15}$$

as required.

Example 4

By definition of a limit, we seek a δ such that for every $\epsilon > 0$, we have $0 < |z - (1 + i)| < \delta \implies |z^2 - 2i| < \epsilon$.

$$\begin{aligned} |z^2 - 2i| &= |z - (1 + i)||z + (1 + i)| \\ &= |z - (1 + i)||z - (1 + i) + (2 + 2i)| \\ &\leq \delta(\delta + 2\sqrt{2}) && \text{(By Triangle Inequality)} \\ &< \epsilon \end{aligned}$$

Where we select δ such that $\delta < \text{the positive solution of } \delta^2 + 2\sqrt{2}\delta - \epsilon = 0 \implies \delta = \frac{-2\sqrt{2} + \sqrt{8 + 4\epsilon}}{2}$.

Example 5

Consider the path $z = iy$, then the limit becomes:

$$\lim_{y \rightarrow 0} \frac{0}{0 + iy} = 0$$

Consider the path $z = x + 0i$, then the limit becomes:

$$\lim_{x \rightarrow 0} \frac{x}{x + 0i} = 1$$

Since the limits along the 2 different paths are different, the limit expression does not exist.

Example 6

1. $f_1(z) = (x + iy)(x^2 + y^2) = (x^3 + xy^2) + i(yx^2 + y^3)$. Then by the Cauchy-Riemann Equations:

$$\frac{\partial u}{\partial x} = 3x^2 + y^2, \quad \frac{\partial u}{\partial y} = 2xy$$

$$\frac{\partial v}{\partial x} = 2xy, \quad \frac{\partial v}{\partial y} = 3y^2 + x^2$$

We have $2xy = -2xy \implies xy = 0$. We also have $3x^2 + y^2 = -3y^2 - x^2 \implies 4x^2 = -4y^2 \implies x^2 = -y^2 \implies x = y = 0$. So the function is differentiable only at $x = y = 0$.

Note that this function is not holomorphic at $z = 0$.

2. $f_2(z) = x^2 + iy^2 \implies 2x = 2y, 0 = -0 \implies x = y$. Hence the function is differentiable $z = x + ix$.
3. Similar to the previous part, we require when $\frac{x}{|x|} = \frac{|y|}{y} \implies xy = |xy|$ (upon noting that the derivative of $|x| = \frac{x}{|x|}$). The above is only true when $xy > 0$. This also means that the function is holomorphic in this region, since it's differentiable on the open set.

Example 7

$u(x, y) = \cos x \cosh y \implies \partial_x^2 u = -\cos x \cosh y, \partial_y^2 u = \cos x \cosh y \implies \partial_x^2 u + \partial_y^2 u = 0$. Hence u is harmonic. The harmonic conjugate is given by solving the CRE's. Let v be the harmonic conjugate so that $\partial_x v = -\cos x \sinh y \implies v(x, y) = -\sin x \sinh y + f(y)$. $\partial_y v = -\sin x \cosh y \implies f'(y) = 0 \implies f(y) = C$. Hence the harmonic conjugate is $v(x, y) = -\sin x \sinh y + C$.

Example 8

Using the same idea as above, you should obtain $v(x, y) = \frac{-y}{x^2 + y^2}$.

Example 9

1. $e^z = 2i \implies x = \log 2i = \ln |2i| + i \arg(2i) = \ln 2 + i\left(\frac{\pi}{2} + 2k\pi\right)$.
2. $\cos z = 3 \implies \frac{e^{iz} + e^{-iz}}{2} = 3$. Thus upon rearrangement, we seek to solve:

$$e^{2iz} - 6e^{iz} + 1 = 0 \implies e^{iz} = \frac{6 \pm \sqrt{36 - 4}}{2} = 3 \pm 2\sqrt{2}$$

Thus $iz = \log(3 \pm 2\sqrt{2}) = \ln |3 \pm 2\sqrt{2}| + i(\arg(3 \pm 2\sqrt{2})) \implies z = -i \ln 3 \pm 2\sqrt{2} + \arg(3 \pm 2\sqrt{2})$.

3. $\cosh z = -4 \implies e^z + e^{-z} = -8 \implies e^{2z} + 8e^z + 1 = 0$. Hence by Quadratic formula, we obtain:

$$e^z = \frac{-8 \pm \sqrt{64 - 4}}{2} = -4 \pm \sqrt{15} \implies z = \ln|-4 \pm \sqrt{15}| + i\arg(-4 \pm \sqrt{15})$$

Example 10

We have $\sin z = i \cos z \implies \sin^2 z = -\cos^2 z \implies \sin^2 z + \cos^2 z = 0$ but this is not valid since we know $\sin^2 z + \cos^2 z \geq 1$.

Example 11

1. Using the principal argument of $\frac{1+\sqrt{3}i}{2} = e^{\frac{\pi}{3}i}$, we can simplify the inside expression to $e^{i\pi}$.

$$\text{pv}(e^{i\pi})^{1-i} = \exp((1-i)\text{Log}(e^{i\pi})) = \exp((1-i)(0+i\pi)) = \exp(i\pi + \pi) = -e^\pi$$

2. $i^i = \exp(i \log i) = \exp(i(0 + i(\frac{\pi}{2} + 2k\pi))) = \exp(-\frac{\pi}{2} - 2k\pi)$, where $k \in \mathbb{Z}$.

3.

$$\lim_{z \rightarrow 0} \exp\left(\frac{1}{z^2} \text{Log}(\cos z)\right) = \exp\left(\lim_{z \rightarrow 0} \frac{\text{Log}(\cos z)}{z^2}\right) = e^{-\frac{1}{2}}$$

Upon using L'Hopital's rule twice.

Example 12

- Note that Log is analytic everywhere except along the negative real axis. So the function $\text{Log}(iz)$ is analytic everywhere except along the positive imaginary axis. (You can get this geometrically).
- Since the limit as $z \rightarrow 0$ doesn't exist, we can discount that. $\text{Log}(z+1)$ is not analytic for $z+1 = x \in \mathbb{R}^-$ so we have $z \in (-\infty, -1]$. Hence g is analytic everywhere except $(-\infty, -1] \cup \{0\} \subseteq \mathbb{R}$.

Example 13

- Note that: $f(z) = \exp(\frac{1}{2}\text{Log}(z+1))$, so the function is analytic everywhere except $z \in (-\infty, -1]$.

2. $f(z) = \exp(\frac{1}{2}\text{Log}(z^2 - 1))$ is analytic except on $z^2 - 1 = x$ for $x \leq 0$, and hence $z = \pm i\sqrt{-x + 1}, x \leq 0$.

Example 14

Solving for z , we obtain:

$$z = \frac{2w}{1 - w}$$

Substituting into the region that we had originally obtained, we simplify the requirement down to:

$$|3w - 1| \leq |w - 1|$$

Which is a circle. One may brute force the rest by substituting $w = x + iy$ and squaring both sides of the inequality to obtain an equation.

Example 15

Let the complex number on the line be given by $z = (2 - 2y) + iy$. Then 3 points on the line are $z = 2, i, 4 - i$, which maps to the points $w = \frac{2-i}{5}, -\frac{i}{2}, \frac{1}{4}$. By constructing perpendicular bisectors to 2 of the lines and finding their intersection point, we arrive at the centre $\frac{1}{8} - \frac{1}{4}i$, and radius of $\frac{\sqrt{5}}{8}$.