



MATH1081 S1 2008 Test 3 v3A

August 19, 2017

These solutions were written by Johann Blanco, typed up by Evgeny Martynov and edited by Allan Loi, Henderson Koh and Aaron Hassan. Please be ethical with this resource. It is for the use of MathSoc members, so do not repost it on other forums or groups without asking for permission. If you appreciate this resource, please consider supporting us by coming to our events and buying our T-shirts! Also, happy studying 😊!

We cannot guarantee that our working is correct, or that it would obtain full marks – please notify us of any errors or typos at unswmathsoc@gmail.com, or on our Facebook page. There are sometimes multiple methods of solving the same question. Remember that in the real class test, you will be expected to explain your steps and working out.

1. (a) The truth tables are given below for each formula (make sure to remember your truth tables for basic things like $P \rightarrow Q$ and $P \wedge Q$ etc.):

$$\sim p \rightarrow (q \wedge p)$$

p	q	$\sim p$	$q \wedge p$	$\sim p \rightarrow (q \wedge p)$
T	T	F	T	T
T	F	F	F	T
F	T	T	F	F
F	F	T	F	F

$$(p \wedge q) \rightarrow q$$

p	q	$p \wedge q$	$(p \wedge q) \rightarrow q$
T	T	T	T
T	F	F	T
F	T	F	T
F	F	F	T

(b) First, it will be easier to see if you produce a combined truth table:

p	q	$\sim p \rightarrow (q \wedge p)$	$(p \wedge q) \rightarrow q$	first \rightarrow second	second \rightarrow first
T	T	T	T	T	T
T	F	T	T	T	T
F	T	F	T	T	F
F	F	F	T	T	F

Recall that for a formula to logically imply another formula, one only needs to look at each row where the first formula produces a T. If every such row also produces a T in the second formula, then this proves that the first formula logically implies the second.

By looking at the truth table, we can see that every time the first formula produces a T (the first two rows), the second formula also produces a T. Hence, the first formula logically implies the second.

On the other hand, to check whether the second formula logically implies the first, we need to look at the rows where the second formula produces a T (every line). However, consider the case where $p = q = F$ (4th line). While the second formula produces a T, the first formula produces a F. Hence, the second formula does **not** logically imply the first.

2. **Theorem:** $\log_6 11$ is irrational.

Proof. We prove by contradiction.

Suppose that $\log_6 11$ is rational, that is, there exist integers p and q such that $\log_6 11 = \frac{p}{q}$. Without loss of generality, we can assume $p, q > 0$.

Then,

$$\begin{aligned}\log_6 11 &= \frac{p}{q} \\ 11 &= 6^{\frac{p}{q}} \\ 11^q &= 6^p\end{aligned}$$

which is a contradiction, as the LHS is always odd and RHS is always even.

Hence, our assumption was false, and it follows that $\log_6 11$ is irrational, thus completing the proof. \square

3. **Theorem:** $q(n) = 11n^2 + 32n$ is a prime number for two integer values of n , and is composite for all other integer values of n .

Proof. First, note that if a number is prime, then it has exactly two factors, which are 1 and the number itself.

Factorising $q(n)$, we obtain $q(n) = n(11n + 32)$. From the definition, for $q(n)$ to be prime, it is necessary that at least one of $n = \pm 1$ or $11n + 32 = \pm 1$ is true.

We now consider the possible cases.

If $n = 1$, then $q(n) = 43$, which is prime.

If $n = -1$, then $q(n) = -21$, which is not prime.

If $11n + 32 = 1$, then $n = \frac{-31}{11} \notin \mathbb{Z}$. This contradicts the constraint that $n \in \mathbb{Z}$ and hence is not possible.

If $11n + 32 = -1$, then $n = -3$, and hence $q(n) = 3$, which is prime.

By exhaustion of all possible cases where $q(n)$ can be prime, we conclude that $q(n)$ is prime in only two cases (namely, $n = 1$ or $n = -3$). For all other values of n , $q(n)$ is divisible by n and $11n + 32$, which are both not equal to 1, implying that it is composite. \square





MATH1081 S2 2008 Test 3 v1A

August 19, 2017

These solutions were written by Johann Blanco, typed up by Evgeny Martynov and edited by Allan Loi, Henderson Koh and Aaron Hassan. Please be ethical with this resource. It is for the use of MathSoc members, so do not repost it on other forums or groups without asking for permission. If you appreciate this resource, please consider supporting us by coming to our events and buying our T-shirts! Also, happy studying 😊!

We cannot guarantee that our working is correct, or that it would obtain full marks – please notify us of any errors or typos at unswmathsoc@gmail.com, or on our Facebook page. There are sometimes multiple methods of solving the same question. Remember that in the real class test, you will be expected to explain your steps and working out.

-
1. (a) The truth tables for each formula are given below:

$$(p \rightarrow (\sim q)) \wedge r$$

p	q	r	$\sim q$	$p \rightarrow (\sim q)$	$(p \rightarrow (\sim q)) \wedge r$
T	T	T	F	F	F
T	T	F	F	F	F
T	F	T	T	T	T
T	F	F	T	T	F
F	T	T	F	T	T
F	T	F	F	T	F
F	F	T	T	T	T
F	F	F	T	T	F

$$q \rightarrow ((\sim p) \wedge r)$$

p	q	r	$\sim p$	$(\sim p) \wedge r$	$q \rightarrow ((\sim p) \wedge r)$
T	T	T	F	F	F
T	T	F	F	F	F
T	F	T	F	F	T
T	F	F	F	F	T
F	T	T	T	T	T
F	T	F	T	F	F
F	F	T	T	T	T
F	F	F	T	F	T

(b) First off, note that the second cannot imply the first when $p = q = r = F$.

On the other hand, the first **does** imply the second, which can be seen from this truth table:

p	q	r	$(p \rightarrow (\sim q)) \wedge r$	$q \rightarrow ((\sim p) \wedge r)$	first \rightarrow second
T	T	T	F	F	T
T	T	F	F	F	T
T	F	T	T	T	T
T	F	F	F	T	T
F	T	T	T	T	T
F	T	F	F	F	T
F	F	T	T	T	T
F	F	F	F	T	T

Hence, the first formula logically implies the second.

2. **Theorem :** If m and n are positive integers, then $m!n! < (m+n)!$

Proof. Let m and n be positive integers.

Now,

$$\begin{aligned}
 m!n! &= (1 \times 2 \times 3 \times \cdots \times m)(1 \times 2 \times 3 \times \cdots \times n) \\
 &< (1 \times 2 \times 3 \times \cdots \times m)(m+1)(m+2)(m+3) \cdots (m+n) \\
 &= (m+n)!
 \end{aligned}$$

as $1 < m+1, 2 < m+2, \dots, n < m+n$.

It follows that $m!n! < (m+n)!$, thus completing the proof. \square

Alternative proof. Let m and n be positive integers. Notice that $\binom{m+n}{m} > 1$ since $m, n \in \mathbb{Z}^+$ so $m, n > 0$.

This is true since this finds us the total number of ways to select m items from a total of

$m + n$ items which is clearly greater than 1 (since $m + n$ is strictly greater than n).
Therefore,

$$\begin{aligned}\binom{m+n}{m} &> 1 \\ \Rightarrow \frac{(m+n)!}{m!n!} &> 1 \\ \Rightarrow m!n! &< (m+n)!,\end{aligned}$$

which proves the result.

3. **Theorem:** If $a \in \mathbb{R}^+$, then the equation $ax = \cos \pi x$ has exactly one solution x such that $0 \leq x \leq 1$.

Proof. Let a be some element of \mathbb{R}^+ . Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f(x) = ax - \cos \pi x$.

Now, $f(0) = -1$ and $f(1) = a + 1$.

Clearly, f is continuous on the interval $[0, 1]$. Thus, as $-1 < 0 < a + 1$, by the intermediate value theorem, there exists at least one number $c \in (0, 1)$ such that $f(c) = 0$.

That is, there is *at least one* solution for $ax = \cos \pi x$ on the specified interval.

Further, on the interval $[0, 1]$,

$$f'(x) = a + \pi \sin \pi x > 0$$

that is, f is monotone increasing, and so it can have at most one root in that interval.

It follows that $ax = \cos \pi x$ has exactly one solution on the interval $[0, 1]$, thus completing the proof. \square

More rigorous proof for the uniqueness condition by contradiction. Suppose that the equation $ax = \cos \pi x$ has two distinct solutions for x in the interval $[0, 1]$. Then f has two distinct roots in the interval $[0, 1]$. Let these roots be x_1 and x_2 .

Since f is continuous and differentiable over the interval $[0, 1]$, f is also continuous over the closed interval $[x_1, x_2]$ and differentiable over the open interval (x_1, x_2) , as $[x_1, x_2]$ and (x_1, x_2) are subsets of $[0, 1]$. By applying Rolle's theorem, there must exist some y in the open interval (x_1, x_2) such that $f'(y) = 0$.

By differentiating the function, we have

$$f'(y) = a + \pi \sin \pi y = 0.$$

However, a and π are positive and $\sin \pi y$ is non-negative for all possible values of y

in $[0, 1]$. This implies that f' is non-zero over the interval $[0, 1]$, which contradicts our assumption that the equation $ax = \cos \pi x$ has two distinct solutions for x in the interval $[0, 1]$. Thus, the equation can only have exactly one solution for x in the interval $[0, 1]$ and this completes the proof.





MATH1081 S1 2009 Test 3 v2A

August 19, 2017

These solutions were written by Johann Blanco, typed up by Evgeny Martynov and edited by Henderson Koh and Aaron Hassan. Please be ethical with this resource. It is for the use of MathSoc members, so do not repost it on other forums or groups without asking for permission. If you appreciate this resource, please consider supporting us by coming to our events and buying our T-shirts! Also, happy studying 😊!

We cannot guarantee that our working is correct or that it would obtain full marks – please notify us of any errors or typos at unswmathsoc@gmail.com, or on our Facebook page. There are sometimes multiple methods of solving the same question. Remember that in the real class test, you will be expected to explain your steps and working out.

1. We have

$$\begin{aligned}
 & p \rightarrow (\sim (q \wedge (\sim p))) \\
 & \Leftrightarrow (\sim p) \vee (\sim (q \wedge (\sim p))) && (\text{as } u \rightarrow v \Leftrightarrow (\sim u) \vee v) \\
 & \Leftrightarrow \sim (p \wedge (q \wedge (\sim p))) && (\text{De Morgan's Law}) \\
 & \Leftrightarrow \sim (p \wedge ((\sim p) \wedge q)) && (\text{Commutative Law}) \\
 & \Leftrightarrow \sim ((p \wedge (\sim p)) \wedge q) && (\text{Associative Law}) \\
 & \Leftrightarrow \sim (\mathbf{c} \wedge q) && (\text{Law of Negation}) \\
 & \Leftrightarrow \sim (\mathbf{c}) && (\text{Domination Law}) \\
 & \Leftrightarrow \mathbf{t}. && (\text{Negation of a Contradiction})
 \end{aligned}$$

2. **Theorem:** If $n \in \mathbb{Z}^+$, then

$$(1 \times 2) + (2 \times 5) + \cdots + n(3n - 1) = n^2(n + 1)$$

Proof. We prove by induction.

When $n = 1$, LHS $= 1 \times 2 = 2$, RHS $= 1^2 \times 2 = 2$, and so the theorem is true for $n = 1$.

Assume the result is true for some particular integer $n \in \mathbb{Z}^+$, that is

$$(1 \times 2) + (2 \times 5) + \cdots + n(3n - 1) = n^2(n + 1).$$

It is required to prove that the result is also true for $n + 1$, that is

$$(1 \times 2) + (2 \times 5) + \cdots + (n + 1)(3n + 2) = (n + 1)^2(n + 2).$$

We have

$$\begin{aligned} \text{LHS} &= (1 \times 2) + \cdots + n(3n - 1) + (n + 1)(3n + 2) && \text{(explicitly writing the second last term)} \\ &= n^2(n + 1) + (n + 1)(3n + 2) && \text{(by the inductive hypothesis)} \\ &= (n + 1)(n^2 + 3n + 2) && \text{(factoring out } (n + 1)) \\ &= (n + 1)(n + 1)(n + 2) && \text{(factorising)} \\ &= (n + 1)^2(n + 2) \\ &= \text{RHS.} \end{aligned}$$

This proves the theorem by induction. □

3. **Theorem:** If $x \in \mathbb{R}$ and $2x^2 - 3 = 0$, then x is irrational.

Proof. We prove by contradiction.

Let $x \in \mathbb{R}$ and $2x^2 - 3 = 0$. Solving, we obtain

$$x^2 = \frac{3}{2}.$$

Suppose that x is rational, that is, it can be written as an irreducible ratio of two integers, $x = \frac{p}{q}$ for some $p, q \in \mathbb{Z} \setminus \{0\}$.

Then, $x^2 = \frac{p^2}{q^2} = \frac{3}{2}$ and thus

$$2p^2 = 3q^2. \tag{1}$$

Clearly, the LHS is even, and so q^2 must be even also. This implies that q is even, and hence $q = 2r$ for some $r \in \mathbb{Z}$.

Substituting this result into (1) and simplifying gives

$$p^2 = 2 \times 3 \times r^2.$$

This shows that p^2 is even, which implies that p must be even. Hence, p and q share a common factor of 2. But since we assumed that p and q form an irreducible fraction, this is a contradiction.

Thus, the initial assumption was wrong.

It follows that x is irrational, and the proof is complete.

□



MATH1081 S2 2009 Test 3 v1A

August 19, 2017

These answers were written by Johann Blanco, typed up by Evgeny Martynov and edited by Henderson Koh and Aaron Hassan. Please be ethical with this resource. It is for the use of MathSoc members, so do not repost it on other forums or groups without asking for permission. If you appreciate this resource, please consider supporting us by coming to our events and buying our T-shirts! Also, happy studying 😊!

We cannot guarantee that our answers are correct, or would obtain full marks – please notify us of any errors or typos at unswmathsoc@gmail.com, or on our Facebook page. There are sometimes multiple methods of solving the same question. Remember that in the real class test, you will be expected to explain your steps and working out.

1. (i) Let

$m = \text{"I earn some money"}$

$h = \text{"I go for a holiday this summer"}$

$w = \text{"I work this summer"}$.

The argument in symbolic form is:

$$\frac{\begin{array}{c} m \rightarrow h \\ h \vee w \end{array}}{\therefore (\sim h) \rightarrow (\sim m) \wedge w}.$$

(ii) Using a truth table, we consider critical rows where the hypotheses are true:

m	h	$\sim h$	w	$m \rightarrow h$	$h \vee w$	$(\sim m) \wedge w$	$(\sim h) \rightarrow (\sim m) \wedge w$
T	T	F	T	T	T	F	T
T	T	F	F	T	T	F	T
T	F	T	T	F	*	*	*
T	F	T	F	F	*	*	*
F	T	F	T	T	T	T	T
F	T	F	F	T	T	F	T
F	F	T	T	T	T	T	T
F	F	T	F	T	F	*	*

Note that the rows we disregard are the ones where at least one of the statements $m \rightarrow h$ and $h \vee w$ are not true, because these are not relevant to our aim. We have highlighted in **green** the rows where the hypotheses are both true and the conclusions in those rows, which we see are also true.

Since we find that the conclusion is true whenever the hypotheses are true, the above argument is logically valid.

2. **Theorem :** Between any two different rational numbers there is another rational number.

Proof. Let x and y be two distinct rational numbers. Suppose without loss of generality that $x < y$.

We claim that $\frac{x+y}{2}$ is a rational number between x and y .

First, we prove that $\frac{x+y}{2}$ is indeed rational. Since x and y are rational, we can write $x = \frac{a}{b}$ and $y = \frac{c}{d}$, where a, b, c and d are integers and $b, d \neq 0$. So, we have

$$\frac{x+y}{2} = \frac{\frac{a}{b} + \frac{c}{d}}{2} = \frac{ad+bc}{2bd},$$

where $ad+bc$ and $2bd$ are integers. As $\frac{x+y}{2}$ can be expressed as a fraction of two integers where $2bd \neq 0$, then it is indeed rational. Now we show that $\frac{x+y}{2}$ is between x and y , that is, $x < \frac{x+y}{2}$ and $\frac{x+y}{2} < y$. We have

$$\begin{aligned} \frac{x+y}{2} - x &= \frac{x+y-2x}{2} \\ &= \frac{y-x}{2} \\ &> 0 && (\text{as } x < y) \\ \Rightarrow x &< \frac{x+y}{2}, \end{aligned}$$

and

$$\begin{aligned}
 y - \frac{x+y}{2} &= \frac{2y - x - y}{2} \\
 &= \frac{y-x}{2} \\
 &> 0 \quad (\text{as } x < y) \\
 \Rightarrow \frac{x+y}{2} &< y.
 \end{aligned}$$

This shows that $\frac{x+y}{2}$ is indeed a rational number between x and y , which proves the theorem. \square

3. **Theorem:** Prove that if n is a positive integer then $4^{2n} + 10n - 1$ is a multiple of 25.

Proof. We prove by induction.

For $n = 1$, $4^{2n} + 10n - 1$ is equal to 25, which is indeed a multiple of 25.

Now, assume that the theorem holds for some positive integer $n = k$, that is $4^{2k} + 10k - 1$ is a multiple of 25, or $4^{2k} = 25m - 10k + 1$ for some integer m .

We will try to prove that the theorem holds when $n = k + 1$, that is $4^{2(k+1)} + 10(k+1) - 1$ is a multiple of 25.

We have

$$\begin{aligned}
 4^{2(k+1)} + 10(k+1) - 1 &= 4^{2k+2} + 10k + 9 \\
 &= 16 \times 4^{2k} + 10k + 9 \\
 &= 16 \times (25m - 10k + 1) + 10k + 9 \quad (\text{induction hypothesis}) \\
 &= 16 \times 25m - 150k + 25 \\
 &= 25(16m - 6k + 1),
 \end{aligned}$$

which is divisible by 25, as $16m - 6k + 1$ is an integer.

Hence, we have proved that the theorem holds when $n = k+1$, whenever it is true for $n = k$ for some integer k . And so, the theorem holds for all positive integers, by mathematical induction. \square



MATH1081 S1 2010 Test 3 v2B

August 19, 2017

These answers were written by Johann Blanco, typed up by Evgeny Martynov and Allan Loi, and edited by Aaron Hassan. Please be ethical with this resource. It is for the use of MathSoc members, so do not repost it on other forums or groups without asking for permission. If you appreciate this resource, please consider supporting us by coming to our events and buying our T-shirts! Also, happy studying 🧐!

We cannot guarantee that our answers are correct or would obtain full marks – please notify us of any errors or typos at unswmathsoc@gmail.com, or on our Facebook page. There are sometimes multiple methods of solving the same question. Remember that in the real class test, you will be expected to explain your steps and working out.

1. We have

$$\begin{aligned}
 & (p \rightarrow q) \wedge (q \rightarrow (\sim p \vee r)) \\
 \Leftrightarrow & ((\sim p) \vee q) \wedge ((\sim q) \vee (\sim p \vee r)) && (\text{as } u \rightarrow v \Leftrightarrow (\sim u) \vee v) \\
 \Leftrightarrow & ((\sim p) \vee q) \wedge ((\sim p) \vee (\sim q \vee r)) && (\text{Associative and Commutative Laws}) \\
 \Leftrightarrow & (\sim p) \vee (q \wedge (\sim q \vee r)) && (\text{Distributive Law}) \\
 \Leftrightarrow & (\sim p) \vee ((q \wedge \sim q) \vee (q \wedge r)) && (\text{Distributive Law}) \\
 \Leftrightarrow & (\sim p) \vee (\mathbf{F} \vee (q \wedge r)) && (\text{Law of Negation}) \\
 \Leftrightarrow & (\sim p) \vee (q \wedge r) && (\text{Identity Law}) \\
 \Leftrightarrow & p \rightarrow (q \wedge r) && (\text{as } u \rightarrow v \Leftrightarrow (\sim u) \vee v)
 \end{aligned}$$

as required.

2. **Theorem:** If n is a positive integer then $(n+1)(n+2)\dots(2n) = 2^n \times 1 \times 3 \times 5 \times \dots \times (2n-1)$.

Proof. We prove this by induction. When $n = 1$, $LHS = 2$ and $RHS = 2$. So the theorem holds for $n = 1$.

Assume now that the theorem holds for some particular integer k , that is

$$(k+1)(k+2)\dots(2k) = 2^k \times 1 \times 3 \times 5 \times \dots \times (2k-1).$$

We will try to prove that the theorem also holds for $n = k+1$, that is

$$(k+2)(k+3)\dots(2(k+1)) = 2^{k+1} \times 1 \times 3 \times 5 \times \dots \times (2k+1).$$

We have

$$\begin{aligned} LHS &= (k+2)(k+3)\dots(2k)(2k+1)(2k+2) \\ &= \frac{(2k+2)(2k+1)}{k+1} \times (k+1)(k+2)\dots(2k) \\ &= \frac{(2k+2)(2k+1)}{k+1} \times 2^k \times 1 \times 3 \times 5 \times \dots \times (2k-1) \\ &\quad \text{(by the inductive hypothesis)} \\ &= \frac{2(k+1)(2k+1)}{k+1} \times 2^k \times 1 \times 3 \times 5 \times \dots \times (2k-1) \\ &= 2^{k+1} \times 1 \times 3 \times 5 \times \dots \times (2k-1) \times (2k+1) \\ &= RHS. \end{aligned}$$

So, we have just proven that the theorem holds for $n = k+1$, whenever it is true for $n = k$. Hence, the theorem holds for all positive integers, by mathematical induction. \square

3. **HINT:** Use a proof by contradiction by first letting $\sqrt{4n-2} = \frac{p}{q}$ where $p, q \in \mathbb{Z}^+$ and are coprime. Here is a sample solution.

Claim: If n is any positive integer then $\sqrt{4n-2}$ is irrational.

Proof. We will prove by contradiction.

Let n be a positive integer. Suppose $\sqrt{4n-2}$ is rational.

Then there exist coprime integers p and $q \neq 0$, such that $\sqrt{4n-2} = \frac{p}{q}$. Squaring both sides yields $4n-2 = \frac{p^2}{q^2}$. Upon multiplying both sides by q^2 , we then get

$$2(2n-1)q^2 = p^2. \tag{1}$$

Since, the left hand side has a factor of 2, then it is even. This implies that p^2 must also

be even. One can then deduce that p is also even, that is there exists some integer r such that $p = 2r$.

Substituting this back into (1) yields

$$2(2n - 1)q^2 = 4r^2 \Leftrightarrow (2n - 1)q^2 = 2r^2.$$

Now, with a similar argument as before, the right hand side is even, which implies that the left hand side must also be even. However, as $2n - 1$ is odd, the only way that the left hand side can be even, is if q^2 is even, which implies that q is even.

However, we assumed that p and q are coprime, which is a contradiction. So our original assumption was incorrect. And so, $\sqrt{4n - 2}$ is indeed irrational. \square

