

UNSW Mathematics Society Presents  
**MATH1081 Workshop**



**Presented by Fred Yan, Ronald Chiang, and Jay Liang**

# Overview I

1. Sets and Functions
2. Integers, Modular Arithmetic and Relations
3. Enumeration and Probability
4. Logic and Proofs
5. Graphs

# 1. Sets and Functions

# 2018 T1 Q1 (i)

- (a) Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .
- (b) Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

# 2018 T1 Q1 (i)

- (a) Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .
- (b) Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

## Size of power sets

The cardinality of a power set  $|P(A)|$  is given by  $2^{|A|}$ .

# 2018 T1 Q1 (i)

- (a) Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .
- (b) Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

## Size of power sets

The cardinality of a power set  $|P(A)|$  is given by  $2^{|A|}$ .

## Size of cartesian product

The cardinality of a cartesian product  $|A \times B|$  is given by  $|A| \times |B|$ .

## a) Solution

Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .

Using  $|X \times Y| = |X| \times |Y|$ , we get

## a) Solution

Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .

Using  $|X \times Y| = |X| \times |Y|$ , we get

$$|P(A) \times P(B)| = |P(A)| \times |P(B)|$$

## a) Solution

Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .

Using  $|X \times Y| = |X| \times |Y|$ , we get

$$\begin{aligned}|P(A) \times P(B)| &= |P(A)| \times |P(B)| \\&= 2^{|A|} \times 2^{|B|}\end{aligned}$$

## a) Solution

Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .

Using  $|X \times Y| = |X| \times |Y|$ , we get

$$\begin{aligned}|P(A) \times P(B)| &= |P(A)| \times |P(B)| \\&= 2^{|A|} \times 2^{|B|} \\&= 2^2 \times 2^3 \\&= 32.\end{aligned}$$

## a) Solution

Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .

Similarly,

$$|P(A \times B)| = 2^{|A \times B|}$$

## a) Solution

Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .

Similarly,

$$\begin{aligned}|P(A \times B)| &= 2^{|A \times B|} \\&= 2^{|A| \times |B|}\end{aligned}$$

## a) Solution

Suppose that  $A$  and  $B$  are sets with  $|A| = 2$  and  $|B| = 3$ .  
Find  $|P(A) \times P(B)|$  and  $|P(A \times B)|$ .

Similarly,

$$\begin{aligned}|P(A \times B)| &= 2^{|A \times B|} \\&= 2^{|A| \times |B|} \\&= 2^{2 \times 3} \\&= 64.\end{aligned}$$

## b) Solution\*

Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

$$\begin{aligned} 2^{|X|+|Y|} &\leq 2^{|X| \times |Y|} \\ |X| + |Y| &\leq |X| \times |Y| \\ |X| = |Y| &= 1 \\ \text{LHS} = 1 + 1 &= 2 \quad \text{RHS} = 1^1 \\ \text{LHS} &\leq \text{RHS} = 1 \end{aligned}$$

## b) Solution\*

Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

Statement is false by counter-example.

$$\begin{aligned} |X| = |Y| &= 1 \\ \text{LHS} &= |P(X) \times P(Y)| \\ &= |P(X)| \times |P(Y)| \end{aligned}$$

## b) Solution\*

Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

Statement is false by counter-example.

$$|X| = |Y| = 1$$

$$\begin{aligned} \text{LHS} &= |P(X) \times P(Y)| \\ &= |P(X)| \times |P(Y)| \\ &= 2^{|X|} \times 2^{|Y|} \end{aligned}$$

$$2^{|X| \times |Y|}$$

$$= 4$$

## b) Solution\*

Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

Statement is false by counter-example.

$$\begin{aligned} \text{RHS} &= |P(X \times Y)| \\ &= 2^{|X \times Y|} \\ &= 2^{|X| \times |Y|} \end{aligned}$$

$\uparrow = 2^{|X|}$   
 $= 2$   
 $LHS = 4$   
 $RHS = 2$   
 $LHS > RHS$

## b) Solution\*

Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

Investigation:

$$\begin{aligned} |P(X) \times P(Y)| \leq |P(X \times Y)| &\iff |P(X)| \times |P(Y)| \leq 2^{|X \times Y|} \\ &\iff 2^{|X|} \times 2^{|Y|} \leq 2^{|X| \times |Y|} \\ &\iff 2^{|X| + |Y|} \leq 2^{|X| \times |Y|} \\ &\iff |X| + |Y| \leq |X| \times |Y|. \end{aligned}$$

What happens when  $|X| = |Y| = 1$ ?

## b) Proof

Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

The statement is false and we'll provide a counterexample:

Let  $|X| = |Y| = 1$ . Then we have

$$\begin{aligned}|P(X) \times P(Y)| &= |P(X)| \times |P(Y)| \\&= 2^{|X|} \times 2^{|Y|} \\&= 2^{|X|+|Y|} \\&= 2^{1+1} \\&= 4.\end{aligned}$$

## b) Proof

Is it true that

$$|P(X) \times P(Y)| \leq |P(X \times Y)|$$

for all non-empty finite sets  $X, Y$ ? Explain your answer.

$$\begin{aligned}|P(X \times Y)| &= 2^{|X \times Y|} \\&= 2^{|X| \times |Y|} \\&= 2^{|X| \times |Y|} \\&= |X| \times |Y| \\&= 1 \times 1 \\&= 1.\end{aligned}$$

Now since  $4 > 1$ , then we have  $|P(X) \times P(Y)| > |P(X \times Y)|$  and the statement is false.  $\square$

## 2017 T1 Q1 (ii)\*

Use the laws of set algebra to simplify  $(A \cap (\overline{A} \cap B)^c) \cup B^c$ .

$$\begin{aligned}&\equiv (A \cap (\overline{A} \cup \overline{B})) \cup B^c && \text{(De Morgan)} \\&\equiv (A \cap \overline{A}) \cup (A \cap \overline{B}) \cup B^c \\&\equiv \emptyset \cup (A \cap B^c) \cup B^c && \text{(Distributivity)} \\&\equiv (A \cap B^c) \cup B^c && \text{(Intersection)} \\&\equiv (A \cap B^c) \cup \emptyset && \text{(Identity)}\end{aligned}$$

# 2017 T1 Q1 (ii)\*

Use the laws of set algebra to simplify  $(A \cap (A \cap B)^c) \cup B^c$ .

$$\equiv (A \cap B^c) \cup B^c$$

$$\equiv A^c \cup B^c \quad (\text{Absorption})$$



## 2017 T1 Q1 (ii)\*

Use the laws of set algebra to simplify  $(A \cap (A \cap B)^c) \cup B^c$ .

$$\begin{aligned}(A \cap (A \cap B)^c) \cup B^c &\equiv (A \cap (A^c \cup B^c)) \cup B^c \text{ (De Morgan's Law)} \\&\equiv (A \cap A^c) \cup (A \cap B^c) \cup B^c \text{ (Distributive Law)} \\&\equiv \emptyset \cup (A \cap B^c) \cup B^c \text{ (Intersection w/ complement)} \\&\equiv (A \cap B^c) \cup B^c \text{ (Identity Law)} \\&\equiv B^c \text{ (Absorption Law)}\end{aligned}$$

## 2016 S2 Q1 (v)

- If  $X$  and  $Y$  are sets with  $|X| = 2$  and  $|Y| = n$ , how many injective functions are there from  $X$  to  $Y$ ? Explain your answer.
- If  $|X| = n$  and  $|Y| = 2$ , how many surjective functions are there from  $X$  to  $Y$ ? Explain your answer.

# 2016 S2 Q1 (v)

- If  $X$  and  $Y$  are sets with  $|X| = 2$  and  $|Y| = n$ , how many injective functions are there from  $X$  to  $Y$ ? Explain your answer.
- If  $|X| = n$  and  $|Y| = 2$ , how many surjective functions are there from  $X$  to  $Y$ ? Explain your answer.

## Injective Function Definition

A function  $f : A \rightarrow B$  is injective if and only if for any  $x_1$  and  $x_2$  in  $A$ ,  $f(x_1) = f(x_2)$  implies that  $x_1 = x_2$ .

In other words, if the outputs are equal then the inputs are equal.

## 2016 S2 Q1 (v)

- If  $X$  and  $Y$  are sets with  $|X| = 2$  and  $|Y| = n$ , how many injective functions are there from  $X$  to  $Y$ ? Explain your answer.
- If  $|X| = n$  and  $|Y| = 2$ , how many surjective functions are there from  $X$  to  $Y$ ? Explain your answer.

### Injective Function Definition

A function  $f : A \rightarrow B$  is injective if and only if for any  $x_1$  and  $x_2$  in  $A$ ,  $f(x_1) = f(x_2)$  implies that  $x_1 = x_2$ .

In other words, if the outputs are equal then the inputs are equal.


### Surjective Function Definition

A function  $f : A \rightarrow B$  is surjective if and only if for any  $y$  in  $B$ , there exists  $x$  in  $A$  such that  $f(x) = y$ .

In other words, every element in the codomain is a function value, i.e.  $f$  of something in  $A$ .

# a) Solution\*


If  $X$  and  $Y$  are sets with  $|X| = 2$  and  $|Y| = n$ , how many injective functions are there from  $X$  to  $Y$ ? Explain your answer.



$$\begin{aligned} & \text{~} \\ & n! \\ & = n(n-1) \end{aligned}$$

## b) Solution\*

If  $|X| = n$  and  $|Y| = 2$ , how many surjective functions are there from  $X$  to  $Y$ ? Explain your answer.



2 cases where  
f is not surjective

$2 \times 2 \times \dots \times 2$

no. of surjective functions  
 $= 2^n - 2$

# Inverse Functions\*

## Theorem

A function  $g : Y \rightarrow X$  is an inverse of  $f : X \rightarrow Y$  if and only if

$$g \circ f = i_x \text{ and } f \circ g = i_y.$$

$$(g \circ f)(x) = x, \forall x \in X$$

$$(f \circ g)(y) = y, \forall y \in Y$$

2017 T1 Q1 (iv)

$$(g \circ f)(x) = g(f(x))$$

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
- Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
- Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- 
- Since the codomain of  $f$  is a subset of the domain of  $g$ ,  $g \circ f$  is defined.  $\text{Dom}(g \circ f) = [1, \infty)$ .  $\text{Codom}(g \circ f) = \mathbb{R}$ .

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- a) Write down the domain and codomain of  $g \circ f$ .
- b) Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
- c) Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.

$$(g \circ f)(x) = x$$

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- b) We know that  $g \circ f : [1, \infty) \rightarrow \mathbb{R}$  is defined. So

$$(g \circ f)(x) = g(f(x))$$

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- b) We know that  $g \circ f : [1, \infty) \rightarrow \mathbb{R}$  is defined. So

$$\begin{aligned}(g \circ f)(x) &= g(f(x)) \\ &= g(\ln x + 1)\end{aligned}$$

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- b) We know that  $g \circ f : [1, \infty) \rightarrow \mathbb{R}$  is defined. So

$$\begin{aligned}(g \circ f)(x) &= g(f(x)) \\&= g(\ln x + 1) \\&= e^{\ln x+1-1}\end{aligned}$$

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- b) We know that  $g \circ f : [1, \infty) \rightarrow \mathbb{R}$  is defined. So

$$\begin{aligned}(g \circ f)(x) &= g(f(x)) \\&= g(\ln x + 1) \\&= e^{\ln x+1-1} \\&= e^{\ln x}\end{aligned}$$

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- b) We know that  $g \circ f : [1, \infty) \rightarrow \mathbb{R}$  is defined. So

$$\begin{aligned}(g \circ f)(x) &= g(f(x)) \\&= g(\ln x + 1) \\&= e^{\ln x+1-1} \\&= e^{\ln x} \\&= x.\end{aligned}$$

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- b) We know that  $g \circ f : [1, \infty) \rightarrow \mathbb{R}$  is defined. So

$$\begin{aligned}(g \circ f)(x) &= g(f(x)) \\&= g(\ln x + 1) \\&= e^{\ln x+1-1} \\&= e^{\ln x} \\&= x.\end{aligned}$$

Since  $(g \circ f)(x) = x$  for  $x \in [1, \infty)$ ,  $g \circ f$  is the identity map on  $[1, \infty)$ .

## 2017 T1 Q1 (iv)

$$\begin{aligned}g \circ f &= i_x \\f \circ g &= i_y\end{aligned}$$

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
- Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
- Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.

2017 T1 Q1 (iv)

Range( $g$ ) =  $(0, \infty)$  Dom( $f$ ) =  $[1, \infty)$

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- c) We have proved in b) that  $g \circ f = i_x$  for all  $x \in [1, \infty)$ . For  $f$  to be the inverse of  $g$ , we also need  $f \circ g = i_y$  for all  $y \in \mathbb{R}$ .

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- a) Write down the domain and codomain of  $g \circ f$ .
  - b) Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - c) Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- c) We have proved in b) that  $g \circ f = i_x$  for all  $x \in [1, \infty)$ . For  $f$  to be the inverse of  $g$ , we also need  $f \circ g = i_y$  for all  $y \in \mathbb{R}$ .

Notice that the range of  $g$  is  $(0, \infty)$  which is not a subset of the domain of  $f$ .

## 2017 T1 Q1 (iv)

Let  $f : [1, \infty) \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \ln x + 1 \quad \text{for all } x \in [1, \infty), \quad g(y) = e^{y-1} \quad \text{for all } y \in \mathbb{R}.$$

- Write down the domain and codomain of  $g \circ f$ .
  - Prove that  $g \circ f$  is the identity map on  $[1, \infty)$ .
  - Is it true that  $f$  is the inverse of  $g$ ? Explain your answer.
- c) We have proved in b) that  $g \circ f = i_x$  for all  $x \in [1, \infty)$ . For  $f$  to be the inverse of  $g$ , we also need  $f \circ g = i_y$  for all  $y \in \mathbb{R}$ .


Notice that the range of  $g$  is  $(0, \infty)$  which is not a subset of the domain of  $f$ .

Hence,  $f \circ g$  is not defined and  $f$  is not the inverse of  $g$ .

## 2018 T2 Q1 (ii)\*

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the function defined by  $f(x, y) = x^2 + y^2$ .

- Suppose that  $S$  is a unit square  $[0, 1] \times [0, 1]$ . Find  $f(S)$ .
- Find and sketch  $f^{-1}([1, 2])$ .




## 2018 T2 Q1 (ii)\*

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the function defined by  $f(x, y) = x^2 + y^2$ .

- Suppose that  $S$  is a unit square  $[0, 1] \times [0, 1]$ . Find  $f(S)$ .
- Find and sketch  $f^{-1}([1, 2])$ .

$$\begin{aligned} & [0, 1] \times \{0\} \\ & [0, 1] \times \{1\} \\ & \{0\} \times [0, 1] \\ & \{1\} \times [0, 1] \end{aligned}$$



## 2018 T2 Q1 (ii)\*

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the function defined by  $f(x, y) = x^2 + y^2$ .

- Suppose that  $S$  is a unit square  $[0, 1] \times [0, 1]$ . Find  $f(S)$ .
- Find and sketch  $f^{-1}([1, 2])$ .

$f(S) = [0, 1]$

1, 2, 3, 4, 5



## 2018 T2 Q1 (ii)\*

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the function defined by  $f(x, y) = x^2 + y^2$ .

- Suppose that  $S$  is a unit square  $[0, 1] \times [0, 1]$ . Find  $f(S)$ .
- Find and sketch  $f^{-1}([1, 2])$ .

a)  $f(S) = [0, 2]$

$$f^{-1}(1), f(x, y) = x^2 + y^2 = 1$$
$$f^{-1}(2), f(x, y) = x^2 + y^2 = 2$$

## 2018 T2 Q1 (ii)\*

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the function defined by  $f(x, y) = x^2 + y^2$ .

- Suppose that  $S$  is a unit square  $[0, 1] \times [0, 1]$ . Find  $f(S)$ .
- Find and sketch  $f^{-1}([1, 2])$ .

a)  $f(S) = [0, 2]$



$$f^{-1}([1, 2]) =$$

$$\{(x, y) \mid 1 \leq x^2 + y^2 \leq 2\}$$


## 2018 T2 Q1 (ii)\*

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the function defined by  $f(x, y) = x^2 + y^2$ .

- Suppose that  $S$  is a unit square  $[0, 1] \times [0, 1]$ . Find  $f(S)$ .
- Find and sketch  $f^{-1}([1, 2])$ .

a)  $f(S) = [0, 2]$

b)  $f(S) = \{(x, y) | 1 \leq x^2 + y^2 \leq 2\}$



A function is bijective if and only if it is both injective and surjective.

A function is bijective if and only if it is both injective and surjective.

Let  $b : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $b(x) = 14x^{13} + 11x^3 + 19x - 16$ .  
 Show that  $b$  is bijective.

Injective


$$b'(x) = 13(14)x^{12} + 33x^2 + 19$$

$$b'(x) > 0, \forall x \in \mathbb{R} \quad > 0$$

$b$  is increasing on  $\mathbb{R}$   
 $b$  is injective.

A function is bijective if and only if it is both injective and surjective.

Let  $b : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $b(x) = 14x^{13} + 11x^3 + 19x - 16$ .  
Show that  $b$  is bijective.



A function is bijective if and only if it is both injective and surjective.

Let  $b : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $b(x) = 14x^{13} + 11x^3 + 19x - 16$ .  
Show that  $b$  is bijective.

$$\lim_{x \rightarrow \infty} b(x) = \infty \quad \lim_{x \rightarrow -\infty} b(x) = -\infty$$

$\forall y \in \mathbb{R}, \exists M > 0$  s.t.

$$x > M \Rightarrow b(x) > y$$

$$x < -M \Rightarrow b(x) < y$$

A function is bijective if and only if it is both injective and surjective.

Let  $b : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $b(x) = 14x^{13} + 11x^3 + 19x - 16$ .  
Show that  $b$  is bijective.

By IVT,  $\exists x_0 \in [n, m]$ ,  
s.t.  $b(x_0) = y, \forall y \in \mathbb{R}$   
 $\text{Range}(b) = \mathbb{R} = \text{codom}(b)$   
Thus,  $b$  is surjective.  
Thus,  $b$  is bijective.

A function is bijective if and only if it is both injective and surjective.

Let  $b : \mathbb{R} \rightarrow \mathbb{R}$  be defined by  $b(x) = 14x^{13} + 11x^3 + 19x - 16$ .  
Show that  $b$  is bijective.

To prove  $b$  is bijective, we have to prove it's both injective and surjective.

Notice  $b'(x) = 182x^{12} + 33x^2 + 19 > 0$ . Since  $b'(x) > 0$  for all  $x \in \mathbb{R}$ ,  $b$  is a strictly increasing function. Hence  $b$  is injective.

Notice that  $b(x) \rightarrow \infty$  as  $x \rightarrow \infty$  and  $b(x) \rightarrow -\infty$  as  $x \rightarrow -\infty$ . Hence, for all  $y \in \mathbb{R}$ , there exists  $M > 0$  such that  $x > M \implies b(x) > y$  and  $x < -M \implies b(x) < y$ . Thus, by the intermediate value theorem, there exists  $x_0 \in [-M, M]$  such that  $b(x_0) = y$  for all  $y \in \mathbb{R}$ .

Hence  $\text{range}(b) = \mathbb{R} = \text{codom}(b)$ . Thus,  $b$  is surjective.

Hence  $b$  is both injective and surjective and thus, bijective.  $\square$

## 2. Integers, Modular Arithmetic and Relations

# 2015 T1 Q2 (i)

•

Evaluate

$$2015^{1082} \pmod{11}.$$

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

We can look at powers of 2015 to find a pattern.

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

We can look at powers of 2015 to find a pattern.  
Let's begin with a power of 1

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

We can look at powers of 2015 to find a pattern.  
Let's begin with a power of 1 where

$$2015^1 \equiv 2 \pmod{11}.$$

## 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

We can look at powers of 2015 to find a pattern.  
Let's begin with a power of 1 where

$$2015^1 \equiv 2 \pmod{11}.$$

Calculator:  $[2015 \div 11] = [-183] = [\times 11] =$

## 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

We can look at powers of 2015 to find a pattern.  
Let's begin with a power of 1 where

$$2015^1 \equiv 2 \pmod{11}.$$

Calculator:  $[2015 \div 11] = [-183] = [\times 11] =$

Now let's try a power of 2 where

$$2015^2 \equiv 2^2 \equiv 4 \pmod{11},$$

## 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

We can look at powers of 2015 to find a pattern.  
Let's begin with a power of 1 where

$$2015^1 \equiv 2 \pmod{11}.$$

Calculator:  $[2015 \div 11] = [-183] = [\times 11] =$

Now let's try a power of 2 where

$$2015^2 \equiv 2^2 \equiv 4 \pmod{11},$$

since, if  $a \equiv b \pmod{m}$ , then  $a^n \equiv b^n \pmod{m}$ .

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

By observing more powers of 2015,

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

By observing more powers of 2015,

$2015^1 \equiv 2 \pmod{11}$	$2015^6 \equiv 9 \pmod{11}$
$2015^2 \equiv 4 \pmod{11}$	$2015^7 \equiv 7 \pmod{11}$
$2015^3 \equiv 8 \pmod{11}$	$2015^8 \equiv 8 \pmod{11}$
$2015^4 \equiv 5 \pmod{11}$	$2015^9 \equiv 6 \pmod{11}$
$2015^5 \equiv 10 \pmod{11}$	$2015^{10} \equiv 1 \pmod{11}$

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

By observing more powers of 2015,

$2015^1 \equiv 2 \pmod{11}$	$2015^6 \equiv 9 \pmod{11}$
$2015^2 \equiv 4 \pmod{11}$	$2015^7 \equiv 7 \pmod{11}$
$2015^3 \equiv 8 \pmod{11}$	$2015^8 \equiv 8 \pmod{11}$
$2015^4 \equiv 5 \pmod{11}$	$2015^9 \equiv 6 \pmod{11}$
$2015^5 \equiv 10 \pmod{11}$	$2015^{10} \equiv 1 \pmod{11}$

we can see that there is a cycle every 10 powers.

# 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

By observing more powers of 2015,

$2015^1 \equiv 2 \pmod{11}$	$2015^6 \equiv 9 \pmod{11}$
$2015^2 \equiv 4 \pmod{11}$	$2015^7 \equiv 7 \pmod{11}$
$2015^3 \equiv 8 \pmod{11}$	$2015^8 \equiv 8 \pmod{11}$
$2015^4 \equiv 5 \pmod{11}$	$2015^9 \equiv 6 \pmod{11}$
$2015^5 \equiv 10 \pmod{11}$	$2015^{10} \equiv 1 \pmod{11}$

we can see that there is a cycle every 10 powers.

Consider

$$1082 \equiv 2 \pmod{10}.$$

## 2015 T1 Q2 (i)

Evaluate

$$2015^{1082} \pmod{11}.$$

By observing more powers of 2015,

$2015^1 \equiv 2 \pmod{11}$	$2015^6 \equiv 9 \pmod{11}$
$2015^2 \equiv 4 \pmod{11}$	$2015^7 \equiv 7 \pmod{11}$
$2015^3 \equiv 8 \pmod{11}$	$2015^8 \equiv 8 \pmod{11}$
$2015^4 \equiv 5 \pmod{11}$	$2015^9 \equiv 6 \pmod{11}$
$2015^5 \equiv 10 \pmod{11}$	$2015^{10} \equiv 1 \pmod{11}$

we can see that there is a cycle every 10 powers.

Consider

$$1082 \equiv 2 \pmod{10}.$$

Hence,

$$2015^{1082} \equiv 2015^2 \equiv 4 \pmod{11}.$$

- a) Find  $\gcd(105, 342)$ .
- b) Solve, or explain why there is no solution to:
  - $\alpha) 105x \equiv 9 \pmod{342}$
  - $\beta) 105x \equiv 8 \pmod{342}$ .

a) Find  $\gcd(105, 342)$ .

# 2015 T1 Q2 (ii)

a) Find  $\gcd(105, 342)$ .

## 2015 T1 Q2 (ii)

a) Find  $\gcd(105, 342)$ .

Remember: For  $a = bq + r$ ,  $\gcd(a, b) = \gcd(b, r)$ .

## 2015 T1 Q2 (ii)

a) Find  $\gcd(105, 342)$ .

Remember: For  $a = bq + r$ ,  $\gcd(a, b) = \gcd(b, r)$ .

To find  $\gcd$ , we can apply the Euclidean Algorithm:

## 2015 T1 Q2 (ii)

a) Find  $\gcd(105, 342)$ .

Remember: For  $a = bq + r$ ,  $\gcd(a, b) = \gcd(b, r)$ .

To find  $\gcd$ , we can apply the Euclidean Algorithm:

$$342 = 105 \times 3 + 27,$$

## 2015 T1 Q2 (ii)

a) Find  $\gcd(105, 342)$ .

Remember: For  $a = bq + r$ ,  $\gcd(a, b) = \gcd(b, r)$ .

To find  $\gcd$ , we can apply the Euclidean Algorithm:

$$342 = 105 \times 3 + 27,$$

$$105 = 27 \times 3 + 24,$$

## 2015 T1 Q2 (ii)

a) Find  $\gcd(105, 342)$ .

Remember: For  $a = bq + r$ ,  $\gcd(a, b) = \gcd(b, r)$ .

To find  $\gcd$ , we can apply the Euclidean Algorithm:

$$342 = 105 \times 3 + 27,$$

$$105 = 27 \times 3 + 24,$$

$$27 = 24 \times 1 + 3,$$

## 2015 T1 Q2 (ii)

a) Find  $\gcd(105, 342)$ .

Remember: For  $a = bq + r$ ,  $\gcd(a, b) = \gcd(b, r)$ .

To find  $\gcd$ , we can apply the Euclidean Algorithm:

$$342 = 105 \times 3 + 27,$$

$$105 = 27 \times 3 + 24,$$

$$27 = 24 \times 1 + 3,$$

$$24 = 3 \times 3.$$

## 2015 T1 Q2 (ii)

a) Find  $\gcd(105, 342)$ .

Remember: For  $a = bq + r$ ,  $\gcd(a, b) = \gcd(b, r)$ .

To find  $\gcd$ , we can apply the Euclidean Algorithm:

$$342 = 105 \times 3 + 27,$$

$$105 = 27 \times 3 + 24,$$

$$27 = 24 \times 1 + 3,$$

$$24 = 3 \times 3.$$

Thus

$$\gcd(105, 342) = \gcd(27, 105) = \gcd(24, 27) = \gcd(3, 24) = 3.$$

b) Solve, or explain why there is no solution to:

- $\alpha)$   $105x \equiv 9 \pmod{342}$ .
- $\beta)$   $105x \equiv 8 \pmod{342}$ .

## Bézout's Identity

Consider

$$ax + by = c,$$

where  $a$ ,  $b$ , and  $c$  are integers, with  $a$  and  $b$  not both zero.

## Bézout's Identity

Consider

$$ax + by = c,$$

where  $a$ ,  $b$ , and  $c$  are integers, with  $a$  and  $b$  not both zero.

The number of integer solutions for  $x$  and  $y$ :

## Bézout's Identity

Consider

$$ax + by = c,$$

where  $a$ ,  $b$ , and  $c$  are integers, with  $a$  and  $b$  not both zero.

The number of integer solutions for  $x$  and  $y$ :

(i) If  $\gcd(a, b)$  does not divide  $c$ , then there are no solutions,

## 2016 T2 Q2 (i)

### Bézout's Identity

Consider

$$ax + by = c,$$

where  $a$ ,  $b$ , and  $c$  are integers, with  $a$  and  $b$  not both zero.

The number of integer solutions for  $x$  and  $y$ :

- (i) If  $\gcd(a, b)$  does not divide  $c$ , then there are no solutions,
- (ii) If  $\gcd(a, b)$  divides  $c$ , then there are  $\gcd(a, b)$  solutions.

## 2016 T2 Q2 (i)

### Bézout's Identity

Consider

$$ax + by = c,$$

where  $a$ ,  $b$ , and  $c$  are integers, with  $a$  and  $b$  not both zero.

The number of integer solutions for  $x$  and  $y$ :

- (i) If  $\gcd(a, b)$  does not divide  $c$ , then there are no solutions,
- (ii) If  $\gcd(a, b)$  divides  $c$ , then there are  $\gcd(a, b)$  solutions.

To check solubility, we refer to Bézout's Identity.

## 2016 T2 Q2 (i)

### Bézout's Identity

Consider

$$ax + by = c,$$

where  $a$ ,  $b$ , and  $c$  are integers, with  $a$  and  $b$  not both zero.

The number of integer solutions for  $x$  and  $y$ :

- (i) If  $\gcd(a, b)$  does not divide  $c$ , then there are no solutions,
- (ii) If  $\gcd(a, b)$  divides  $c$ , then there are  $\gcd(a, b)$  solutions.

To check solubility, we refer to Bézout's Identity.

Note:

$$ax + by = c \Leftrightarrow ax \equiv c \pmod{b}.$$

- b) Solve, or explain why there is no solution to:
- $105x \equiv 9 \pmod{342}$ .

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:
- $105x \equiv 9 \pmod{342}$ .

We know

$$105x \equiv 9 \pmod{342} \Leftrightarrow 105x + 342y = 9$$

for some integer  $y$ .

- b) Solve, or explain why there is no solution to:
- $105x \equiv 9 \pmod{342}$ .

We know

$$105x \equiv 9 \pmod{342} \Leftrightarrow 105x + 342y = 9$$

for some integer  $y$ .

Since  $\gcd(105, 342) = 3$  divides 9, there are 3 integer solutions for  $x$ .  
(From Bézout's Identity)

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Dividing both side by the greatest common divisor,

$$105x + 342y = 9 \Rightarrow 35x + 114y = 3. \textcolor{red}{•}$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Dividing both side by the greatest common divisor,

$$105x + 342y = 9 \Rightarrow \underline{35x} + \underline{114y} = 3.$$

Through the Euclidean Algorithm:

$$\begin{aligned} 114 &= 3(35) + \underline{9}, \\ 35 &= 3(\underline{9}) + \underline{8}, \\ 9 &= 1(\underline{8}) + \underline{1}, \\ 8 &= 8(1). \end{aligned}$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:
- a)  $105x \equiv 9 \pmod{342}$ .

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:
- $105x \equiv 9 \pmod{342}$ .

From the Extended Euclidean Algorithm, the previous equations can be rearranged to have 1 be expressed in terms of 35 and 114, ie.

## 2015 T2 Q2 (ii)

b) Solve, or explain why there is no solution to:

a)  $105x \equiv 9 \pmod{342}$ .

From the Extended Euclidean Algorithm, the previous equations can be rearranged to have 1 be expressed in terms of 35 and 114, ie.

$$1 = \underline{9 - 8},$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

From the Extended Euclidean Algorithm, the previous equations can be rearranged to have 1 be expressed in terms of 35 and 114, ie.

$$\begin{aligned} 1 &= 9 - \underline{8}, \\ &= 9 - (35 - 3(9)), \end{aligned}$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

From the Extended Euclidean Algorithm, the previous equations can be rearranged to have 1 be expressed in terms of 35 and 114, ie.

$$\begin{aligned} 1 &= 9 - 8, \\ &= 9 - \underline{(35 - 3(9))}, \\ &= (114 - 3(35)) - (35 - 3(114 - 3(35))), \end{aligned}$$

## 2015 T2 Q2 (ii)

b) Solve, or explain why there is no solution to:

a)  $105x \equiv 9 \pmod{342}$ .

From the Extended Euclidean Algorithm, the previous equations can be rearranged to have 1 be expressed in terms of 35 and 114, ie.

$$\begin{aligned} 1 &= \underline{9 - 8}, \\ &= 9 - (35 - 3(9)), \\ &= (114 - 3(35)) - (35 - 3(114 - 3(35))), \\ &= 35(-13) + 114(4). \end{aligned}$$

## 2015 T2 Q2 (ii)

b) Solve, or explain why there is no solution to:

a)  $105x \equiv 9 \pmod{342}$ .

From the Extended Euclidean Algorithm, the previous equations can be rearranged to have 1 be expressed in terms of 35 and 114, ie.

$$\begin{aligned} 1 &= 9 - 8, \\ &= 9 - (35 - 3(9)), \\ &= (114 - 3(35)) - (35 - 3(114 - 3(35))), \\ &= 35(-13) + 114(4). \end{aligned}$$

Multiplying both sides by 3,

$$3 = 35(-39) + 114(12) = 35x + 114y.$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:
- a)  $105x \equiv 9 \pmod{342}$ .

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Equating coefficients in

$$35(-39) + 114(12) = 35x + 114y,$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Equating coefficients in

$$35(-39) + 114(12) = 35x + 114y,$$

we find that  $x = -39$  is an integer solution.

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Equating coefficients in

$$35(-39) + 114(12) = 35x + 114y,$$

we find that  $x = -39$  is an integer solution.

However, solutions must be  $0 \leq x < 342$ . (342 from original mod)

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Equating coefficients in

$$35(-39) + 114(12) = 35x + 114y$$

we find that  $x = -39$  is an integer solution.

However, solutions must be  $0 \leq x < \underline{342}$ . (342 from original mod)

We can find more solutions by adding 114 to our  $x$ .

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Equating coefficients in

$$35(-39) + 114(12) = 35x + 114y,$$

we find that  $x = -39$  is an integer solution.

However, solutions must be  $0 \leq x < 342$ . (342 from original mod)

We can find more solutions by adding 114 to our  $x$ .

Thus,

$$x = -39 + 114$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Equating coefficients in

$$35(-39) + 114(12) = 35x + 114y,$$

we find that  $x = -39$  is an integer solution.

However, solutions must be  $0 \leq x < 342$ . (342 from original mod)

We can find more solutions by adding 114 to our  $x$ .

Thus,

$$x = -39 + 114, -39 + 2(114)$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
α)  $105x \equiv 9 \pmod{342}$ .

Equating coefficients in

$$35(-39) + 114(12) = 35x + 114y,$$

we find that  $x = -39$  is an integer solution.

However, solutions must be  $0 \leq x < 342$ . (342 from original mod)

We can find more solutions by adding 114 to our  $x$ .

Thus,

$$x = -39 + 114, -39 + 2(114), -39 + 3(114) = 75, 189, 303.$$

- b) Solve, or explain why there is no solution to:  
β)  $105x \equiv 8 \pmod{342}$ .



## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
β)  $105x \equiv 8 \pmod{342}$ .

We know

$$105x \equiv 8 \pmod{342} \Leftrightarrow 105x + 342y = 8,$$

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
β)  $105x \equiv 8 \pmod{342}$ .

We know

$$105x \equiv 8 \pmod{342} \Leftrightarrow 105x + 342y = 8,$$

for some integer  $y$ , but  $\gcd(105, 342) = 3$  does not divide 8.

## 2015 T2 Q2 (ii)

- b) Solve, or explain why there is no solution to:  
β)  $105x \equiv 8 \pmod{342}$ .

We know

$$105x \equiv 8 \pmod{342} \Leftrightarrow 105x + 342y = 8,$$

for some integer  $y$ , but  $\gcd(105, 342) = 3$  does not divide 8.

Hence, by Bézout's Identity, there are no solutions for

$$105x \equiv 8 \pmod{342}.$$

## 2016 T2 Q1 (vii)

Suppose  $\gcd(a, m) = d$ .

Show that if  $x$  is a solution of  $ax \equiv b \pmod{m}$ , then so is  $x + m/d$ .

## 2016 T2 Q1 (vii)

Suppose  $\gcd(a, m) = d$ .

Show that if  $x$  is a solution of  $ax \equiv b \pmod{m}$ , then so is  $x + m/d$ .

## 2016 T2 Q1 (vii)

Suppose  $\gcd(a, m) = d$ .

Show that if  $x$  is a solution of  $ax \equiv b \pmod{m}$ , then so is  $x + m/d$ .

Suppose that

$$ax \equiv b \pmod{m}.$$

## 2016 T2 Q1 (vii)

Suppose  $\gcd(a, m) = d$ .

Show that if  $x$  is a solution of  $ax \equiv b \pmod{m}$ , then so is  $x + m/d$ .

Suppose that

$$ax \equiv b \pmod{m}.$$

We need to prove that

$$\underline{a(x + m/d)} \equiv b \pmod{m}.$$

## 2016 T2 Q1 (vii)

Suppose  $\gcd(a, m) = d$ .

Show that if  $x$  is a solution of  $ax \equiv b \pmod{m}$ , then so is  $x + m/d$ .

Suppose that

$$ax \equiv b \pmod{m}.$$

We need to prove that

$$a(x + m/d) \equiv b \pmod{m}.$$

Consider

$$a(x + m/d) = ax + m(a/d).$$

## 2016 T2 Q1 (vii)

Suppose  $\gcd(a, m) = d$ .

Show that if  $x$  is a solution of  $ax \equiv b \pmod{m}$ , then so is  $x + m/d$ .

Suppose that

$$ax \equiv b \pmod{m}.$$

We need to prove that

$$a(x + m/d) \equiv b \pmod{m}.$$

Consider

$$a(x + m/d) = ax + m(a/d).$$

Since  $\gcd(a, m) = d$ ,  $a/d$  must be an integer.

## 2016 T2 Q1 (vii)

Suppose  $\gcd(a, m) = d$ .

Show that if  $x$  is a solution of  $ax \equiv b \pmod{m}$ , then so is  $x + m/d$ .

Suppose that

$$ax \equiv b \pmod{m}.$$

We need to prove that

$$a(x + m/d) \equiv b \pmod{m}.$$

Consider

$$a(x + m/d) = ax + m(a/d).$$

Since  $\gcd(a, m) = d$ ,  $a/d$  must be an integer.

Thus

$$a(x + m/d) \equiv ax + \underline{m}(a/d)$$

## 2016 T2 Q1 (vii)

Suppose  $\gcd(a, m) = d$ .

Show that if  $x$  is a solution of  $ax \equiv b \pmod{m}$ , then so is  $x + m/d$ .

Suppose that

$$\underline{ax \equiv b \pmod{m}}.$$

We need to prove that

$$a(x + m/d) \equiv b \pmod{m}.$$

Consider

$$a(x + m/d) = ax + m(a/d).$$

Since  $\gcd(a, m) = d$ ,  $a/d$  must be an integer.

Thus

$$\underline{a(x + m/d)} \equiv \underline{ax + m(a/d)} \equiv \underline{ax} \equiv \underline{b} \pmod{m}.$$

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- Find integers  $k$  and  $l$  such that  $1039k + 2013l = 1$ .
- What is the general solution to  $1039x + 2013y = 4$ ?

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- Find integers  $k$  and  $l$  such that  $1031k + 2013l = 1$ .

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- Find integers  $k$  and  $l$  such that  $1031k + 2013l = 1$ .

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- Find integers  $k$  and  $l$  such that  $1031k + 2013l = 1$ .

Remember: The inverse  $a^{-1}$  is a number such that  $aa^{-1} \equiv 1 \pmod{m}$ .

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- a) Find integers  $k$  and  $l$  such that  $1031k + 2013l = 1$ .

Remember: The inverse  $a^{-1}$  is a number such that  $aa^{-1} \equiv 1 \pmod{m}$ .

We know that

$$1039 \times 31 \equiv 1 \pmod{2013} \Leftrightarrow 1039(31) + 2013y = 1$$

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- a) Find integers  $k$  and  $l$  such that  $\underline{\underline{1031k + 2013l = 1}}$ .

Remember: The inverse  $a^{-1}$  is a number such that  $aa^{-1} \equiv 1 \pmod{m}$ .

We know that

$$1039 \times 31 \equiv 1 \pmod{2013} \Leftrightarrow 1039(31) + 2013\underline{y} = 1$$

for some integer  $y$ . This equation is strikingly similar to the question.

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- Find integers  $k$  and  $l$  such that  $1031k + 2013l = 1$ .

Remember: The inverse  $a^{-1}$  is a number such that  $aa^{-1} \equiv 1 \pmod{m}$ .

We know that

$$1039 \times 31 \equiv 1 \pmod{2013} \Leftrightarrow 1039(31) + 2013y = 1$$

for some integer  $y$ . This equation is strikingly similar to the question.

Solving for  $y$  we see that  $y = \frac{1 - 1039(31)}{2013} = -16$ .

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- Find integers  $k$  and  $l$  such that  $1031k + 2013l = 1$ .

Remember: The inverse  $a^{-1}$  is a number such that  $aa^{-1} \equiv 1 \pmod{m}$ .

We know that

$$1039 \times 31 \equiv 1 \pmod{2013} \Leftrightarrow 1039(31) + 2013y = 1$$

for some integer  $y$ . This equation is strikingly similar to the question.

Solving for  $y$  we see that  $y = \frac{1 - 1039(31)}{2013} = -16$ .

Therefore

$$\underline{\hspace{10em}} 1039(31) + 2013(-16) = 1 = \underline{\hspace{10em}} 1031k + 2013l. \underline{\hspace{10em}}$$

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- a) Find integers  $k$  and  $l$  such that  $1031k + 2013l = 1$ .

Remember: The inverse  $a^{-1}$  is a number such that  $aa^{-1} \equiv 1 \pmod{m}$ .

We know that

$$1039 \times 31 \equiv 1 \pmod{2013} \Leftrightarrow 1039(31) + 2013y = 1$$

for some integer  $y$ . This equation is strikingly similar to the question.

Solving for  $y$  we see that  $y = \frac{1 - 1039(31)}{2013} = -16$ .

Therefore

$$1039(31) + 2013(-16) = 1 = 1031k + 2013l.$$

Comparing coefficients, we see that  $k = 31$  and  $l = -16$ .

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

From a), we can multiply both sides by 4 to find that

$$1039(31) + 2013(-16) = 1 \Rightarrow 1039(124) + 2013(-64) = 4.$$

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $\underline{1039x + 2013y = 4}$ ?

From a), we can multiply both sides by 4 to find that

$$1039(31) + 2013(-16) = 1 \Rightarrow 1039(124) + 2013(-64) = 4.$$

This can be equated to the equation in the question where

$$1039(124) + 2013(-64) = 4 = 1039x + 2013y.$$

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

From a), we can multiply both sides by 4 to find that

$$1039(31) + 2013(-16) = 1 \Rightarrow 1039(124) + 2013(-64) = 4.$$

This can be equated to the equation in the question where

$$1039(124) + 2013(-64) = 4 = 1039x + 2013y.$$

Thus, possible solutions are  $x = 124$  and  $y = -64$ .

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

From a), we can multiply both sides by 4 to find that

$$1039(31) + 2013(-16) = 1 \Rightarrow 1039(124) + 2013(-64) = 4.$$

This can be equated to the equation in the question where

$$1039(124) + 2013(-64) = 4 = 1039x + 2013y.$$

Thus, possible solutions are  $x = 124$  and  $y = -64$ .

Hence, the general solution is  $x = 124 + 2013z$  for some integer  $z$ .

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

Consider  $1039(124 + 2013z) = 1039(124) + 1039(2013z)$ , then

$$1039(124 + 2013z) - 1039(2013z) + 2013(-64) = 4,$$

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

Consider  $1039(124 + 2013z) = 1039(124) + 1039(2013z)$ , then

$$1039(124 + 2013z) - 1039(2013z) + 2013(-64) = 4,$$

$$1039(124 + 2013z) + 2013(-64 - 1039z) = 4,$$

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

- b) What is the general solution to  $1039x + 2013y = 4$ ?

Consider  $1039(124 + 2013z) = 1039(124) + 1039(2013z)$ , then

$$1039(124 + 2013z) - 1039(2013z) + 2013(-64) = 4,$$

$$1039(124 + 2013z) + 2013(-64 - 1039z) = 4,$$

$$1039(124 + 2013z) + 2013(-64 - 1039z) = 1039x + 2013y.$$

## 2012 T2 Q2 (ii)

Given that  $1039^{-1} \equiv 31 \pmod{2013}$ :

b) What is the general solution to  $1039x + 2013y = 4$ ?

Consider  $1039(124 + 2013z) = 1039(124) + 1039(2013z)$ , then

$$1039(124 + 2013z) - 1039(2013z) + 2013(-64) = 4,$$

$$1039(124 + 2013z) + 2013(-64 - 1039z) = 4,$$

$$1039(124 + 2013z) + 2013(-64 - 1039z) = 1039x + 2013y.$$

Hence, from equating coefficients, the general solution is

$x = 124 + 2013z$  and  $y = -64 - 1039z$  for some integer  $z$ .

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- Prove that  $\sim$  is an equivalence relation.
- Give a geometric description of the equivalence class with  $(1,0)$ .
- Is it true that all equivalence classes are infinite in size? Explain.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

## Properties and Types of Relations

Let  $R$  be a relation defined on the set  $A$

## Properties and Types of Relations

Let  $R$  be a relation defined on the set  $A$

- (i) Reflexive: All elements of the relation relate to themselves.  
ie.  $xRx$  for all  $x \in A$ .

## Properties and Types of Relations

Let  $R$  be a relation defined on the set  $A$

- (i) Reflexive: All elements of the relation relate to themselves.  
ie.  $xRx$  for all  $x \in A$ .
- (ii) Symmetrical: Elements that relate must relate both ways.  
ie. If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

## Properties and Types of Relations

Let  $R$  be a relation defined on the set  $A$

(i) Reflexive: All elements of the relation relate to themselves.

ie.  $xRx$  for all  $x \in A$ .

(ii) Symmetrical: Elements that relate must relate both ways.

ie. If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

(iii) Anti-Symm: Elements only relate both ways with themselves.

ie. If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

## Properties and Types of Relations

Let  $R$  be a relation defined on the set  $A$

(i) Reflexive: All elements of the relation relate to themselves.

ie.  $xRx$  for all  $x \in A$ .

(ii) Symmetrical: Elements that relate must relate both ways.

ie. If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

(iii) Anti-Symm: Elements only relate both ways with themselves.

ie. If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

(iv) Transitive: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

## Properties and Types of Relations

Let  $R$  be a relation defined on the set  $A$

(i) Reflexive: All elements of the relation relate to themselves.

ie.  $xRx$  for all  $x \in A$ .

(ii) Symmetrical: Elements that relate must relate both ways.

ie. If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

(iii) Anti-Symm: Elements only relate both ways with themselves.

ie. If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

(iv) Transitive: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Equivalence Relation: Reflexive, Symmetrical, Transitive.

## Properties and Types of Relations

Let  $R$  be a relation defined on the set  $A$

(i) Reflexive: All elements of the relation relate to themselves.

ie.  $xRx$  for all  $x \in A$ .

(ii) Symmetrical: Elements that relate must relate both ways.

ie. If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

(iii) Anti-Symm: Elements only relate both ways with themselves.

ie. If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

(iv) Transitive: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Equivalence Relation: Reflexive, Symmetrical, Transitive.

Partial Order: Reflexive, Anti-Symmetrical, Transitive.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

Let  $a_1$  and  $a_2$  be real numbers.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

Let  $a_1$  and  $a_2$  be real numbers. Consider

$$a_1^2 + a_2^2 = a_1^2 + a_2^2.$$

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

a) Prove that  $\sim$  is an equivalence relation.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

Let  $a_1$  and  $a_2$  be real numbers. Consider

$$a_1^2 + a_2^2 = a_1^2 + a_2^2.$$

Therefore

$$(a_1, a_2) \sim (a_1, a_2).$$

where  $(a_1, a_2)$  is a part of the set  $\mathbb{R} \times \mathbb{R}$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

a) Prove that  $\sim$  is an equivalence relation.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

Let  $a_1$  and  $a_2$  be real numbers. Consider

$$a_1^2 + a_2^2 = a_1^2 + a_2^2.$$

Therefore

$$(a_1, a_2) \sim (a_1, a_2).$$

where  $(a_1, a_2)$  is a part of the set  $\mathbb{R} \times \mathbb{R}$ .

Hence, the relation  $\sim$  is reflexive.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove symmetry: If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove symmetry: If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

Let  $a_1, a_2, b_1$  and  $b_2$  be real numbers.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove symmetry: If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

Let  $a_1, a_2, b_1$  and  $b_2$  be real numbers. Suppose that  $(a_1, a_2) \sim (b_1, b_2)$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove symmetry: If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

Let  $a_1, a_2, b_1$  and  $b_2$  be real numbers. Suppose that  $(a_1, a_2) \sim (b_1, b_2)$ .  
By definition

$$a_1^2 + a_2^2 = b_1^2 + b_2^2$$

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove symmetry: If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

Let  $a_1, a_2, b_1$  and  $b_2$  be real numbers. Suppose that  $(a_1, a_2) \sim (b_1, b_2)$ .  
By definition

$$a_1^2 + a_2^2 = b_1^2 + b_2^2 \Rightarrow b_1^2 + b_2^2 = a_1^2 + a_2^2.$$

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove symmetry: If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

Let  $a_1, a_2, b_1$  and  $b_2$  be real numbers. Suppose that  $(a_1, a_2) \sim (b_1, b_2)$ . By definition

$$a_1^2 + a_2^2 = b_1^2 + b_2^2 \Rightarrow b_1^2 + b_2^2 = a_1^2 + a_2^2.$$

Therefore  $(b_1, b_2) \sim (a_1, a_2)$ , where  $(a_1, a_2)$  and  $(b_1, b_2)$  are in  $\mathbb{R} \times \mathbb{R}$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove symmetry: If  $xRy$ , then  $yRx$  for all  $x, y \in A$ .

Let  $a_1, a_2, b_1$  and  $b_2$  be real numbers. Suppose that  $(a_1, a_2) \sim (b_1, b_2)$ . By definition

$$a_1^2 + a_2^2 = b_1^2 + b_2^2 \Rightarrow b_1^2 + b_2^2 = a_1^2 + a_2^2.$$

Therefore  $(b_1, b_2) \sim (a_1, a_2)$ , where  $(a_1, a_2)$  and  $(b_1, b_2)$  are in  $\mathbb{R} \times \mathbb{R}$ .

Hence, the relation  $\sim$  is symmetrical.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $a_1, a_2, b_1, b_2, c_1$  and  $c_2$  be real numbers.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $a_1, a_2, b_1, b_2, c_1$  and  $c_2$  be real numbers.

Suppose that  $(a_1, a_2) \sim (b_1, b_2)$  and  $(b_1, b_2) \sim (c_1, c_2)$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $a_1, a_2, b_1, b_2, c_1$  and  $c_2$  be real numbers.

Suppose that  $(a_1, a_2) \sim (b_1, b_2)$  and  $(b_1, b_2) \sim (c_1, c_2)$ . By definition,

$$a_1^2 + a_2^2 = b_1^2 + b_2^2 \text{ and } b_1^2 + b_2^2 = c_1^2 + c_2^2.$$



## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $a_1, a_2, b_1, b_2, c_1$  and  $c_2$  be real numbers.

Suppose that  $(a_1, a_2) \sim (b_1, b_2)$  and  $(b_1, b_2) \sim (c_1, c_2)$ . By definition,

$$a_1^2 + a_2^2 = b_1^2 + b_2^2 \text{ and } b_1^2 + b_2^2 = c_1^2 + c_2^2.$$

Therefore  $(a_1, a_2) \sim (c_1, c_2)$ , since  $a_1^2 + a_2^2 = b_1^2 + b_2^2 = c_1^2 + c_2^2$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

a) Prove that  $\sim$  is an equivalence relation.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $a_1, a_2, b_1, b_2, c_1$  and  $c_2$  be real numbers.

Suppose that  $(a_1, a_2) \sim (b_1, b_2)$  and  $(b_1, b_2) \sim (c_1, c_2)$ . By definition,

$$a_1^2 + a_2^2 = b_1^2 + b_2^2 \text{ and } b_1^2 + b_2^2 = c_1^2 + c_2^2.$$

Therefore  $(a_1, a_2) \sim (c_1, c_2)$ , since  $a_1^2 + a_2^2 = b_1^2 + b_2^2 = c_1^2 + c_2^2$ .

Hence, the relation  $\sim$  is transitive.

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

We know that the relation  $\sim$  is reflexive, symmetrical and transitive.

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- a) Prove that  $\sim$  is an equivalence relation.

We know that the relation  $\sim$  is reflexive, symmetrical and transitive.

Hence, the relation  $\sim$  is an equivalence relation.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- b) Give a geometric description of the equivalence class with  $(1,0)$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- b) Give a geometric description of the equivalence class with  $(1,0)$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- b) Give a geometric description of the equivalence class with  $(1,0)$ .

Equivalence Classes:  $[a] = \{x \in A \mid xRa\}$  where  $a$  is an element of  $A$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- b) Give a geometric description of the equivalence class with  $(1,0)$ .

Equivalence Classes:  $[a] = \{x \in A \mid xRa\}$  where  $a$  is an element of  $A$ .

Consider

$$[(1, 0)] = \{(a_1, a_2) \in \mathbb{R} \times \mathbb{R} \mid (a_1, a_2) \sim (1, 0)\}.$$

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- b) Give a geometric description of the equivalence class with  $(1,0)$ .

Equivalence Classes:  $[a] = \{x \in A \mid xRa\}$  where  $a$  is an element of  $A$ .

Consider

$$[(1, 0)] = \{(a_1, a_2) \in \mathbb{R} \times \mathbb{R} \mid (a_1, a_2) \sim (1, 0)\}.$$

By definition,  $\underline{a_1^2} + \underline{a_2^2} = 1^2 + 0^2 = 1$  for the real numbers  $\underline{a_1}$  and  $\underline{a_2}$

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- b) Give a geometric description of the equivalence class with  $(1,0)$ .

Equivalence Classes:  $[a] = \{x \in A \mid xRa\}$  where  $a$  is an element of  $A$ .

Consider

$$[(1, 0)] = \{(a_1, a_2) \in \mathbb{R} \times \mathbb{R} \mid (a_1, a_2) \sim (1, 0)\}.$$

By definition,  $a_1^2 + a_2^2 = 1^2 + 0^2 = 1$  for the real numbers  $a_1$  and  $a_2$ .

Hint: Let  $a_1 = x$  and  $a_2 = y$ . Thus,  $x^2 + y^2 = 1$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- b) Give a geometric description of the equivalence class with  $(1,0)$ .

Equivalence Classes:  $[a] = \{x \in A \mid xRa\}$  where  $a$  is an element of  $A$ .

Consider

$$[(1,0)] = \{(a_1, a_2) \in \mathbb{R} \times \mathbb{R} \mid (a_1, a_2) \sim (1,0)\}.$$

By definition,  $a_1^2 + a_2^2 = 1^2 + 0^2 = 1$  for the real numbers  $a_1$  and  $a_2$ .

Hint: Let  $a_1 = x$  and  $a_2 = y$ . Thus,  $x^2 + y^2 = 1$ .

The equivalence class with  $(1,0)$  is the set of ordered pairs containing all the coordinates on the unit circle.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- c) Is it true that all equivalence classes are infinite in size? Explain.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

Thus,  $a_1^2 = -a_2^2$  where  $a_1^2 \geq 0$  and  $a_2^2 \geq 0$  since squares are non-negative.

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

Thus,  $a_1^2 = -a_2^2$  where  $a_1^2 \geq 0$  and  $a_2^2 \geq 0$  since squares are non-negative.

If  $a_1^2 > 0$ , then  $-a_2^2 > 0$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

- c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

Thus,  $a_1^2 = -a_2^2$  where  $a_1^2 \geq 0$  and  $a_2^2 \geq 0$  since squares are non-negative.

If  $a_1^2 > 0$ , then  $-a_2^2 > 0$ . Therefore,  $\underline{a_2^2 < 0}$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

Thus,  $a_1^2 = -a_2^2$  where  $a_1^2 \geq 0$  and  $a_2^2 \geq 0$  since squares are non-negative.

If  $a_1^2 > 0$ , then  $-a_2^2 > 0$ . Therefore,  $a_2^2 < 0$ . This contradicts  $a_2^2 \geq 0$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

Thus,  $a_1^2 = -a_2^2$  where  $a_1^2 \geq 0$  and  $a_2^2 \geq 0$  since squares are non-negative.

If  $\underline{a_1^2 > 0}$ , then  $-a_2^2 > 0$ . Therefore,  $a_2^2 < 0$ . This contradicts  $\underline{a_2^2 \geq 0}$ .

We can also use this argument for the case that  $\underline{a_2^2 > 0}$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

Thus,  $a_1^2 = -a_2^2$  where  $a_1^2 \geq 0$  and  $a_2^2 \geq 0$  since squares are non-negative.

If  $a_1^2 > 0$ , then  $-a_2^2 > 0$ . Therefore,  $a_2^2 < 0$ . This contradicts  $a_2^2 \geq 0$ .

We can also use this argument for the case that  $a_2^2 > 0$ .

Thus,  $a_1^2 = a_2^2 = 0$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

Thus,  $a_1^2 = -a_2^2$  where  $a_1^2 \geq 0$  and  $a_2^2 \geq 0$  since squares are non-negative.

If  $a_1^2 > 0$ , then  $-a_2^2 > 0$ . Therefore,  $a_2^2 < 0$ . This contradicts  $a_2^2 \geq 0$ .

We can also use this argument for the case that  $a_2^2 > 0$ .

Thus,  $a_1^2 = a_2^2 = 0$ . Therefore, there is only 1 element for  $[(0, 0)]$ :  $(0, 0)$ .

## 2017 T1 Q2 (iii)

Let  $\sim$  be the relation on the set  $\mathbb{R} \times \mathbb{R}$  defined by

$$(a_1, a_2) \sim (b_1, b_2) \text{ if and only if } a_1^2 + a_2^2 = b_1^2 + b_2^2.$$

c) Is it true that all equivalence classes are infinite in size? Explain.

This is not true. Consider the equivalence class  $[(0, 0)]$ .

By definition,  $a_1^2 + a_2^2 = 0^2 + 0^2 = 0$  for some real numbers  $a_1$  and  $a_2$ .

Thus,  $a_1^2 = -a_2^2$  where  $a_1^2 \geq 0$  and  $a_2^2 \geq 0$  since squares are non-negative.

If  $a_1^2 > 0$ , then  $-a_2^2 > 0$ . Therefore,  $a_2^2 < 0$ . This contradicts  $a_2^2 \geq 0$ .

We can also use this argument for the case that  $a_2^2 > 0$ .

Thus,  $a_1^2 = a_2^2 = 0$ . Therefore, there is only 1 element for  $[(0, 0)]$ :  $(0, 0)$ . Hence, there exists an equivalence class which has a size of 1.

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.



## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.


## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

To find all equivalence classes, we need all the

$$[a] = \{x \in \mathbb{Z} \mid x \star a\} \text{ such that } \bigcup [a] = \mathbb{Z}.$$


## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

To find all equivalence classes, we need all the

$$[a] = \{x \in \mathbb{Z} \mid x \star a\} \text{ such that } \bigcup [a] = \mathbb{Z}.$$

As we are in the world of modulo 4, all integers can be represented by 0, 1, 2 and 3. We just need to check each of these.

• • • •

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

$$\text{For } x \equiv 0 \pmod{4} \Rightarrow x^2 \equiv 0 \pmod{4}.$$

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 0 \pmod{4} \Rightarrow x^2 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 0 \pmod{4}$  are in  $[0]$ .

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 0 \pmod{4} \Rightarrow x^2 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 0 \pmod{4}$  are in [0]. These are all the numbers in the form  $4n$ , where  $n$  is some integer ie. ..., -8, -4, 0, 4, ...

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 0 \pmod{4} \Rightarrow x^2 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 0 \pmod{4}$  are in [0]. These are all the numbers in the form  $4n$ , where  $n$  is some integer ie. ..., -8, -4, 0, 4, ...

For  $x \equiv 1 \pmod{4} \Rightarrow x^2 \equiv 1 \pmod{4}$

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 0 \pmod{4} \Rightarrow x^2 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 0 \pmod{4}$  are in [0]. These are all the numbers in the form  $4n$ , where  $n$  is some integer ie. ..., -8, -4, 0, 4, ...

For  $x \equiv 1 \pmod{4} \Rightarrow x^2 \equiv 1 \pmod{4}$  Thus all integers where  $a \equiv 1 \pmod{4}$  are in [1].

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 0 \pmod{4} \Rightarrow x^2 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 0 \pmod{4}$  are in [0]. These are all the numbers in the form  $4n$ , where  $n$  is some integer ie. ..., -8, -4, 0, 4,...

For  $x \equiv 1 \pmod{4} \Rightarrow x^2 \equiv 1 \pmod{4}$  Thus all integers where  $a \equiv 1 \pmod{4}$  are in [1]. These are all the numbers in the form  $1+4n$ , where  $n$  is some integer ie. ..., -7, -3, 1, 5,...

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 2 \pmod{4} \Rightarrow x^2 \equiv 4 \equiv 0 \pmod{4}$ .

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 2 \pmod{4} \Rightarrow x^2 \equiv 4 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 2 \pmod{4}$  are in  $[0]$ .

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 2 \pmod{4} \Rightarrow x^2 \equiv 4 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 2 \pmod{4}$  are in [0]. These are all the numbers in the form  $2+4n$ , where  $n$  is some integer ie. ..., -6, -2, 2, 6, ...

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 2 \pmod{4} \Rightarrow x^2 \equiv 4 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 2 \pmod{4}$  are in [0]. These are all the numbers in the form  $2+4n$ , where  $n$  is some integer ie. ..., -6, -2, 2, 6, ...

For  $x \equiv 3 \pmod{4} \Rightarrow x^2 \equiv 9 \equiv 1 \pmod{4}$ .

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 2 \pmod{4} \Rightarrow x^2 \equiv 4 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 2 \pmod{4}$  are in [0]. These are all the numbers in the form  $2+4n$ , where  $n$  is some integer ie. ..., -6, -2, 2, 6, ...

For  $x \equiv 3 \pmod{4} \Rightarrow x^2 \equiv 9 \equiv 1 \pmod{4}$ . Thus all integers where  $a \equiv 3 \pmod{4}$  is in [1].

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

For  $x \equiv 2 \pmod{4} \Rightarrow x^2 \equiv 4 \equiv 0 \pmod{4}$ . Thus all integers where  $a \equiv 2 \pmod{4}$  are in [0]. These are all the numbers in the form  $2+4n$ , where n is some integer ie. ..., -6, -2, 2, 6,...

For  $x \equiv 3 \pmod{4} \Rightarrow x^2 \equiv 9 \equiv 1 \pmod{4}$ . Thus all integers where  $a \equiv 3 \pmod{4}$  is in [1]. These are all the numbers in the form  $3+4n$ , where n is some integer ie. ..., -5, -1, 3, 7,...

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

Thus, we know that

$$[0] = \{..., -8, -4, 0, 4, ...\} \cup \{..., -6, -2, 2, 6, ...\}$$

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

Thus, we know that

$$\begin{aligned}[0] &= \{\dots, -8, -4, 0, 4, \dots\} \cup \{\dots, -6, -2, 2, 6, \dots\} \\ &= \{\dots - 6, -4, -2, 0, 2, 4, 6\} = \{2j \mid j \in \mathbb{Z}\}\end{aligned}$$

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

Thus, we know that

$$\begin{aligned}[0] &= \{\dots, -8, -4, 0, 4, \dots\} \cup \{\dots, -6, -2, 2, 6, \dots\} \\ &= \{\dots - 6, -4, -2, 0, 2, 4, 6\} = \{2j \mid j \in \mathbb{Z}\}\end{aligned}$$

$$[1] = \{\dots, -7, -3, 1, 5, \dots\} \cup \{\dots, -5, -1, 3, 7, \dots\}$$



## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

Thus, we know that

$$\begin{aligned}[0] &= \{\dots, -8, -4, 0, 4, \dots\} \cup \{\dots, -6, -2, 2, 6, \dots\} \\ &= \{\dots - 6, -4, -2, 0, 2, 4, 6\} = \{2j \mid j \in \mathbb{Z}\}\end{aligned}$$

$$\begin{aligned}[1] &= \{\dots, -7, -3, 1, 5, \dots\} \cup \{\dots, -5, -1, 3, 7, \dots\} \\ &= \{\dots - 3, -1, 1, 3, 5, 7\} = \{2k + 1 \mid k \in \mathbb{Z}\}\end{aligned}$$

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

Thus, we know that

$$\begin{aligned}[0] &= \{\dots, -8, -4, 0, 4, \dots\} \cup \{\dots, -6, -2, 2, 6, \dots\} \\ &= \{\dots - 6, -4, -2, 0, 2, 4, 6\} = \{2j \mid j \in \mathbb{Z}\}\end{aligned}$$

$$\begin{aligned}[1] &= \{\dots, -7, -3, 1, 5, \dots\} \cup \{\dots, -5, -1, 3, 7, \dots\} \\ &= \{\dots - 3, -1, 1, 3, 5, 7\} = \{2k + 1 \mid k \in \mathbb{Z}\}\end{aligned}$$

Since [0] contains all even integers and [1] contains all odd integers,  
 $[0] \cup [1] = \mathbb{Z}$ .

## 2016 T2 Q2 (ii)

Let  $\star$  be the relation on the set of integers  $\mathbb{Z}$  defined by

$$a \star b \text{ if and only if } a^2 \equiv b^2 \pmod{4}.$$

Given that  $\star$  is an equivalence relation, find its equivalence classes.

Thus, we know that

$$\begin{aligned}[0] &= \{\dots, -8, -4, 0, 4, \dots\} \cup \{\dots, -6, -2, 2, 6, \dots\} \\ &= \{\dots - 6, -4, -2, 0, 2, 4, 6\} = \{2j \mid j \in \mathbb{Z}\}\end{aligned}$$

$$\begin{aligned}[1] &= \{\dots, -7, -3, 1, 5, \dots\} \cup \{\dots, -5, -1, 3, 7, \dots\} \\ &= \{\dots - 3, -1, 1, 3, 5, 7\} = \{2k + 1 \mid k \in \mathbb{Z}\}\end{aligned}$$

Since [0] contains all even integers and [1] contains all odd integers,  $[0] \cup [1] = \mathbb{Z}$ . Hence, [0] and [1] are all possible equivalence classes.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ . 

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

- a) Prove that  $|$  is a partial order.
- b) Draw a Hasse diagram for the partially ordered set  $(A, |)$ .
- c) Find, if they exist, all
  - $\alpha$ ) minimal elements.
  - $\beta$ ) least elements.
  - $\gamma$ ) maximal elements.
  - $\delta$ ) greatest elements.
- d) Find two elements of  $A$  that do not have a greatest lower bound.  
Explain why that is.

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

Let  $x$  be a number in the set  $A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

Let  $x$  be a number in the set  $A$ .

The number  $x$  is known to divide  $x$  as  $x \neq 0$ , since  $0 \notin A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

Let  $x$  be a number in the set  $A$ .

The number  $x$  is known to divide  $x$  as  $x \neq 0$ , since  $0 \notin A$ .

Therefore,  $x|x$  where  $x \in A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

To prove reflexivity:  $xRx$  for all  $x \in A$ .

Let  $x$  be a number in the set  $A$ .

The number  $x$  is known to divide  $x$  as  $x \neq 0$ , since  $0 \notin A$ .

Therefore,  $x|x$  where  $x \in A$ .

Hence, the relation  $|$  is reflexive.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove anti-symmetry: If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove anti-symmetry: If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

Let  $x$  and  $y$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|x$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove anti-symmetry: If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

Let  $x$  and  $y$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|x$ .

By definition,  $x$  divides  $y$ , meaning  $x \leq y$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove anti-symmetry: If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

Let  $x$  and  $y$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|x$ .

By definition,  $x$  divides  $y$ , meaning  $x \leq y$ . Similarly, since  $y$  divides  $x$ ,  $y \leq x$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove anti-symmetry: If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

Let  $x$  and  $y$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|x$ .

By definition,  $x$  divides  $y$ , meaning  $x \leq y$ . Similarly, since  $y$  divides  $x$ ,  $y \leq x$ . Combining these inequalities gives us:  $x \leq y \leq x$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove anti-symmetry: If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

Let  $x$  and  $y$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|x$ .

By definition,  $x$  divides  $y$ , meaning  $x \leq y$ . Similarly, since  $y$  divides  $x$ ,  $y \leq x$ . Combining these inequalities gives us:  $x \leq y \leq x$ .

Therefore by sandwich theorem,  $x = y$  where  $x, y \in A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove anti-symmetry: If  $xRy$  and  $yRx$ , then  $x = y$  for all  $x, y \in A$ .

Let  $x$  and  $y$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|x$ .

By definition,  $x$  divides  $y$ , meaning  $x \leq y$ . Similarly, since  $y$  divides  $x$ ,  $y \leq x$ . Combining these inequalities gives us:  $x \leq y \leq x$ .

Therefore by sandwich theorem,  $x = y$  where  $x, y \in A$ .

Hence, the relation  $|$  is anti-symmetrical.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $x, y$  and  $z$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|z$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $x, y$  and  $z$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|z$ .

By definition,  $y = ax$  and  $z = by$  for some integers  $a$  and  $b$ .

Substituting this value of  $y$  into the second equation gives us  $z = abx$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .  
a) Prove that  $|$  is a partial order.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $x, y$  and  $z$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|z$ .

By definition,  $y = ax$  and  $z = by$  for some integers  $a$  and  $b$ .

Substituting this value of  $y$  into the second equation gives us  $z = abx$ .

Since  $ab$  is an integer, it is shown that  $x|z$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

To prove transitivity: If  $xRy$  and  $yRz$ , then  $xRz$  for all  $x, y, z \in A$ .

Let  $x, y$  and  $z$  be numbers in the set  $A$ . Suppose that  $x|y$  and  $y|z$ .

By definition,  $y = ax$  and  $z = by$  for some integers  $a$  and  $b$ .

Substituting this value of  $y$  into the second equation gives us  $z = abx$ .

Since  $ab$  is an integer, it is shown that  $x|z$ .

Hence, the relation  $|$  is transitive.

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

Thus, the relation  $|$  is reflexive, anti-symmetrical and transitive.

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

a) Prove that  $|$  is a partial order.

Thus, the relation  $|$  is reflexive, anti-symmetrical and transitive.

Hence, the relation  $|$  is a partial order.

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

b) Draw a Hasse diagram for the partially ordered set  $(A, |)$ .


2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

b) Draw a Hasse diagram for the partially ordered set  $(A, |)$ .

Let's draw this up!



## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) Find, if they exist, all

- $\alpha)$  minimal elements.
- $\beta)$  least elements.
- $\gamma)$  maximal elements.
- $\delta)$  greatest elements.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) α) Find, if they exist, all minimal elements.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) α) Find, if they exist, all minimal elements.

Minimal Elements: There exists no element that relates to the minimal elements other than itself.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) α) Find, if they exist, all minimal elements.

Minimal Elements: There exists no element that relates to the minimal elements other than itself.

In this case, the minimal elements are the primes in  $A$ : 2, 3, 5.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) α) Find, if they exist, all minimal elements.

Minimal Elements: There exists no element that relates to the minimal elements other than itself.

In this case, the minimal elements are the primes in  $A$ : 2, 3, 5.

From the diagram, we know that all composites have an element that relate to them.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) α) Find, if they exist, all minimal elements.

Minimal Elements: There exists no element that relates to the minimal elements other than itself.

In this case, the minimal elements are the primes in  $A$ : 2, 3, 5.

From the diagram, we know that all composites have an element that relate to them. Since primes only have the factors 1 and itself, there are no elements that divide 2, 3 and 5 in  $A$  other than itself.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) α) Find, if they exist, all minimal elements.

**Minimal Elements:** There exists no element that relates to the minimal elements other than itself.

In this case, the minimal elements are the primes in  $A$ : 2, 3, 5.

From the diagram, we know that all composites have an element that relate to them. Since primes only have the factors 1 and itself, there are no elements that divide 2, 3 and 5 in  $A$  other than itself. This is because 1 is not in  $A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c)  $\beta$ ) Find, if they exist, all least elements.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c)  $\beta$ ) Find, if they exist, all least elements.

Least Elements: The element must relate to every other element.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) β) Find, if they exist, all least elements.

Least Elements: The element must relate to every other element.

Let  $a$  be the least element.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c)  $\beta$ ) Find, if they exist, all least elements.

Least Elements: The element must relate to every other element.

Let  $a$  be the least element. Thus,  $a|2$  and  $a|3$ , since 2 and 3 are elements of  $A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c)  $\beta$ ) Find, if they exist, all least elements.

Least Elements: The element must relate to every other element.

Let  $a$  be the least element. Thus,  $a|2$  and  $a|3$ , since 2 and 3 are elements of  $A$ . Since 2 and 3 are coprime,  $a$  must be 1, but 1 is not in  $A$ .

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c)  $\beta$ ) Find, if they exist, all least elements.

Least Elements: The element must relate to every other element.

Let  $a$  be the least element. Thus,  $a|2$  and  $a|3$ , since 2 and 3 are elements of  $A$ . Since 2 and 3 are coprime,  $a$  must be 1, but 1 is not in  $A$ . Therefore there exists no element of  $A$  that relates to 2 and 3.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c)  $\beta$ ) Find, if they exist, all least elements.

Least Elements: The element must relate to every other element.

Let  $a$  be the least element. Thus,  $a|2$  and  $a|3$ , since 2 and 3 are elements of  $A$ . Since 2 and 3 are coprime,  $a$  must be 1, but 1 is not in  $A$ . Therefore there exists no element of  $A$  that relates to 2 and 3.

Hence, there are no least elements.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) γ) Find, if they exist, all maximal elements.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) γ) Find, if they exist, all maximal elements.

Maximal Elements: There are no elements that the maximal elements relate to other than itself.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) γ) Find, if they exist, all maximal elements.

Maximal Elements: There are no elements that the maximal elements relate to other than itself.

From the diagram we see that 210 is the only element that does not relate to another.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) γ) Find, if they exist, all maximal elements.

Maximal Elements: There are no elements that the maximal elements relate to other than itself.

From the diagram we see that 210 is the only element that does not relate to another.

Hence, the only maximal element is 210.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) δ) Find, if they exist, all greatest elements.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) δ) Find, if they exist, all greatest elements.

Greatest Elements: Every element relates to the greatest element.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) δ) Find, if they exist, all greatest elements.

Greatest Elements: Every element relates to the greatest element.

We know that every element in  $A$  divides 210. Thus every element relates to 210.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

c) δ) Find, if they exist, all greatest elements.

Greatest Elements: Every element relates to the greatest element.

We know that every element in  $A$  divides 210. Thus every element relates to 210.

Hence, the greatest element is 210.

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

d) Find two elements of  $A$  that do not have a greatest lower bound.

Explain why that is.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

d) Find two elements of  $A$  that do not have a greatest lower bound.

Explain why that is.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

d) Find two elements of  $A$  that do not have a greatest lower bound.

Explain why that is.

Given two elements, the lower bounds are all the elements that relate to both. Thus, the greatest lower bound is the greatest element from these lower bounds.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

d) Find two elements of  $A$  that do not have a greatest lower bound.

Explain why that is.

Given two elements, the lower bounds are all the elements that relate to both. Thus, the greatest lower bound is the greatest element from these lower bounds.

Consider the greatest lower bound of 2 and 3.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

d) Find two elements of  $A$  that do not have a greatest lower bound.

Explain why that is.

Given two elements, the lower bounds are all the elements that relate to both. Thus, the greatest lower bound is the greatest element from these lower bounds.

Consider the greatest lower bound of 2 and 3. This would be an element that relates to both i.e. dividing both 2 and 3.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

d) Find two elements of  $A$  that do not have a greatest lower bound.

Explain why that is.

Given two elements, the lower bounds are all the elements that relate to both. Thus, the greatest lower bound is the greatest element from these lower bounds.

Consider the greatest lower bound of 2 and 3. This would be an element that relates to both i.e. dividing both 2 and 3. Since 2 and 3 are coprime, the only number possible which could relate to both 2 and 3 is 1.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

d) Find two elements of  $A$  that do not have a greatest lower bound.

Explain why that is.

Given two elements, the lower bounds are all the elements that relate to both. Thus, the greatest lower bound is the greatest element from these lower bounds.

Consider the greatest lower bound of 2 and 3. This would be an element that relates to both i.e. dividing both 2 and 3. Since 2 and 3 are coprime, the only number possible which could relate to both 2 and 3 is 1. We know that 1 is not in  $A$  and therefore there are no elements which relate to both 2 and 3.

## 2013 T1 Q2 (i)

Suppose that  $A = \{2, 3, 5, 6, 15, 30, 35, 70, 105, 210\}$ .

Consider the relation  $|$  on set  $A$  defined by  $a|b$  if and only if  $a$  divides  $b$ .

d) Find two elements of  $A$  that do not have a greatest lower bound.

Explain why that is.

Given two elements, the lower bounds are all the elements that relate to both. Thus, the greatest lower bound is the greatest element from these lower bounds.


Consider the greatest lower bound of 2 and 3. This would be an element that relates to both i.e. dividing both 2 and 3. Since 2 and 3 are coprime, the only number possible which could relate to both 2 and 3 is 1. We know that 1 is not in  $A$  and therefore there are no elements which relate to both 2 and 3.

Since no lower bounds exist, 2 and 3 can't have a greatest lower bound.

### 3. Enumeration and Probability

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

2016 T2 Q4 v)



## Pigeonhole Principle

Given that there are  $n + 1$  pigeons sorted across  $n$  pigeonholes then there is at least one pigeonhole with two or more pigeons.

## Pigeonhole Principle

Given that there are  $n + 1$  pigeons sorted across  $n$  pigeonholes then there is at least one pigeonhole with two or more pigeons.

Generalised Pigeonhole Principle:

With  $n$  pigeons and  $k$  pigeonholes, there is at least one pigeonhole with  $\left\lceil \frac{n}{k} \right\rceil$  or more pigeons.

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

The midpoint of  $(a, b)$  and  $(c, d)$  is

$$\left( \frac{a+c}{2}, \frac{b+d}{2} \right).$$

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

The midpoint of  $(a, b)$  and  $(c, d)$  is

$$\left( \frac{a+c}{2}, \frac{b+d}{2} \right).$$

To have the average of two integers be an integer, they must both either be odd or both be even.

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

The midpoint of  $(a, b)$  and  $(c, d)$  is

$$\left( \frac{a+c}{2}, \frac{b+d}{2} \right).$$

To have the average of two integers be an integer, they must both either be odd or both be even.

Therefore, we need to have  $a$  and  $c$  with  $b$  and  $d$  be the same parity.

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

2016 T2 Q4 v)

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

Coordinates are represented by two numbers,  $x$  and  $y$ .

2016 T2 Q4 v)

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

Coordinates are represented by two numbers,  $x$  and  $y$ .

All possible combinations of parity for  $x$  and  $y$  are given by

(odd, odd), (even, odd), (even, even), (odd, even).

2016 T2 Q4 v)

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

Coordinates are represented by two numbers,  $x$  and  $y$ .

All possible combinations of parity for  $x$  and  $y$  are given by

(odd, odd), (even, odd), (even, even), (odd, even).

Since there are 5 distinct points and only 4 possible cases of parity

## 2016 T2 Q4 v)

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

Coordinates are represented by two numbers,  $x$  and  $y$ .

All possible combinations of parity for  $x$  and  $y$  are given by

(odd, odd), (even, odd), (even, even), (odd, even).

Since there are 5 distinct points and only 4 possible cases of parity, by pigeonhole principle, there must exist 2 points which share parity for both coordinates.

2016 T2 Q4 v)

Let  $(x_i, y_i)$  for  $i = 1, 2, 3, 4, 5$  be 5 distinct points in the plane with integer coordinates. Prove that at least one midpoint of two such points has integer coordinates.

Coordinates are represented by two numbers,  $x$  and  $y$ .

All possible combinations of parity for  $x$  and  $y$  are given by

(odd, odd), (even, odd), (even, even), (odd, even).

Since there are 5 distinct points and only 4 possible cases of parity, by pigeonhole principle, there must exist 2 points which share parity for both coordinates.

These two points would make a midpoint with integer coordinates.

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- a) with no further restriction?
- b) with each child getting at least one ball and exactly 5 lollipops?

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- a) with no further restriction?

## 2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- a) with no further restriction?

Two problems:

- α) Distribute 80 identical balls among 8 children.

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

a) with no further restriction?

Two problems:

- $\alpha$ ) Distribute 80 identical balls among 8 children.
- $\beta$ ) Distribute 40 distinct lollipops among 8 children.

## 2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- a) with no further restriction?

Two problems:

- $\alpha$ ) Distribute 80 identical balls among 8 children.
- $\beta$ ) Distribute 40 distinct lollipops among 8 children.

We can get the answer to a) by multiplying the answers of each sub-problem since it would cover all combinations of distributing 80 identical balls and 40 distinct lollipops among 9 children.

2019 T3 Q3 i)

a) Distribute 80 identical balls among 8 (distinct) children.

# 2019 T3 Q3 i)

a) Distribute 80 identical balls among 8 (distinct) children.


We can represent the amount of balls each child gets by  $x_i$ .

2019 T3 Q3 i)

a) Distribute 80 identical balls among 8 (distinct) children.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$



## 2019 T3 Q3 i)

a) Distribute 80 identical balls among 8 (distinct) children.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a non-negative integer.

## 2019 T3 Q3 i)

a) Distribute 80 identical balls among 8 (distinct) children.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a non-negative integer.

The general solution for  $n$  identical objects and  $r$  distinct boxes is

$$\underline{\underline{\binom{n+r-1}{r-1}}}$$

## 2019 T3 Q3 i)

a) Distribute 80 identical balls among 8 (distinct) children.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a non-negative integer.

The general solution for  $n$  identical objects and  $r$  distinct boxes is

$$\binom{n+r-1}{r-1} \Rightarrow \binom{80+8-1}{8-1} = \binom{87}{7},$$

## 2019 T3 Q3 i)

a) Distribute 80 identical balls among 8 (distinct) children.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a non-negative integer.

The general solution for  $n$  identical objects and  $r$  distinct boxes is

$$\binom{n+r-1}{r-1} \Rightarrow \binom{80+8-1}{8-1} = \binom{87}{7},$$

where there are  $r = 8$  children (boxes) and  $n = 80$  balls (objects).

# 2019 T3 Q3 i)

a) Distribute 80 identical balls among 8 (distinct) children.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a non-negative integer.

The general solution for  $n$  identical objects and  $r$  distinct boxes is

$$\binom{n+r-1}{r-1} \Rightarrow \binom{80+8-1}{8-1} = \boxed{\binom{87}{7}},$$

where there are  $r = 8$  children (boxes) and  $n = 80$  balls (objects).

This formula only works when  $x_i \geq 0$  i.e. can be 0

$\beta)$  Distribute 40 distinct lollipops among 8 children.

$\beta)$  Distribute 40 distinct lollipops among 8 children.

The general solution for  $n$  distinct objects and  $r$  distinct boxes is

$$\bullet \overset{r^n}{\bullet}$$

$\beta)$  Distribute 40 distinct lollipops among 8 children.

The general solution for  $n$  **distinct objects** and  $r$  **distinct boxes** is

$$r^n \Rightarrow 8^{40}$$

$\beta)$  Distribute 40 distinct lollipops among 8 children.

The general solution for  $n$  **distinct objects** and  $r$  **distinct boxes** is

$$r^n \Rightarrow 8^{40}$$

where there are  $r = 8$  children (boxes) and  $n = 40$  lollipops (objects).

$\beta)$  Distribute 40 distinct lollipops among 8 children.

The general solution for  $n$  **distinct objects** and  $r$  **distinct boxes** is

$$r^n \Rightarrow 8^{40}$$

where there are  $r = 8$  children (boxes) and  $n = 40$  lollipops (objects).

The reasoning behind this is the fact that each distinct object has  $r$  choices to go to.

$\beta)$  Distribute 40 distinct lollipops among 8 children.

The general solution for  $n$  **distinct objects** and  $r$  **distinct boxes** is

$$r^n \Rightarrow 8^{40}$$

where there are  $r = 8$  children (boxes) and  $n = 40$  lollipops (objects).

The reasoning behind this is the fact that each distinct object has  $r$  choices to go to. In this case, each lollipop has a choice from 8 children to go to.

β) Distribute 40 distinct lollipops among 8 children.

The general solution for  $n$  **distinct objects** and  $r$  **distinct boxes** is

$$r^n \Rightarrow 8^{40}$$

where there are  $r = 8$  children (boxes) and  $n = 40$  lollipops (objects).

The reasoning behind this is the fact that each distinct object has  $r$  choices to go to. In this case, each lollipop has a choice from 8 children to go to. 40 of these lollipops gives us  $8^{40}$  by multiplication principle.

## 2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- a) with no further restriction?

2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

a) with no further restriction?

Since we know that a) has

$$\binom{87}{7}$$

different ways to distribute the 80 identical balls

2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

a) with no further restriction?

Since we know that  $\alpha$ ) has

$$\binom{87}{7}$$

different ways to distribute the 80 identical balls and  $\beta$ ) has

$$8^{40}$$

different ways to distribute the 40 distinct lollipops.

2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

a) with no further restriction?

Since we know that  $\alpha$ ) has

$$\binom{87}{7}$$

different ways to distribute the 80 identical balls and  $\beta$ ) has

$$8^{40}$$

different ways to distribute the 40 distinct lollipops.

Hence, there are

$$\binom{87}{7} \times 8^{40}$$

different ways to distribute the objects among 8 children.

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

b) with each child getting at least one ball and exactly 5 lollipops?

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- b) with each child getting at least one ball and exactly 5 lollipops?

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

b) with each child getting at least one ball and exactly 5 lollipops?

Two problems:

a) Distribute 80 identical balls among 8 children

## 2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- b) with each child getting at least one ball and exactly 5 lollipops?

Two problems:

- a) Distribute 80 identical balls among 8 children

AND each child must get at least one ball. 

## 2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- b) with each child getting at least one ball and exactly 5 lollipops?

Two problems:

- $\alpha$ ) Distribute 80 identical balls among 8 children

AND each child must get at least one ball.

- $\beta$ ) Distribute 40 distinct lollipops among 8 children

## 2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- b) with each child getting at least one ball and exactly 5 lollipops?

Two problems:

- $\alpha$ ) Distribute 80 identical balls among 8 children

AND each child must get at least one ball.

- $\beta$ ) Distribute 40 distinct lollipops among 8 children

AND each child gets exactly 5 lollipops.



2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

b) with each child getting at least one ball and exactly 5 lollipops?

Two problems:

$\alpha$ ) Distribute 80 identical balls among 8 children

AND each child must get at least one ball.

$\beta$ ) Distribute 40 distinct lollipops among 8 children

AND each child gets exactly 5 lollipops.

Again, we can get the answer to b) by multiplying  $\alpha$  and  $\beta$ ).

- a) Distribute 80 identical balls among 8 children  
AND each child must get at least one ball.

2019 T3 Q3 i)

- a) Distribute 80 identical balls among 8 children  
AND each child must get at least one ball.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$



where all  $x_i$  is a positive integer.

2019 T3 Q3 i)

- a) Distribute 80 identical balls among 8 children  
AND each child must get at least one ball.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a positive integer.

The general solution for  $n$  identical objects and  $r$  distinct boxes is

$$\binom{n-1}{r-1}$$

## 2019 T3 Q3 i)

- a) Distribute 80 identical balls among 8 children  
AND each child must get at least one ball.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a positive integer.

The general solution for  $n$  identical objects and  $r$  distinct boxes is

$$\binom{n-1}{r-1} \Rightarrow \binom{80-1}{8-1} = \binom{79}{7},$$

## 2019 T3 Q3 i)

- a) Distribute 80 identical balls among 8 children  
AND each child must get at least one ball.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a positive integer.

The general solution for  $n$  identical objects and  $r$  distinct boxes is

$$\binom{n-1}{r-1} \Rightarrow \binom{80-1}{8-1} = \binom{79}{7},$$

where there are  $r = 8$  children (boxes) and  $n = 80$  balls (objects).

### 2019 T3 Q3 i)

- a) Distribute 80 identical balls among 8 children  
AND each child must get at least one ball.

We can represent the amount of balls each child gets by  $x_i$ . Thus

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 + x_8 = 80,$$

where all  $x_i$  is a positive integer.

The general solution for  $n$  identical objects and  $r$  distinct boxes is

$$\binom{n-1}{r-1} \Rightarrow \binom{80-1}{8-1} = \binom{79}{7},$$

where there are  $r = 8$  children (boxes) and  $n = 80$  balls (objects).

This formula only works when  $x_i$  must be positive i.e. can't be 0.

2019 T3 Q3 i)

- β) Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

2019 T3 Q3 i)

- $\beta)$  Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We can have each child choose 5 lollipops from the pile.

## 2019 T3 Q3 i)

- β) Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We can have each child choose 5 lollipops from the pile.

This means that the first child has  ${}^{40}C_5$  ways to choose.

## 2019 T3 Q3 i)

- β) Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We can have each child choose 5 lollipops from the pile.

This means that the first child has  ${}^{40}C_5$  ways to choose.

After every choice, there are 5 less lollipops so the second person has  ${}^{40-5}C_5$  ways to choose.

## 2019 T3 Q3 i)

- β) Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We can have each child choose 5 lollipops from the pile.

This means that the first child has  ${}^{40}C_5$  ways to choose.

After every choice, there are 5 less lollipops so the second person has  ${}^{40-5}C_5$  ways to choose.

We can do this for each child to obtain the expression

## 2019 T3 Q3 i)

- $\beta)$  Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We can have each child choose 5 lollipops from the pile.

This means that the first child has  ${}^{40}C_5$  ways to choose.

After every choice, there are 5 less lollipops so the second person has  ${}^{40-5}C_5$  ways to choose.

We can do this for each child to obtain the expression

$$\binom{40}{5} \times \binom{35}{5} \times \dots \times \binom{10}{5} \times \binom{5}{5},$$

## 2019 T3 Q3 i)

- $\beta)$  Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We can have each child choose 5 lollipops from the pile.

This means that the first child has  ${}^{40}C_5$  ways to choose.

After every choice, there are 5 less lollipops so the second person has  ${}^{40-5}C_5$  ways to choose.

We can do this for each child to obtain the expression

$$\binom{40}{5} \times \binom{35}{5} \times \dots \times \binom{10}{5} \times \binom{5}{5},$$

i.e. how many ways you can distribute 40 lollipops so each child gets 5.

$\beta)$  Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

## 2019 T3 Q3 i)

$\beta)$  Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We know that

$$\binom{n}{r} = \frac{n!}{(n-r)!r!}.$$

β) Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We know that

$$\binom{n}{r} = \frac{n!}{(n-r)!r!}.$$

Thus,

$$\binom{40}{5} \times \binom{35}{5} \times \dots \times \binom{10}{5} \times \binom{5}{5} = \frac{40!}{\cancel{35!} \cancel{30!} \dots \cancel{10!} \cancel{5!}} \times \frac{35!}{\cancel{35!} \cancel{30!} \dots \cancel{10!} \cancel{5!}} \times \dots \times \frac{10!}{\cancel{10!} \cancel{5!}} \times \frac{5!}{\cancel{5!} \cancel{0!}}$$

## 2019 T3 Q3 i)

β) Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We know that

$$\binom{n}{r} = \frac{n!}{(n-r)!r!}.$$

Thus,

$$\begin{aligned}\binom{40}{5} \times \binom{35}{5} \times \dots \times \binom{10}{5} \times \binom{5}{5} &= \frac{40!}{35!5!} \times \frac{35!}{30!5!} \times \dots \times \frac{10!}{5!5!} \times \frac{5!}{0!5!} \\ &= \frac{40!}{35!5!} \times \frac{35!}{30!5!} \times \dots \times \frac{10!}{5!5!} \times \frac{5!}{0!5!}\end{aligned}$$

2019 T3 Q3 i)

β) Distribute 40 distinct lollipops among 8 children  
AND each child gets exactly 5 lollipops.

We know that

$$\binom{n}{r} = \frac{n!}{(n-r)!r!}.$$

Thus,

$$\begin{aligned}\binom{40}{5} \times \binom{35}{5} \times \dots \times \binom{10}{5} \times \binom{5}{5} &= \frac{40!}{35!5!} \times \frac{35!}{30!5!} \times \dots \times \frac{10!}{5!5!} \times \frac{5!}{0!5!} \\ &= \frac{40!}{35!5!} \times \frac{35!}{30!5!} \times \dots \times \frac{10!}{5!5!} \times \frac{5!}{0!5!} \\ &= \boxed{\frac{40!}{(5!)^8}} \text{ ways to distribute.}\end{aligned}$$

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

- b) with each child getting at least one ball and exactly 5 lollipops?

2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

b) with each child getting at least one ball and exactly 5 lollipops?

Since we know that  $\alpha$ ) has

$$\binom{79}{7}$$

different ways to distribute the 80 identical balls

2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

b) with each child getting at least one ball and exactly 5 lollipops?

Since we know that  $\alpha$ ) has

$$\binom{79}{7}$$

different ways to distribute the 80 identical balls and  $\beta$ ) has

$$\frac{40!}{(5!)^8}$$

different ways to distribute the 40 distinct lollipops.

2019 T3 Q3 i)

How many different ways are there to distribute 80 identical balls and 40 distinct lollipops among 8 children:

b) with each child getting at least one ball and exactly 5 lollipops?

Since we know that  $\alpha$ ) has

$$\binom{79}{7}$$

different ways to distribute the 80 identical balls and  $\beta$ ) has

$$\frac{40!}{(5!)^8}$$

different ways to distribute the 40 distinct lollipops.

Hence, there are

$$\binom{79}{7} \times \frac{40!}{(5!)^8}$$

different ways to distribute the objects among 8 children.

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

We want to make it so each term can be represented by consecutive integers.

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

We want to make it so each term can be represented by consecutive integers. Being odd restricts this as you cannot have a range like 0,1,2,3,... and must adhere to 1,3,5,....

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

We want to make it so each term can be represented by consecutive integers. Being odd restricts this as you cannot have a range like 0,1,2,3,... and must adhere to 1,3,5,....

To fix this, we can rewrite  $x_i = 2y_i + 1$ , where  $y_i$  is an integer.

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

We want to make it so each term can be represented by consecutive integers. Being odd restricts this as you cannot have a range like 0,1,2,3,... and must adhere to 1,3,5,....

To fix this, we can rewrite  $x_i = 2y_i + 1$ , where  $y_i$  is an integer. We know  $x_i \geq 1$  as it cannot be 0 since it is odd.

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

We want to make it so each term can be represented by consecutive integers. Being odd restricts this as you cannot have a range like 0,1,2,3,... and must adhere to 1,3,5,....

To fix this, we can rewrite  $x_i = 2y_i + 1$ , where  $y_i$  is an integer. We know  $x_i \geq 1$  as it cannot be 0 since it is odd. Thus  $\underline{2y_i + 1 \geq 1 \Rightarrow y_i \geq 0}$ .

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

We want to make it so each term can be represented by consecutive integers. Being odd restricts this as you cannot have a range like 0,1,2,3,... and must adhere to 1,3,5,....

To fix this, we can rewrite  $x_i = 2y_i + 1$ , where  $y_i$  is an integer. We know  $x_i \geq 1$  as it cannot be 0 since it is odd. Thus  $2y_i + 1 \geq 1 \Rightarrow y_i \geq 0$ .

Since  $x_i \leq 7$ ,  $2y_i + 1 \leq 7$ . Hence,  $y_i \leq 3$ .

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

Using this substitution of  $x_i = 2y_i + 1$ ,

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

Using this substitution of  $x_i = 2y_i + 1$ ,

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36$$

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

Using this substitution of  $x_i = 2y_i + 1$ ,

$$\begin{aligned} x_1 + x_2 + x_3 + x_4 + x_5 + x_6 &= 36 \\ (2y_1 + 1) + (2y_2 + 1) + \dots + (2y_6 + 1) &= \end{aligned}$$

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

Using this substitution of  $x_i = 2y_i + 1$ ,

$$\begin{aligned} x_1 + x_2 + x_3 + x_4 + x_5 + x_6 &= 36 \\ (2y_1 + 1) + (2y_2 + 1) + \dots + (2y_6 + 1) &= \\ 2(y_1 + y_2 + y_3 + y_4 + y_5 + y_6) + 6 &= 36 \end{aligned}$$


2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

Using this substitution of  $x_i = 2y_i + 1$ ,

$$\begin{aligned} x_1 + x_2 + x_3 + x_4 + x_5 + x_6 &= 36 \\ (2y_1 + 1) + (2y_2 + 1) + \dots + (2y_6 + 1) &= \\ \underline{\bullet} 2(y_1 + y_2 + y_3 + y_4 + y_5 + y_6) + \underline{\bullet} 6 &= 36 \quad \bullet \\ y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= \underline{\bullet} 15. \end{aligned}$$

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?

2017 T1 Q4 iii)

Consider the equation

$$x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 36,$$

where  $x_1, x_2, x_3, x_4, x_5, x_6$  are non-negative integers.

How many solutions are there with all the  $x_i$  being odd and  $x_i \leq 7$ ?



Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

Instead we can look at the Total - Complement.

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

Instead we can look at the Total - Complement. We know that the amount of solutions for all  $y_i \leq 3$  is equivalent to the total number of solutions minus the solutions where some  $y_i > 3 \Rightarrow y_i \geq 4$ .

2017 T1 Q4 iii)

Consider the equation

$$\underline{y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15},$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

Instead we can look at the Total - Complement. We know that the amount of solutions for all  $y_i \leq 3$  is equivalent to the total number of solutions minus the solutions where some  $y_i > 3 \Rightarrow y_i \geq 4$ .

From the formula before, with no restrictions there are a total of

$$\binom{15 + 6 - 1}{6 - 1} = \binom{20}{5}$$

solutions for the equation above.

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

To find all the solutions for the complement, where some  $y_i \geq 4$

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

To find all the solutions for the complement, where some  $y_i \geq 4$ , we need to find out when there is one such  $y_i$ , there are two such  $y_i$  and there are three such  $y_i$ .

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

To find all the solutions for the complement, where some  $y_i \geq 4$ , we need to find out when there is one such  $y_i$ , there are two such  $y_i$  and there are three such  $y_i$ . The reason there cannot be four is because there are only 15 objects so there cannot be four or more  $y_i \geq 4$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For one such  $y_i$ , we can consider the case that  $y_1 \geq 4$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For one such  $y_i$ , we can consider the case that  $y_1 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$  and for every other  $y_i = a_i$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For one such  $y_i$ , we can consider the case that  $y_1 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$  and for every other  $y_i = a_i$ . This give us

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15$$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For one such  $y_i$ , we can consider the case that  $y_1 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$  and for every other  $y_i = a_i$ . This give us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + a_2 + a_3 + a_4 + a_5 + a_6 &= 15 \end{aligned}$$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For one such  $y_i$ , we can consider the case that  $y_1 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$  and for every other  $y_i = a_i$ . This give us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + a_2 + a_3 + a_4 + a_5 + a_6 &= 15 \\ a_1 + a_2 + a_3 + a_4 + a_5 + a_6 &= 11, \end{aligned}$$

where  $a_i \geq 0$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For one such  $y_i$ , we can consider the case that  $y_1 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$  and for every other  $y_i = a_i$ . This give us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + a_2 + a_3 + a_4 + a_5 + a_6 &= 15 \\ a_1 + a_2 + a_3 + a_4 + a_5 + a_6 &= 11, \end{aligned}$$

where  $a_i \geq 0$ . Therefore the total amount of solutions is  ${}^{16}C_5$ .

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For two such  $y_i$ , we can consider the case that  $y_1, y_2 \geq 4$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For two such  $y_i$ , we can consider the case that  $y_1, y_2 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$  and for every other  $y_i = a_i$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For two such  $y_i$ , we can consider the case that  $y_1, y_2 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$  and for every other  $y_i = a_i$ . This gives us

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15$$

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For two such  $y_i$ , we can consider the case that  $y_1, y_2 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$  and for every other  $y_i = a_i$ . This gives us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + (a_2 + 4) + a_3 + a_4 + a_5 + a_6 &= 15 \end{aligned}$$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For two such  $y_i$ , we can consider the case that  $y_1, y_2 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$  and for every other  $y_i = a_i$ . This gives us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + (a_2 + 4) + a_3 + a_4 + a_5 + a_6 &= 15 \\ \underline{\underline{a_1 + a_2 + a_3 + a_4 + a_5 + a_6}} &= 7, \end{aligned}$$

where  $a_i \geq 0$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For two such  $y_i$ , we can consider the case that  $y_1, y_2 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$  and for every other  $y_i = a_i$ . This gives us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + (a_2 + 4) + a_3 + a_4 + a_5 + a_6 &= 15 \\ a_1 + a_2 + a_3 + a_4 + a_5 + a_6 &= 7, \end{aligned}$$

where  $a_i \geq 0$ . Therefore the total amount of solutions is  ${}^{12}C_5$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For three such  $y_i$ , we can consider the case that  $y_1, y_2, y_3 \geq 4$ .

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For three such  $y_i$ , we can consider the case that  $y_1, y_2, y_3 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$ ,  $y_3 = a_3 + 4$  and for every other  $y_i = a_i$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For three such  $y_i$ , we can consider the case that  $y_1, y_2, y_3 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$ ,  $y_3 = a_3 + 4$  and for every other  $y_i = a_i$ . This gives us

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15$$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For three such  $y_i$ , we can consider the case that  $y_1, y_2, y_3 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$ ,  $y_3 = a_3 + 4$  and for every other  $y_i = a_i$ . This gives us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + (a_2 + 4) + (a_3 + 4) + a_4 + a_5 + a_6 &= 15 \end{aligned}$$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For three such  $y_i$ , we can consider the case that  $y_1, y_2, y_3 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$ ,  $y_3 = a_3 + 4$  and for every other  $y_i = a_i$ . This gives us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + (a_2 + 4) + (a_3 + 4) + a_4 + a_5 + a_6 &= 15 \\ a_1 + a_2 + a_3 + a_4 + a_5 + a_6 &= 3, \end{aligned}$$

where  $a_i \geq 0$ .

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For three such  $y_i$ , we can consider the case that  $y_1, y_2, y_3 \geq 4$ . Let's make the substitution that  $y_1 = a_1 + 4$ ,  $y_2 = a_2 + 4$ ,  $y_3 = a_3 + 4$  and for every other  $y_i = a_i$ . This gives us

$$\begin{aligned} y_1 + y_2 + y_3 + y_4 + y_5 + y_6 &= 15 \\ (a_1 + 4) + (a_2 + 4) + (a_3 + 4) + a_4 + a_5 + a_6 &= 15 \\ a_1 + a_2 + a_3 + a_4 + a_5 + a_6 &= 3, \end{aligned}$$

where  $a_i \geq 0$ . Therefore the total amount of solutions is  ${}^8C_5$ .

## The Inclusion-Exclusion Principle

$$\begin{aligned}|A_1 \cup A_2 \cup \dots \cup A_n| &= |A_1| + |A_2| + \dots + |A_n| \\&\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_{n-1} \cap A_n| \\&\quad + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n| + \dots \\&\quad \dots \\&\quad + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|\end{aligned}$$

## The Inclusion-Exclusion Principle

$$\begin{aligned}|A_1 \cup A_2 \cup \dots \cup A_n| &= |A_1| + |A_2| + \dots + |A_n| \\&\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_{n-1} \cap A_n| \\&\quad + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n| + \dots \\&\quad \dots \\&\quad + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|\end{aligned}$$

In our case,  $A_i$  represents when  $y_i \geq 4$ .

2017 T1 Q4 iii)

## The Inclusion-Exclusion Principle

$$\begin{aligned}|A_1 \cup A_2 \cup \dots \cup A_n| &= |A_1| + |A_2| + \dots + |A_n| \\&\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_{n-1} \cap A_n| \\&\quad + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n| + \dots \\&\quad \dots \\&\quad + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|\end{aligned}$$

In our case,  $A_i$  represents when  $y_i \geq 4$ . So  $A_1$  is when  $y_1 \geq 4$

## The Inclusion-Exclusion Principle

$$\begin{aligned}|A_1 \cup A_2 \cup \dots \cup A_n| &= |A_1| + |A_2| + \dots + |A_n| \\&\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_{n-1} \cap A_n| \\&\quad + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n| + \dots \\&\quad \dots \\&\quad + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|\end{aligned}$$

In our case,  $A_i$  represents when  $y_i \geq 4$ . So  $A_1$  is when  $y_1 \geq 4$  and  $A_1 \cap A_2$  is when both  $y_1, y_2 \geq 4$ .

2017 T1 Q4 iii)

## The Inclusion-Exclusion Principle

$$\begin{aligned}|A_1 \cup A_2 \cup \dots \cup A_n| &= |A_1| + |A_2| + \dots + |A_n| \\&\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_{n-1} \cap A_n| \\&\quad + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n| + \dots \\&\quad \dots \\&\quad + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|\end{aligned}$$

In our case,  $A_i$  represents when  $y_i \geq 4$ . So  $A_1$  is when  $y_1 \geq 4$  and  $A_1 \cap A_2$  is when both  $y_1, y_2 \geq 4$ .

For us,  $|A_1| = |A_2|$

2017 T1 Q4 iii)

## The Inclusion-Exclusion Principle

$$\begin{aligned}|A_1 \cup A_2 \cup \dots \cup A_n| &= |A_1| + |A_2| + \dots + |A_n| \\&\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_{n-1} \cap A_n| \\&\quad + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n| + \dots \\&\quad \dots \\&\quad + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|\end{aligned}$$

In our case,  $A_i$  represents when  $y_i \geq 4$ . So  $A_1$  is when  $y_1 \geq 4$  and  $A_1 \cap A_2$  is when both  $y_1, y_2 \geq 4$ .

For us,  $|A_1| = |A_2|$  since we can follow the same process for  $A_i$  that we did with  $A_1$  to derive the same amount of solutions.

## The Inclusion-Exclusion Principle

$$\begin{aligned}|A_1 \cup A_2 \cup \dots \cup A_n| &= |A_1| + |A_2| + \dots + |A_n| \\&\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_{n-1} \cap A_n| \\&\quad + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n| + \dots \\&\quad \dots \\&\quad + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|\end{aligned}$$

In our case,  $A_i$  represents when  $y_i \geq 4$ . So  $A_1$  is when  $y_1 \geq 4$  and  $A_1 \cap A_2$  is when both  $y_1, y_2 \geq 4$ .

For us,  $|A_1| = |A_2|$  since we can follow the same process for  $A_i$  that we did with  $A_1$  to derive the same amount of solutions. We can also do the same for  $A_i \cap A_j$  and  $A_i \cap A_j \cap A_k$ .

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For all cases of some  $y_i \geq 4$ , from inclusion-exclusion principle, we have

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For all cases of some  $y_i \geq 4$ , from inclusion-exclusion principle, we have

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_6| &= |A_1| + |A_2| + \dots + |A_6| \\ &\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_5 \cap A_6| \\ &\quad + |A_1 \cap A_2 \cap A_3| + \dots + |A_4 \cap A_5 \cap A_6| \end{aligned}$$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For all cases of some  $y_i \geq 4$ , from inclusion-exclusion principle, we have

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_6| &= \cancel{|A_1|} + \cancel{|A_2|} + \dots + \cancel{|A_6|} && \text{Red arrow pointing to } \cancel{|A_1|} \\ &\quad - \cancel{|A_1 \cap A_2|} - \cancel{|A_1 \cap A_3|} - \dots - \cancel{|A_5 \cap A_6|} && \text{Red arrow pointing to } \cancel{|A_1 \cap A_2|} \\ &\quad + \cancel{|A_1 \cap A_2 \cap A_3|} + \dots + \cancel{|A_4 \cap A_5 \cap A_6|} && \text{Red arrow pointing to } \cancel{|A_1 \cap A_2 \cap A_3|} \\ &= {}^6C_1|A_1| - {}^6C_2|A_1 \cap A_2| + {}^6C_3|A_1 \cap A_2 \cap A_3| \end{aligned}$$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For all cases of some  $y_i \geq 4$ , from inclusion-exclusion principle, we have

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_6| &= |A_1| + |A_2| + \dots + |A_6| \\ &\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_5 \cap A_6| \\ &\quad + |A_1 \cap A_2 \cap A_3| + \dots + |A_4 \cap A_5 \cap A_6| \\ &= {}^6C_1|A_1| - {}^6C_2|A_1 \cap A_2| + {}^6C_3|A_1 \cap A_2 \cap A_3| \\ &= {}^6C_1{}^16C_5 - {}^6C_2{}^12C_5 + {}^6C_3{}^8C_5 \end{aligned}$$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For all cases of some  $y_i \geq 4$ , from inclusion-exclusion principle, we have

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_6| &= |A_1| + |A_2| + \dots + |A_6| \\ &\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_5 \cap A_6| \\ &\quad + |A_1 \cap A_2 \cap A_3| + \dots + |A_4 \cap A_5 \cap A_6| \\ &= {}^6C_1|A_1| - {}^6C_2|A_1 \cap A_2| + {}^6C_3|A_1 \cap A_2 \cap A_3| \\ &= {}^6C_1{}^16C_5 - {}^6C_2{}^12C_5 + {}^6C_3{}^8C_5 \end{aligned}$$

Hence, we have  ${}^{20}C_5$

2017 T1 Q4 iii)

Consider the equation

$$y_1 + y_2 + y_3 + y_4 + y_5 + y_6 = 15,$$

where  $y_i \geq 0$  is an integer. How many solutions for all  $y_i \leq 3$ ?

For all cases of some  $y_i \geq 4$ , from inclusion-exclusion principle, we have

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_6| &= |A_1| + |A_2| + \dots + |A_6| \\ &\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_5 \cap A_6| \\ &\quad + |A_1 \cap A_2 \cap A_3| + \dots + |A_4 \cap A_5 \cap A_6| \\ &= {}^6C_1|A_1| - {}^6C_2|A_1 \cap A_2| + {}^6C_3|A_1 \cap A_2 \cap A_3| \\ &= {}^6C_1{}^16C_5 - {}^6C_2{}^12C_5 + {}^6C_3{}^8C_5 \end{aligned}$$

Hence, we have  ${}^{20}C_5 - ({}^6C_1{}^16C_5 - {}^6C_2{}^12C_5 + {}^6C_3{}^8C_5) = 86$  solutions.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

- a) How many such words are there altogether?
- b) How many words in  $S$  contain the sub-words JULIA or TONY?
- c) How many words in  $S$  contain all five vowels?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

a) How many such words are there altogether?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

a) How many such words are there altogether?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

a) How many such words are there altogether?

We have 26 letters where repetition is not allowed.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

- a) How many such words are there altogether?

We have 26 letters where repetition is not allowed. This is just choosing 14 letters from the 26 and then ordering them.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

a) How many such words are there altogether?

We have 26 letters where repetition is not allowed. This is just choosing 14 letters from the 26 and then ordering them.

Hence, within the set  $S$ , there are  ${}^{26}P_{14}$  words altogether.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

These words hint at the use of inclusion-exclusion

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

These words hint at the use of inclusion-exclusion such that

$$|A_{JULIA} \cup A_{TONY}| = |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}|$$

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

These words hint at the use of inclusion-exclusion such that

$$|A_{JULIA} \cup A_{TONY}| = |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}|$$

where  $A_{JULIA}$  are the words that contain JULIA and  $A_{TONY}$  are the words that contain TONY.

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

These words hint at the use of inclusion-exclusion such that

$$|A_{JULIA} \cup A_{TONY}| = |A_{JULIA}| + |A_{TONY}| - \underline{|A_{JULIA} \cap A_{TONY}|}$$

where  $A_{JULIA}$  are the words that contain JULIA and  $A_{TONY}$  are the words that contain TONY.

Thus, to find how many words with JULIA or TONY

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

These words hint at the use of inclusion-exclusion such that

$$|A_{JULIA} \cup A_{TONY}| = |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}|$$

where  $A_{JULIA}$  are the words that contain JULIA and  $A_{TONY}$  are the words that contain TONY.

Thus, to find how many words with JULIA or TONY, we need all the words with JULIA

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

These words hint at the use of inclusion-exclusion such that

$$|A_{JULIA} \cup A_{TONY}| = |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}|$$

where  $A_{JULIA}$  are the words that contain JULIA and  $A_{TONY}$  are the words that contain TONY.

Thus, to find how many words with JULIA or TONY, we need all the words with JULIA, all the with with TONY

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

These words hint at the use of inclusion-exclusion such that

$$|A_{JULIA} \cup A_{TONY}| = |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}|$$

where  $A_{JULIA}$  are the words that contain JULIA and  $A_{TONY}$  are the words that contain TONY.

Thus, to find how many words with JULIA or TONY, we need all the words with JULIA, all the with with TONY and all the words that have both JULIA and TONY as subwords.

---

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA, consider

J U L I A - - - - -  
• • • • .

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA, consider

[ J U L I A ] -----

where the letters for JULIA have been chosen, so 9 spaces are left

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA, consider

[ J U L I A ] -----

where the letters for JULIA have been chosen, so 9 spaces are left from the remaining 21 letters in the alphabet (no repetition).

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA, consider

J [ I U L I A ] -----

where the letters for JULIA have been chosen, so 9 spaces are left from the remaining 21 letters in the alphabet (no repetition). This gives us  ${}^{21}C_9$  ways to choose these letters.

To arrange these, we want to preserve the order of JULIA

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA, consider

[ J U L I A ] -----

where the letters for JULIA have been chosen, so 9 spaces are left from the remaining 21 letters in the alphabet (no repetition). This gives us  ${}^{21}C_9$  ways to choose these letters.

To arrange these, we want to preserve the order of JULIA, so we can treat this as a letter

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA, consider

[ J U L I A ] -----

where the letters for JULIA have been chosen, so 9 spaces are left from the remaining 21 letters in the alphabet (no repetition). This gives us  ${}^{21}C_9$  ways to choose these letters.

To arrange these, we want to preserve the order of JULIA, so we can treat this as a letter, giving us 10 objects and  $10!$  arrangements.

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA, consider

[ J U L I A ] -----

where the letters for JULIA have been chosen, so 9 spaces are left from the remaining 21 letters in the alphabet (no repetition). This gives us  ${}^{21}C_9$  ways to choose these letters.

To arrange these, we want to preserve the order of JULIA, so we can treat this as a letter, giving us 10 objects and  $10!$  arrangements.

Hence, there are  ${}^{21}C_9 \times 10!$  words with JULIA.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with TONY, consider

[ T O N Y ] -----

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with TONY, consider

[ T O N Y ] -----

where the letters for TONY have been chosen, so 10 spaces are left from the remaining 22 letters in the alphabet (no repetition). This gives us  ${}^{22}C_{10}$  ways to choose these letters.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with TONY, consider

[ T O N Y ] -----

where the letters for TONY have been chosen, so 10 spaces are left from the remaining 22 letters in the alphabet (no repetition). This gives us  ${}^{22}C_{10}$  ways to choose these letters.

To arrange these, we want to preserve the order of TONY, so we can treat this as a letter, giving us 11 objects and  $11!$  arrangements.

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with TONY, consider

[ T O N Y ] -----

where the letters for TONY have been chosen, so 10 spaces are left from the remaining 22 letters in the alphabet (no repetition). This gives us  ${}^{22}C_{10}$  ways to choose these letters.

To arrange these, we want to preserve the order of TONY, so we can treat this as a letter, giving us 11 objects and  $11!$  arrangements.

Hence, there are  ${}^{22}C_{10} \times 11!$  words with TONY.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA and TONY

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA and TONY, consider

[ J U L I A ] [ T O N Y ] \_ \_ \_ \_

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA and TONY, consider

[ J U L I A ] [ T O N Y ] -----

where the letters for JULIA and TONY have been chosen, so 5 spaces are left from the remaining 17 letters in the alphabet (no repetition). This gives us  ${}^{17}C_5$  ways to choose these letters.

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA and TONY, consider

[ J U L I A ] [ T O N Y ] -----

where the letters for JULIA and TONY have been chosen, so 5 spaces are left from the remaining 17 letters in the alphabet (no repetition). This gives us  ${}^{17}C_5$  ways to choose these letters.

We want to preserve both JULIA and TONY

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA and TONY, consider

[ J U L I A ] [ T O N Y ] \_ \_ \_ \_

where the letters for JULIA and TONY have been chosen, so 5 spaces are left from the remaining 17 letters in the alphabet (no repetition). This gives us  ${}^{17}C_5$  ways to choose these letters.

We want to preserve both JULIA and TONY so they are both individual letters.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA and TONY, consider

[ J U L I A ] [ T O N Y ] \_ \_ \_ \_

where the letters for JULIA and TONY have been chosen, so 5 spaces are left from the remaining 17 letters in the alphabet (no repetition). This gives us  ${}^{17}C_5$  ways to choose these letters.

We want to preserve both JULIA and TONY so they are both individual letters. This gives us 7 objects and  $7!$  arrangements.

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

To find the words with JULIA and TONY, consider

[ J U L I A ] [ T O N Y ] \_ \_ \_ \_

where the letters for JULIA and TONY have been chosen, so 5 spaces are left from the remaining 17 letters in the alphabet (no repetition). This gives us  ${}^{17}C_5$  ways to choose these letters.

We want to preserve both JULIA and TONY so they are both individual letters. This gives us 7 objects and  $7!$  arrangements.

Hence, there are  ${}^{17}C_5 \times 7!$  words with JULIA and TONY.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

From inclusion-exclusion principle,

$$|A_{JULIA} \cup A_{TONY}| = |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}|$$

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

From inclusion-exclusion principle,

$$\begin{aligned}|A_{JULIA} \cup A_{TONY}| &= |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}| \\&= {}^{21}C_9 \times 10! + {}^{22}C_{10} \times 11! - {}^{17}C_5 \times 7!,\end{aligned}$$

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

From inclusion-exclusion principle,

$$\begin{aligned}|A_{JULIA} \cup A_{TONY}| &= |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}| \\&= {}^{21}C_9 \times 10! + {}^{22}C_{10} \times 11! - {}^{17}C_5 \times 7!,\end{aligned}$$

where  $A_{JULIA}$  are the words that contain JULIA and  $A_{TONY}$  are the words that contain TONY.

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

b) How many words in  $S$  contain the sub-words JULIA or TONY?

From inclusion-exclusion principle,

$$\begin{aligned}|A_{JULIA} \cup A_{TONY}| &= |A_{JULIA}| + |A_{TONY}| - |A_{JULIA} \cap A_{TONY}| \\&= {}^{21}C_9 \times 10! + {}^{22}C_{10} \times 11! - {}^{17}C_5 \times 7!,\end{aligned}$$

where  $A_{JULIA}$  are the words that contain JULIA and  $A_{TONY}$  are the words that contain TONY.

Hence, there are  ${}^{21}C_9 \times 10! + {}^{22}C_{10} \times 11! - {}^{17}C_5 \times 7!$  such words.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

c) How many words in  $S$  contain all five vowels?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

c) How many words in  $S$  contain all five vowels?

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

c) How many words in  $S$  contain all five vowels?

Suppose that all 5 vowels have already been chosen for the word.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

c) How many words in  $S$  contain all five vowels?

Suppose that all 5 vowels have already been chosen for the word. This leaves us with 9 more letters for the 14-letter word and a choice from 21 letters left over, since there is no repetition.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

c) How many words in  $S$  contain all five vowels?

Suppose that all 5 vowels have already been chosen for the word. This leaves us with 9 more letters for the 14-letter word and a choice from 21 letters left over, since there is no repetition.

Thus, there are only  ${}^{21}C_9$  ways to choose the rest of the letters.

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

c) How many words in  $S$  contain all five vowels?

Suppose that all 5 vowels have already been chosen for the word. This leaves us with 9 more letters for the 14-letter word and a choice from 21 letters left over, since there is no repetition.

Thus, there are only  ${}^{21}C_9$  ways to choose the rest of the letters.

Now, we just need to arrange the 14 letters that have been chosen, which gives us  $14!$  arrangements. This is because the vowels don't need to be in a particular order.

2013 T1 Q4 ii)

Let  $S$  be the set of all 14-letter words from the 26 letters of the English alphabet, where no letter is used more than once.

c) How many words in  $S$  contain all five vowels?

Suppose that all 5 vowels have already been chosen for the word. This leaves us with 9 more letters for the 14-letter word and a choice from 21 letters left over, since there is no repetition.

Thus, there are only  ${}^{21}C_9$  ways to choose the rest of the letters.

Now, we just need to arrange the 14 letters that have been chosen, which gives us  $14!$  arrangements. This is because the vowels don't need to be in a particular order.

Hence, there are  ${}^{21}C_9 \times 14!$  words in  $S$  with all 5 vowels.

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

- a) Find the probability that June and Wayne are dealt all 13 spades.
- b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

Probability: 
$$\frac{\text{Number of Cases with Requirement}}{\text{Total Cases}}$$

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

Probability: 
$$\frac{\text{Number of Cases with Requirement}}{\text{Total Cases}}$$

Across June and Wayne, there are 26 cards dealt.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

Probability: 
$$\frac{\text{Number of Cases with Requirement}}{\text{Total Cases}}$$

Across June and Wayne, there are 26 cards dealt. Thus, the total cases is given by choosing 26 cards from the deck of 52:  ${}^{52}C_{26}$ .

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

Probability: 
$$\frac{\text{Number of Cases with Requirement}}{\text{Total Cases}}$$

Across June and Wayne, there are 26 cards dealt. Thus, the total cases is given by choosing 26 cards from the deck of 52:  ${}^{52}C_{26}$ .

The cases where all 13 spades are dealt means that 13 of the 26 slots are taken by the spades.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

Probability: 
$$\frac{\text{Number of Cases with Requirement}}{\text{Total Cases}}$$

Across June and Wayne, there are 26 cards dealt. Thus, the total cases is given by choosing 26 cards from the deck of 52:  ${}^{52}C_{26}$ .

The cases where all 13 spades are dealt means that 13 of the 26 slots are taken by the spades. Thus, there are 13 more cards to be chosen from the 39 remaining cards.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

Probability:  $\frac{\text{Number of Cases with Requirement}}{\text{Total Cases}}$

Across June and Wayne, there are 26 cards dealt. Thus, the total cases is given by choosing 26 cards from the deck of 52:  ${}^{52}C_{26}$ .

The cases where all 13 spades are dealt means that 13 of the 26 slots are taken by the spades. Thus, there are 13 more cards to be chosen from the 39 remaining cards. Meaning the number of cases with 13 spades chosen is given by  ${}^{39}C_{13}$ .

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

a) Find the probability that June and Wayne are dealt all 13 spades.

Probability:  $\frac{\text{Number of Cases with Requirement}}{\text{Total Cases}}$

Across June and Wayne, there are 26 cards dealt. Thus, the total cases is given by choosing 26 cards from the deck of 52:  ${}^{52}C_{26}$ .

The cases where all 13 spades are dealt means that 13 of the 26 slots are taken by the spades. Thus, there are 13 more cards to be chosen from the 39 remaining cards. Meaning the number of cases with 13 spades chosen is given by  ${}^{39}C_{13}$ .

Hence the probability is given by  ${}^{39}C_{13}/{}^{52}C_{26}$ .

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain **at least one** complete suite. (13 spades/hearts/clubs/diamonds)

We know, from a). that there are  $\underline{52C_{26}}$  total cases.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain **at least one** complete suite. (13 spades/hearts/clubs/diamonds)

We know, from a). that there are  ${}^{52}C_{26}$  total cases.

This is another application of inclusion-exclusion principle.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain **at least one** complete suite. (13 spades/hearts/clubs/diamonds)

We know, from a). that there are  ${}^{52}C_{26}$  total cases.

This is another application of inclusion-exclusion principle. From a) we also know that the cases to get a single complete suite is  ${}^{39}C_{13}$ .

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain **at least one** complete suite. (13 spades/hearts/clubs/diamonds)

We know, from a). that there are  ${}^{52}C_{26}$  total cases.

This is another application of inclusion-exclusion principle. From a) we also know that the cases to get a single complete suite is  ${}^{39}C_{13}$ .

There is only one way to get 13 spades and hearts dealt since there are only 26 cards between June and Wayne.

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain **at least one** complete suite. (13 spades/hearts/clubs/diamonds)

We know, from a). that there are  ${}^{52}C_{26}$  total cases.

This is another application of inclusion-exclusion principle. From a) we also know that the cases to get a single complete suite is  ${}^{39}C_{13}$ .

There is only one way to get 13 spades and hearts dealt since there are only 26 cards between June and Wayne.

Since there are only 26 cards, there is no way to get 3 or more complete suites as that would require 39 or more cards to be dealt between June and Wayne.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

Now we can apply the inclusion-exclusion principle:

$$\begin{aligned}|A_S \cup A_H \cup A_C \cup A_D| &= |A_S| + |A_H| + |A_C| + |A_D| \\&\quad - |A_S \cap A_H| - \dots - |A_C \cap A_D|\end{aligned}$$

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

Now we can apply the inclusion-exclusion principle:

$$\begin{aligned} |A_S \cup A_H \cup A_C \cup A_D| &= |A_S| + |A_H| + |A_C| + |A_D| \\ &\quad - |A_S \cap A_H| - \dots - |A_C \cap A_D| \cancel{+} \\ &= {}^4C_1 \times |A_S| - {}^4C_2 \times |A_S \cap A_H| \end{aligned}$$

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

Now we can apply the inclusion-exclusion principle:

$$\begin{aligned}|A_S \cup A_H \cup A_C \cup A_D| &= |A_S| + |A_H| + |A_C| + |A_D| \\&\quad - |A_S \cap A_H| - \dots - |A_C \cap A_D| \\&= {}^4C_1 \times |A_S| - {}^4C_2 \times |A_S \cap A_H| \\&= {}^4C_1 \times {}^{39}C_{13} - {}^4C_2 \times 1\end{aligned}$$

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

Now we can apply the inclusion-exclusion principle:

$$\begin{aligned}|A_S \cup A_H \cup A_C \cup A_D| &= |A_S| + |A_H| + |A_C| + |A_D| \\&\quad - |A_S \cap A_H| - \dots - |A_C \cap A_D| \\&= {}^4C_1 \times |A_S| - {}^4C_2 \times |A_S \cap A_H| \\&= {}^4C_1 \times {}^{39}C_{13} - {}^4C_2 \times 1 \\&= 4 \times {}^{39}C_{13} - {}^4C_2,\end{aligned}$$

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)

Now we can apply the inclusion-exclusion principle:

$$\begin{aligned}|A_S \cup A_H \cup A_C \cup A_D| &= |A_S| + |A_H| + |A_C| + |A_D| \\&\quad - |A_S \cap A_H| - \dots - |A_C \cap A_D| \\&= {}^4C_1 \times |A_S| - {}^4C_2 \times |A_S \cap A_H| \\&= {}^4C_1 \times {}^{39}C_{13} - {}^4C_2 \times 1 \\&= 4 \times {}^{39}C_{13} - {}^4C_2,\end{aligned}$$

where  $A_S$  represents where there is a complete suite of spades etc.

2012 T1 Q4 iii)

Four friends, June, Alex, Wayne, Joe, are playing a hand of bridge. A standard pack of 52 playing cards is dealt, giving each player 13 cards.

b) Find the probability that the cards dealt to June and Wayne contain at least one complete suite. (13 spades/hearts/clubs/diamonds)


Now we can apply the inclusion-exclusion principle:

$$\begin{aligned}|A_S \cup A_H \cup A_C \cup A_D| &= |A_S| + |A_H| + |A_C| + |A_D| \\&\quad - |A_S \cap A_H| - \dots - |A_C \cap A_D| \\&= {}^4C_1 \times |A_S| - {}^4C_2 \times |A_S \cap A_H| \\&= {}^4C_1 \times {}^{39}C_{13} - {}^4C_2 \times 1 \\&= 4 \times {}^{39}C_{13} - {}^4C_2,\end{aligned}$$

where  $A_S$  represents where there is a complete suite of spades etc.

Hence, the probability is  $(4 \times {}^{39}C_{13} - {}^4C_2)/{}^{52}C_{26}$


A straight path of width 2 units is to be laid using 1-unit by 2-unit paving slabs. Here is an example of a path of length 8 units.



Let  $a_n$  be the number of ways to lay a path of width 2 and length  $n$ .

- Find  $a_1$ ,  $a_2$  and  $a_3$ .
- Obtain a recurrence relation for  $a_n$ . Explain your answer.  
(You do NOT need to solve this recurrence relation.)

A straight path of width 2 units is to be laid using 1-unit by 2-unit paving slabs. Here is an example of a path of length 8 units.



Let  $a_n$  be the number of ways to lay a path of width 2 and length  $n$ .


a) Find  $a_1$ ,  $a_2$  and  $a_3$ .

2018 T2 Q4 v)


A straight path of width 2 is laid using 1 by 2 paving slabs. Let  $a_n$  be the number of ways to lay a path of width 2 and length  $n$ .

a) Find  $a_1$ ,  $a_2$  and  $a_3$ .

Let's draw this up!



A straight path of width 2 units is to be laid using 1-unit by 2-unit paving slabs. Here is an example of a path of length 8 units.



Let  $a_n$  be the number of ways to lay a path of width 2 and length  $n$ .

b) Obtain a recurrence relation for  $a_n$ . Explain your answer.

(You do NOT need to solve this recurrence relation.)

A straight path of width 2 is laid using 1 by 2 paving slabs. Let  $a_n$  be the number of ways to lay a path of width 2 and length  $n$ .

b) Obtain a recurrence relation for  $a_n$ . Explain your answer.

(You do NOT need to solve this recurrence relation.)

A straight path of width 2 is laid using 1 by 2 paving slabs. Let  $a_n$  be the number of ways to lay a path of width 2 and length  $n$ .

- b) Obtain a recurrence relation for  $a_n$ . Explain your answer.  
(You do NOT need to solve this recurrence relation.)

The recurrence relation for  $a_n$  is

$$a_n = a_{n-1} + a_{n-2}$$


A straight path of width 2 is laid using 1 by 2 paving slabs. Let  $a_n$  be the number of ways to lay a path of width 2 and length  $n$ .

- b) Obtain a recurrence relation for  $a_n$ . Explain your answer.  
(You do NOT need to solve this recurrence relation.)

The recurrence relation for  $a_n$  is

$$a_n = a_{n-1} + a_{n-2}.$$

We know this is true since we can make a path of length  $n$  from a path of length  $n - 1$  by adding a single block vertically to all  $n - 1$  cases.  
Hence, the  $a_{n-1}$ .

A straight path of width 2 is laid using 1 by 2 paving slabs. Let  $a_n$  be the number of ways to lay a path of width 2 and length  $n$ .

- b) Obtain a recurrence relation for  $a_n$ . Explain your answer.  
(You do NOT need to solve this recurrence relation.)

The recurrence relation for  $a_n$  is

$$a_n = a_{n-1} + a_{n-2}.$$

We know this is true since we can make a path of length  $n$  from a path of length  $n - 1$  by adding a single block vertically to all  $n - 1$  cases. Hence, the  $a_{n-1}$ .

There is actually another way to make a path of length  $n$  which is adding two horizontal blocks to make a 2 by 2 and adding it to all  $n - 2$  cases. Hence, the  $a_{n-2}$ .

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

2019 T3 Q3 iv)

## Definitions for Recurrence Relations

A general recurrence relation of order  $k$  has the form

$$a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = \underline{f(n)}$$

2019 T3 Q3 iv)

## Definitions for Recurrence Relations

A general recurrence relation of order  $k$  has the form

$$a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = f(n)$$

where  $f(n)$  and  $c_i$  are functions of  $n$

## Definitions for Recurrence Relations

A general recurrence relation of order  $k$  has the form

$$a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = f(n)$$

where  $f(n)$  and  $c_i$  are functions of  $n$

Some properties:

## Definitions for Recurrence Relations

A general recurrence relation of order  $k$  has the form

$$a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = f(n)$$

where  $f(n)$  and  $c_i$  are functions of  $n$

Some properties:

a) Linear: All  $c_i$  are constants.

## Definitions for Recurrence Relations

A general recurrence relation of order  $k$  has the form

$$a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = f(n)$$

where  $f(n)$  and  $c_i$  are functions of  $n$

Some properties:

- a) Linear: All  $c_i$  are constants.
- b) Homogeneous:  $f(n) = 0$ .

## Definitions for Recurrence Relations

A general recurrence relation of order  $k$  has the form

$$a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = f(n)$$

where  $f(n)$  and  $c_i$  are functions of  $n$

Some properties:

- a) Linear: All  $c_i$  are constants.
- b) Homogeneous:  $f(n) = 0$ .
- c) Non-Homogeneous:  $f(n)$  is not 0.

## Definitions for Recurrence Relations

A general recurrence relation of order  $k$  has the form

$$a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = f(n)$$

where  $f(n)$  and  $c_i$  are functions of  $n$

Some properties:

- a) Linear: All  $c_i$  are constants.
- b) Homogeneous:  $f(n) = 0$ .
- c) Non-Homogeneous:  $f(n)$  is not 0.

Solutions for Linear, Homogeneous, Order 1:  $a_n = r a_{n-1}$



## Definitions for Recurrence Relations

A general recurrence relation of order  $k$  has the form

$$a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = f(n)$$

where  $f(n)$  and  $c_i$  are functions of  $n$

Some properties:

- a) Linear: All  $c_i$  are constants.
- b) Homogeneous:  $f(n) = 0$ .
- c) Non-Homogeneous:  $f(n)$  is not 0.

Solutions for Linear, Homogeneous, Order 1:  $a_n = r a_{n-1}$

The solution is  $a_n = A r^n$  for some constant  $A$ .

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ .

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ . Let the solutions to  $r^2 + pr + q = 0$  (characteristic polynomial) be  $\alpha$  and  $\beta$ .

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ . Let the solutions to  $r^2 + pr + q = 0$  (characteristic polynomial) be  $\alpha$  and  $\beta$ .

If  $\alpha$  and  $\beta$  are real and non-equal

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ . Let the solutions to  $r^2 + pr + q = 0$  (characteristic polynomial) be  $\alpha$  and  $\beta$ .

If  $\alpha$  and  $\beta$  are real and non-equal, the general solution is

$$a_n = A\alpha^n + B\beta^n.$$

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ . Let the solutions to  $r^2 + pr + q = 0$  (characteristic polynomial) be  $\alpha$  and  $\beta$ .

If  $\alpha$  and  $\beta$  are real and non-equal, the general solution is

$$a_n = A\alpha^n + B\beta^n.$$

If  $\alpha$  and  $\beta$  are real and equal

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ . Let the solutions to  $r^2 + pr + q = 0$  (characteristic polynomial) be  $\alpha$  and  $\beta$ .

If  $\alpha$  and  $\beta$  are real and non-equal, the general solution is

$$a_n = A\alpha^n + B\beta^n.$$

If  $\alpha$  and  $\beta$  are real and equal, the general solution is

$$a_n = A\alpha^n + Bn\beta^n.$$

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ . Let the solutions to  $r^2 + pr + q = 0$  (characteristic polynomial) be  $\alpha$  and  $\beta$ .

If  $\alpha$  and  $\beta$  are real and non-equal, the general solution is

$$a_n = A\alpha^n + B\beta^n.$$

If  $\alpha$  and  $\beta$  are real and equal, the general solution is

$$a_n = A\alpha^n + Bn\beta^n.$$

If  $\alpha$  and  $\beta$  are complex conjugates  $R \exp \pm i\theta$

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ . Let the solutions to  $r^2 + pr + q = 0$  (characteristic polynomial) be  $\alpha$  and  $\beta$ .

If  $\alpha$  and  $\beta$  are real and non-equal, the general solution is

$$a_n = A\alpha^n + B\beta^n.$$

If  $\alpha$  and  $\beta$  are real and equal, the general solution is

$$a_n = A\alpha^n + Bn\beta^n.$$

If  $\alpha$  and  $\beta$  are complex conjugates  $R \exp \pm i\theta$ , the general solution is

$$a_n = R^n(A \cos(n\theta) + B \sin(n\theta)).$$

2019 T3 Q3 iv)

## Solutions to Linear, Homogeneous, Order 2

Consider the recurrence  $a_n + pa_{n-1} + qa_{n-2} = 0$ . Let the solutions to  $r^2 + pr + q = 0$  (characteristic polynomial) be  $\alpha$  and  $\beta$ .

If  $\alpha$  and  $\beta$  are real and non-equal, the general solution is

$$a_n = A\alpha^n + B\beta^n.$$

If  $\alpha$  and  $\beta$  are real and equal, the general solution is

$$a_n = A\alpha^n + Bn\beta^n.$$

If  $\alpha$  and  $\beta$  are complex conjugates  $R \exp \pm i\theta$ , the general solution is

$$a_n = R^n(A \cos(n\theta) + B \sin(n\theta)).$$

Here  $A$  and  $B$  are some constants.

2019 T3 Q3 iv)

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .  $h_n$  is the solution if the recurrence was homogeneous and  $p_n$  is known as a particular solution.

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .  $h_n$  is the solution if the recurrence was homogeneous and  $p_n$  is known as a particular solution.

$p_n$  can be guessed by mirroring the form of  $f(n)$ :

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .  $h_n$  is the solution if the recurrence was homogeneous and  $p_n$  is known as a particular solution.

$p_n$  can be guessed by mirroring the form of  $f(n)$ :

a) Constant: Constant.

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .  $h_n$  is the solution if the recurrence was homogeneous and  $p_n$  is known as a particular solution.

$p_n$  can be guessed by mirroring the form of  $f(n)$ :

- a) Constant: Constant.
- b) Polynomial: Polynomial of the same degree.

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .  $h_n$  is the solution if the recurrence was homogeneous and  $p_n$  is known as a particular solution.

$p_n$  can be guessed by mirroring the form of  $f(n)$ :

- a) Constant: Constant.
- b) Polynomial: Polynomial of the same degree.
- c) Exponential: Constant  $\times$  Exponential of same base.

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .  $h_n$  is the solution if the recurrence was homogeneous and  $p_n$  is known as a particular solution.

$p_n$  can be guessed by mirroring the form of  $f(n)$ :

- a) Constant: Constant.
- b) Polynomial: Polynomial of the same degree.
- c) Exponential: Constant  $\times$  Exponential of same base.
- d) Product of Previous: Product of corresponding guesses.

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .  $h_n$  is the solution if the recurrence was homogeneous and  $p_n$  is known as a particular solution.

$p_n$  can be guessed by mirroring the form of  $f(n)$ :

- a) Constant: Constant.
- b) Polynomial: Polynomial of the same degree.
- c) Exponential: Constant  $\times$  Exponential of same base.
- d) Product of Previous: Product of corresponding guesses.

A guess does not work if there are shared terms in  $\underline{h_n}$  and  $\underline{p_n}$ .

2019 T3 Q3 iv)

## Solutions to Non-Homogeneous

In this case,  $a_n = h_n + p_n$ .  $h_n$  is the solution if the recurrence was homogeneous and  $p_n$  is known as a particular solution.

$p_n$  can be guessed by mirroring the form of  $f(n)$ :

- a) Constant: Constant.
- b) Polynomial: Polynomial of the same degree.
- c) Exponential: Constant  $\times$  Exponential of same base.
- d) Product of Previous: Product of corresponding guesses.

A guess does not work if there are shared terms in  $h_n$  and  $p_n$ . If that is the case, multiply  $p_n$  by  $n$  to form a new guess.

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

This is a non-homogeneous linear second-order recurrence.

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

This is a non-homogeneous linear second-order recurrence.  
This means that the solution for this relation is in the form

$$\underline{a_n = h_n + p_n},$$

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

This is a non-homogeneous linear second-order recurrence.  
This means that the solution for this relation is in the form

$$a_n = h_n + p_n,$$

where  $h_n$  is the solution if it were homogeneous.

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

This is a non-homogeneous linear second-order recurrence.  
This means that the solution for this relation is in the form

$$a_n = h_n + p_n,$$

where  $h_n$  is the solution if it were homogeneous.

The homogeneous recurrence for this relation is  $a_n - 6a_{n-1} + 9a_{n-2} = 0$ .

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

This is a non-homogeneous linear second-order recurrence.  
This means that the solution for this relation is in the form

$$a_n = h_n + p_n,$$

where  $h_n$  is the solution if it were homogeneous.

The homogeneous recurrence for this relation is  $a_n - 6a_{n-1} + 9a_{n-2} = 0$ .  
To do this we look at  $r^2 - 6r + 9 = 0$  where the solutions are  $r = \underline{3, 3}$ .

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

This is a non-homogeneous linear second-order recurrence.  
This means that the solution for this relation is in the form

$$a_n = h_n + p_n,$$

where  $h_n$  is the solution if it were homogeneous.

The homogeneous recurrence for this relation is  $a_n - 6a_{n-1} + 9a_{n-2} = 0$ .  
To do this we look at  $r^2 - 6r + 9 = 0$  where the solutions are  $r = 3, 3$ .

Hence, the general solution  $h_n$  is given by  $A(3^n) + Bn(3^n)$  since the  
solutions of the  $r$  quadratic are the same.

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

$p_n$  is a particular solution which relates to the function that the relation is equal to:  $5n3^n$ .

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

$p_n$  is a particular solution which relates to the function that the relation is equal to:  $5n3^n$ . We want to mirror this function, which is a linear and an exponential with base 3

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = \underline{5n3^n}.$$

$p_n$  is a particular solution which relates to the function that the relation is equal to:  $5n3^n$ . We want to mirror this function, which is a linear and an exponential with base 3 so a possible

$$p_n = \underline{(cn + d)3^n} = d(3^n) + cn(3^n) \text{ for some real numbers } \underline{c} \text{ and } \underline{d}.$$

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

$p_n$  is a particular solution which relates to the function that the relation is equal to:  $5n3^n$ . We want to mirror this function, which is a linear and an exponential with base 3 so a possible  
 $p_n = (cn + d)3^n = d(3^n) + cn(3^n)$  for some real numbers  $c$  and  $d$ .

A  $p_n$  only works if it does not share terms with  $h_n$ .

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

$p_n$  is a particular solution which relates to the function that the relation is equal to:  $5n3^n$ . We want to mirror this function, which is a linear and an exponential with base 3 so a possible  
 $p_n = (cn + d)3^n = d(3^n) + cn(3^n)$  for some real numbers  $c$  and  $d$ .

A  $p_n$  only works if it does not share terms with  $h_n$ . Since  $cn(3^n)$  is the same as  $Bn(3^n)$ , this  $p_n$  does not work.

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

$p_n$  is a particular solution which relates to the function that the relation is equal to:  $5n3^n$ . We want to mirror this function, which is a linear and an exponential with base 3 so a possible  
 $p_n = (cn + d)3^n = d(3^n) + cn(3^n)$  for some real numbers  $c$  and  $d$ .

A  $p_n$  only works if it does not share terms with  $h_n$ . Since  $cn(3^n)$  is the same as  $Bn(3^n)$ , this  $p_n$  does not work.

A new  $p_n$  can be found by multiplying  $n$  to the old one.

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

$p_n$  is a particular solution which relates to the function that the relation is equal to:  $5n3^n$ . We want to mirror this function, which is a linear and an exponential with base 3 so a possible

$$p_n = (cn + d)3^n = d(3^n) + cn(3^n) \text{ for some real numbers } c \text{ and } d.$$

A  $p_n$  only works if it does not share terms with  $h_n$ . Since  $cn(3^n)$  is the same as  $Bn(3^n)$ , this  $p_n$  does not work.

A new  $p_n$  can be found by multiplying  $n$  to the old one. Our new  
 $p_n = n(cn + d)3^n = dn(3^n) + cn^2(3^n)$

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

$p_n$  is a particular solution which relates to the function that the relation is equal to:  $5n3^n$ . We want to mirror this function, which is a linear and an exponential with base 3 so a possible

$$p_n = (cn + d)3^n = d(3^n) + cn(3^n) \text{ for some real numbers } c \text{ and } d.$$

A  $p_n$  only works if it does not share terms with  $h_n$ . Since  $cn(3^n)$  is the same as  $Bn(3^n)$ , this  $p_n$  does not work.

A new  $p_n$  can be found by multiplying  $n$  to the old one. Our new  $p_n = n(cn + d)3^n = \underline{dn(3^n)} + cn^2(3^n)$  has the term  $dn(3^n)$  which is also the same as  $Bn(3^n)$  so it also does not work.

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

Instead, we use  $p_n = n^2(cn + d)3^n$ .

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

Instead, we use  $p_n = n^2(cn + d)3^n$ . To find the constants of our  $p_n$  you use  $p_n$  in place of  $a_n$

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

Instead, we use  $p_n = n^2(cn + d)3^n$ . To find the constants of our  $p_n$  you use  $p_n$  in place of  $a_n$ , which gives us

$$(cn^3 + dn^2)3^n - 6(c(n-1)^3 + d(n-1)^2)3^{n-1}$$

$$\underline{\underline{+9(c(n-2)^3 + d(n-2)^2)3^{n-2}}} = 5n3^n.$$

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

Instead, we use  $p_n = n^2(cn + d)3^n$ . To find the constants of our  $p_n$  you use  $p_n$  in place of  $a_n$ , which gives us

$$\begin{aligned}(cn^3 + dn^2)3^n - 6(c(n-1)^3 + d(n-1)^2)3^{n-1} \\+ 9(c(n-2)^3 + d(n-2)^2)3^{n-2} = 5n3^n.\end{aligned}$$

Simplifying this expression gets us equations from coefficients:

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

Instead, we use  $p_n = n^2(cn + d)3^n$ . To find the constants of our  $p_n$  you use  $p_n$  in place of  $a_n$ , which gives us

$$\begin{aligned}(cn^3 + dn^2)3^n - 6(c(n-1)^3 + d(n-1)^2)3^{n-1} \\ + 9(c(n-2)^3 + d(n-2)^2)3^{n-2} = 5n3^n.\end{aligned}$$

Simplifying this expression gets us equations from coefficients:

$$6cn - 6c + 2d = 5n + 0 \Rightarrow \underline{\underline{6c = 5}} \text{ and } \underline{\underline{2d - 6c = 0}}.$$

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

Instead, we use  $p_n = n^2(cn + d)3^n$ . To find the constants of our  $p_n$  you use  $p_n$  in place of  $a_n$ , which gives us

$$\begin{aligned}(cn^3 + dn^2)3^n - 6(c(n-1)^3 + d(n-1)^2)3^{n-1} \\+ 9(c(n-2)^3 + d(n-2)^2)3^{n-2} = 5n3^n.\end{aligned}$$

Simplifying this expression gets us equations from coefficients:

$$6cn - 6c + 2d = 5n + 0 \Rightarrow 6c = 5 \text{ and } 2d - 6c = 0.$$

Solving these,  $c = \frac{5}{6}$  and  $d = \frac{5}{2}$ .

2019 T3 Q3 iv)

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

Instead, we use  $p_n = n^2(cn + d)3^n$ . To find the constants of our  $p_n$  you use  $p_n$  in place of  $a_n$ , which gives us

$$\begin{aligned}(cn^3 + dn^2)3^n - 6(c(n-1)^3 + d(n-1)^2)3^{n-1} \\ + 9(c(n-2)^3 + d(n-2)^2)3^{n-2} = 5n3^n.\end{aligned}$$

Simplifying this expression gets us equations from coefficients:

$$6cn - 6c + 2d = 5n + 0 \Rightarrow 6c = 5 \text{ and } 2d - 6c = 0.$$

Solving these,  $c = \frac{5}{6}$  and  $d = \frac{5}{2}$ . Therefore,  $p_n = n^2 \left( \frac{5n}{6} + \frac{5}{2} \right) 3^n$ .

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

From our previous work we have shown that

$$\begin{cases} h_n = A(3^n) + Bn(3^n), \\ p_n = n^2 \left( \frac{5n}{6} + \frac{5}{2} \right) 3^n. \end{cases}$$

Obtain the general solution to the recurrence relation

$$a_n - 6a_{n-1} + 9a_{n-2} = 5n3^n.$$

From our previous work we have shown that

$$\begin{cases} h_n = A(3^n) + Bn(3^n), \\ p_n = n^2 \left( \frac{5n}{6} + \frac{5}{2} \right) 3^n. \end{cases}$$

Hence, the general solution is

$$a_n = A(3^n) + Bn(3^n) + n^2 \left( \frac{5n}{6} + \frac{5}{2} \right) 3^n.$$

## 4. Logic and Proofs

# Mathematical Induction

Prove by mathematical induction that  $3^n > n^3$  for all integers  $n \geq 4$ .

# Mathematical Induction

Prove by mathematical induction that  $3^n > n^3$  for all integers  $n \geq 4$ .

Proof: Let  $P(n)$  be the statement  $3^n > n^3$  where  $n \in \mathbb{Z}^+$  and  $n \geq 4$ .

Step 1: We have to check our base case  $P(4)$  is true.

# Mathematical Induction

Prove by mathematical induction that  $3^n > n^3$  for all integers  $n \geq 4$ .

Proof: Let  $P(n)$  be the statement  $3^n > n^3$  where  $n \in \mathbb{Z}^+$  and  $n \geq 4$ .

Step 1: We have to check our base case  $P(4)$  is true.

We have that  $3^4 = 81$  and  $4^3 = 64$ .

# Mathematical Induction

Prove by mathematical induction that  $3^n > n^3$  for all integers  $n \geq 4$ .

Proof: Let  $P(n)$  be the statement  $3^n > n^3$  where  $n \in \mathbb{Z}^+$  and  $n \geq 4$ .

Step 1: We have to check our base case  $P(4)$  is true.

We have that  $3^4 = 81$  and  $4^3 = 64$ .

So  $3^4 > 4^3$  and  $P(4)$  is true.

# Mathematical Induction

Prove by mathematical induction that  $3^n > n^3$  for all integers  $n \geq 4$ .

Step 2:

Assume  $P(k)$  is true for all integers  $k \geq 4$ .  
i.e  $3^k > k^3$ .

# Mathematical Induction

Prove by mathematical induction that  $3^n > n^3$  for all integers  $n \geq 4$ .

Step 2:

Assume  $P(k)$  is true for all integers  $k \geq 4$ .  
i.e  $3^k > k^3$ .

We now have to prove  $P(k + 1)$  is also true.  
i.e  $3^{k+1} > (k + 1)^3$ .

# Mathematical Induction\*

Prove by mathematical induction that  $3^n > n^3$  for all integers  $n \geq 4$ .

Assume  $3^k > k^3$ .

RTP:  $3^{k+1} > (k+1)^3 = k^3 + 3k^2 + 3k + 1$ .

$$\begin{aligned} 3^{k+1} &= 3 \cdot 3^k && \rightarrow = k^3 + 3k^2 + 3k + 17k \\ &> 3k^3 && \text{(by assumption)} \\ &= k^3 + 2k^3 && \geq k^3 + 3k^2 + 3k + 1 \\ &= k^3 + 2k \cdot k^2 && = (k+1)^3 \quad (k \geq 4) \\ &\geq k^3 + 8k^2 && \quad (k \geq 4) \\ &= k^3 + 3k^2 + 5k^2 \\ &\geq k^3 + 3k^2 + 20k && \quad (k \geq 4) \end{aligned}$$

# Mathematical Induction

Prove by mathematical induction that  $3^n > n^3$  for all integers  $n \geq 4$ .

$$\begin{aligned}3^{k+1} &= 3(3^k) \\&> 3k^3 \text{ (by induction hypothesis)} \\&= k^3 + 2k^3 \\&\geq k^3 + 8k^2 \text{ (since } k \geq 4\text{)} \\&= k^3 + 3k^2 + 5k^2 \\&\geq k^3 + 3k^2 + 20k \text{ (since } k \geq 4\text{)} \\&= k^3 + 3k^2 + 3k + 17k \\&\geq k^3 + 3k^2 + 3k + 1 \text{ (since } k \geq 4\text{)} \\&= (k+1)^3.\end{aligned}$$

Hence  $P(k+1)$  is true if  $P(k)$  is true. Thus, by the principle of mathematical induction,  $P(n)$  is true for all integers  $n \geq 4$ .  $\square$

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

Assume for a contradiction that  $\sqrt[3]{3} + \sqrt{2}$  is rational.

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

Assume for a contradiction that  $\sqrt[3]{3} + \sqrt{2}$  is rational.

Let  $\sqrt[3]{3} + \sqrt{2} = \frac{a}{b}$  where  $a, b \in \mathbb{Z}^+$ . Then we have

$$\sqrt[3]{3} = \frac{a}{b} - \sqrt{2}$$

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

Assume for a contradiction that  $\sqrt[3]{3} + \sqrt{2}$  is rational.

Let  $\sqrt[3]{3} + \sqrt{2} = \frac{a}{b}$  where  $a, b \in \mathbb{Z}^+$ . Then we have

$$\sqrt[3]{3} = \frac{a}{b} - \sqrt{2}$$

$$3 = \left( \frac{a}{b} - \sqrt{2} \right)^3$$

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

Assume for a contradiction that  $\sqrt[3]{3} + \sqrt{2}$  is rational.

Let  $\sqrt[3]{3} + \sqrt{2} = \frac{a}{b}$  where  $a, b \in \mathbb{Z}^+$ . Then we have

$$\sqrt[3]{3} = \frac{a}{b} - \sqrt{2}$$

$$3 = \left( \frac{a}{b} - \sqrt{2} \right)^3$$

$$3 = \frac{a^3}{b^3} - \frac{3\sqrt{2}a^2}{b^2} + \frac{12a}{b} - 2\sqrt{2}$$

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

Assume for a contradiction that  $\sqrt[3]{3} + \sqrt{2}$  is rational.

Let  $\sqrt[3]{3} + \sqrt{2} = \frac{a}{b}$  where  $a, b \in \mathbb{Z}^+$ . Then we have

$$\sqrt[3]{3} = \frac{a}{b} - \sqrt{2}$$

$$3 = \left( \frac{a}{b} - \sqrt{2} \right)^3$$

$$3 = \frac{a^3}{b^3} - \frac{3\sqrt{2}a^2}{b^2} + \frac{12a}{b} - 2\sqrt{2}$$

$$3b^3 = a^3 - 3\sqrt{2}a^2b + 12ab^2 - 2\sqrt{2}b^3$$

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

Assume for a contradiction that  $\sqrt[3]{3} + \sqrt{2}$  is rational.

Let  $\sqrt[3]{3} + \sqrt{2} = \frac{a}{b}$  where  $a, b \in \mathbb{Z}^+$ . Then we have

$$\sqrt[3]{3} = \frac{a}{b} - \sqrt{2}$$

$$3 = \left( \frac{a}{b} - \sqrt{2} \right)^3$$

$$3 = \frac{a^3}{b^3} - \frac{3\sqrt{2}a^2}{b^2} + \frac{12a}{b} - 2\sqrt{2}$$

$$3b^3 = a^3 - 3\sqrt{2}a^2b + 12ab^2 - 2\sqrt{2}b^3$$

$$\sqrt{2}(3a^2b + 2b^3) = a^3 - 3b^3 + 12ab^2$$

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

Assume for a contradiction that  $\sqrt[3]{3} + \sqrt{2}$  is rational.

Let  $\sqrt[3]{3} + \sqrt{2} = \frac{a}{b}$  where  $a, b \in \mathbb{Z}^+$ . Then we have

$$\sqrt[3]{3} = \frac{a}{b} - \sqrt{2}$$

$$3 = \left( \frac{a}{b} - \sqrt{2} \right)^3$$

$$3 = \frac{a^3}{b^3} - \frac{3\sqrt{2}a^2}{b^2} + \frac{12a}{b} - 2\sqrt{2}$$

$$3b^3 = a^3 - 3\sqrt{2}a^2b + 12ab^2 - 2\sqrt{2}b^3$$

$$\sqrt{2}(3a^2b + 2b^3) = a^3 - 3b^3 + 12ab^2$$

$$\sqrt{2} = \frac{a^3 - 3b^3 + 12ab^2}{3a^2b + 2b^3}$$

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

$$\begin{aligned}\sqrt{2} &= \frac{a^3 - 3b^3 + 12ab^2}{3a^2b + 2b^3} \\ \sqrt{2} &= \frac{A}{B} \text{ (where A and B are integers).}\end{aligned}$$

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

$$\begin{aligned}\sqrt{2} &= \frac{a^3 - 3b^3 + 12ab^2}{3a^2b + 2b^3} \\ \sqrt{2} &= \frac{A}{B} \text{ (where A and B are integers).}\end{aligned}$$

Hence  $\sqrt{2}$  is rational.

# Contradiction

Prove  $\sqrt[3]{3} + \sqrt{2}$  is irrational.

$$\sqrt{2} = \frac{a^3 - 3b^3 + 12ab^2}{3a^2b + 2b^3}$$

$$\sqrt{2} = \frac{A}{B} \text{ (where A and B are integers).}$$

Hence  $\sqrt{2}$  is rational.

But this is a contradiction as  $\sqrt{2}$  is irrational and our assumption is false. Thus,  $\sqrt[3]{3} + \sqrt{2}$  is irrational.  $\square$

## Lemma

$\sqrt{2}$  is irrational.

# Contrapositive

## Definition

$$p \rightarrow q \equiv \sim q \rightarrow \sim p.$$

# Contrapositive

## Definition

$$p \rightarrow q \equiv \sim q \rightarrow \sim p.$$

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

# Contrapositive

## Definition

$$p \rightarrow q \equiv \sim q \rightarrow \sim p.$$

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

## Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $n$  is not prime, then  $2^n - 1$  is not prime.

# Contrapositive

## Definition

$$p \rightarrow q \equiv \sim q \rightarrow \sim p.$$

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

## Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $n$  is not prime, then  $2^n - 1$  is not prime.

Let  $n \in \mathbb{Z}^+$ . Suppose  $n$  is not prime. Then  $n$  is either 1 or composite.

Case 1:  $n = 1$ .

# Contrapositive

## Definition

$$p \rightarrow q \equiv \sim q \rightarrow \sim p.$$

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

## Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $n$  is not prime, then  $2^n - 1$  is not prime.

Let  $n \in \mathbb{Z}^+$ . Suppose  $n$  is not prime. Then  $n$  is either 1 or composite.

### Case 1: $n = 1$ .

If  $n = 1$  then we have  $2^n - 1 = 2^1 - 1 = 1$  which is not prime. Hence,  $2^n - 1$  is not prime for  $n = 1$ .

# Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

Case 2:  $n = ab$  where  $a, b \in \mathbb{Z}^+$  and  $a, b > 1$ .

# Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

Case 2:  $n = ab$  where  $a, b \in \mathbb{Z}^+$  and  $a, b > 1$ .

If  $n = ab$ , then we have

$$2^n - 1 = 2^{ab} - 1$$

# Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

Case 2:  $n = ab$  where  $a, b \in \mathbb{Z}^+$  and  $a, b > 1$ .

If  $n = ab$ , then we have

$$\begin{aligned}2^n - 1 &= 2^{ab} - 1 \\&= (2^a)^b - 1\end{aligned}$$

# Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

Case 2:  $n = ab$  where  $a, b \in \mathbb{Z}^+$  and  $a, b > 1$ .

If  $n = ab$ , then we have

$$\begin{aligned}2^n - 1 &= 2^{ab} - 1 \\&= (2^a)^b - 1 \\&= (2^a - 1)(1 + 2^a + (2^a)^2 + \dots + (2^a)^{b-1})\\&\quad \textcolor{blue}{\geq 1} \quad \textcolor{blue}{\geq 1}\end{aligned}$$

# Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

Case 2:  $n = ab$  where  $a, b \in \mathbb{Z}^+$  and  $a, b > 1$ .

If  $n = ab$ , then we have

$$\begin{aligned}2^n - 1 &= 2^{ab} - 1 \\&= (2^a)^b - 1 \\&= (2^a - 1)(1 + 2^a + (2^a)^2 + \dots + (2^a)^{b-1}) \\&= AB\end{aligned}$$

Where  $A, B \in \mathbb{Z}^+$  and  $A, B > 1$ .

# Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

Case 2:  $n = ab$  where  $a, b \in \mathbb{Z}^+$  and  $a, b > 1$ .

If  $n = ab$ , then we have

$$\begin{aligned}2^n - 1 &= 2^{ab} - 1 \\&= (2^a)^b - 1 \\&= (2^a - 1)(1 + 2^a + (2^a)^2 + \dots + (2^a)^{b-1}) \\&= AB\end{aligned}$$

Where  $A, B \in \mathbb{Z}^+$  and  $A, B > 1$ .

Hence when  $n$  is composite,  $2^n - 1$  is not prime.

# Contrapositive

For all  $n \in \mathbb{Z}^+$ , prove that if  $2^n - 1$  is prime, then  $n$  is prime.

Case 2:  $n = ab$  where  $a, b \in \mathbb{Z}^+$  and  $a, b > 1$ .

If  $n = ab$ , then we have

$$\begin{aligned}2^n - 1 &= 2^{ab} - 1 \\&= (2^a)^b - 1 \\&= (2^a - 1)(1 + 2^a + (2^a)^2 + \dots + (2^a)^{b-1}) \\&= AB\end{aligned}$$

Where  $A, B \in \mathbb{Z}^+$  and  $A, B > 1$ .

Hence when  $n$  is composite,  $2^n - 1$  is not prime.

Thus in both cases, the contrapositive statement is true. Hence the original statement is true for all  $n \in \mathbb{Z}^+$ .  $\square$

# 2017 T1 Q3 (iv)

Absolute value function

$$|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0. \end{cases}$$

# 2017 T1 Q3 (iv)

Absolute value function

$$|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0. \end{cases}$$

Prove that if  $x$  is a real number then

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

# 2017 T1 Q3 (iv)

Absolute value function

$$|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0. \end{cases}$$

Prove that if  $x$  is a real number then

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

The absolute value is a strong indicator that we need cases!!!

# 2017 T1 Q3 (iv)

Absolute value function

$$|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0. \end{cases}$$

Prove that if  $x$  is a real number then

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

The absolute value is a strong indicator that we need cases!!! Notice that

$$|x - 3| = \begin{cases} x - 3 & x \geq 3 \\ -(x - 3) & x < 3. \end{cases}$$

# 2017 T1 Q3 (iv)

Absolute value function

$$|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0. \end{cases}$$

Prove that if  $x$  is a real number then

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

The absolute value is a strong indicator that we need cases!!! Notice that

$$|x - 3| = \begin{cases} x - 3 & x \geq 3 \\ -(x - 3) & x < 3. \end{cases}$$

So lets take cases when  $x \geq 3$  and  $x < 3$ .

2017 T1 3) (iv)\*

Prove that if  $x$  is a real number then

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

Case 1:  $x \geq 3$

$$\begin{aligned} & 2x^2 - 9x + 12 - x|x - 3| \\ &= 2x^2 - 9x + 12 - x(x - 3) \\ &= 2x^2 - 9x + 12 - x^2 + 3x \\ &\leq x^2 - 6x + 12 \\ &= \underline{(x-3)^2 + 3} \\ &\geq 0 \end{aligned}$$

Hence, when

$$x \geq 3,$$

$$x|x - 3| \leq 2x^2 - 9x + 12$$

2017 T1 3) (iv)\*

Prove that if  $x$  is a real number then

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

Case 2:  $x < 3$

$$\begin{aligned} & 2x^2 - 9x + 12 - x|x - 3| \\ &= 2x^2 - 9x + 12 - x(3 - x) \quad \left. \begin{array}{l} \text{Hence for } x < 3, \\ |x - 3| \leq 3 - x \end{array} \right\} \\ &= 2x^2 - 9x + 12 - 3x + x^2 \quad \left. \begin{array}{l} \text{Hence, for all} \\ x \in \mathbb{R}, \\ |x - 3| \leq 3 - x \end{array} \right\} \\ &= 3x^2 - 12x + 12 \\ &= 3(x^2 - 4x + 4) \\ &= 3(x - 2)^2 \\ &\geq 0 \end{aligned}$$

2017 T1 3) (iv)

Prove that if  $x$  is a real number then

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

Suppose  $x \in \mathbb{R}$ .

Case 1: For the first case, we take  $x \geq 3$ . Now consider

$$\begin{aligned} 2x^2 - 9x + 12 - x|x - 3| &= 2x^2 - 9x + 12 - x(x - 3) \\ &= 2x^2 - 9x + 12 - x^2 + 3x \\ &= x^2 - 6x + 12 \\ &= (x - 3)^2 + 3 \\ &\geq 0 \end{aligned}$$

Hence for  $x \geq 3$ ,  $2x^2 - 9x + 12 - x|x - 3| \geq 0$  implying that

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

2017 T1 3) (iv)

Prove that if  $x$  is a real number then

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

Case 2: For the second case, we take  $x < 3$ . Now consider

$$\begin{aligned} 2x^2 - 9x + 12 - x|x - 3| &= 2x^2 - 9x + 12 + x(x - 3) \\ &= 2x^2 - 9x + 12 + x^2 - 3x \\ &= 3x^2 - 12x + 12 \\ &= 3(x - 2)^2 \\ &\geq 0 \end{aligned}$$

Hence for  $x < 3$ ,  $2x^2 - 9x + 12 - x|x - 3| \geq 0$  implying that

$$x|x - 3| \leq 2x^2 - 9x + 12.$$

Thus,  $x|x - 3| \leq 2x^2 - 9x + 12$ . for all  $x \in \mathbb{R}$ .

For a sequence of integers  $\{a_n\}_{n=1}^{\infty}$ , we say that  $a_n$  is eventually even if and only if

$$\exists M \in \mathbb{Z} \quad \forall n > M \quad a_n \text{ is even.}$$

- Write symbolically the statement " $a_n$  is not eventually even", simplifying your answer so that the negation symbols are not used.
- Prove that the sequence given by

$$a_n = 475 + \left\lfloor \frac{28n + 995}{n} \right\rfloor$$

is not eventually even.

## a) Solution

For a sequence of integers  $\{a_n\}_{n=1}^{\infty}$ , we say that  $a_n$  is eventually even if and only if

$$\exists M \in \mathbb{Z} \quad \forall n > M \quad a_n \text{ is even.}$$

- a) Write symbolically the statement " $a_n$  is not eventually even", simplifying your answer so that the negation symbols are not used.

Let  $P$  be the statement " $a_n$  is eventually even".

## a) Solution

For a sequence of integers  $\{a_n\}_{n=1}^{\infty}$ , we say that  $a_n$  is eventually even if and only if

$$\exists M \in \mathbb{Z} \quad \forall n > M \quad a_n \text{ is even.}$$

- a) Write symbolically the statement " $a_n$  is not eventually even", simplifying your answer so that the negation symbols are not used.

Let  $P$  be the statement " $a_n$  is eventually even". The statement " $a_n$  is not eventually even" can be represented by

$$\sim P \equiv \sim (\exists M \in \mathbb{Z} \quad \underline{\forall n > M} \quad a_n \text{ is even})$$

## a) Solution

For a sequence of integers  $\{a_n\}_{n=1}^{\infty}$ , we say that  $a_n$  is eventually even if and only if

$$\exists M \in \mathbb{Z} \quad \forall n > M \quad a_n \text{ is even.}$$

- a) Write symbolically the statement " $a_n$  is not eventually even", simplifying your answer so that the negation symbols are not used.

Let  $P$  be the statement " $a_n$  is eventually even". The statement " $a_n$  is not eventually even" can be represented by

$$\begin{aligned}\sim P &\equiv \sim (\exists M \in \mathbb{Z} \quad \forall n > M \quad a_n \text{ is even}) \\ &\equiv \quad \forall M \in \mathbb{Z} \quad \exists n > M \quad a_n \text{ is odd.}\end{aligned}$$

## b) Proof\*

b) Prove that the sequence given by

$$a_n = 475 + \left\lfloor \frac{28n + 995}{n} \right\rfloor$$

is not eventually even.  $\sim P \equiv \forall M \in \mathbb{Z} \quad \exists n > M \quad a_n \text{ is odd.}$

$$\begin{aligned} a_n &= 475 + \left\lfloor \frac{28n}{n} + \frac{995}{n} \right\rfloor \quad \left| \begin{array}{l} 0 \leq \frac{995}{n} < 1 \\ n > M \end{array} \right. \\ &= 475 + \left\lfloor 28 + \frac{995}{n} \right\rfloor \quad \left| \begin{array}{l} n > 995 \\ n = M^2 + 995 \end{array} \right. \\ \text{If } 0 \leq \frac{995}{n} < 1, \quad & \left| \begin{array}{l} n = M^2 + 995 \\ > 995, M \end{array} \right. \\ \left\lfloor 28 + \frac{995}{n} \right\rfloor &= 28 \\ a_n &= 475 + 28 = 503 \end{aligned}$$

## b) Proof\*

b) Prove that the sequence given by

$$a_n = 475 + \left\lfloor \frac{28n + 995}{n} \right\rfloor$$

is not eventually even.  $\sim P \equiv \forall M \in \mathbb{Z} \quad \exists n > M \quad a_n \text{ is odd.}$

Let  $M \in \mathbb{Z}$ , choose  $n = M^2 + 996 > M$ .

$$\begin{aligned} a_n &= 475 + \left\lfloor 28 + \frac{995}{M^2 + 996} \right\rfloor \\ &= 475 + 28 \quad \left( 0 < \frac{995}{M^2 + 996} < 1 \right) \\ &= 503 \end{aligned}$$

Thus  $a_n$  is odd. So  $a_n$  is not eventually even.  $\square$

## b) Proof

For a sequence of integers  $\{a_n\}_{n=1}^{\infty}$ , we say that  $a_n$  is eventually even if and only if

$$\exists M \in \mathbb{Z} \quad \forall n > M \quad a_n \text{ is even.}$$

b) Prove that the sequence given by

$$a_n = 475 + \left\lfloor \frac{28n + 995}{n} \right\rfloor$$

is not eventually even.

To prove  $a_n$  is not eventually even, we have to prove  $\sim P$  is true.

We are required to prove  $\forall M \in \mathbb{Z} \quad \exists n > M \quad a_n \text{ is odd.}$

Notice  $a_n = 475 + \left\lfloor 28 + \frac{995}{n} \right\rfloor$ . We can make  $a_n = 503$  which is odd if  $0 \leq \frac{995}{n} < 1$ . So we need  $n > 995$  and  $n > M$ . So we can choose  $n = M^2 + 996$  which will guarantee  $n > 995$  and  $n > M$ .

## b) Proof

Proof: Let  $M \in \mathbb{Z}$ . Choose  $n = M^2 + 996 > M$ . Then we have

$$\begin{aligned} a_n &= 475 + \left\lfloor \frac{28n + 995}{n} \right\rfloor \\ &= 475 + \left\lfloor 28 + \frac{995}{n} \right\rfloor \\ &= 475 + \left\lfloor 28 + \frac{995}{M^2 + 996} \right\rfloor \\ &= 475 + 28 \text{ (since } 0 < \frac{995}{M^2 + 996} < 1\text{)} \\ &= 503. \end{aligned}$$

Hence,  $a_n$  is odd and  $\sim P$  is true. Thus,  $a_n$  is not eventually even.  $\square$

# 2016 T1 Q1\*

i) Construct a truth table for

$$(p \wedge q) \rightarrow (\underline{q \rightarrow p}).$$

Is this a tautology, contradiction or contingency? Give reasons.

P	q	$P \wedge q$	$q \rightarrow P$	$(P \wedge q) \rightarrow (q \rightarrow P)$
T	T	T	T	T
T	F	F	T	T
F	T	F	F	T
F	F	F	T	T

we have a tautology.

# 2016 T1 Q1\*

i) Construct a truth table for

$$(p \wedge q) \rightarrow (q \rightarrow p).$$

Is this a tautology, contradiction or contingency? Give reasons.

$p$	$q$	$p \wedge q$	$q \rightarrow p$	$(p \wedge q) \rightarrow (q \rightarrow p)$
T	T	T	T	T
T	F	F	T	T
F	T	F	F	T
F	F	F	T	T

Since  $(p \wedge q) \rightarrow (q \rightarrow p)$  is always true, this is a tautology.

# 2016 T1 Q1\*

ii) Suppose you have a complete truth table for  $n$  propositions  $p_1, p_2, \dots, p_n$ .

In how many of the  $2^n$  lines of the truth table is  $p_1 \rightarrow p_n$  true?

$p_1$	$p_n$	$p_1 \rightarrow p_n$
$\frac{2^n}{2}$	F	$F \rightarrow$ anything always TRUE
$\frac{2^n}{2}$	T	$T \rightarrow T$ True
$\frac{2^n}{4}$	F	$T \rightarrow F$ False

total lines:

$$\frac{2^n}{2} + \frac{2^n}{4}$$

$$= 2 \cdot 2^{n-2} + 2^{n-2}$$

$$= 3 \cdot 2^{n-2}$$

# 2016 T1 Q1\*

ii) Suppose you have a complete truth table for  $n$  propositions  $p_1, p_2, \dots, p_n$ .

In how many of the  $2^n$  lines of the truth table is  $p_1 \rightarrow p_n$  true?

Notice initially, half of  $p_1$  is true and the other half is false.

Case 1:  $p_1$  is false.

If  $p_1$  is false, then  $p_1 \rightarrow p_n$  will be true regardless of  $p_n$ . This can occur in  $\frac{2^n}{2} = 2^{n-1}$  ways.

Case 2:  $p_1$  is true.

If  $p_1$  is true, then  $p_1 \rightarrow p_n$  is only true when  $p_n$  is false. Notice that within the true  $p_1$ , half of them will correspond to a true  $p_n$ . Hence, this can occur in  $\frac{2^n}{2} = 2^{n-2}$  ways.

Hence the total number of ways is  $2(2^{n-2}) + 2^{n-2} = 3(2^{n-2})$  ways.

## 2016 T1 3) (iii)\*

Consider the following arguments:

If

- (1) I will have a good holiday ~~only if~~ I have both time and money.
- (2) If I work then I will have no time.
- (3) If I do not work, then I will have no money.
- (4) Therefore, I will not have a good holiday.

~~only if~~

$g \rightarrow (t \wedge m)$

$w \rightarrow \sim t$

$\sim w \rightarrow \sim m$

$\sim g$

- a) Express this argument in symbolic logical form using the labels:

$g = \text{"I will have a good holiday"}$

$t = \text{"I have time"}$

$m = \text{"I have money"}$

$w = \text{"I work"}$ .

## a) Solution

Consider the following arguments:

- (1) I will have a good holiday only if I have both time and money.
- (2) If I work then I will have no time.
- (3) If I do not work, then I will have no money.
- (4) Therefore, I will not have a good holiday.

$$P_1 \ g \rightarrow (t \wedge m)$$

$$P_2 \ w \rightarrow \sim t$$

$$P_3 \sim w \rightarrow \sim m$$

$$P_4 \sim g$$

2016 T1 3) (iii)\*

b) Show that  $P_4$  can be logically deduced from the previous statements.

$$\begin{array}{l} P_1 g \rightarrow (t \wedge m) \equiv \neg(t \wedge m) \rightarrow \neg g \equiv (\underbrace{\neg t \vee \neg m}) \rightarrow \neg g \\ P_2 w \rightarrow \neg t \\ P_3 \neg w \rightarrow \neg m \quad \text{if or } \neg m \text{ is true} \\ P_4 \neg g \end{array}$$

$$\begin{aligned} & P_1 \wedge P_2 \wedge P_3 \\ &= (\underline{w \rightarrow \neg t}) \wedge (\underline{\neg w \rightarrow \neg m}) \wedge ((\neg t \vee \neg m) \rightarrow \neg g) \\ &= (\underbrace{\neg t \vee \neg m}) \wedge (\underbrace{(\neg t \vee \neg m)}_{\text{(case exhaustion)}} \rightarrow \neg g) \\ &\Rightarrow \neg g \quad (\text{modus ponens}) \\ \text{Hence, } P_4 \text{ can be logically deduced.} \end{aligned}$$

2016 T1 3) (iii)\*

b) Show that  $P_4$  can be logically deduced from the previous statements.

$$P_1 \ g \rightarrow (t \wedge m)$$

$$P_2 \ w \rightarrow \sim t$$

$$P_3 \sim w \rightarrow \sim m$$

$$P_4 \sim g$$

Consider  $P_1 \wedge P_2 \wedge P_3$

$$\equiv (w \rightarrow \sim t) \wedge (\sim w \rightarrow \sim m) \wedge (g \rightarrow (t \wedge m))$$

$$\equiv (w \vee \sim w) \wedge (w \rightarrow \sim t) \wedge (\sim w \rightarrow \sim m) \wedge (g \rightarrow (t \wedge m)) \text{ (Union)}$$

$$\implies (\sim t \vee \sim m) \wedge (g \rightarrow (t \wedge m)) \text{ (Case exhaustion)}$$

$$\equiv (\sim t \vee \sim m) \wedge ((\sim t \vee \sim m) \rightarrow \sim g) \text{ (Contrapositive)}$$

$$\implies \sim g \text{ (Modus ponens)}$$

$$\equiv P_4.$$

Hence,  $P_4$  can be logically deduced.

## 2017 T2 Q3 (v)\*

Use induction to show that for all integers  $n \geq 4$ , there exists  $a, b \in \mathbb{N}$  such that  $n = 2a + 5b$ .

Let  $P(n)$  be statement " $\exists a, b \in \mathbb{N}$  s.t.  $n = 2a + 5b$ ,  $\forall n \geq 4$ ".

S1  $P(4)$ :  $4 = 2(2) + 5(0)$ ,  $2, 0 \in \mathbb{N}$

So  $P(4)$  is true.

S2 Assume  $P(k)$  is true,  $\forall k \geq 4$   
RTP:  $P(k+1)$  is also true.

## 2017 T2 Q3 (v)\*

Use induction to show that for all integers  $n \geq 4$ , there exists  $a, b \in \mathbb{N}$  such that  $n = 2a + 5b$ .

## 2017 T2 Q3 (v)\*

Use induction to show that for all integers  $n \geq 4$ , there exists  $\underline{a}, \underline{b} \in \mathbb{N}$  such that  $n = 2a + 5b$ .

Assume  $\exists a, b \in \mathbb{N}$ ,  $b = \underline{2a+5b}$ ,

RTP  $\exists A, B \in \mathbb{N}$ ,  $b+2 = \underline{2A+5B}$ ,

$$\text{LHS} = b+2$$

$$= 2a+5b+2 \quad (\text{by assumption})$$


$$= 2\underline{(a+1)} + 5b$$

$$= 2\underline{A} + 5\underline{B}, \quad A, B \in \mathbb{N}$$

$P(b+2)$  is true if  $P(b)$  is true.

## 2017 T2 Q3 (v)\*

Use induction to show that for all integers  $n \geq 4$ , there exists  $a, b \in \mathbb{N}$  such that  $n = 2a + 5b$ .



$$P(5): 5 = 2(0) + 5(1), \quad 0, 1 \in \mathbb{N}$$

$P(5)$  is true.

## 2017 T2 Q3 (v)\*

Use induction to show that for all integers  $n \geq 4$ , there exists  $a, b \in \mathbb{N}$  such that  $n = 2a + 5b$ .

Let  $P(n)$  be the statement  $n = 2a + 5b$  for some  $a, b \in \mathbb{N}$ .

Step 1: Prove  $P(4)$  and  $P(5)$  is true.

For  $P(4)$ , we have  $P(4) = 4 = 2(2) + 5(0)$  where  $0, 2 \in \mathbb{N}$ . Hence,  $P(4)$  is true.

For  $P(5)$ , we have  $P(5) = 5 = 2(0) + 5(1)$  where  $0, 1 \in \mathbb{N}$ . Hence,  $P(5)$  is true.

## 2017 T2 Q3 (v)

Use induction to show that for all integers  $n \geq 4$ , there exists  $a, b \in \mathbb{N}$  such that  $n = 2a + 5b$ .

Step 2:

Assume  $P(k)$  is true for all integers  $k \geq 4$ .

i.e There exists  $a, b \in \mathbb{N}$  such that  $k = 2a + 5b$ .

We are now required to prove  $P(k + 2)$  is also true.

i.e There exists  $A, B \in \mathbb{N}$  such that  $k + 2 = 2A + 5B$ .

$$\begin{aligned} k + 2 &= 2a + 5b + 2 \text{ (from induction hypothesis)} \\ &= 2(a + 1) + 5b \\ &= 2A + 5B \end{aligned}$$

where  $A = a + 1 \in \mathbb{N}$  and  $B = b \in \mathbb{N}$ . Hence,  $P(k + 2)$  is true.

## 2017 T2 Q3 (v)


Use induction to show that for all integers  $n \geq 4$ , there exists  $a, b \in \mathbb{N}$  such that  $n = 2a + 5b$ .

Step 3:

Hence, because  $P(4)$  and  $P(5)$  is true from step 1 and  $P(k + 2)$  is true if  $P(k)$  is true from step 2, then by the principle of mathematical induction,  $P(n)$  is true for all integers  $n \geq 4$ .  $\square$

## 5. Graphs

# Types of paths and circuits



## Definition: Euler Path


A path that passes through every edge of the graph  $G$  exactly once.

\* A connected graph  $G$  has an Euler path if and only if there are exactly 2 vertices of odd degree.

## Definition: Euler Circuit

A circuit that passes through every edge of the graph  $G$  exactly once.

\* A connected graph  $G$  has an Euler circuit if and only if there are only even vertex degrees.



# Types of paths and circuits



## Definition: Hamilton Path

A path that passes through every vertex of the graph  $G$  exactly once.

No easy criteria for Hamilton's.

A Hamilton path or circuit uses at most 2 edges incident to each vertex.

## Definition: Hamilton Circuit

A circuit that passes through every vertex of the graph  $G$  exactly once except for the first and last vertex.

\* If a graph  $G$  has a Hamilton circuit, then the circuit must include all edges incident with vertices of degree 2.


\* Let  $G$  be a connected and simple graph with  $n \geq 3$  vertices, such that each vertex has degree at least  $n/2$ . Then  $G$  has a Hamilton circuit.

# Example 1

b) simple

6 vertices total

$$\frac{6}{2} = 3$$



2018 S2 Q2 iii)

Let  $G$  denote the graph shown above.

- a) Does  $G$  have an Euler path? Explain your answer. ✓
- b) Does  $G$  have a Hamilton circuit? Explain your answer. ✓
- c) Is  $G$  bipartite? Explain your answer.

# Example 1 Solutions

a)

$G$  does have an Euler path. Why?

# Example 1 Solutions

a)

$G$  does have an Euler path. Why?

$G$  is connected. ✓

$G$  has exactly two odd vertices at  $c$  and  $f$ . ✓

# Example 1 Solutions

a)

$G$  does have an Euler path. Why?

$G$  is connected.

$G$  has exactly two odd vertices at  $c$  and  $f$ .

b)

$G$  does have a Hamilton circuit. Why?

# Example 1 Solutions

a)

$G$  does have an Euler path. Why?

$G$  is connected.

$G$  has exactly two odd vertices at  $c$  and  $f$ .

b)

$G$  does have a Hamilton circuit. Why?

$G$  is connected, simple and has more than 3 vertices.

$G$  has 6 vertices and each vertex has degree greater than  $\frac{6}{2} = 3$ .


(c)

We will return to c) later.

## Example 2

c)  $K$

connected  
every vertex  
has only even  
degrees.



2012 S2 Q2 iii) c) and d)

Let the graph above be  $K$ .

- c) Show that  $K$  has an Euler circuit. ✓
- d) Show that  $K$  has no Hamiltonian circuit. ✗

## Example 2 Solutions

c)

$K$  is connected.

$K$  has only vertices with even degrees.

## Example 2 Solutions

c)

$K$  is connected.

$K$  has only vertices with even degrees.

d)

To show this, we shall combine two different results, as well as a proof by contradiction.

## Example 2 Solutions

c)

$K$  is connected.

$K$  has only vertices with even degrees.

d)

To show this, we shall combine two different results, as well as a proof by contradiction.

Firstly, we have that, if  $K$  did contain a Hamiltonian circuit, it should use up all edges incident with vertices with degree of 2. In  $K$ , this would be the vertices:  $A, C, D, F$ .

This means edges  $AB, DB, FB, CB$  must be used.

## Example 2 Solutions

c)

$K$  is connected.

$K$  has only vertices with even degrees.

d)

To show this, we shall combine two different results, as well as a proof by contradiction.


Firstly, we have that, if  $K$  did contain a Hamiltonian circuit, it should use up all edges incident with vertices with degree of 2. In  $K$ , this would be the vertices:  $A, C, D, F$ .

This means edges  $AB, DB, FB, CB$  must be used.

Secondly, we have that a Hamiltonian circuit uses at most 2 edges incident to any vertex.

Thus, the vertex  $B$  leads to a contradiction as  $B$  has all 4 of its edges being used up.

# Bipartite Graph




Definition: Bipartite Graph

A graph  $G$  is bipartite if there are two disjoint sets  $V_1$  and  $V_2$  such that  $V_1 \cup V_2 = V$  and every edge of  $G$  connects a vertex in  $V_1$  to a vertex in  $V_2$ .

In loose terms, a graph is bipartite if every vertex in one subgroup only 'bridges' over to vertices in the other subgroup and not to its own.

## Example 3



Return to 2018 S2 Q2 iii) c)


Let  $G$  denote the graph shown above.

- Does  $G$  have an Euler path? Explain your answer.
- Does  $G$  have a Hamilton circuit? Explain your answer.
- Is  $G$  bipartite? Explain your answer.

## Example 3 Solutions

c)

$G$  cannot be bipartite. Why?



$H$  cannot be bipartite.

Because in  $V_2$ ,

the vertex  $b$  is connected  
to  $e$  and  $f$  within  
the same subset.

## Example 3 Solutions

c)

$G$  cannot be bipartite. Why?

Let's try splitting the vertices into two groups!

# Planar Graphs



## Definition: Planar Graph


A graph is planar if you can draw it (in 2D!) without crossing edges.

## Theorems for Planar Graphs

If  $G$  is a connected planar graph with  $e$  edges and  $v \geq 3$  vertices, then

(1)  $e \leq 3v - 6$ ,

(2)  $e \leq 2v - 4$  (if  $G$  has no cycles of length 3).



# Other Useful Theorems/Lemmas

## Definition: Euler's Formula

If  $G$  is a connected planar graph with  $e$  edges and  $v$  vertices, and if  $r$  is the number of regions in a planar representation of  $G$ , then,

$$v - e + r = 2.$$

## Definition: Handshaking Lemma

For any graph with vertices  $V$  and edges  $E$ :


$$2|E| = \sum_{v \in V} \deg(v).$$

total vertex degrees

For any planar graph with vertices  $V$  and regions  $R$ :


$$2|E| = \sum_{r \in R} \deg(r).$$

total region degrees.

## Example 4

$$V=7, R=?$$
$$e = ? \quad V - e + R = 2$$

2014 S2 Q2 ii)

A connected planar graph has vertices of degrees 6, 5, 5, 2, 2, 1, 1.

- How many regions does the graph have?
- Draw an example of such a planar graph.

$$\text{total vertex degrees} = 6+5+5+2+2+1+1 \\ = 22.$$

# Example 4 Solutions

a)

We are given the number of vertices and degrees of each vertex. We are trying to find number of regions. What can we do?

# Example 4 Solutions

a)

We are given the number of vertices and degrees of each vertex. We are trying to find number of regions. What can we do?

Notice we have most components of Euler's formula, and furthermore, Handshaking Lemma can be used to determine the missing piece, the number of edges.

## Example 4 Solutions

a)


We are given the number of vertices and degrees of each vertex. We are trying to find number of regions. What can we do?

Notice we have most components of Euler's formula, and furthermore, Handshaking Lemma can be used to determine the missing piece, the number of edges.

22

By Handshaking Lemma,  $2e = 6 + 5 + 5 + 2 + 2 + 1 + 1 \implies e = 11$ .

Finally, by Euler's formula,  $7 - 11 + r = 2 \implies r = 6$ .


$$\begin{matrix} v & e & r \end{matrix}$$


## Example 4 Solutions

b)

Let's try draw it out!

$$\Gamma = 7, \quad e = 11, \quad r = 6.$$

## Example 5

2014 S1 Q2 iv)

- a) State Euler's Formula for a connected planar graph having  $v$  vertices,  $e$  edges and  $r$  regions.  $V - e + r = 2$
- b) Show that if  $G$  is a connected planar simple graph with  $v \geq 3$  vertices, then  $e \leq 3v - 6$ .
- c) Hence show that a connected planar simple graph with  $v \geq 3$  vertices has at least one vertex of degree less than or equal to 6.

# Example 5 Solutions

a)

$$v - e + r = 2.$$

# Example 5 Solutions

a)

$$v - e + r = 2.$$

b)

The proof uses a few niche results. I will briefly describe it here, but it is more so knowing how to use the  $e \leq 3v - 6$  theorem that it more important.

Typically, the theorem will be used to prove that a graph is **not** planar!

## Example 5 Solutions

a)

$$v - e + r = 2.$$

b)

The proof uses a few niche results. I will briefly describe it here, but it is more so knowing how to use the  $e \leq 3v - 6$  theorem that it more important.

Typically, the theorem will be used to prove that a graph is **not** planar!  $G$  is simple, connected, planar and has  $v \geq 3$ , therefore, each of its regions must have at least a degree of 3." (1)

Handshaking Lemma tells us that  $2e =$  total region degrees. (2)

$$\text{So, } 2e \geq 3r \implies 2e \geq 3(2 + e - v).$$

$$(1)+(2) \quad 2e \geq 6 + 3e - 3v$$

$2e = \text{total region degrees} \geq 3r$  each region has at least deg. number of regions

## Example 5 Solutions

'at least 1 vertex has degree  $\leq 6'$

c)

Clearly, trying to brute force this will be rather hard considering that 'at least' accounts for a lot of possibilities. Instead, we should consider proof by contradiction, as this turns the 'at least' statement into 'all vertices have degree greater than 6'.

Assume that all vertices have degree greater than 6.

## Example 5 Solutions

c)

Clearly, trying to brute force this will be rather hard considering that 'at least' accounts for a lot of possibilities. Instead, we should consider proof by contradiction, as this turns the 'at least' statement into 'all vertices have degree greater than 6'.

Recall the other Handshaking Lemma.  $2e = \text{total vertex degrees}$ .

Thus, by our assumption that 'all vertices have degree greater than 6',  
 $2e \geq 6v \implies e \geq 3v$ .

This leads to a contradiction, as  $e \leq 3v - 6$ .

$2e \geq 6v \rightarrow e \geq 3v$ .  
each vertex  $\nearrow$  number of vertices  
has at least deg of 6.

↑  
this must  
actually be  
false.

# Isomorphism and Homeomorphism

## Definition: Isomorphism

Let  $G$  and  $G'$  be graphs with vertices  $V$  and  $V'$ , and edges  $E$  and  $E'$ . Then  $G$  is isomorphic to  $G'$ , if and only if there are two bijections  $f : V \rightarrow V'$  and  $g : E \rightarrow E'$ , such that  $e$  is incident with  $v$  in  $G$  if and only if  $g(e)$  is incident with  $f(v)$  in  $G'$ .



Loosely speaking, this means that the two graphs are the same except for how the vertices are labelled.

## Important Property of Isomorphic Graphs

If two graphs are isomorphic, they must have the same invariant properties.

Examples of invariant properties include: number of vertices and edges, total degree, number of vertices of a certain degree, etc.


example of isomorphism.



# Isomorphism and Homeomorphism

## Definition: Homeomorphism

Two graphs are homeomorphic if and only if they can both be obtained from a common graph by elementary subdivisions.




## Definition: Kuratowski's Theorem

A graph is planar if and only if it does not contain a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$ .




## Example 6

$$V=6$$
$$e=9$$



$$V=6$$
$$e=9$$



2018 S1 Q2 iv) c)

- c) Are  $G_1$  and  $G_2$  isomorphic? Explain why.

# Example 6 Solutions

c)

Usually, unless it's trivial to determine the isomorphism, you should begin by trying to **disprove** the existence of an isomorphism.

# Example 6 Solutions


c)

Usually, unless it's trivial to determine the isomorphism, you should begin by trying to **disprove** the existence of an isomorphism.

Run through some common invariants and see if there's any discrepancy between the two graphs.

In this case, notice that in  $G_2$ , vertex  $B$  has degree 3, whereas on  $G_1$ , no vertices have degree 3.

## Example 7



2012 S2 Q2 iv)

- State Kuratowski's theorem characterising non-planar graphs.
- Show that the above graph is not planar.

## Example 7 Solutions

a)

A graph is planar if and only if it does not contain a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$  (as shown in lecture slides).

A graph is non-planar iff it does contain a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$

## Example 7 Solutions

a)

A graph is planar if and only if it does not contain a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$  (as shown in lecture slides).

Taking its contrapositive we get, a graph is **non-planar** if and only if it **does contain** a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$ .

# Example 7 Solutions

a)

A graph is planar if and only if it does not contain a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$  (as shown in lecture slides).

Taking its contrapositive we get, a graph is **non-planar** if and only if it **does contain** a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$ .

b)

The idea would be to determine whether the above graph is a subgraph of  $K_{3,3}$  or  $K_5$ .

# Example 7 Solutions

a)

A graph is planar if and only if it does not contain a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$  (as shown in lecture slides).

Taking its contrapositive we get, a graph is **non-planar** if and only if it **does contain** a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$ .

b)

The idea would be to determine whether the above graph is a subgraph of  $K_{3,3}$  or  $K_5$ .

This means we should in a sense, do the opposite of elementary subdivisions.

This is because rather than taking  $K_{3,3}$  or  $K_5$  and breaking it down to a subgraph, we are starting at the potential subgraph and are trying to build it back up to being a  $K_{3,3}$  or  $K_5$ .

## Example 7 Solutions



b)

Let's note some key features of  $K_{3,3}$  and  $K_5$ . A  $K_{3,3}$  has 6 vertices and 9 edges, whereas a  $K_5$  has 5 vertices and 10 edges.

# Example 7 Solutions

b)

Let's note some key features of  $K_{3,3}$  and  $K_5$ . A  $K_{3,3}$  has 6 vertices and 9 edges, whereas a  $K_5$  has 5 vertices and 10 edges.

The graph we have right now has 7 vertices and 11 edges. It seems it would be easier to attain the  $K_{3,3}$  graph from our graph, than to try attain  $K_5$ .

# Example 7 Solutions

b)

Let's note some key features of  $K_{3,3}$  and  $K_5$ . A  $K_{3,3}$  has 6 vertices and 9 edges, whereas a  $K_5$  has 5 vertices and 10 edges.

The graph we have right now has 7 vertices and 11 edges. It seems it would be easier to attain the  $K_{3,3}$  graph from our graph, than to try attain  $K_5$ .


We are trying to seek a change to our graph that leads to a decrease of 1 vertex and 2 edges. Let's try remove vertex  $E$ , and replace the detour path  $FED$  with just  $FD$ .

Let's draw it out and see if we can form a  $K_{3,3}$ !

# Adjacency Matrix

## Definition: Adjacency Matrix

Let  $G$  be a graph with an ordered listing of vertices  $v_1, v_2, \dots, v_n$ . The adjacency matrix of  $G$  is the  $n \times n$  matrix  $A = [a_{ij}]$  with  $a_{ij}$  being the number of edges connecting  $v_i$  and  $v_j$ .


$$A = \begin{pmatrix} v_1 & v_2 \\ v_1 & v_2 \end{pmatrix} \quad \begin{matrix} v_1 \\ v_2 \end{matrix}$$


## Powers of the Adjacency Matrix

Given the  $A^k$  matrix, the  $i$ th row and  $j$ th column value of this matrix graphically represents the number of walks between  $v_i$  and  $v_j$  that is of length  $k$ .

$$A^6 = \begin{pmatrix} v_1 & v_2 \\ v_1 & v_2 \end{pmatrix}$$

there are 4 walks of length 6 from  $v_1$  to  $v_2$ .



## Example 8

2013 S1 Q2 ii)

A graph  $H$  on the vertices  $A, B, C, D$  (in that order) has adjacency matrix:

$$M = \begin{pmatrix} 1 & 2 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix}$$


- Draw the graph  $H$ .
- $M^3$  has the number 12 in position 1, 2 (that is first row, second column). What does that mean in terms of the graph?

## Example 8 Solutions

a)

Let's try draw it out!

$$M = \begin{pmatrix} A & B & C & D \\ A & 1 & 2 & 0 & 0 \\ B & 2 & 0 & 1 & 0 \\ C & 0 & 1 & 0 & 0 \\ D & 0 & 0 & 0 & 2 \end{pmatrix}$$



## Example 8 Solutions

$$M^3 = \begin{pmatrix} - & 12 & - & - \\ - & - & - & - \\ - & - & - & - \\ - & - & - & - \end{pmatrix}$$

b)

Recall that the  $n$ th power of an adjacency matrix conveys information regarding the number of possible walks from one vertex to another that passes through  $n$  edges.

there are 12 walks from A to B  
of length 3.

# Example 8 Solutions

b)

Recall that the  $n$ th power of an adjacency matrix conveys information regarding the number of possible walks from one vertex to another that passes through  $n$  edges.

So for  $H$ , position 1, 2 refers to vertices  $A, B$ . This means that there are 12 possible walks from  $A$  to  $B$  that has length of 3.

Let's head back to our graph and see what these walks may look like.

## Example 9

2012 S1 Q2 iii)



Consider the graph  $G$  consisting of the vertices of a cube and its edges.

- a) Draw a planar representation of this graph, labelled with vertices 1, 2, 3, 4, 5, 6, 7, 8 so that the path 123456781 is a Hamiltonian circuit, and so that 1 and 6 correspond to opposite vertices in the cube.
- b) Is the graph  $G$  bipartite?
- c) With the above labelling, write down the first four rows of the adjacency matrix  $A$  of the graph.
- d) Without computing the matrix product  $A^3$ , compute the (1, 3) entry and the (1, 6) entry of  $A^3$ .

## Example 9 Solutions

a)

Let's try draw it! Keep in mind that 'Hamiltonian' means passing through all vertices exactly once.



b)

Try this yourselves.

## Example 9 Solutions

c)

Let's first define the adjacency matrix to have rows and columns ordered 1, 2, 3, ..., 8.

Refer back to our graph drawn above. We can fill in the entries one by one.

$$A = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 \\ 2 & & & & & & & \\ 3 & & & & & & & \\ 4 & & & & & & & \\ \vdots & & & & & & & \\ 8 & & & & & & & \end{pmatrix}$$

The matrix is an 8x8 adjacency matrix. The columns are labeled 1 through 8 at the top. The rows are labeled 1 through 8 on the left. The matrix entries are filled in as follows:

- Row 1: 0, 1, 0, 1, 0, 0, 0, 1
- Row 2: 1, 0, 1, 0, 0, 0, 0, 1
- Row 3: 0, 0, 0, 0, 0, 0, 0, 0
- Row 4: 0, 0, 0, 0, 0, 0, 0, 0
- Row 5: 0, 0, 0, 0, 0, 0, 0, 0
- Row 6: 0, 0, 0, 0, 0, 0, 0, 0
- Row 7: 0, 0, 0, 0, 0, 0, 0, 0
- Row 8: 0, 0, 0, 0, 0, 0, 0, 0

Red annotations highlight the following:

- The row and column indices 1 through 8 are written above the matrix and to the left of the matrix respectively.
- The entry in the second row, second column (labeled 2, 2) is circled in red.
- The entry in the second row, third column (labeled 2, 3) is circled in red.
- The entry in the second row, fourth column (labeled 2, 4) is circled in red.
- The entry in the second row, fifth column (labeled 2, 5) is circled in red.
- The entry in the second row, sixth column (labeled 2, 6) is circled in red.
- The entry in the second row, seventh column (labeled 2, 7) is circled in red.
- The entry in the second row, eighth column (labeled 2, 8) is circled in red.

# Example 9 Solutions

c)

Let's first define the adjacency matrix to have rows and columns ordered 1, 2, 3, ..., 8.

Refer back to our graph drawn above. We can fill in the entries one by one.

$$A = \begin{pmatrix} 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}.$$

## Example 9 Solutions

d)

Again, recall what the  $n$ th power of a matrix means.

Rather than calculating the powers of the 8 by 8 matrix, we can instead do this question more intuitively and graphically.

$$A^3 = \begin{pmatrix} 1 & & & & & & & \\ & 3 & & & & & & \\ & & 6 & & & & & \\ & & & 1 & & & & \\ & & & & 1 & & & \\ & & & & & 1 & & \\ & & & & & & 1 & \\ & & & & & & & 1 \end{pmatrix}$$

# Example 9 Solutions

d)

Again, recall what the  $n$ th power of a matrix means.

Rather than calculating the powers of the 8 by 8 matrix, we can instead do this question more intuitively and graphically.

We are aiming to find the number of possible walks for vertex 1 to 3 and vertex 1 to 6 that are exactly of length 3.

Let's refer back to our graph and see if we can visually determine the solution.

# Example 9 Solutions

d)

Again, recall what the  $n$ th power of a matrix means.

Rather than calculating the powers of the 8 by 8 matrix, we can instead do this question more intuitively and graphically.

We are aiming to find the number of possible walks for vertex 1 to 3 and vertex 1 to 6 that are exactly of length 3.

Let's refer back to our graph and see if we can visually determine the solution.

Turns out the (1, 3) entry is 0 and the (1, 6) entry is 6.

# Weighted Trees

## Definition: Tree


A tree is a simple connected graph with no circuits.

## Definition: Spanning Tree

A spanning tree is a sub-tree that contains every vertex of graph.


## Definition: Weighted Tree

A weighted tree is a tree whose edges have been given numbers called weights. The weight of an edge  $e$  is denoted by  $w(e)$ .



# Kruskal's Algorithm

$n$  vertices  
 $n-1$  edges




## Definition: Kruskal's Algorithm

Kruskal's Algorithm can be used to find the minimal spanning tree. This means the spanning tree with lowest sum of weights.

1. Start with  $T$ , the empty tree.
2. Sort the edges by increasing weight.
3. Going down the list, add an edge to  $T$  if and only if it does not form a circuit with the existing edges.
4. Continue until  $T$  has  $n - 1$  edges.

## Example 10



2013 S2 Q2 v)

- a) Use Kruskal's algorithm to construct a minimal spanning tree  $T$  for the above weighted graph. Make a table showing the details of each step in your application of the algorithm.

# Example 10 Solutions

a)

Recall Kruskal's Algorithm. We shall begin with an empty tree and build our way up to a minimal spanning tree by constructing a table with the weighted edges.

Edge	Weight	Chosen?
$BE$	4	✓
$DE$	4	✓
$AF$	5	✓
$EF$	5	✓
$AE$	6	✗
$BC$	6	✓
$BD$	6	
$CD$	6	
$DF$	6	

T

A

C

B

4

E

S

D

F

# Dijkstra's Algorithm




## Definition: Dijkstra's Algorithm

Dijkstra's Algorithm can be used to find the minimal spanning tree with respect to a particular vertex. This means, we can for example obtain the tree such that all the walks from a particular vertex  $A$  to any other vertex is the least in terms of weight.

1. Start with the subgraph  $T$ , containing only the given vertex,  $A$ .
2. Consider all edges  $e$  incident to  $A$ .
3. Of these edges, choose an edge  $e$  with the lowest weight and note down the vertex,  $B$ , it leads to.
4. Add edge  $e$  and  $B$  to  $T$ .
5. Continue until  $T$  contains all the vertices of  $G$ .

## Example 11



2014 S1 Q2 v)

- a) Use Dijkstra's algorithm to find a spanning tree that gives the shortest paths from A to every other vertex of the graph. Make a table showing the details of each step in your application of the algorithm.

# Example 11 Solutions

a)

Recall Dijkstra's algorithm. We start at the assigned vertex, in this case,  $A$ , and we build up the tree by constructing a table to consider the potential edge candidates.

Edge candidates	Next edge	Next vertex
$AE(1), AD(2), AB(6)$	$AE(1)$	$E$
$AB(6), \cancel{AD}(2), EB(4), ED(4), EC(4)$	$AD(2)$	$D$
$AB(6), DC(6), EB(4), EC(4)$	$EB(4)$	$B$
<u><math>BC(6), DC(6), EC(4)</math></u>	<u><math>EC(4)</math></u>	<u><math>C</math></u>

T

