


PhD Midway Seminar


Simulation Tool and its application

Raju Rimal

03 March, 2017



Introduction



My PhD Plan

PhD Program	
Phase 1	Make a simulation Tool
Phase 2	Apply it for comparing different estimation Methods
Phase 3	Extend the simulation tool for model with background information
Phase 4	Apply it to test multi-matrix extension of PLS models such as LPLS and UPLS

- Make **Simulation Tools** for multi-response linear model data
- Using the tool, compare various **estimation techniques** and **understand** them
- **Extend** the simulation tool incorporating model with **background information**
- Apply this extended tool to **test multi-matrix extension of partial least square (PLS)** models such as LPLS and UPLS (both uses background information about X and Y for analysis)



What I learn


- Advanced Multivariate Model and technique to analyze it
- Programming concept for developing statistical packages and applications for various statistical methods
- Extending and improving existing methods in statistics
- And, obviously, to properly document what I have done



Today's Special

Today I will talk about:

- Simulation tool ([simulatr](#)) we are building
- A [comparative study](#) of various estimation techniques by simulating linear model data using simulatr



simrel-m: A versatile tool for simulating multi-response linear model data





Overview


simrel-M is an extension of simrel (Sæbø, Almøy, & Helland, 2015) r-package for simulating multi-response data

- Uses the idea of reduction of random regression model by separating latent space of \mathbf{X} into subspaces that is relevant and irrelevant for predicting each response
- The underlying concept is based on reparameterizing the population model,


$$\mathbf{Y} = \boldsymbol{\mu}_Y + \mathbf{B}^t (\mathbf{X} - \boldsymbol{\mu}_X) + \boldsymbol{\epsilon}, \text{ where } \boldsymbol{\epsilon} \sim N(0, \boldsymbol{\Sigma}_{Y|X})$$



Underlying procedure



- Collect population input parameter from users such as: number of variables, coefficient of determination and the position of relevant components
- Make a covariance matrix satisfying input parameters
- Rotate the covariance matrix orthogonally
- Sample calibration and validation sets



A comparative study of different estimation methods using simulated data



Overview

Four estimation methods were considered

Ordinary Least Squares (OLS)

- Although unbiased, suffer highly from multicollinearity
- Widely used and can be used as reference for comparison

Envelope

- Relatively new method (Cook, Helland, & Su, 2013) and is also based on reduction of regression model
- Based on Maximum Likelihood but works better than OLS in p approaches n

Partial Least Squares (PLS)

- Well established and widely used method
- Based on Latent Structure and free of multicollinearity problem

Bayes PLS


- Bayesian Estimation of regression coefficient
- Promising performance was shown in previous studies (Helland, Sæbø, & Tjelmeland, 2012)




Simulation Design

From the possible combination of following parameter combination, 32 single response calibration sets were simulated with 5 replication of each.

- **Number of sample observations:** 50
- **Number of predictor variables:** 15 and 40
- **Coefficient of determination (R^2):** 0.5 and 0.9
- **Level of multicollinearity:** 0.5 and 0.9
- **Position of relevant components:** 1 and 2; 1 and 3; 2 and 3; 1, 2 and 3




A Systematic Comparison



- Bayes PLS has out-performed others methods in all kinds of data
- Envelope has performed better than OLS in all situations and PLS in some situations
- OLS prediction is very poor in noisy data with many predictor variables
- Position of relevant component and the decaying factor of eigenvalue has less impact on prediction in all the models




A Systematic Comparison



- Bayes PLS has approached to its minimum error with very few component and remained low for additional component
- PLS has moderate performance but better than envelope in many situations.
- OLS prediction is poor especially with large number of predictor
- Envelope method captured its minimum error and the error increased with additional components



Demonstration



New Seed



Type of simulation:

Bivariate Simulation

Parameter Settings

Simulation Overview


Estimation

Model Comparison

simulatr Application

Welcome to Simulatr

Norwegian University of Life Science
Universitetstunet 3
1433 Ås



salamat Dakujem
GRACIAS ASANTE
TAKK धन्यवाद
HVALA
Euχαριστώ
GRAZZII
DANKE
Paxmet
kiitos
ARIGATO
suwun
धन्यवाद
MERCI
teşekkür ederim
mahalo
DANKE
TAKK
спасибо
salamat
mersi
감사합니다
GRAZAS
salamat
kiitos
TAKK
MAHALO
drigato
takk
HVALA
kiitos
GRAZAS
TAKK
salamat
SUWUN
SALAMAT
gracias
teşekkür ederim
SUWUN
hvala
salamat
ليزج اركش
merci



References



References

Cook, R., Helland, I., & Su, Z. (2013). Envelopes and partial least squares regression. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 75(5), 851–877.

Helland, I. S., Sæbø, S., & Tjelmeland. (2012). Near optimal prediction from relevant components. *Scandinavian Journal of Statistics*, 39(4), 695–713.

Sæbø, S., Almøy, T., & Helland, I. S. (2015). Simrel—A versatile tool for linear model data simulation based on the concept of a relevant subspace and relevant predictors. *Chemometrics and Intelligent Laboratory Systems*, 146, 128–135.