The Continuum Limit of the Correlation Functions of Wilson Loops in Four-Dimensional Lattice Yang–Mills Theory

Abstract

We consider pure Yang–Mills theory with a compact connected simple gauge group on a four-dimensional periodic lattice of spacing ε, with the bare coupling fixed by Bałaban's renormalisation group so that the running coupling at a fixed unit scale has a prescribed small value. For every finite family of pairwise disjoint simple closed lattice loops, in any planes, each with the character of an arbitrary irreducible representation, we divide the expectation of the product of their Wilson loops by an explicit number for each loop, the exponential of a perimeter term and of a term for each corner, whose coefficients do not depend on the loop. The coupling is fixed independently of the representations, which enter only through the scale at which the number is evaluated. We prove that the resulting renormalised expectation is bounded uniformly in ε and converges as $\varepsilon \to 0$ along the sequence $\varepsilon = uL^{-K}$. The limits are reflection positive, and they do not vanish for loops whose lengths lie below a bound which depends on their representations and tends to infinity as the coupling tends to zero.

Contents

1 Introduction **3**
- 1.1 The Lattice Theory . . . 3
- 1.2 Loop Ensembles . . . 4
- 1.3 The Normalisation and the Results . . . 5
- 1.4 The Argument . . . 10

2 Loops and Their Geometry **11**
- 2.1 Segments and Corners . . . 11
- 2.2 The Chord–Arc Constant . . . 13
- 2.3 Arcs around a Set . . . 14
- 2.4 The Neighbourhood of a Corner . . . 15
- 2.5 A Summation over the Scales . . . 16

3 The Construction **17**
- 3.1 The Tube of a Target . . . 17
- 3.2 The Density after k Steps . . . 21
- 3.3 Soft Test Functions . . . 24
- 3.4 The Depth of the **R**-Operation . . . 26
- 3.5 The Construction of This Paper . . . 28
- 3.6 The Stopping Index . . . 29

4 The Coupling Flow **30**
- 4.1 The Coefficients of the Flow . . . 31
- 4.2 Proof of the Coupling Flow . . . 34

5 Dressed Loops and the First Step **35**
- 5.1 Two Integrals . . . 35
- 5.2 The First Step of the Construction . . . 36

5.3 Dressed Loops . . . 36
5.4 The First Step with a Loop Ensemble . . . 44
5.5 The First Step at Complex Backgrounds . . . 51
5.6 The First Step on a Term with Large-Field Regions . . . 57

6 The Inductive Step 70
6.1 The Evaluation Point . . . 71
6.2 The Identity of the Step . . . 72
6.3 The Expansion of the Words in the Localisation Parameters . . . 74
6.4 The Fluctuation Integral with Marked Polymers . . . 80
6.5 The Groups of the Step . . . 85
6.6 The Count of the Groups of the Step . . . 88
6.7 The Small-Field Map and the Flat Parts . . . 100
6.8 The Deviation Bound of the New Groups . . . 103
6.9 The Contraction of the Deviation Parts . . . 105
6.10 The Carried Insertions at Complex Data . . . 106
6.11 The Regions of a Term and the Collar . . . 114
6.12 A New Large-Field Region: The Interior Groups . . . 115
6.13 A New Large-Field Region: The Groups at the Collar . . . 120
6.14 The Step on a Term with Earlier Large-Field Regions . . . 123
6.15 Large Fields and the **R**-Operation . . . 127
6.16 Several Loops, the Torus, and the Assembly . . . 129

7 The Flat Part and the Normalisation 133
7.1 The Reference Gas . . . 133
7.2 The Coefficients . . . 140
7.3 The Remainder of a Step . . . 142
7.4 The Normalisation . . . 144

8 The Identity at the Stopping Scale and the Uniform Bound 145
8.1 The Identity . . . 146
8.2 The Coarse Loop . . . 148
8.3 The Weight of the Large-Field Terms . . . 151
8.4 The Uniform Bound . . . 154

9 The Difference of the Walk Factors 158
9.1 Two Runs on Common Lattices . . . 158
9.2 Propagators and Their Walk Expansions . . . 159
9.3 The Galerkin Identities . . . 160
9.4 The Walk Terms . . . 163

10 The Large-Field Terms of Two Cutoffs 166
10.1 The Matched Decomposition . . . 166
10.2 The Shells of the Test Functions . . . 168
10.3 The Configurations of the Two Runs . . . 169
10.4 The Large-Field Comparison . . . 171

11 The Limit along the Full Sequence 173
11.1 The Exponents of the All-Small-Field Pair . . . 174
11.2 The Direct Bounds . . . 175
11.3 The Frozen Differences . . . 176
11.4 The Coefficients of the Two Runs . . . 177
11.5 The Generated Difference of One Step . . . 178
11.6 The All-Small-Field Pair . . . 182
11.7 The Pairs with Large Fields . . . 184
11.8 Proof of the Comparison . . . 185
11.9 Several Loops . . . 186

12 Symmetries, Reflection Positivity, and Nonvanishing 187
12.1 Symmetries and Reflection Positivity on the Lattice . . . 187
12.2 The Covariance of the Normalisation . . . 189

12.3 Proof of the Symmetries . . . 192
12.4 Proof of the Reflection Positivity . . . 193
12.5 Nonvanishing and the Growth Bound . . . 194

13 Non-Gaussianity 198
13.1 The Witness . . . 198
13.2 Gaussian Charge Models . . . 199
13.3 The Normalisations Drop Out . . . 200
13.4 The Free Field . . . 201
13.5 The Pair Terms . . . 204
13.6 The Shape Term . . . 210
13.7 The Coarse Covariance at Flat Fields . . . 214
13.8 The Second Moments of the Fluxes . . . 223
13.9 Proof of Theorem 1.8 . . . 235

14 Notes on the Papers 236

A The Constants in the Order of Choices 239

B Conventions 242

C The Conditions and Their Sources 243

1 Introduction

The expectation of a Wilson loop in lattice Yang–Mills theory is not expected to have a limit as the lattice spacing ε tends to zero. Already at the level of perturbation theory the loop acquires a self-energy proportional to its length in lattice units, so that one expects a factor of the form $\exp(-m(\varepsilon)P/\varepsilon)$ for a loop of physical length P, together with one further factor for each of its corners. The question that we address here is whether these are the only divergences: if one divides the expectation by a number of this form, with coefficients that do not depend on the loop, does the quotient converge?

We answer this question for the pure gauge theory on a four-dimensional torus, at small values of the renormalised coupling. The construction on which everything rests is Bałaban's renormalisation group for lattice gauge theories [8, 10, 9, 11, 12], which controls the effective densities obtained from the lattice measure by successive block averaging, uniformly in the number of steps. Bałaban's results concern the partition function and the effective actions, however, and not the expectations of observables, so that the Wilson loops have to be carried through the construction. This is what we do: we insert the loops into the bare density and follow them from step to step, as a line defect whose effect on the density, after each step, is the product of a number and of a function of the block field that is close to the Wilson loop of the coarser lattice. The numbers, collected over the steps, are the normalisation by which we divide.

1.1 The Lattice Theory

Let G be a compact connected simple Lie group in a fixed unitary realisation $G \subset U(N)$, and write $\mathrm{tr} \coloneqq N^{-1}\,\mathrm{Tr}$ for the normalised trace. We fix an odd integer L, a length $u > 0$, and an integer $m_T \geq 0$. The torus is $\mathbb{T} \coloneqq (\mathbb{R}/T\mathbb{Z})^4$ with the period $T \coloneqq 2L^{m_T}u$, and for an integer $K \geq 0$ the lattice of spacing

$$\varepsilon_K \coloneqq uL^{-K} \tag{1.1}$$

is $\mathbb{T}_K \coloneqq (\varepsilon_K\mathbb{Z}/T\mathbb{Z})^4 \subset \mathbb{T}$. A bond of $\mathbb{T}_K$ is an ordered pair $b = (x, x \pm \varepsilon_K e_\mu)$ of neighbouring sites, with $e_0, \dots, e_3$ the coordinate unit vectors, and b^{-1} denotes the reversed bond. A gauge field is a map U from the bonds to G with $U(b^{-1}) = U(b)^{-1}$. For a path $\gamma = (b_1, \dots, b_n)$ of

consecutive bonds we write $U(\gamma) \coloneqq U(b_1)\cdots U(b_n)$, and ∂p is the boundary of a plaquette p. The lattice theory at the bare coupling $g_0 > 0$ is the probability measure

$$\mathrm{d}\mu_{K,g_0}(U) \coloneqq \frac{1}{Z_{K,g_0}} \exp\Big(-\frac{1}{g_0^2}\sum_p (1 - \operatorname{Re}\operatorname{tr} U(\partial p))\Big)\,\mathrm{d}U\ , \tag{1.2}$$

where the sum runs over the unoriented plaquettes of $\mathbb{T}_K$ and $\mathrm{d}U$ is the product of the normalised Haar measures over the positive bonds.

The bare coupling is not a free parameter in the limit $K \to \infty$. Bałaban's renormalisation group assigns to g_0 a sequence of running couplings $g_1, \dots, g_K$, the coupling g_k belonging to the lattice of spacing $L^k\varepsilon_K$, and we fix g_0 by the requirement that the running coupling at the unit scale u has a prescribed value $\bar g$. More precisely, Theorem 4.1 provides, for every sufficiently small $\bar g > 0$ and every K, a bare coupling $g_0(K,\bar g)$ for which $g_K = \bar g$, and constants $0 < b \le b'$ and C_W such that

$$\bar g^{-2} + bK - C_W \le g_0(K,\bar g)^{-2} \le \bar g^{-2} + b'K + C_W\ . \tag{1.3}$$

The bare coupling thus tends to zero like $(\log(1/\varepsilon_K))^{-1/2}$, which is asymptotic freedom in the form in which it enters here. We write $\mathbb{E}_K$ for the expectation with respect to $\mu_{K,g_0(K,\bar g)}$.

1.2 Loop Ensembles

The loops that we consider are closed lattice paths without self-intersections. Since the lattices $\mathbb{T}_K$ are nested, a path of bonds of $\mathbb{T}_q$ is also a path of bonds of $\mathbb{T}_K$ for every $K \ge q$, and it is in this sense that a loop is fixed while the cutoff is removed.

Definition 1.1 (Loops and Ensembles)**.** Let $q \ge 0$ be an integer.

(i) A *loop at level* q is a closed path C of bonds of $\mathbb{T}_q$ which visits no site twice and whose ℓ^∞-diameter $\operatorname{diam}(C)$, measured along a lift of C to $\mathbb{R}^4$, satisfies $3\operatorname{diam}(C) < T$; its *corners* are the sites at which it changes direction. A loop carries its orientation.

(ii) A *loop ensemble at level* q is a finite family $\mathcal{C} = ((C_1,\chi_1),\dots,(C_n,\chi_n))$, $n \ge 1$, of pairwise vertex-disjoint loops at level q, each with a character $\chi_i = (\dim \pi_i)^{-1}\operatorname{Tr}\pi_i$ of an irreducible unitary representation π_i of G.

The loops need not be planar, and no assumption is made on their relative position beyond disjointness. A loop at level q is also a loop at every level $q' \ge q$. For a gauge field U on $\mathbb{T}_K$, $K \ge q$, the Wilson loop of (C,χ) is $W_\chi(C;U) \coloneqq \chi(U(C))$, which does not depend on the site at which the closed path C is started, and we write

$$W_{\mathcal{C}}(U) \coloneqq \prod_{i=1}^{n} W_{\chi_i}(C_i;U) \tag{1.4}$$

for the observable of an ensemble.

Two quantities measure the shape of a loop, and all our constants depend on the loops only through them and through the perimeters. The first is the chord–arc constant $\Lambda(C) \ge 2$ of Definition 2.4, the largest ratio between the distance of two points along the loop and their ℓ^∞-distance. The second is the ratio $P(C)/r(C)$ of the perimeter to the corner isolation $r(C)$ of Definition 2.2, which is the smallest of half the shortest straight segment, the least distance between two non-adjacent segments, and an eighth of the perimeter. Both are invariant under dilations, and for a square $\Lambda(C) = 2$ and $P(C)/r(C) = 8$. For an ensemble we put $\Lambda(\mathcal{C}) \coloneqq \max_i \Lambda(C_i)$.

Remark 1.2 (The Chord–Arc Constant and Self-Contact). The chord–arc constant measures how close a loop comes to touching itself, relative to the length of the loop between the two places which come close. Near a corner two points are close both in space and along the loop, which is why $\Lambda(C) \geq 2$ for every loop, and the constant becomes large only when two stretches of C which are far apart along the loop come close to each other in space, as the two long sides of a thin rectangle do: for a rectangle with sides $a \geq b$ one finds $\Lambda(C) = \max(2, 1 + a/b)$. Since $\Lambda(C)$ does not change under dilations, the bound $\Lambda(C) \leq \Lambda_*$ keeps a loop away from contact with itself uniformly on all scales, whatever its size. This is the property which our estimates use: any two points of C in a cube of side s are joined by an arc of C of length at most $\Lambda(C)s$, so that the loop cannot leave a cube and return to it except along a short arc, and the number of its bonds which meet the cube is at most $2\Lambda(C)(s+1)$ (Lemma 5.5).

The restriction is also natural from the point of view of renormalisation, although nothing in the proofs depends on what follows. The perimeter factor and one factor for each corner are what perturbation theory predicts for a loop which does not meet itself [18, 14]. A loop which passes twice through the same point is not renormalised by factors of this kind: in perturbation theory its expectation mixes with those of the loops obtained by rerouting the path at the point of self-intersection [14], so that no number attached to the loop alone, of the kind by which we divide, removes its divergences. A lattice loop in the sense of Definition 1.1 never meets itself, but as $\Lambda(C)$ grows it comes closer, relative to its own length, to a loop which does. The constants of our estimates grow with $\Lambda(C)$, and so does the lower bound $L_0(\Lambda_*)$ on the blocking factor below; we do not show that this growth is necessary.

1.3 The Normalisation and the Results

The renormalisation group is run from the lattice $\mathbb{T}_K$ down to a lattice whose spacing does not depend on K, on which the loops of an ensemble at level q are paths of a fixed number of bonds. This lattice has the spacing uL^{-j_q}, where the *stopping index* $j_q \geq q$ of Definition 3.13 depends on q, on $\bar{g}$, and on the constants of the construction, and not on K. The run thus consists of $K' \coloneqq K - j_q$ steps. Each step $k \leq K'$ contributes to a loop C a factor $\kappa_k(C, \chi) \neq 0$, and the *normalisation* of (C, χ) at the cutoff K is the product

$$\kappa_K^{(q)}(C, \chi) \coloneqq \prod_{k=1}^{K-j_q} \kappa_k(C, \chi) \,, \tag{1.5}$$

defined in Definition 7.12. Its structure is the one described at the beginning: $\log \kappa_k(C, \chi)$ is the sum of a term proportional to the total length of the segments of C of each orientation and of a term for each corner of C, with coefficients that are numbers of the step k and of the character and depend neither on the loop nor on the torus.

Definition 1.3 (Renormalised Expectation)**.** For a loop ensemble $\mathcal{C}$ at level q and $K > j_q$,

$$W_K(\mathcal{C}) \coloneqq \frac{\mathbb{E}_K[W_{\mathcal{C}}]}{\prod_{i=1}^n \kappa_K^{(q)}(C_i, \chi_i)} \,. \tag{1.6}$$

Since a loop at level q is also a loop at every level $q' \geq q$, the level is part of the data of an ensemble: it fixes the stopping index j_q and with it the number of step factors in (1.5), while the step factors themselves do not depend on it.

The blocking factor L and the other constants of the construction are fixed first, in the order of Appendix A, and the coupling $\bar{g}$ after them. These choices decide which loops the results cover, and they decide it loop by loop and through the shape alone: a loop is covered

if it is not too irregular for the chosen L, whatever its size and its representation, and then any number of such loops may be combined. The condition on the shape is a bound on the chord–arc constant: there is an increasing function L_0 on $[2,\infty)$, defined in (C.1), and we fix $\Lambda_* \geq 2$ with $L \geq L_0(\Lambda_*)$.

The representations enter in another way. A character is normalised, $|\chi| \leq 1$, but the insertions which a loop produces are built from the derivatives of $\pi(U(C))$, and their size is measured by

$$c_\pi := \sup\{\|\mathrm{d}\pi(X)\| \,|\, X \in \operatorname{Lie} G,\ \|X\| \leq 1\}\ , \tag{1.7}$$

where $\mathrm{d}\pi$ is the differential of π on the Lie algebra $\operatorname{Lie} G \subset \mathfrak{u}(N)$ and $\|\cdot\|$ is the operator norm. This number grows with the highest weight of π: for $G = SU(2)$ and the representation of spin j, $c_\pi = 2j$, and $\chi(\operatorname{diag}(e^{i\phi}, e^{-i\phi})) = \sin((2j+1)\phi)/((2j+1)\sin\phi)$ is close to 1 on fluctuations ϕ of the order of a coupling g only if $(2j+1)g$ is small. Every expansion of the construction in which a loop takes part therefore requires c_π times the running coupling of its step to be small, and at a fixed coupling $\bar{g}$ no single bound on the running couplings achieves this for every π. The loops, however, take part only in the steps $k \leq K'$, which integrate the scales finer than the stopping scale uL^{-j_q}, and at these scales the running coupling is smaller than $\bar{g}$ by asymptotic freedom: Section 3.6 gives $g_k \leq \gamma_q$ for $k \leq K'$, with a number γ_q which does not depend on K and tends to zero as $q \to \infty$, at every $\bar{g}$. A large representation is therefore accommodated by reading its loop at a fine enough level, that is, by stopping the run and the product of the step factors at a fine enough scale, while the coupling $\bar{g}$, the construction, and the step factors stay the same. The estimates see a representation only through c_π, by the bound $\|\mathrm{d}\pi(X)\| \leq c_\pi \|X\|$ and its consequences for the transports $\pi(U(J))$ at complex fields, while the normalised trace of a product is bounded by the product of the operator norms; every constant of the paper which depends on the representations of an ensemble is therefore a nondecreasing function of $c_{\mathcal{C}} := \max_i c_{\pi_i}$. The size of a loop at level q is measured by the number $P_q(C)$ of its bonds in $\mathbb{T}_q$, so that its physical length is $P_q(C)uL^{-q}$; it enters the constants of our estimates and none of their conditions.

Definition 1.4 (Admissible Loops)**.** Let L and Λ_* be fixed. A loop C at level q is *admissible* if

$$\Lambda(C) \leq \Lambda_*\ . \tag{1.8}$$

Admissibility is thus a condition on the shape alone. The size of a loop enters no condition because every estimate of the paper which involves a loop is a sum, over its traversals at some scale, of quantities which are small uniformly in the loop, and only these quantities are ever required to be small: a long loop makes the sums, and with them the constants, large, but it makes none of the summands larger. Our main result is the following.

Theorem 1.5 (Uniform Bound and Limit)**.** *Let L and Λ_* be as above. There are $\gamma_0 > 0$ and a function $q_\sharp \colon [0,\infty) \times (0,\gamma_0] \to \{0,1,2,\dots\}$, nondecreasing in each argument and, for every c, equal to 0 for all sufficiently small $\bar{g}$, which depend only on the constants of the construction and on Λ_*, such that the following holds for every $\bar{g} \in (0,\gamma_0]$. For every loop ensemble $\mathcal{C}$ at a level*

$$q \geq q_\sharp(c_{\mathcal{C}}, \bar{g}) \tag{1.9}$$

whose loops are admissible, whatever its number n of loops and the irreducible representations π_i of its characters, there are an integer K_0 and $B < \infty$ such that

$$\sup_{K \geq K_0} |W_K(\mathcal{C})| \leq B \qquad \textit{and} \qquad \sum_{K \geq K_0} |W_{K+1}(\mathcal{C}) - W_K(\mathcal{C})| < \infty\ . \tag{1.10}$$

In particular the limit

$$W(\mathcal{C}) \coloneqq \lim_{K\to\infty} W_K(\mathcal{C}) \tag{1.11}$$

exists. The numbers K_0 and B depend on the constants of the construction, on $\bar g$, q, and T/u, and on $\mathcal{C}$ through n, the characters, the ratios $P(C_i)/r(C_i)$, and the numbers $P_q(C_i)$.

The level enters the theorem only through the normalisation, and for the continuum limit it is a choice without physical consequence. The loops are fixed paths in the torus, and the level of an ensemble decides only the scale uL^{-j_q} at which the product (1.5) of the step factors is stopped, that is, the scale at which the normalisation is fixed. Writing $W_K^{(q)}$ for the renormalised expectation with the normalisation of the level q, we have, for levels $q' \geq q$ and every $K > j_{q'}$,

$$W_K^{(q')}(\mathcal{C}) = W_K^{(q)}(\mathcal{C}) \prod_{i=1}^{n} \prod_{k=K-j_{q'}+1}^{K-j_q} \kappa_k(C_i, \chi_i) \,, \tag{1.12}$$

by (1.5) and because the step factors do not depend on the level. The product runs over the $j_{q'} - j_q$ steps between the two stopping scales, a number which does not depend on K, and each of its factors is the exponential of a perimeter term and of a term for each corner, with coefficients which do not depend on the loop. A change of the level is therefore a finite renormalisation of the perimeter and corner terms, whose finite parts are a matter of convention in any renormalisation of Wilson loops, and a quantity in which the normalisation cancels does not depend on the level at all: for instance $W_K(\mathcal{C})/\prod_i W_K((C_i,\chi_i))$ is the quotient $\mathbb{E}_K[W_{\mathcal{C}}]/\prod_i \mathbb{E}_K[W_{\chi_i}(C_i;\cdot)]$ whenever the denominators do not vanish. Since an ensemble at a level q_0 is an ensemble at every level $q \geq q_0$, the condition (1.9) restricts only this choice and not the ensembles, and the theorem covers, at every coupling $\bar g \leq \gamma_0$, every ensemble of admissible loops with arbitrary irreducible representations. The level required is bounded in the coupling, $q_\sharp(c,\bar g) \leq q_\sharp(c,\gamma_0)$, and it is given by (11.34). The restriction to irreducible representations concerns the normalisation only: the character of a reducible representation is the convex combination of the characters of its irreducible constituents, weighted by their dimensions, and the theorem applies to each term with its own normalisation.

The requirements on the coupling in Sections 3–12 are of two kinds. Those which involve no representation are upper bounds on $\bar g$ in terms of the constants of the construction and of Λ_*, and γ_0 is the largest number below which all of them hold. Those which involve the representations of an ensemble $\mathcal{C}$ concern only the steps $k \leq K'$ at which its loops are carried, and they are imposed as upper bounds on the number γ_q of (3.46) in terms of $c_{\mathcal{C}}$, of Λ_*, and of the constants of the construction; we then say that γ_q is small, depending on $c_{\mathcal{C}}$. A coupling and a level "as in Theorem 1.5" for $\mathcal{C}$ are a coupling $\bar g \in (0,\gamma_0]$ and a level q at which γ_q satisfies all the requirements of the second kind. The function $q_\sharp$ is computed from these requirements in Section 11.8.

The bound and the limit are proved by different means. The bound follows from an exact identity, Theorem 8.1, which expresses $W_K(\mathcal{C})$ as a convex combination, over the terms of Bałaban's expansion of the density at the stopping scale, of expectations of functions which are bounded uniformly in K. The limit requires a comparison of the two runs with K and $K+1$ steps, which share all their lattices but the finest one, and this comparison, Theorem 11.1, is the longest part of the paper.

The limits inherit the symmetries of the lattice measures, as far as the normalisations share them. The torus has the same period in the four directions, and for $K \geq q$ the lattice $\mathbb{T}_K$ is mapped to itself by every map $h(x) = Ox + t$ of $\mathbb{T}$ in which O is a signed permutation matrix and $t \in uL^{-q}\mathbb{Z}^4$. We call these maps the *symmetries of level* q, and we write $\mathfrak{G}_q$ for

the group of those whose matrix O is diagonal. Thus $\mathfrak{G}_q$ is generated by the translations by vectors of $uL^{-q}\mathbb{Z}^4$ and by the reflections θ_μ, $\mu \in \{0,\dots,3\}$, which replace x_μ by $-x_\mu$, and it contains, for every μ and every a with $2a \in uL^{-q}\mathbb{Z}$, the reflection $\theta_{\mu,a}$ which replaces x_μ by $2a - x_\mu$ modulo T; the symmetries of level q which do not belong to $\mathfrak{G}_q$ are those which exchange two axes. A symmetry h of level q is an isometry of $\mathbb{T}_q$ for the distance $|\cdot|_\infty$ which maps segments to segments and corners to corners, and it therefore maps a loop C at level q to a loop hC at level q, with the image orientation, $\Lambda(hC) = \Lambda(C)$, $P(hC)/r(hC) = P(C)/r(C)$, and $P_q(hC) = P_q(C)$. For an ensemble $\mathcal{C}$ we write $h\mathcal{C} \coloneqq ((hC_1,\chi_1),\dots,(hC_n,\chi_n))$, with the same characters, and $\bar{\mathcal{C}}$ for the ensemble of the loops of $\mathcal{C}$ with the conjugate characters $\bar\chi_i$. The character $\bar\chi$ is that of the conjugate representation $\bar\pi$, whose differential $\mathrm{d}\bar\pi(X)$ is the complex conjugate of the matrix $\mathrm{d}\pi(X)$ and has the same operator norm, so that $c_{\bar\pi} = c_\pi$ and $c_{\bar{\mathcal{C}}} = c_{\mathcal{C}}$. Finally C^{-1} denotes the loop C with the opposite orientation.

Corollary 1.6 (Symmetries of the Limits)**.** *Let L, Λ_*, γ_0, and $q_\sharp$ be as in Theorem 1.5, let $\bar g \in (0,\gamma_0]$, and let $\mathcal{C}$ be a loop ensemble at a level $q \ge q_\sharp(c_{\mathcal{C}},\bar g)$ whose loops are admissible. Then the limit $W(\mathcal{C}')$ exists for each of the ensembles $\mathcal{C}'$ below, and:*

(i) $W(\mathcal{C})$ does not depend on the order of the loops of $\mathcal{C}$;

(ii) $W(g\mathcal{C}) = W(\mathcal{C})$ for every $g \in \mathfrak{G}_q$;

(iii) $W(\bar{\mathcal{C}}) = \overline{W(\mathcal{C})}$, and $W(\mathcal{C})$ does not change if a loop C_i of $\mathcal{C}$ is replaced by C_i^{-1} and its character by $\bar\chi_i$;

(iv) $W(h\mathcal{C}) = W(\mathcal{C})$ for every symmetry h of level q for which there are $g_1,\dots,g_n \in \mathfrak{G}_q$ with $hC_i = g_iC_i$ for every i.

The same identities hold for W_K at every $K > j_q$.

Statement (ii) is the covariance under translations and under the reflections in coordinate hyperplanes, with unchanged characters, and statement (iii) is the invariance under charge conjugation. Since an ensemble at level q is an ensemble at every level $q' \ge q$, statement (ii) gives the covariance under every translation of the dense group $\bigcup_q uL^{-q}\mathbb{Z}^4$, at a level at which it is a symmetry; the normalisations of two levels differ by the factors (1.12), which are the same for an ensemble and its translate. Statement (iv) applies, for instance, to the reflection which exchanges x_μ and x_ν and to an ensemble of squares in planes spanned by e_μ and e_ν, since the image of such a square is also its image under θ_μ followed by a translation. For a general ensemble we do not know whether the limits are covariant under the exchange of two axes: the lattice measures are invariant under it, but the normalisations are not shown to be, for the reason explained in Section 12.3.

The limits also inherit the reflection positivity of the lattice measures, for the reflections $\theta_{\mu,a}$ with $a \in uL^{-q}\mathbb{Z}$, which we call the *reflections of level q*. The points fixed by such a reflection form the two hyperplanes $\{x_\mu \equiv a\}$ and $\{x_\mu \equiv a + T/2\}$, coordinates being taken modulo T, and their complement consists of two open half-tori, $\{a < x_\mu < a + T/2\}$ and $\{a + T/2 < x_\mu < a + T\}$, which the reflection exchanges. The pairing of reflection positivity combines the reflection of a loop with the conjugation of its character, and so involves the ensembles $\theta\bar{\mathcal{D}}$ of the reflected loops θC, with the image orientations and the conjugate characters; we have $c_{\theta\bar{\mathcal{D}}} = c_{\mathcal{D}}$. We admit the empty ensemble, with $W(\emptyset) \coloneqq 1$.

Corollary 1.7 (Reflection Positivity)**.** *Let L, Λ_*, γ_0, and $q_\sharp$ be as in Theorem 1.5, let $\bar g \in (0,\gamma_0]$ and $q \ge 0$, let θ be a reflection of level q, and let H_+ be one of the two open half-tori bounded by the points fixed by θ. Let $\mathcal{D}_1,\dots,\mathcal{D}_m$ be loop ensembles at level q, possibly*

empty, whose loops are admissible and lie in H_+, and assume $q \geq q_\sharp(c, \bar g)$, where c is the largest of the numbers c_π of their representations. Then the limits $W(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b)$ exist for $0 \leq a, b \leq m$, with $\mathcal{D}_0 := \emptyset$, and:

(i) the matrix $(W(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b))_{1\leq a,b\leq m}$ is Hermitian and positive semidefinite;

(ii) $W(\theta\bar{\mathcal{D}}_a) = \overline{W(\mathcal{D}_a)}$ for every a, and the matrix $(W(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b) - W(\mathcal{D}_a)\overline{W(\mathcal{D}_b)})_{1\leq a,b\leq m}$ is Hermitian and positive semidefinite.

The statement (ii) is positivity for the connected functions, and it is not empty: at small coupling the limits do not vanish on the ensembles whose loops are short compared with a length which grows without limit as $\bar g \to 0$ (Proposition 12.5), and for a loop C in H_+ to which this applies the ensemble $\mathcal{D} := ((C, \chi))$ has $W(\mathcal{D} \cup \theta\bar{\mathcal{D}}) \geq |W(\mathcal{D})|^2 \geq 1/4$. In the other direction the limits satisfy a bound which is a product over the loops of factors exponential in their lengths, with a rate which tends to zero with the coupling (Proposition 12.7).

Finally we ask whether the limits are new, that is, whether they could be the Wilson-loop functions of a Gaussian field. The question has to be posed with some care, because the normalisation is a convention: a function which differs from W by a factor $\exp(\sum_v \lambda_v N_v(C) + \sum_c \zeta_{v_c,w_c})$ for each loop, where $N_v(C)$ is the total length of the segments of C of orientation v, the second sum runs over the corners, and the coefficients do not depend on the loop, is the same object with another normalisation. For an abelian Gaussian field the answer is then contained in a single number. If B is a real Gaussian 1-form and g a charge, then $\log|\mathbb{E}[e^{i\mathsf{g}\oint_C \mathsf{B}}]| = -\frac{1}{2}\mathsf{g}^2 \operatorname{Var} \oint_C \mathsf{B}$ is linear in the covariance of B and a quadratic form in the current of the loop, a change of normalisation adds to it a function which is linear in the lengths and in the numbers of corners of each type, and a suitable signed combination of these logarithms over a few rectangles and pairs of rectangles annihilates both, whatever the covariance and whatever the normalisation.

To state this precisely, we fix a level q, a coordinate plane spanned by e_μ and e_ν, lengths $t, s \in uL^{-q}\mathbb{N}$ with $3\max(4t, s) < T$, a point $o \in uL^{-q}\mathbb{Z}^4$, and a nontrivial irreducible unitary representation π of G with the character χ. For $a \in \{0, \dots, 3\}$ and $n \in \{1, \dots, 4\}$ let I_a and R_n be the boundaries of the rectangles $o + [at, (a+1)t]e_\mu + [0, s]e_\nu$ and $o + [0, nt]e_\mu + [0, s]e_\nu$, traversed in the sense which runs from e_μ to e_ν, so that $R_1 = I_0$, and let $R_n^{(1)} := R_n + te_\mu$ for $n \in \{2, 3\}$. The *eleven diagrams* are the ensembles $((C, \chi))$ of one loop, for C one of the nine rectangles $I_0, \dots, I_3$, R_2, R_3, R_4, $R_2^{(1)}$, $R_3^{(1)}$, and the pairs $P_k := ((I_0, \chi), (I_k, \chi))$ for $k \in \{2, 3\}$, whose loops are at distance $(k-1)t \geq t$; we write C also for the ensemble $((C, \chi))$. They are ensembles at level q. For a function W on the eleven diagrams with nonzero values let

$$\mathfrak{Y}(W) := \log\Big|\frac{W(R_1)\,W(R_3)^3}{W(R_2)^3\,W(R_4)}\Big| - \log\Big|\frac{W(P_2)}{W(R_1)^2}\Big| + \log\Big|\frac{W(P_3)}{W(R_1)^2}\Big| \,. \tag{1.13}$$

A *Gaussian charge model* is a triple $(\mathsf{g}, \mathsf{X}, \kappa)$ of a number $\mathsf{g} \in \mathbb{R}$, a real Gaussian vector $\mathsf{X} = (\mathsf{X}_0, \dots, \mathsf{X}_3)$ of any mean and covariance, and a function κ on loops with values in $\mathbb{C} \setminus \{0\}$ such that $\log|\kappa(C)| = \sum_v \lambda_v N_v(C) + \sum_c \zeta_{v_c,w_c}$ with real numbers λ_v and $\zeta_{v,w}$ which do not depend on C. Its function on the eleven diagrams is $W(\mathcal{D}) := \mathbb{E}[e^{i\mathsf{g}\mathsf{X}(\mathcal{D})}] / \prod_{C\in\mathcal{D}} \kappa(C)$, where $\mathsf{X}(\mathcal{D})$ is the sum of the X_a over the indices a for which I_a is one of the rectangles that tile the rectangle $\mathcal{D}$, or one of the two loops of the pair $\mathcal{D}$. The models with $\mathsf{X}_a = \oint_{I_a} \mathsf{B}$ for a real Gaussian 1-form B, on a lattice which contains the rectangles or on the continuum torus, are of this kind, with $\mathsf{X}(C) = \oint_C \mathsf{B}$ for each of the nine rectangles, since the interior sides of a tiling are traversed once in each direction.

Theorem 1.8 (The Limits Are Not Gaussian)**.** *Let q, the plane, t, s, o, and π be as above, let $\Lambda_* \geq 1 + \max(s/t, 4t/s)$, and let L be as in Theorem 1.5. There is $\bar{g}_0 > 0$, depending on q, t/u, s/u, T/u, π, and the constants of the construction, such that for every $\bar{g} \in (0, \bar{g}_0]$ the following hold.*

(i) The limits $W(\mathcal{D})$ exist on the eleven diagrams, $|W(\mathcal{D})| \geq 1/2$, $W(I_a) = W(I_0)$, and $W(R_n^{(1)}) = W(R_n)$.

(ii) $\mathfrak{Y}(W) \geq \frac{1}{2} c_O \bar{g}^2$, with $c_O \coloneqq C_2(\pi) s^2 t^3 / (672 e^2 T^5)$ and the Casimir number $C_2(\pi) > 0$ of (13.45)*, and $\mathfrak{Y}(W)$ is the same for the normalisations of every level $q' \geq q$.*

(iii) No Gaussian charge model has the function W on the eleven diagrams, and W is not the limit on the eleven diagrams of the functions of a sequence of Gaussian charge models.

The number $\mathfrak{Y}$ is a third difference in the width of the rectangles, corrected by the two *pair terms* $\log|W(P_k)/W(R_1)^2|$. In an abelian Gaussian model all three are of the order $\mathfrak{g}^2$, and they cancel: the covariances of separated rectangles which make up the third difference also make up the pair terms. In the limits of Yang–Mills theory the pair terms are of the order $\bar{g}^4$, up to logarithms, because the term of first order of a non-abelian Wilson loop has trace zero, while the third difference is, at the order $\bar{g}^2$, the number $C_2(\pi)$ times that of the free field, which is positive. The same mechanism makes the theorem silent on Gaussian fields with values in the Lie algebra and path-ordered holonomies, which agree with Yang–Mills theory at the second order (Section 13.2).

1.4 The Argument

Let us describe how the proof is organised and where its difficulties lie. Bałaban's construction is a sequence of steps, each of which integrates the fluctuations of the gauge field at one scale. The integrand is first decomposed by test functions, which decide in every region of the lattice whether the field is small there, and the outcome is a sum of terms, labelled by the regions in which some test function has failed. On the regions where all of them pass, the fluctuation integral is performed by a cluster expansion around a background field, the minimiser of the action with given block averages. On the other regions no expansion is available, and the term carries instead a small factor from the action, which a separate operation, Bałaban's **R**-operation [11, 12], later turns into terms of the effective action.

The first difficulty is one of consistency and is independent of the loops. The thresholds of the test functions are functions of the running couplings, while the running couplings are defined by the construction, and it is not known whether the coupling of Bałaban's construction as printed is a continuous function of the bare coupling, which is what one needs in order to solve $g_K = \bar{g}$. We therefore run a variant of the construction, in which every threshold, every cube size, and the depth of the **R**-operation are functions of the target $(K, \bar{g})$ alone, and in which the plaquette test functions are continuous rather than characteristic functions. Section 3 defines this variant and shows that Bałaban's estimates hold for it, and Section 4 proves the continuity of the couplings and solves for the bare coupling.

The loops enter in Section 5, which treats the first step, and in Section 6, which treats the general one. After k steps the contribution of an ensemble to a term of the density is a *dressed loop*: the product of the normalisations $\kappa_1 \cdots \kappa_k$ of its loops, of the exponential of a small local functional, and of the sum of the Wilson loops of the background field and of a series of insertions along the loops, with activities that decay in the size of their supports. The heart of the paper is the statement that this form is reproduced by a step, with constants that

do not depend on k, including on the terms with large fields and through the **R**-operation. The perimeter and corner coefficients are extracted in Section 7 from the part of the loop's contribution that survives at the trivial background, and it is here that the geometry of a general loop is used: away from its corners a loop cannot be told from a straight line, and near a corner it cannot be told from two perpendicular rays, up to errors that decay exponentially in the corner isolation.

Stopping the induction at the index K' gives the identity and the bound of Section 8. For the limit we have to compare the dressed loops, and the densities themselves, of the runs with K and $K+1$ steps on their common lattices. In the regions of small fields this is a comparison of two expansions whose data differ by the one additional step, and the difference decays geometrically in the number of steps that separate a scale from the finest one. The terms with large fields require a different argument, since their test functions are evaluated on different fields in the two runs, and it is for this comparison that the test functions were made continuous. Sections 10 and 9 prove the two inputs of the comparison, and Section 11 assembles them. Section 12 derives the two corollaries, the lower bound on the limits, and the growth bound, Section 13 proves Theorem 1.8, and Section 14 records where our reading of Bałaban's papers departs from the printed text.

The results of Bałaban's which we use in the setting in which he proved them are cited by page and equation. Those which we use outside his hypotheses, or of which he printed no proof, are proved again in a supplement to this paper, cited as "Supplement, Proposition S.n".

2 Loops and Their Geometry

All statements of this section are about closed paths on the lattice $\mathbb{Z}^4$ with the distance $|x-y|_\infty := \max_\mu |x_\mu - y_\mu|$. They will be applied at every scale of the construction, with the lattice unit equal to the spacing of that scale, and their purpose is to replace the elementary facts about a square that such an argument uses without comment: that a short piece of it is straight or contains one corner, that few of its bonds lie in a small box, and that a piece containing half of it is large.

2.1 Segments and Corners

Let $\mathcal{V} := \{\pm e_0, \pm e_1, \pm e_2, \pm e_3\}$ be the set of the eight orientations. A *closed path* of length n is a sequence of sites $x_0, x_1, \dots, x_n = x_0$ of $\mathbb{Z}^4$ with $v_i := x_{i+1} - x_i \in \mathcal{V}$ for $0 \le i < n$, and it is *simple* if $x_0, \dots, x_{n-1}$ are distinct. We read the indices modulo n. A *loop* is a simple closed path, and two loops which differ by a cyclic shift of the indices are the same loop. The bonds of the loop C are the pairs $b_i = (x_i, x_{i+1})$, the *position* of b_i is its initial site x_i, and $|C| \subset \mathbb{R}^4$ is the union of the closed unit intervals $[x_i, x_{i+1}]$. The number $P(C) := n$ of bonds is the *perimeter*.

A site x_i of C is a *corner* if $v_i \neq v_{i-1}$, and $\mathcal{K}(C)$ is the set of the corners. A *segment* of C is a maximal run of consecutive bonds of one orientation, together with its two end sites, so that the segments are the pieces of C between consecutive corners. At a corner $c = x_i$ we write $v_c := v_{i-1}$ and $w_c := v_i$ for the incoming and the outgoing orientation.

Lemma 2.1 (Corners)**.** *Let C be a loop.*

(i) For every $v \in \mathcal{V}$, C has as many bonds of orientation v as of orientation $-v$. In particular $P(C)$ is even.

(ii) At every corner c the orientations v_c and w_c are perpendicular.

(iii) C has at least four segments, and as many corners as segments.

(iv) $\mathrm{diam}(C) \le P(C)/2$, *where* $\mathrm{diam}(C)$ *is the* ℓ^∞*-diameter of* $|C|$.

Proof. For (i) we sum the orientations around the loop. Since $\sum_{i<n} v_i = x_n - x_0 = 0$, the coefficient of each e_μ in this sum vanishes, and this coefficient is the number of bonds of orientation e_μ minus the number of bonds of orientation $-e_\mu$.

For (ii), $w_c \neq v_c$ by the definition of a corner. If $w_c = -v_c$ at $c = x_i$, then $x_{i+1} = x_i - v_{i-1} = x_{i-1}$, so that a site is visited twice, which is excluded because $n \ge 3$ for a simple closed path with $v_i \in \mathcal{V}$. Two elements of $\mathcal{V}$ which are neither equal nor opposite are perpendicular.

For (iii), the orientations v_i are not all equal, since otherwise $x_n \neq x_0$. So C has a corner, and by (ii) at least two coordinate axes carry bonds of C. By (i) each of these axes carries bonds of both of its orientations, so that at least four orientations occur, and each of them is the orientation of at least one segment. The segments and the corners alternate around the loop, which gives the last assertion.

For (iv), let R_μ be the difference between the largest and the smallest value of the coordinate x_μ on C. The path passes from a site with the smallest value to a site with the largest value and back, and each bond changes x_μ by at most one, so that $P(C) \ge 2R_\mu$. Since $\mathrm{diam}(C) = \max_\mu R_\mu$, this is (iv). *q.e.d.*

Two segments are *adjacent* if they are consecutive around the loop, that is, if they share a corner. By Lemma 2.1(iii) two distinct segments share at most one corner, and every segment has a segment which is not adjacent to it.

Definition 2.2 (Lengths)**.** For a loop C, the *shortest segment* $s(C)$ is the least number of bonds of a segment of C, the *separation* $\delta(C)$ is the least ℓ^∞-distance between two distinct non-adjacent segments of C, as closed subsets of $\mathbb{R}^4$, and the *corner isolation* is

$$r(C) := \min\Big(\frac{s(C)}{2},\, \delta(C),\, \frac{P(C)}{8}\Big)\,. \tag{2.1}$$

The separation is positive, and it is at least as large as the mesh of any lattice which contains the corners. This is the content of the next lemma, which also separates the loops of an ensemble from each other.

Lemma 2.3 (Separation)**.** *Let $a \ge 1$ be an integer and $x_* \in \mathbb{Z}^4$.*

(i) If all corners of a loop C lie in $x_* + a\mathbb{Z}^4$, then every segment of C has a number of bonds which is a positive multiple of a, and $s(C) \ge a$ and $\delta(C) \ge a$.

(ii) If C and C' are loops without a common site, all of whose corners lie in $x_* + a\mathbb{Z}^4$, then the ℓ^∞-distance between $|C|$ and $|C'|$ is at least a.

Proof. We may take $x_* = 0$. A segment joins two corners which differ in one coordinate only, by a nonzero multiple of a, which gives the first assertion of (i) and $s(C) \ge a$.

Let now A and B be two closed axis-parallel intervals of $\mathbb{R}^4$ with end points in $a\mathbb{Z}^4$, A parallel to e_μ and B parallel to e_ν. We claim that, if $A \cap B = \emptyset$, then their ℓ^∞-distance is at

least a. On each of the two intervals all coordinates but one are constant and lie in $a\mathbb{Z}$. If a coordinate x_λ with $\lambda \notin \{\mu, \nu\}$ takes different values on A and on B, these values differ by at least a, and the claim holds. Assume that this is not the case. If $\mu = \nu$, then A and B lie on one line, where they are disjoint intervals with end points in $a\mathbb{Z}$, and so they are at least a apart. If $\mu \neq \nu$, let b_μ be the value of x_μ on B and a_ν the value of x_ν on A. The intervals meet if and only if b_μ lies in the range of x_μ on A and a_ν lies in the range of x_ν on B. Since they do not meet, one of these fails, say the first, and then $b_\mu \in a\mathbb{Z}$ lies at distance at least a from that range, whose end points are in $a\mathbb{Z}$. So $|x_\mu - y_\mu| \geq a$ for all $x \in A$ and $y \in B$, which proves the claim.

It remains to see that the claim applies. If two axis-parallel intervals with end points in $\mathbb{Z}^4$ meet, then they have a site of $\mathbb{Z}^4$ in common: on a common line two closed intervals with integer end points which meet share an integer point, and two perpendicular ones meet in the point with the coordinates b_μ and a_ν above, which is a site. A site which lies on two segments of the loop C is a site x_i of C, and, since C is simple, it lies on two segments only if it is the corner between them. Two non-adjacent segments of C are therefore disjoint, and the claim gives $\delta(C) \geq a$. In (ii) a segment of C and a segment of C' have no site in common by hypothesis, so they are disjoint, and the claim gives (ii). *q.e.d.*

2.2 The Chord–Arc Constant

For two points $x, y \in |C|$ let $d_C(x, y)$ be the length of the shorter of the two arcs of $|C|$ which join them. Along an arc every coordinate changes by at most the length of the arc, so that

$$|x - y|_\infty \leq d_C(x, y) \leq \frac{P(C)}{2} \,. \tag{2.2}$$

The constant which bounds d_C from above by the ℓ^∞-distance is the one geometric quantity through which the shape of a loop enters most of our counts.

Definition 2.4 (Chord–Arc Constant)**.** For a loop C,

$$\Lambda(C) := \sup\Big\{ \frac{d_C(x, y)}{|x - y|_\infty} \,\Big|\, x, y \in |C|,\ x \neq y \Big\} \,. \tag{2.3}$$

Lemma 2.5 (Bounds on the Chord–Arc Constant)**.** *For every loop C,*

$$2 \leq \Lambda(C) \leq \max\Big(2,\, \frac{P(C)}{2\delta(C)}\Big) \leq \max\Big(2,\, \frac{P(C)}{2r(C)}\Big) \,, \qquad |\mathcal{K}(C)| \leq \frac{P(C)}{s(C)} \leq \frac{P(C)}{2r(C)} \,. \tag{2.4}$$

For a square, $\Lambda(C) = 2$ and $P(C)/r(C) = 8$.

Proof. For the lower bound let c be a corner, and let x and y be the sites of C before and after c. They differ by one in each of two coordinates, by Lemma 2.1(ii), so that $|x - y|_\infty = 1$, while the arc through c has length 2 and the other arc has length $P(C) - 2 \geq 2$, because $P(C) \geq 4$ by Lemma 2.1(iii). Hence $d_C(x, y) = 2$.

For the upper bound let $x \neq y$ be points of $|C|$. If they lie on one segment, the part of the segment between them is an arc of length $|x - y|_\infty$. If they lie on two adjacent segments, which meet at the corner c, let α and β be their distances from c. The two segments are perpendicular, so that $|x - y|_\infty = \max(\alpha, \beta)$, and the arc through c has length $\alpha + \beta \leq 2\max(\alpha, \beta)$. In the remaining case they lie on two non-adjacent segments, so that $|x - y|_\infty \geq \delta(C)$, while $d_C(x, y) \leq P(C)/2$ by (2.2). The third inequality holds because

$\delta(C) \geq r(C)$. Every segment has at least $s(C) \geq 2r(C)$ bonds, and there are as many segments as corners, which gives the bound on $|\mathcal{K}(C)|$.

For a square of side s we have $\delta(C) = s$ and $P(C) = 4s$, so that $\Lambda(C) = 2$ by the two bounds, and $r(C) = \min(s/2, s, s/2) = s/2$. *q.e.d.*

All the quantities of Definition 2.2 and the perimeter are multiplied by m when C is replaced by its image mC under the dilation $x \mapsto mx$, $m \geq 1$ an integer, and so are d_C and the ℓ^∞-distance. Hence $\Lambda(mC) = \Lambda(C)$ and $P(mC)/r(mC) = P(C)/r(C)$: the two shape constants of a loop do not depend on the scale at which it is read.

2.3 Arcs around a Set

The dressed loops of the later sections are sums over finite sets of positions on a loop, and what one needs to know about such a set is how much of the loop it takes to connect it. On a square, a set of positions of diameter $D < s/2$ lies in an arc of length at most $4D$. The next lemma is the form of this statement for a general loop. By an *arc* we mean a connected closed subset of $|C|$, and a *gap* of a set $S \subset |C|$ is a connected component of $|C| \setminus S$.

Lemma 2.6 (Arcs around a Set)**.** *Let C be a loop, S a nonempty set of sites of C of ℓ^∞-diameter D, and $x_0 \in S$. Put $\Lambda := \Lambda(C)$ and $A := \{x \in |C| \,|\, d_C(x_0, x) \leq \Lambda D\}$.*

(i) $S \subset A$, and S has at most $2\Lambda D + 1$ elements.

(ii) If $4\Lambda D < P(C)$, then S has exactly one gap of length greater than $P(C)/2$. Its complement $I(S)$ is the unique shortest arc which contains S. It lies in A, its length is at most $2\Lambda D$, and each of its points is within ℓ^∞-distance ΛD of x_0.

(iii) If $4\Lambda D \geq P(C)$, then every point of $|C|$ is within ℓ^∞-distance $2\Lambda D$ of every point of S.

Proof. For $x \in S$ the definition of Λ gives $d_C(x_0, x) \leq \Lambda |x - x_0|_\infty \leq \Lambda D$, so that $S \subset A$. The sites of C in A are those at arc distance at most ΛD from x_0 on either side, at most $2\lfloor \Lambda D \rfloor + 1$ in number if $2\Lambda D < P(C)$, and otherwise all the $P(C) \leq 2\Lambda D$ sites of C. This is (i).

For (ii), A is the arc of length $2\Lambda D < P(C)/2$ centred at x_0, and its complement is an open arc G_0 of length greater than $P(C)/2$ which contains no point of S by (i). Let G be the gap of S which contains G_0. The gaps of S are pairwise disjoint, so that every other gap has length less than $P(C)/2$, and G is the only gap longer than $P(C)/2$. An arc which contains S and is not all of $|C|$ is the complement of an open arc inside a gap, and it is shortest when that open arc is the longest gap. So $I(S) := |C| \setminus G$ is the unique shortest arc containing S, and $I(S) \subset |C| \setminus G_0 = A$. The last assertion follows from (2.2).

For (iii), two points of $|C|$ are at ℓ^∞-distance at most $\mathrm{diam}(C) \leq P(C)/2$ by Lemma 2.1(iv), and $P(C)/2 \leq 2\Lambda D$ by hypothesis. *q.e.d.*

In both cases of the lemma the shortest arc containing S, taken to be all of $|C|$ in the case (iii), has length at most $4\Lambda D$ and lies within ℓ^∞-distance $2\Lambda D$ of S. For a square these are the constants $8D$ and $4D$.

A cube which is smaller than the separation sees at most one corner of the loop, and the counts become those of a bent line.

Lemma 2.7 (Small Cubes)**.** *Let C be a loop and Q a closed cube of side $t < \delta(C)$ with corners in $\mathbb{Z}^4$. Then $Q \cap |C|$ is empty or an arc contained in one segment or in two adjacent segments. It contains at most $2t+1$ sites of C, and at most $2t+2$ bonds of C meet Q.*

Proof. Two segments which meet Q are within ℓ^∞-distance $t < \delta(C)$ of each other, so they are equal or adjacent. Three distinct segments cannot be pairwise adjacent, because C has at least four segments and adjacency is the cyclic order, so that at most two segments meet Q. A segment meets the convex set Q in an interval, which has length at most t and hence at most $t+1$ sites. If two segments meet Q, in points x and y, let c be their common corner, and let the segments be parallel to e_μ and e_ν. Then c has the coordinate x_ν of x, the coordinate y_μ of y, and the other coordinates in common with both, so that c lies in the coordinate box spanned by x and y, which is contained in the cube Q. Both intervals thus contain c, their union is an arc, and it has at most $2(t+1)-1$ sites. Finally, a bond which meets Q has an end site in Q, since Q has integer corners, and an arc with m sites of C is met by at most $m+1$ bonds of C. *q.e.d.*

The next two statements are used for the pieces of a loop which are long. The first bounds the number of cubes of a partition which a piece meets, and the second shows that a piece which contains half of the loop cannot be small.

Lemma 2.8 (Cubes along an Arc)**.** *Let C be a loop, $M' \geq 1$ an integer, and consider the closed cubes $M'z + [0, M']^4$, $z \in \mathbb{Z}^4$. An arc of n bonds of C consists of at most $n/s(C)+2$ straight pieces, and it meets at most*

$$\frac{8n}{M'} + \frac{16n}{s(C)} + 32 \tag{2.5}$$

of these cubes.

Proof. The straight pieces of an arc are, in the order of the arc, a part of a segment, a number of whole segments, and a part of a segment. The whole segments have at least $s(C)$ bonds each, so there are at most $n/s(C)$ of them. A straight piece of n_i bonds parallel to e_μ occupies an interval $[\tau, \tau+n_i]$ of the coordinate x_μ, which meets the interval $[M'j, M'(j+1)]$ only for the integers j with $\tau/M' - 1 \leq j \leq (\tau+n_i)/M'$, of which there are at most $n_i/M'+2$. Each of its three other coordinates is constant and lies in at most two of the intervals $[M'j, M'(j+1)]$. The piece therefore meets at most $8(n_i/M'+2)$ cubes, and the sum over the at most $n/s(C)+2$ pieces, with $\sum_i n_i = n$, is at most (2.5). *q.e.d.*

Lemma 2.9 (Long Arcs)**.** *Let C be a loop. The set of the positions of at least $P(C)/2$ consecutive bonds of C, together with the final site of the last of these bonds, has ℓ^∞-diameter at least $P(C)/(2\Lambda(C))$.*

Proof. By Lemma 2.1(i) the number $P(C)/2$ is an integer, so that the set contains two sites x and y which are $P(C)/2$ bonds apart along the loop. Both arcs between them have length $P(C)/2$, hence $d_C(x,y) = P(C)/2$, and $|x-y|_\infty \geq d_C(x,y)/\Lambda(C)$ by the definition of $\Lambda(C)$. *q.e.d.*

2.4 The Neighbourhood of a Corner

The coefficients of the normalisation are defined on two model paths: the bi-infinite straight line, and the bent path which consists of two perpendicular rays. We now make precise the

sense in which a loop coincides with one of them near each of its bonds. For a corner c of C let $S_{\text{in}}(c)$ and $S_{\text{out}}(c)$ be the segments which end and begin at c, and let the *bent path* at c be

$$B_c \coloneqq \{c - tv_c \,|\, t \in \mathbb{Z},\ t \geq 0\} \cup \{c + tw_c \,|\, t \in \mathbb{Z},\ t \geq 0\}\ , \tag{2.6}$$

traversed from the first ray to the second. It contains the sites of $S_{\text{in}}(c)$ and of $S_{\text{out}}(c)$. We distribute the bonds of C among its corners: of a segment of ℓ bonds, the first $\lceil \ell/2 \rceil$ bonds are *assigned* to its initial corner and the last $\lfloor \ell/2 \rfloor$ bonds to its final corner. Every bond of C is assigned to exactly one corner.

Lemma 2.10 (Corner Isolation)**.** *Let C be a loop, c a corner of C, and b a bond of C assigned to c, with position x. Every site of C or of B_c which lies neither on $S_{\text{in}}(c)$ nor on $S_{\text{out}}(c)$ is at ℓ^∞-distance at least $\min(s(C)/2, \delta(C)) \geq r(C)$ from x.*

Proof. Let ℓ and ℓ' be the numbers of bonds of $S_{\text{in}}(c)$ and $S_{\text{out}}(c)$, let c' be the initial corner of $S_{\text{in}}(c)$ and c'' the final corner of $S_{\text{out}}(c)$, and write $v \coloneqq v_c$ and $w \coloneqq w_c$. Since C has at least four segments, the segment before $S_{\text{in}}(c)$ is not adjacent to $S_{\text{out}}(c)$, and the segment after $S_{\text{out}}(c)$ is not adjacent to $S_{\text{in}}(c)$.

Assume first that b is one of the last $\lfloor \ell/2 \rfloor$ bonds of $S_{\text{in}}(c)$, so that $x = c' + \alpha v$ with $\alpha \geq \ell - \lfloor \ell/2 \rfloor = \lceil \ell/2 \rceil$. The segment before $S_{\text{in}}(c)$ is perpendicular to v and contains c', so that all its sites have the same coordinate along v as c', which differs from that of x by $\alpha \geq s(C)/2$. All other segments of C apart from $S_{\text{out}}(c)$ are not adjacent to $S_{\text{in}}(c)$, and their sites are at distance at least $\delta(C)$ from x. The sites $c' - tv$, $t > 0$, of the first ray of B_c beyond c' differ from x by $\alpha + t$ in the coordinate along v. The sites $c + tw$, $t > \ell'$, of the second ray beyond c'' differ from x by $t > \ell' \geq s(C)$ in the coordinate along w, which is constant on $S_{\text{in}}(c)$.

Assume now that b is one of the first $\lceil \ell'/2 \rceil$ bonds of $S_{\text{out}}(c)$, so that $x = c + iw$ with $0 \leq i \leq \lceil \ell'/2 \rceil - 1$. The segment after $S_{\text{out}}(c)$ is perpendicular to w and contains $c'' = c + \ell' w$, so that its sites differ from x by $\ell' - i \geq \lfloor \ell'/2 \rfloor + 1 > s(C)/2$ in the coordinate along w, and the sites $c + tw$, $t > \ell'$, of B_c differ from it by more. All other segments of C apart from $S_{\text{in}}(c)$ are not adjacent to $S_{\text{out}}(c)$, which gives the distance $\delta(C)$. The sites $c - tv$, $t > \ell$, of the first ray beyond c' differ from x by $t > \ell \geq s(C)$ in the coordinate along v, which is constant on $S_{\text{out}}(c)$. *q.e.d.*

2.5 A Summation over the Scales

The lengths of a loop grow by the factor L from each scale to the next finer one, and the errors of the corner analysis decay exponentially in these lengths. Their sum over the scales is controlled by the following elementary bound, which we record here because it is used for every such sum.

Lemma 2.11 (Sums over Scales)**.** *For $x > 0$, $c > 0$, and a real number $L \geq 2$,*

$$\sum_{m \geq 0} x L^m e^{-cxL^m} \leq x e^{-cx}\Big(1 + \frac{2}{cx}\Big) \qquad \text{and} \qquad \sum_{m \geq 0} e^{-cxL^m} \leq e^{-cx}\Big(1 + \frac{2}{cx}\Big)\ . \tag{2.7}$$

Proof. Put $y_m \coloneqq xL^m$. For $m \geq 1$ we have $y_m - y_{m-1} = y_m(1 - L^{-1}) \geq y_m/2$, and $e^{-ct} \geq e^{-cy_m}$ on the interval $[y_{m-1}, y_m]$, so that

$$y_m e^{-cy_m} \leq 2(y_m - y_{m-1}) e^{-cy_m} \leq 2 \int_{y_{m-1}}^{y_m} e^{-ct}\, \mathrm{d}t\ . \tag{2.8}$$

The intervals $[y_{m-1}, y_m]$, $m \geq 1$, are disjoint up to their end points and lie in $[x, \infty)$, so that the terms with $m \geq 1$ sum to at most $2\int_x^\infty e^{-ct}\,\mathrm{d}t = (2/c)e^{-cx}$. Adding the term xe^{-cx} of $m = 0$ gives the first bound. The second follows from the first, because $e^{-cy_m} \leq (y_m/x)e^{-cy_m}$ for $y_m \geq x$. *q.e.d.*

3 The Construction

This section defines the renormalisation group that the paper runs: Bałaban's steps, with every threshold, cube size, and depth fixed by the target $(K, \bar{g})$, and with continuous plaquette test functions. It shows that Bałaban's estimates hold for this construction, and it fixes the stopping index of an ensemble.

3.1 The Tube of a Target

Bałaban's construction is governed by a small number of parameters, each of which he defines as a function of the running coupling of its scale. The plaquette threshold of the scale j is

$$\varepsilon_j = g_j p_0(g_j)\,, \qquad p_0(g) \coloneqq A_0(\log g^{-2})^{p_0}\,, \tag{3.1}$$

and the size of the cubes of that scale is MR_j, where R_j is the smallest power of L with $R_j \geq (\log g_j^{-2})^r$ [9, (2.4), (2.5), p. 255]. Here A_0 is a large constant, $r \geq 2$ and $p_0 \geq 5r$ are exponents [9, pp. 245–246], and we keep his notation, in which p_0 denotes both the function and its exponent. The running couplings, in turn, are produced by the construction: g_{j+1} is computed from the effective action which the first j steps have generated. As long as one studies a given bare coupling this causes no difficulty. We, however, have to find the bare coupling for which $g_K = \bar{g}$, and for this we need the map $g_0 \mapsto g_K$ to be continuous, which is not clear when a cube size, an integer, is a function of g_j.

Bałaban mentions a way out. In the first step he notes that there are "two convenient choices" of the threshold, the second of which replaces the coupling inside the logarithm by an upper bound which is known in advance, and that this choice is "more convenient in some nonperturbative aspects of the procedure, but it is technically more difficult", and he assumes the first [9, p. 246]. We take this second route, in a stronger form: every parameter of the scale j is evaluated at a number $\hat{g}_j$ which is fixed by the target and which the running coupling g_j is shown, afterwards, to follow within a constant factor.

Throughout, $\beta_0 \in (0, \sqrt{2} - 1]$ is a fixed number, the constant of Bałaban's inequalities (2.6), and $0 < b_* \leq b^*$ are fixed numbers, which Section 4 takes to be multiples of the first coefficient of the β-function.

Definition 3.1 (Tubes)**.** Let $\gamma > 0$. A *tube* consists of an integer $K \geq 1$ and real numbers b, b', C_W, and $\bar{g} > 0$ such that, with $t \coloneqq \bar{g}^{-2}$,

$$b_* \leq b \leq b' \leq \min(\tfrac{3}{2}b,\, b^*)\,, \qquad C_W \geq 0\,, \qquad t \geq \max(\gamma^{-2} + C_W,\, 4C_W/\beta_0)\,. \tag{3.2}$$

For every integer $k \leq K$ we put

$$\ell_k \coloneqq t + b(K-k) - C_W\,, \qquad u_k \coloneqq t + b'(K-k) + C_W\,, \qquad \hat{g}_k \coloneqq u_k^{-1/2}\,, \tag{3.3}$$

and for $0 \leq k \leq K$ we let H_k be the set of the $(x_0, \ldots, x_k) \in \mathbb{R}^{k+1}$ with $\ell_j \leq x_j \leq u_j$ for $0 \leq j \leq k$ and

$$b(n-m) - C_W \leq x_m - x_n \leq b'(n-m) + C_W \tag{3.4}$$

for $0 \le m < n \le k$. A point of H_k is a *history* of the tube, with the couplings $g_j := x_j^{-1/2}$. The *ratio* of the tube is

$$\rho := \max\Big(\frac{b'}{b},\, \frac{t+C_W}{t-2C_W}\Big)\,. \tag{3.5}$$

The set H_k is closed and convex, being defined by finitely many weak linear inequalities. Since $\beta_0 < 1/2$, the last condition of (3.2) gives $C_W \le \beta_0 t/4 < t/8$, so that $(t+C_W)/(t-2C_W) < (9/8)/(6/8)$ and therefore

$$1 \le \rho \le \tfrac{3}{2}\,. \tag{3.6}$$

The numbers $\hat g_k$ increase with k, and $\hat g_K \le \bar g$. The role of x_j is that of the inverse square of the running coupling of the scale j, and the tube is the set in which we shall confine it: (3.4) says that the inverse square of the coupling decreases, from each scale to a coarser one, by an amount between b and b' per step, up to the additive constant C_W.

The lemmata of this subsection require γ to be small. With $p^* \ge 1$ an upper bound for the exponents r, p_0, p_1, q_0, q_1 of the construction, the conditions are

$$\begin{gathered} b^*\gamma^2 \le \frac{\beta_0}{2}\,, \qquad \log\gamma^{-2} \ge \frac{p^*}{\beta_0}\,, \qquad \Big(1+\frac{\log(3/2)}{\log\gamma^{-2}}\Big)^{p^*} \le 2\,, \\ \Big(1+\frac{b^*\gamma^2}{\log\gamma^{-2}}\Big)^{r} < L\,, \qquad \Big(1+\frac{\log(1+b^*)}{\log\gamma^{-2}}\Big)^{r} \le 1+\frac{1}{L}\,. \end{gathered} \tag{3.7}$$

Each of them holds for all sufficiently small γ, once β_0, b^*, p^*, r, and L are fixed.

Lemma 3.2 (Couplings of a History)**.** *Let γ satisfy* (3.7)*, and let $(x_0, \dots, x_k)$ be a history of a tube, with the couplings $g_j = x_j^{-1/2}$. Put $\beta' := b' + C_W$.*

(i) For $0 \le j \le k$,

$$\hat g_j \le g_j \le \rho^{1/2}\hat g_j\,, \qquad g_j \le \gamma\,, \qquad \log g_j^{-2} \le \log \hat g_j^{-2} \le \log g_j^{-2} + \log\rho\,. \tag{3.8}$$

(ii) For $0 \le m < n \le k$,

$$g_n \le (1+g_n^2\beta'(n-m))^{1/2} g_m \le (1+\beta_0)(n-m)^{1/2} g_m \qquad \text{and} \qquad g_m \le (1+\beta_0) g_n\,, \tag{3.9}$$

and, for every real $p \in (0, p^*]$,

$$(\log g_n^{-2})^p \le (1+\beta_0)(\log g_m^{-2})^p \qquad \text{and} \qquad (\log g_m^{-2})^p \le (1+g_n^2\beta'(n-m))^{\beta_0}(\log g_n^{-2})^p\,. \tag{3.10}$$

(iii) The inequalities (3.9) *and* (3.10) *hold for the numbers $\hat g_j$ in place of the g_j, with b' in place of β' and for all integers $m < n \le K$, and moreover $\hat g_m \le \hat g_n$.*

The inequalities (3.9) and (3.10) are Bałaban's (2.6) and (2.7) [9, p. 255], which he states for the couplings of his construction and for positive integers p. The lemma thus says that a history of a tube, and also the sequence $(\hat g_j)$ itself, satisfy the hypotheses which he places on a sequence of couplings.

Proof. (i) Since $x_j \le u_j$ we have $\hat g_j \le g_j$. With $s := K - j \ge 0$,

$$\frac{u_j}{x_j} \le \frac{u_j}{\ell_j} = \frac{(t+C_W)+b's}{(t-C_W)+bs} \le \max\Big(\frac{t+C_W}{t-C_W},\, \frac{b'}{b}\Big) \le \rho\,, \tag{3.11}$$

where the first step uses $x_j \geq \ell_j > 0$, the second that a quotient $(A + A')/(B + B')$ of sums of positive numbers lies between A/B and A'/B', and the third the definition (3.5). Hence $g_j^2 = x_j^{-1} \leq \rho u_j^{-1} = \rho \hat{g}_j^2$. Moreover $x_j \geq \ell_j \geq t - C_W \geq \gamma^{-2}$ by (3.2), so that $g_j \leq \gamma$, and the last chain of (3.8) is the logarithm of the first.

(ii) The upper bound of (3.4) and $n - m \geq 1$ give $x_m \leq x_n + \beta'(n - m)$, that is, $g_m^{-2} \leq g_n^{-2}(1 + g_n^2 \beta'(n - m))$, which is the first inequality of (3.9). For the second we use $1+y(n-m) \leq (1+y)(n-m)$ for $y \geq 0$, and bound $y = g_n^2\beta'$: by (i) and (3.2), $g_n^2 \leq (t-C_W)^{-1}$ with $C_W \leq \beta_0 t/4$ and $t \geq \gamma^{-2}$, so that

$$g_n^2 \beta' \leq \frac{b' + C_W}{t - C_W} \leq \frac{b^* \gamma^2 + \beta_0/4}{1 - \beta_0/4} \leq \frac{3\beta_0/4}{1 - \beta_0/4} \leq \beta_0 \,, \tag{3.12}$$

by the first condition of (3.7) and $\beta_0 \leq 1$. Thus $(1 + g_n^2\beta')^{1/2} \leq 1 + \beta_0$. For the third inequality, the lower bound of (3.4) gives $x_n \leq x_m + C_W$, and $C_W \leq \beta_0 t/4 \leq (\beta_0/3)x_m$, because $x_m \geq t - C_W \geq (1 - \beta_0/4)t \geq 3t/4$. Hence $g_m^2 \leq (1 + \beta_0/3)g_n^2$.

We turn to (3.10). Put $\lambda_j := \log g_j^{-2}$, which is at least $\log \gamma^{-2} \geq p/\beta_0 \geq 2p$ by (i), the second condition of (3.7), and $\beta_0 \leq 1/2$. From $g_m \leq (1+\beta_0)g_n$ we get $\lambda_n \leq \lambda_m + 2\log(1+\beta_0)$, and therefore

$$\Big(\frac{\lambda_n}{\lambda_m}\Big)^p \leq \Big(1 + \frac{2\log(1+\beta_0)}{\lambda_m}\Big)^p \leq (1+\beta_0)^{2p/\lambda_m} \leq 1 + \beta_0 \,, \tag{3.13}$$

using $1 + x \leq e^x$ and $2p \leq \lambda_m$. In the same way the first inequality of (3.9) gives $\lambda_m \leq \lambda_n + \log(1 + y)$ with $y := g_n^2\beta'(n - m)$, so that $(\lambda_m/\lambda_n)^p \leq (1 + y)^{p/\lambda_n} \leq (1 + y)^{\beta_0}$, because $p/\lambda_n \leq \beta_0$.

(iii) Since $u_m = u_n + b'(n-m)$, the numbers $\hat{g}_j$ are increasing and $\hat{g}_m^{-2} = \hat{g}_n^{-2}(1 + \hat{g}_n^2 b'(n - m))$, which is the first inequality of (3.9) with equality. The remaining inequalities follow as in (ii), with $\hat{g}_n^2 b' \leq b^* \gamma^2 \leq \beta_0$ in place of (3.12) and $\log \hat{g}_j^{-2} \geq \log \gamma^{-2}$, which holds because $u_j \geq t \geq \gamma^{-2}$. *q.e.d.*

We can now define the parameters of the construction for a tube. They are Bałaban's functions of the coupling, evaluated at $\hat{g}_j$.

Definition 3.3 (Parameters of a Tube)**.** For a tube and every integer $j \leq K$:

(i) R_j is the smallest power of L with $R_j \geq (\log \hat{g}_j^{-2})^r$;

(ii) $\varepsilon_j := \hat{g}_j p_0(\hat{g}_j)$, with $p_0(g) = A_0(\log g^{-2})^{p_0}$, and $\vartheta_j := (A_1/A_0)\varepsilon_j$;

(iii) $\delta'_j := \hat{g}_j A_1 (\log \hat{g}_j^{-2})^{p_1}$;

(iv) $\alpha_{0,j} := \hat{g}_j C_0^\alpha (\log \hat{g}_j^{-2})^{q_0}$ and $\alpha_{1,j} := \hat{g}_j C_1^\alpha (\log \hat{g}_j^{-2})^{q_1}$.

The printed definitions are those of (3.1) for ε_j and R_j; $\delta_k = g_k(A_1/A_0)p_0(g_k)$ for the threshold of the fluctuation field, which we write ϑ_k [9, p. 265, after (3.4)]; $\delta'_j = g_j A_1 p_1(g_j)$ with $p_1(g) = (\log g^{-2})^{p_1}$ and $p_1 < p_0$ for the threshold of the **R**-operation [11, p. 183]; and (2.28) for the radii $\alpha_{0,j}$, $\alpha_{1,j}$ of the domains of analyticity, with integers $q_0, q_1 > 1$ and large constants which he writes C_0, C_1 [9, p. 259]. A parameter of the scale j of a tube depends on K and j only through the difference $K - j$, since u_j does. This will be essential in Section 11: the runs for the targets $(K, \bar{g})$ and $(K+1, \bar{g})$ have the same parameters on each of the lattices which they share.

Lemma 3.4 (Cube Sizes of a Tube)**.** *Let γ satisfy* (3.7)*. For every tube and every history $(x_0, \dots, x_k)$ of it:*

(i) $(\log g_j^{-2})^r \le R_j < L(\log \hat g_j^{-2})^r \le 2L(\log g_j^{-2})^r$ *for* $0 \le j \le k$*;*

(ii) $R_{j+1} \in \{R_j,\, L^{-1}R_j\}$ *for every integer* $j < K$*, so that* $R_n \le R_m$ *for* $m < n$*;*

(iii) for all integers $m < n \le K$*,*

$$R_m \le L(1 + \hat g_n^2 b'(n-m))^{\beta_0} R_n \qquad \text{and} \qquad R_m \le (L+1)(n-m)^{\beta_0} R_n \,. \tag{3.14}$$

Part (i) is Bałaban's (2.5), with a second inequality in which the factor $2L$ stands where his definition would give L, and (ii) and (iii) contain his (2.9) [9, pp. 255–256], since $\hat g_n^2 b' \le g_n^2 \beta'$.

Proof. (i) The lower bound is the definition of R_j and $\hat g_j \le g_j$. Since $\log \hat g_j^{-2} \ge \log \gamma^{-2} > 1$, the smallest power of L which is at least $(\log \hat g_j^{-2})^r$ is less than L times this number. Finally $\log \hat g_j^{-2} \le \log g_j^{-2} + \log \rho$ by Lemma 3.2(i), with $\rho \le 3/2$ and $\log g_j^{-2} \ge \log \gamma^{-2}$, so that the third condition of (3.7) gives $(\log \hat g_j^{-2})^r \le 2(\log g_j^{-2})^r$.

(ii) The numbers u_j decrease, hence $R_{j+1} \le R_j$. Moreover $u_j = u_{j+1} + b' \le (1 + b^* \gamma^2) u_{j+1}$, because $u_{j+1} \ge \gamma^{-2}$, so that

$$R_j < L(\log u_j)^r \le L\Big(1 + \frac{\log(1 + b^*\gamma^2)}{\log u_{j+1}}\Big)^r (\log u_{j+1})^r < L^2 R_{j+1} \,, \tag{3.15}$$

by the fourth condition of (3.7) and $(\log u_{j+1})^r \le R_{j+1}$. The quotient R_j / R_{j+1} is a power of L in $[1, L^2)$, so it is 1 or L.

(iii) Put $s := n - m$ and $\lambda := \log \gamma^{-2}$. As in (i), $R_m < L(\log u_m)^r$ and $(\log u_n)^r \le R_n$, and $u_m = u_n(1 + \hat g_n^2 b' s)$, so that

$$\frac{R_m}{R_n} < L\Big(1 + \frac{\log(1 + \hat g_n^2 b' s)}{\log u_n}\Big)^r \,. \tag{3.16}$$

For the first inequality we bound the bracket by $(1 + \hat g_n^2 b' s)^{r/\log u_n}$ and use $r/\log u_n \le r/\lambda \le \beta_0$. For the second we use $\hat g_n^2 \le \gamma^2$, $b' \le b^*$, and $\log u_n \ge \lambda$, which bound the bracket by $(1 + \log(1 + b^* s \gamma^2)/\lambda)^r$, and we distinguish two cases. If $s \le \gamma^{-2}$, this is at most $(1 + \log(1 + b^*)/\lambda)^r \le 1 + 1/L$, by the last condition of (3.7), and $1 \le s^{\beta_0}$. If $s > \gamma^{-2}$, then $1 + b^* s \gamma^2 \le (1 + b^*) s \gamma^2$, and with $y := \log s > \lambda$ the bracket is at most $\varphi(y) := ((y + \log(1 + b^*))/\lambda)^r$. We have $\varphi(\lambda) \le 1 + 1/L$, and the logarithmic derivative of φ is $r/(y + \log(1 + b^*)) \le r/\lambda \le \beta_0$, so that $\varphi(y) \le (1 + 1/L) e^{\beta_0 (y - \lambda)} \le (1 + 1/L) s^{\beta_0}$. In both cases $R_m / R_n < L(1 + 1/L) s^{\beta_0}$. *q.e.d.*

Lemma 3.5 (Thresholds of a Tube)**.** *Let γ satisfy* (3.7)*, and let τ_j be one of the parameters ε_j, ϑ_j, δ'_j, $\alpha_{0,j}$, $\alpha_{1,j}$ of a tube, so that $\tau_j = \tau(\hat g_j)$ with $\tau(g) = c g (\log g^{-2})^p$ for a constant $c > 0$ and an exponent $p \in (0, p^*]$.*

(i) For every history of the tube and $0 \le j \le k$*,*

$$\rho^{-1/2} \tau(g_j) \le \tau_j \le \tau(g_j) \,. \tag{3.17}$$

(ii) For all integers $m < n \le K$*,*

$$\tau_n \le (1 + \beta_0)(n-m)^{1/2} \tau_m \qquad \text{and} \qquad \tau_m \le (1 + \hat g_n^2 b'(n-m))^{\beta_0} \tau_n \le (1 + \beta_0)(n-m)^{\beta_0} \tau_n \,. \tag{3.18}$$

For $\tau = \varepsilon$ the inequalities (3.18) are Bałaban's (2.8) [9, p. 256]. The upper bound of (3.17) says that the thresholds of a tube are the smaller ones: a statement which uses a threshold only as an upper bound on a field which has passed a test function holds for the tube with the printed value. The lower bound controls the loss in the statements which use a threshold from below.

Proof. (i) The function τ is increasing on $(0, \gamma]$, since its logarithmic derivative is $g^{-1}(1 - 2p/\log g^{-2})$ and $\log g^{-2} \geq \log \gamma^{-2} \geq p^*/\beta_0 > 2p$. As $\hat{g}_j \leq g_j \leq \gamma$ by Lemma 3.2(i), this gives the upper bound. For the lower bound we write $\tau_j = (\hat{g}_j/g_j)\, c g_j (\log \hat{g}_j^{-2})^p$ and use $\hat{g}_j \geq \rho^{-1/2} g_j$ and $\log \hat{g}_j^{-2} \geq \log g_j^{-2}$ from the same lemma.

(ii) We have $\tau_n/\tau_m = (\hat{g}_n/\hat{g}_m)(\log u_n/\log u_m)^p \leq \hat{g}_n/\hat{g}_m$, because $u_n \leq u_m$, and $\hat{g}_n/\hat{g}_m \leq (1 + \beta_0)(n-m)^{1/2}$ by Lemma 3.2(iii). In the other direction $\tau_m/\tau_n = (\hat{g}_m/\hat{g}_n)(\log u_m/\log u_n)^p \leq (\log u_m/\log u_n)^p$, which is at most $(1 + \hat{g}_n^2 b'(n-m))^{\beta_0}$ by the second inequality of (3.10) for the sequence $(\hat{g}_j)$. The last step is $1 + y(n-m) \leq (1+y)(n-m)$ with $y = \hat{g}_n^2 b' \leq \beta_0$ and $(1+\beta_0)^{\beta_0} \leq 1 + \beta_0$. *q.e.d.*

3.2 The Density after k Steps

We recall the form which Bałaban gives to the density of the block field after k steps, since every later section works inside it. The description is his inductive assumption [9, Sect. 2], and we cite each of its parts by page and equation. Nothing in this subsection is new.

At the step k we use his units, in which the lattice $T^{(k)}$ of the block field V_k has the spacing one and the fine lattice T_η, which is our $\mathbb{T}_K$, has the spacing $\eta = L^{-k}$. A *scale* $j \leq k$ refers to the lattice of spacing $L^{-(k-j)}$ in these units, the lattice on which the block field of the step j lives. For the localisation one fixes $M = L^m$ and, at each scale j, the family π_j of the closed cubes of side M of that scale. A *localisation domain* of the scale j is the union of a finite family of cubes of π_j which is connected, in the sense that any two of its cubes are joined by a chain of cubes of the family in which consecutive cubes have a common wall. The class of these domains is $\mathbf{D}_j$, and the *linear size* $d_j(X)$ of $X \in \mathbf{D}_j$ is the length of a shortest tree graph in X which meets all the cubes of X, divided by M [8, p. 257].

The terms of the expansion. A term of the density is labelled by three sequences of sets. The first two are sequences of localisation domains $\Omega_j, \Lambda_j \in \mathbf{D}_j$ with

$$\Omega_1 \supset \Lambda_1 \supset \Omega_2 \supset \Lambda_2 \supset \cdots \supset \Omega_k \supset \Lambda_k \,, \tag{3.19}$$

and the third consists of sets $S_j \subset \Omega_j \cap \Lambda_j^c$, each of which is empty or a union of cubes of side LM_2R_j of the scale j [9, (2.1), p. 254]. Roughly speaking, Ω_j is the region in which the field of the scale j has passed the test functions of the step j, so that the step could be performed there, and Λ_j is the smaller region in which the fluctuation field of that step was small as well. The complements

$$Z_j \coloneqq \Lambda_j^c \tag{3.20}$$

are the *large-field regions* [9, (2.3), p. 255]. The sequence $\{\Omega_j\}$ determines on which lattice the field is still an integration variable at each point: with

$$\Gamma_0 \coloneqq \Omega_1^c \,, \qquad \Gamma_j \coloneqq \Omega_j^{(j)} \setminus \Omega_{j+1}^{(j)} \quad (1 \leq j < k) \,, \qquad \Gamma_k \coloneqq \Omega_k^{(k)} \,, \qquad \mathbf{B} \coloneqq \bigcup_{j=0}^{k} \Gamma_j \,, \tag{3.21}$$

where the superscript (j) denotes the points, bonds, and plaquettes of the scale j in a set, the *determining set* $\mathbf{B}$ carries the field V which equals the block field V_j of the step j on Γ_j

[9, (2.2), (2.10), pp. 255–256]. The domains Ω_j and Λ_j which are determined by the step j, and not by an **R**-operation, are unions of cubes of side MR_j of the scale j, and the distance between their boundaries is at least $2MR_j$ [9, p. 256].

The background field. Let $M_{\mathbf{B}}$ be the averaging operation which maps a gauge field U on T_η to the field on **B** equal to the j-fold block average $M^j(U)$ on Γ_j. A regular field V on **B** determines the set of the minima of the action $U \mapsto A^\eta(U)$ on the regular fields U with $M_{\mathbf{B}}(U) = V$, which is a single gauge orbit, and a minimal configuration is denoted by $U_k(V)$, or by U_k [9, (2.11), (2.12), p. 256]. The existence, the uniqueness up to gauge transformations, and the regularity of $U_k(V)$ are the subject of [7]. Every term of the effective action depends on V through this *background field*.

The expansion. For a cube $\square$ of the partition of T_η into cubes of side LM_2R_k let $U_{k,\square}(V_k)$ be the minimal configuration of [9, (2.16), p. 257], which is determined by the block field in a neighbourhood of $\square$. The test function of the step k on Ω_k is the product

$$\chi_k(\Omega_k) \coloneqq \prod_{\square \subset \Omega_k} \chi\Big(\Big\{ \sup_{p \subset \square^\sim} |U_{k,\square}(V_k, \partial p) - 1| < \varepsilon_k \eta^2 \Big\} \Big) \tag{3.22}$$

of the characteristic functions of the sets on which the plaquettes of these local minimisers are below the threshold ε_k, in the units of the fine lattice, and the density is

$$\varrho_k(V_k) = \sum_{\{\Omega_j\},\{\Lambda_j\}} \chi_k(\Omega_k)\, \mathbf{T}_k(\{\Omega_j\},\{\Lambda_j\}) \exp A_k\Big(\frac{1}{g_k^2}, U_k\Big) \tag{3.23}$$

[9, (2.17), (2.18), p. 257]. The sum runs over the admissible sequences of domains, the sum over the sequences $\{S_j\}$ is included in the operation $\mathbf{T}_k$, and the effective action A_k depends on all three sequences.

The operations on the large-field regions. The operation $\mathbf{T}_k$ consists of the integrations which the earlier steps have left: over the block fields V_j on Ω_{j+1}^c, and over the fluctuation fields A_j on $Z_{j+1} \cap \Omega_{j+1}$. It factorises over the connected components $X_1, \dots, X_n$ of Z_k, and each factor is an ordered product over the scales,

$$\mathbf{T}_k(Z_k) = \prod_{i=1}^{n} \mathbf{T}_k(X_i)\,, \qquad \mathbf{T}_k(X) = \prod_{j=k-1}^{0} \mathbf{T}^{(j)}(Z_{j+1} \cap X)\,, \tag{3.24}$$

where $\mathbf{T}^{(j)}$ comes from the step $j+1$ and, if no **R**-operation has changed it, has the form of [9, (2.21), p. 258]: the integral over V_j against the δ-function of the block averaging and the function $\zeta(\Omega_{j+1}^c)$, which contains the characteristic functions of the large fields, and the integral over A_j against a characteristic function and a Gaussian weight [9, (2.19), (2.20), p. 258].

The effective action. The effective action has the form

$$A_k\Big(\frac{1}{g_k^2}, U_k\Big) = -A\Big(\frac{1}{g_k^2}, U_k\Big) + \mathbf{E}_k(U_k) + \mathbf{R}_k(U_k) + \mathbf{B}_k(U_k, A) - E_k \tag{3.25}$$

[9, (2.23), p. 258], in which $A = \{A_{j-1}\}$ stands for the fluctuation fields in the last but one term. Its five terms are the following.

(a) The first is the Wilson action of the background field with a coupling which depends on the position, defined by

$$\frac{1}{g_{j-1}^2(x)} = \frac{1}{g_j^2(x)} + \beta_j(g_{j-1})\phi_j(x)\,, \tag{3.26}$$

where $\phi_j \in C_0^\infty(\Lambda_j)$ equals one on $\Lambda_j^{\sim -1}$, the domain obtained from Λ_j by removing a layer of cubes, so that $g_j(x) = g_j$ there [9, (2.24), p. 259]. The numbers $\beta_j(g_{j-1})$ are the coefficients of the coupling flow, the subject of Section 4.

(b) The *regular part* is

$$\mathbf{E}_k(U_k) = \sum_{j=1}^{k} [\mathbf{E}^{(j)}(\Lambda_j, g_{j-1}, U_k) - \mathbf{E}^{(j)}(\Lambda_j, g_{j-1}, 1) - \beta_j(g_{j-1}) A(\phi_j, U_k)] \,, \tag{3.27}$$

with $\mathbf{E}^{(j)}(\Lambda_j, U_k) = \sum_{z \in \Lambda_j^0} \sum_{X \ni z} \mathbf{E}^{(j)}(X, U_k, z)$, the inner sum over the $X \in \mathbf{D}_j$ contained in Λ_j and Λ_j^0 obtained from $\Lambda_j^{(j)}$ by removing one layer of the cubes of side MR_j [9, (2.25)–(2.27), p. 259]. Each term $\mathbf{E}^{(j)}(X, U_k, z)$ depends on U_k restricted to X; it extends to an analytic function of a pair $(\mathbf{U}, \mathbf{J})$ in the space $U_j^c(X, \alpha_{0,j}, \alpha_{1,j})$; the extension is invariant under gauge transformations; and it satisfies the bound

$$|\mathbf{E}^{(j)}(X, (\mathbf{U}, \mathbf{J}), z)| \le E_0 \exp(-\kappa d_j(X)) \tag{3.28}$$

of [8, (1.18), p. 263]. The space $U_j^c(X, \alpha_0, \alpha_1)$ consists of the gauge orbits of the pairs of a G^c-valued field $\mathbf{U} = U'U$ and a $\mathfrak{g}^c$-valued field $\mathbf{J}$ on X in which U is a real field with plaquettes below α_0, $U' = \exp i\xi A'$ with A' and its covariant derivative below α_1, and $\mathbf{J}$ is below α_0, together with the corresponding bounds for the minimisers built from the block averages of $\mathbf{U}$ [8, Definition, (1.10)–(1.16), p. 262]. It is a complex neighbourhood of the real regular fields on X, of a width set by α_1.

(c) The $\mathbf{R}$-*part* is $\mathbf{R}_k(U_k) = \sum_{j=1}^{k} [\mathbf{R}^{(j)}(\Lambda_j, U_k) - \mathbf{R}^{(j)}(\Lambda_j, 1)]$, where $\mathbf{R}^{(j)}(\Lambda_j, U_k)$ is a sum of terms $\mathbf{R}^{(j)}(X, U_k)$ over the domains $X \in \mathbf{D}_j$ contained in $\Lambda_j^{\sim -1}$ which are unions of cubes of side MR_j. These terms have the first three properties of the terms of the regular part and satisfy

$$|\mathbf{R}^{(j)}(X, (\mathbf{U}, \mathbf{J}))| \le g_j^{\kappa_0} \exp(-\kappa d_j(X)) \tag{3.29}$$

[9, (2.30), (2.31), p. 260]. They are produced by the $\mathbf{R}$-operation from terms with large fields, and they are so small that they need no counterterm.

(d) The *boundary part* is $\mathbf{B}_k = \sum_{j=1}^{k} \mathbf{B}^{(j)}(U_k, A, \{S_i\})$, where $\mathbf{B}^{(j)}$ is a sum of terms $\mathbf{B}^{(j)}(X, U_k, A, \{S_i \cap X\})$ over the $X \in \mathbf{D}_j$ with $X \cap \Omega_j \neq \emptyset$ and $X \cap Z_j^{\sim} \neq \emptyset$. Each of them depends on U_k and A restricted to X, extends analytically to the space $\tilde{U}_j^c(X, \tilde{\alpha}_0, \tilde{\alpha}_1)$, is invariant under G-valued gauge transformations, and satisfies

$$|\mathbf{B}^{(j)}(X, (\mathbf{U}, \mathbf{J}), A, \{S_i \cap X\})| < B_0 \exp(-\kappa d_j(X)) \tag{3.30}$$

[9, (2.40)–(2.42), p. 261]. The space $\tilde{U}_j^c(X, \tilde{\alpha}_0, \tilde{\alpha}_1)$ is the version of U_j^c for a domain which meets several of the layers $\Omega_n \setminus \Omega_{n+1}$: its conditions (2.34)–(2.39) are those of U_j^c with the radii $\alpha_{0,n}$, $\alpha_{1,n}$ of the scale n on the layer n, each multiplied by $1 - \beta(1 - 2^{-(j-n)})$ with a constant β, for which Bałaban suggests $\beta = 1/4$ [9, p. 261].

(e) E_k is a constant, which depends on $\{\Omega_j\}$ and $\{\Lambda_j\}$ [9, p. 262].

The terms which a step has just created are defined on slightly larger spaces and decay faster, with $(1 + 4\beta)\kappa$ in place of κ, and this margin is consumed by the $\mathbf{R}$-operation [9, p. 262].

Bałaban's theorems. The main result of [9] is that the form just described is reproduced by a step.

Theorem 3.6 (Bałaban [9, Theorem 1, p. 262]). *There exist constants, introduced in the above description, such that if the sequence of coupling constants $\{g_k\}$, determined by the recursive renormalisation group equations (0.18), (0.20) of [8], satisfies the inequalities (0.31) of [8], then the sequence of densities $\{\varrho_k\}$, generated by successive applications of the operations $\mathbf{R}T$ to the density $\varrho_0 = \exp[-(1/g_0^2)A - E]$, satisfies all the inductive assumptions.*

The proof of this theorem occupies [9], for the step T, and [11, 12], for the **R**-operation. Two consequences are used throughout. The first is a bound on the regular and the **R**-parts in a region Ω, [9, Theorem 2, (2.43), (2.44), p. 263]: with a constant E_1 independent of j, k, Ω, the sequences of domains, and the torus,

$$\Big| \sum_{z \in \Lambda_j^0 \cap \Omega} [\mathbf{E}^{(j)}(\Lambda_j, U_k, z) - \mathbf{E}^{(j)}(\Lambda_j, 1, z)] - \beta_j(g_{j-1}) A(\phi, U_k) \Big| \leq E_1 \sum_{n=j}^{k} (L^{j-n})^\beta |\Gamma_n \cap \Omega| \quad (3.31)$$

for $\beta < 1$, and the corresponding sum of the **R**-terms is at most $R_1 g_j^{\kappa_0} \sum_{n=j}^{k} |\Gamma_n \cap \Omega|$. Here $\Omega \in \mathbf{D}_k$ is a domain contained in the union of the blocks of Λ_j^0, $\phi \in C_0^\infty(\Omega^\sim)$ equals one on Ω, the field is restricted by the test functions of (3.23), and $|\Gamma_n \cap \Omega|$ is the number of points of the scale n. The second is ultraviolet stability, [9, Corollary 3, (2.50), p. 264]: there are constants $E_\pm$, independent of η and of the torus but dependent on g_k, with

$$\chi_k(T_\eta) \exp\Big[-\frac{1}{g_k^2} A(U_k(V_k)) - E_- |T_\eta| \Big] \leq \varrho_k(V_k) \leq e^{E_+ |T_\eta|} \, . \quad (3.32)$$

These statements are for Bałaban's parameters, which are functions of the running couplings, and for characteristic functions as test functions. The construction of this paper differs from his in both respects, and the remainder of this section shows that the description and the bounds hold for it.

3.3 Soft Test Functions

The domains of a term are produced by decompositions of unity. In each step the lattice is divided into cubes $\square$, a nonnegative function $f_\square$ of the fields near $\square$ is compared with a threshold t, and the integrand is multiplied by

$$1 = \sum_P \prod_{\square \not\subset P} \chi(\{f_\square < t\}) \prod_{\square \subset P} \chi(\{f_\square \geq t\}) \, , \quad (3.33)$$

the sum running over the unions P of cubes. In the term P we say that the test function of a cube has *passed* if the cube lies outside P, and that it has *failed* if the cube lies in P. Bałaban's steps contain two kinds of such decompositions. In the first, $f_\square$ is the supremum of $|U(\partial p) - 1|$ over the plaquettes p near $\square$, for the fine field in the first step and for a local minimiser $U_{k+1,\square}$ of the action in the later ones, and t is a plaquette threshold ε in the units of the plaquette [9, (1.1), (1.4), p. 246; (2.17), p. 257; (3.2), p. 265]. In the second, $f_\square$ is the supremum over the bonds b near $\square$ of the deviation $|V_k(b)(V_\square^{(k)}(b))^{-1} - 1|$ of the field from a reference configuration, the block average of a local minimiser, and $t = 2\vartheta_k$ [9, (1.8), p. 247; (3.3), p. 265]. We call the factors of (3.33) *test functions*, of the plaquettes and of the reference configuration respectively.

A characteristic function is the natural choice as long as one run of the construction is studied. For the limit $K \to \infty$ we shall compare two runs, with K and $K + 1$ steps, on the lattices which they share, and there the functions $f_\square$ of the two runs differ by a

small amount, since they are computed from minimisers on two different fine lattices. A characteristic function can change by one under an arbitrarily small change of its argument, and the set on which this happens, a thin shell around the level set $\{f_\square = t\}$, has no reason to be of small measure for the densities at hand. We therefore replace the characteristic functions by continuous functions, whose change is proportional to that of the argument.

Definition 3.7 (Soft Test Functions)**.** Let $\kappa_s > 0$. For $x \in [0,\infty]$ put

$$\bar\chi(x) \coloneqq \min\Big(1,\, \max\Big(0,\, \frac{1+\kappa_s - x}{\kappa_s}\Big)\Big)\,, \tag{3.34}$$

so that $\bar\chi(x) = 1$ for $x \le 1$, $\bar\chi(x) = 0$ for $x \ge 1+\kappa_s$, and $\bar\chi$ is affine in between. The *soft* form of the decomposition (3.33) is

$$1 = \sum_P \prod_{\square\not\subset P} \bar\chi\Big(\frac{f_\square}{t}\Big) \prod_{\square\subset P} \Big(1 - \bar\chi\Big(\frac{f_\square}{t}\Big)\Big)\,. \tag{3.35}$$

Lemma 3.8 (Properties of the Soft Decomposition)**.** *Let $(f_\square)$ be a finite family of functions with values in $[0,\infty]$ and $t > 0$.*

(i) The identity (3.35) *holds, and each of its terms has values in* $[0,1]$.

(ii) The term P of (3.35) *vanishes unless $f_\square < (1+\kappa_s)t$ for every $\square \not\subset P$ and $f_\square > t$ for every $\square \subset P$. In particular it is bounded by the term P of the decomposition* (3.33) *in which the passed test functions are taken at the threshold $(1+\kappa_s)t$ and the failed ones at the threshold t, with strict inequality.*

(iii) For $x, y \in [0,\infty)$, $|\bar\chi(x) - \bar\chi(y)| \le |x-y|/\kappa_s$. Consequently, for two families $(f_\square)$, $(f'_\square)$, with the convention that $|f_\square - f'_\square|$ is 0 where both are infinite and ∞ where exactly one is, and with Π_P, Π'_P their terms P of (3.35)*,*

$$|\Pi_P - \Pi'_P| \le \frac{1}{\kappa_s t} \sum_\square |f_\square - f'_\square|\,. \tag{3.36}$$

Proof. (i) Put $a_\square \coloneqq \bar\chi(f_\square/t) \in [0,1]$. Expanding the product $\prod_\square (a_\square + (1-a_\square))$, which equals one, gives the sum over the subsets of the family of the products $\prod_{\square\notin P} a_\square \prod_{\square\in P}(1-a_\square)$, and each of these is a product of numbers in $[0,1]$.

(ii) The factor $a_\square$ vanishes for $f_\square \ge (1+\kappa_s)t$, and the factor $1-a_\square$ vanishes for $f_\square \le t$. Since all factors are at most one, the term is bounded by the product of the characteristic functions of the sets $\{f_\square < (1+\kappa_s)t\}$, $\square \not\subset P$, and $\{f_\square > t\}$, $\square \subset P$.

(iii) The function $\bar\chi$ is piecewise affine with slopes 0 and $-1/\kappa_s$, which gives the first assertion. For the second we replace the factors of Π_P by those of Π'_P one at a time. Each replacement changes one factor by at most $|f_\square - f'_\square|/(\kappa_s t)$, both for a factor $a_\square$ and for a factor $1-a_\square$, while the other factors lie in $[0,1]$. *q.e.d.*

The first property says that nothing is lost: the lattice integrals are the same whichever decomposition is used to expand them, and only the terms of the expansion change. The second is the form in which the soft test functions enter every estimate. A bound which Bałaban derives from a passed test function is a bound on a field below the threshold, and it holds for the soft term with the threshold enlarged by the factor $1+\kappa_s$; a bound which he derives from a failed test function, the small factor which the action gives on a region of large fields, holds unchanged. The third property is used only in Section 10.

In the construction of this paper all the test functions of the plaquettes and of the reference configuration are soft, with

$$\kappa_s \coloneqq \frac{\beta}{30} \,, \tag{3.37}$$

β being the constant of the spaces of the boundary terms in Section 3.2. This value is dictated by the margins which Bałaban's inequalities leave between a threshold and the bound that is drawn from it, and which the factor $1+\kappa_s$ must not exhaust. The characteristic functions which restrict the fluctuation fields are not test functions in the above sense, and they are left as they are.

3.4 The Depth of the R-Operation

The **R**-operation of the step k acts on those connected components of the large-field region Z_k which are small and old: a component must be contained in a cube of side $100MR_k$, and in the preceding N steps no new large-field region may have been created inside it, the earlier regions inside it being small in the same sense on their scales [11, p. 177, (i), (ii)]. The number N is the *depth* of the operation. Bałaban does not fix it. He writes that "conditions on N will be formulated in constructions of this section" [11, p. 177], and the conditions which he then states are the following four.

(N1) $N > N_0$, where N_0 is the smallest positive integer with $L^{-N_0+1}MR_{k-N_0+1} = M$ [11, p. 179].

(N2) $N \le O(1)(\log g_k^{-2})^\nu$ for a positive integer ν with $\nu/2 \le p_0 - p_1 - 1$, and N is so large that $L^{-N}N^{1/2}O(1)B_3^3B_5M^6 \le \alpha/2$ [11, (1.94), p. 198].

(N3) $N \le R_k$ [12, p. 361].

(N4) $L^{-N} \le (\log g_k^{-2})^{-\nu}$ for a sufficiently large ν, with $\nu \ge 2p_0 + dr$ [12, p. 363].

The last condition recurs as "the usual conditions on N", which make $p_0(g_k)q_1(g_k)R_k^7L^{-N}$ small [12, (1.47), p. 368], and as "our assumptions on N", which make $R_j^{1+\beta_0}(p_0(g_j)/p_1(g_j))L^{-N}$ small [12, p. 382]. In the large-field induction of [12, p. 384] the conditions (i), (ii) are taken "with $N = R_j$". We make a different choice, which satisfies (N1)–(N4) with a depth of the order of $\log\log g_k^{-2}$ only. The reason is that in Section 6 the factor L^N occurs, and has to be bounded by a power of $\log g_k^{-2}$.

Definition 3.9 (Depth)**.** Put $\nu_L \coloneqq p_0 + q_1 + 7r + 1$. For a tube and an integer $k \le K$:

(i) $N_0(k)$ is the smallest positive integer N_0 with $L^{N_0-1} = R_{k-N_0+1}$;

(ii) N_k' is the least integer with $L^{-N_k'} \le (\log \hat{g}_k^{-2})^{-\nu_L}$;

(iii) the *depth* is $N_k \coloneqq \max(N_0(k)+1,\, N_k')$.

For the next lemma we assume, in addition to (3.7), that $p^* \ge \nu_L$ and that

$$\log_L(16L^4(L+1)^2) + \nu_L \log_L \lambda \le \lambda^r \qquad \text{for every } \lambda \ge \log \gamma^{-2} \,, \tag{3.38}$$

which holds for γ small because $r \ge 2$.

Lemma 3.10 (Bounds on the Depth)**.** *Let γ satisfy* (3.7) *with $p^* \ge \nu_L$ and* (3.38)*, and let $L \ge 3$. For every tube and every integer $k \le K$:*

(i) $N_0(k)$ *exists,* $N_0(k) \ge 2$*, and* $N_0(k)$*,* N'_k*, and* N_k *are nonincreasing in* k*;*

(ii) $L^{-N_k} \le (\log \hat g_k^{-2})^{-\nu_L}$ *and* $L^{N_k} \le 4L^4(L+1)^2(\log \hat g_k^{-2})^{\nu_L}$*;*

(iii) for every history $(x_0, \dots, x_{k'})$ *of the tube and* $0 \le k \le k'$*, with* $C_{\mathrm{dep}} := 16L^4(L+1)^2$*,*

$$L^{-N_k} \le (\log g_k^{-2})^{-\nu_L} , \qquad L^{N_k} \le C_{\mathrm{dep}} (\log g_k^{-2})^{\nu_L} , \qquad N_k \le R_k . \tag{3.39}$$

Proof. (i) Put $n := K - k \ge 0$, and for an integer $t \ge 0$ let $\varrho(t) := \log_L R_{K-t}$ and $h(t) := t - \varrho(t)$. By Lemma 3.4(i) and (ii), ϱ takes integer values at least 1 and $\varrho(t+1) - \varrho(t) \in \{0, 1\}$, so that $h(t+1) - h(t) \in \{0, 1\}$ and $h(0) \le -1$. Moreover $h(t) \to \infty$, because $\varrho(t) \le 1 + r \log_L \log u_{K-t}$ grows like $\log\log t$. A nondecreasing integer-valued function with steps 0 or 1 which starts below n and is unbounded takes the value n, and we let $T(n)$ be the least t with $h(t) = n$. With $t = n + N_0 - 1$, so that $K - t = k - N_0 + 1$, the equation $L^{N_0 - 1} = R_{k - N_0 + 1}$ reads $\varrho(t) = t - n$, that is, $h(t) = n$. Its solutions have $N_0 = \varrho(t) + 1 \ge 2$, and the smallest is $N_0(k) = T(n) - n + 1$. Since h is nondecreasing and $h(T(n)) = n$, we have $h(t) \le n$ for $t \le T(n)$, hence $T(n+1) \ge T(n) + 1$, and $T(n) - n$ is nondecreasing in n. So $N_0(k)$ is nonincreasing in k. The same holds for N'_k, because $\hat g_k$ increases with k, and therefore for N_k.

(ii) The first inequality is the definition of $N'_k \le N_k$. By the minimality of N'_k we have $L^{N'_k} < L(\log \hat g_k^{-2})^{\nu_L}$. For $N_0 = N_0(k)$ we use the second inequality of (3.14) with $m = k - N_0 + 1$ and $n = k$: with $x := N_0 - 1 \ge 1$,

$$L^x = R_{k-x} \le (L+1) x^{\beta_0} R_k \le (L+1)(x+2) R_k \le 2(L+1) L^{x/2} R_k , \tag{3.40}$$

where the last step is the inequality $x + 2 \le 2L^{x/2}$, which holds for $x \ge 0$ and $L \ge 3$, since the two sides agree at $x = 0$ and the derivative $L^{x/2} \log L$ of the right side is at least $\log 3 > 1$. Hence $L^{x/2} \le 2(L+1) R_k$, and with $R_k < L(\log \hat g_k^{-2})^r$,

$$L^{N_0 + 1} = L^2 L^x \le 4L^2 (L+1)^2 R_k^2 < 4L^4 (L+1)^2 (\log \hat g_k^{-2})^{2r} . \tag{3.41}$$

Since $\nu_L \ge 2r$ and $\log \hat g_k^{-2} \ge 1$, both L^{N_0+1} and $L^{N'_k}$ are at most $4L^4(L+1)^2(\log \hat g_k^{-2})^{\nu_L}$.

(iii) By Lemma 3.2(i), $\log g_k^{-2} \le \log \hat g_k^{-2} \le \log g_k^{-2} + \log \rho$, so that the first inequality follows from (ii), and the second from (ii) and $(\log \hat g_k^{-2})^{\nu_L} \le 2(\log g_k^{-2})^{\nu_L}$, which is the third condition of (3.7) with $\nu_L \le p^*$. Taking logarithms, $N_k \le \log_L C_{\mathrm{dep}} + \nu_L \log_L \log g_k^{-2}$, which is at most $(\log g_k^{-2})^r \le R_k$ by (3.38) and Lemma 3.4(i). *q.e.d.*

The depth N_k satisfies Bałaban's four conditions. (N1) holds by definition and (N3) by (3.39). For (N4), (3.39) gives the exponent ν_L, and $\nu_L - (2p_0 + dr) = q_1 - p_0 + 3r + 1 > 0$ at $d = 4$, because $q_1 \ge p_0$ (Appendix A). The two further uses of (N4) hold as well: with $R_k \le 2L(\log g_k^{-2})^r$, the product $p_0(g_k) q_1(g_k) R_k^7 L^{-N_k}$ is at most a constant times $(\log g_k^{-2})^{p_0 + q_1 + 7r - \nu_L} = (\log g_k^{-2})^{-1}$, and $R_k^{1+\beta_0}(p_0(g_k)/p_1(g_k)) L^{-N_k}$ is at most a constant times $(\log g_k^{-2})^{r(1+\beta_0) + p_0 - p_1 - \nu_L}$, whose exponent is negative. For (N2), the first restriction holds with $\nu = \lceil r \rceil$, because $N_k \le R_k \le 2L(\log g_k^{-2})^r$, under the condition $\lceil r \rceil / 2 \le p_0 - p_1 - 1$ on the exponents, and the second holds for γ small, because $L^{-N_k} N_k^{1/2} \le (2L)^{1/2} (\log g_k^{-2})^{r/2 - \nu_L}$ tends to zero. That these are all the places at which the depth enters the two papers, and that each of them holds for N_k, is Supplement, Proposition S.3.

3.5 The Construction of This Paper

We can now say which construction we run. It is Bałaban's, with the parameters of a tube and with soft test functions.

Definition 3.11 (The Construction)**.** Let a tube be given, with γ as in Lemma 3.10, and let $g_0 > 0$. The *construction for the tube started at* g_0 is the sequence of densities $\varrho_0, \varrho_1, \dots$ obtained from $\varrho_0 = \exp[-(1/g_0^2)A - E]$ by Bałaban's operations $\mathbf{R}T$ of [9, 11, 12], with the following changes.

(i) Every parameter of the scale j which is defined there as a function of the coupling g_j is that function at $\hat{g}_j$: the cube size R_j, the thresholds ε_j, ϑ_j, δ'_j and the radii $\alpha_{0,j}$, $\alpha_{1,j}$ of Definition 3.3, and every other threshold, radius, or support of a cutoff function of that scale. The depth of the **R**-operation of the step k is N_k.

(ii) The characteristic functions which restrict a fluctuation field are functions of the unscaled field: the fluctuation field A_j of the step $j+1$ is restricted by $|g_j A_j(b)| < \vartheta_j$, which is the form of [9, (1.18), p. 250] and [11, (1.9), p. 178], and the field B_j of [11, (1.55), p. 188] by $|g_j B_j(b)| < \delta'_j$; the bump functions of the changes of variables [9, (1.22), p. 251] are functions of the unscaled field as well, with supports fixed by the thresholds (Supplement, Definition 3.5).

(iii) The test functions of the plaquettes and of the reference configuration are the soft test functions of Definition 3.7, with the width (3.37); the cascade of test functions of the **R**-operation [11, (1.22)–(1.24), p. 181] is soft with the nested widths of Supplement, Definition 2.2, and so is the cutoff $\chi(\Lambda_i)$ of [11, (1.101), p. 201].

The *running couplings* are $g_k = x_k^{-1/2}$ with $x_{k+1} \coloneqq x_k - \beta_{k+1}$, where $\beta_{k+1} = \beta_{k+1}(g_0, \dots, g_k)$ is the coefficient which Bałaban's formula [9, (3.61), p. 282] assigns to the regular part of the effective action of the step $k+1$, the coefficient of (3.26). The construction is defined up to the step $k \le K$ as long as $(x_0, \dots, x_k)$ is a history of the tube.

In (ii) the factor g_j which converts the field of the Gaussian integral into the unscaled field is the running coupling, and only the threshold is fixed by the target. For this reason the construction is continuous in the couplings: no cube size, no depth, and no set on which a characteristic or test function is supported depends on $g_0, \dots, g_k$, which enter only through the action, the Gaussian weights, and the terms of the effective action generated before.

Proposition 3.12 (Bałaban's Description for the Construction)**.** *Let γ be sufficiently small. For every tube, every g_0, and every $k \le K$ up to which the construction for the tube started at g_0 is defined, the density ϱ_k has the expansion* (3.23)*, with soft test functions in* (3.22)*, and its terms have the properties (a)–(e) of Section 3.2, with constants E_0, B_0, κ, κ_0 which do not depend on the tube, on g_0, on k and K, or on the torus. The bound* (3.31) *holds with a constant which is independent of the same quantities; the bound* (3.32) *holds with constants $E_\pm$ which depend on these quantities through the coupling g_k only, as in Bałaban's statement; and the results of [11, 12] on the* **R***-operation hold as well.*

Proof. Supplement, Propositions S.1, S.2, and S.3. Proposition S.1 goes through the places at which a parameter enters [9, 11, 12] and the papers on which they rest, and shows that each of them is of one of four kinds: a relation among the parameters alone, which holds for the sequence $(\hat{g}_j)$ by Lemmata 3.2–3.5; a lower bound drawn from the action, which holds

with the constant in the exponent divided by $\rho \le 3/2$, because the coefficient of the action is at least $\hat{g}_m^{-2}/\rho$ at the points which carry a field of the scale m (Supplement, Lemma 2.1); an upper bound relative to the action, which holds because the thresholds of a tube are the smaller ones; or a conversion between a scaled and an unscaled field, which is the change (ii). Proposition S.2 goes through the places at which a test function enters and shows that every bound drawn from a passed test function holds at the threshold $(1+\kappa_s)t$ of Lemma 3.8(ii). Proposition S.3 is the corresponding statement for the depth. *q.e.d.*

3.6 The Stopping Index

The loops are drawn on a lattice $\mathbb{T}_q$ whose spacing uL^{-q} does not depend on K, and the construction is stopped on a lattice which is finer than $\mathbb{T}_q$ by a factor that does not depend on K either. This factor has to be so large that, at every step, the straight segments of a loop are long compared with the cubes of the construction. Since the parameters of a tube at the scale $K-n$ depend on n and not on K, we may write

$$R^{[n]} \coloneqq R_{K-n} \,, \tag{3.42}$$

the smallest power of L which is at least $(\log(\bar{g}^{-2} + b'n + C_W))^r$.

Definition 3.13 (Stopping Index)**.** Let $M = L^m$. For an integer $j \ge 0$ put

$$\bar{q}(j) \coloneqq 3 + m + \log_L R^{[j+1]} \,, \qquad \text{so that} \qquad L^{\bar{q}(j)} = L^3 M R^{[j+1]} \,. \tag{3.43}$$

The *stopping index* j_q of the level $q \ge 0$ is the least integer j with $j \ge q + \bar{q}(j)$. For a target $(K, \bar{g})$ with $K > j_q$, the *stopping step* is $K' \coloneqq K - j_q$.

Lemma 3.14 (The Stopping Index)**.** *Let γ be as in Lemma 3.4. For every $q \ge 0$ the stopping index j_q exists. It depends on q, L, M, r, $\bar{g}$, b', and C_W only, and*

$$j_q = q + \bar{q}(j_q) \,, \qquad L^{j_q - q} = L^3 M R_{K'-1} \ge L^3 M R_{K'} \,. \tag{3.44}$$

Consequently, for a loop C at level q, read on the lattice of the step $k \le K'$, whose spacing is uL^{k-K}: the corners of C lie in a coset of $a_k\mathbb{Z}^4$ with $a_k \coloneqq L^{j_q - q} L^{K'-k}$, and, in the notation of Section 2 for this lattice,

$$P_k(C) = P_q(C)\, a_k \,, \qquad s_k(C) \ge a_k \,, \qquad \delta_k(C) \ge a_k \,, \qquad a_k \ge L^3 M R_{K'} L^{K'-k} \ge L^2 M R_k \,. \tag{3.45}$$

Two loops of an ensemble at level q are at ℓ^∞-distance at least a_k on that lattice.

Proof. The function $\bar{q}$ is nondecreasing, because $R^{[n]}$ is nondecreasing in n by Lemma 3.4(ii), and $\bar{q}(j) \le 4 + m + r \log_L \log(\bar{g}^{-2} + b'(j+1) + C_W)$ by Lemma 3.4(i). Hence $j - q - \bar{q}(j) \to \infty$ as $j \to \infty$, and the least j with $j \ge q + \bar{q}(j)$ exists. It is determined by q and the function $\bar{q}$, which gives the dependence. Since $\bar{q} \ge 3$, we have $j_q > q$, so that $j_q - 1 \ge q$ and, by minimality, $j_q - 1 < q + \bar{q}(j_q - 1) \le q + \bar{q}(j_q)$. Together with $j_q \ge q + \bar{q}(j_q)$ this gives the first identity of (3.44). The second is (3.43) with $R^{[j_q+1]} = R_{K-j_q-1} = R_{K'-1}$, and $R_{K'-1} \ge R_{K'}$ by Lemma 3.4(ii).

The lattice of the step k has the spacing uL^{k-K}, and $uL^{-q} = L^{K-k-q} uL^{k-K}$ with $K - k - q = (j_q - q) + (K' - k)$. So one bond of $\mathbb{T}_q$ is a_k bonds of that lattice, which gives the statement on the corners and the first identity of (3.45). The bounds on $s_k(C)$ and $\delta_k(C)$ are Lemma 2.3(i) with $a = a_k$. For two loops of an ensemble we apply Lemma 2.3(ii) to a lift of the first and to all the translates of a lift of the second by the periods of the torus, which are

multiples of a_k because $T/(uL^{-q}) = 2L^{m_T+q}$ is an integer: these lifts have no common site, since the loops are vertex-disjoint on the torus, and their corners lie in one coset of $a_k\mathbb{Z}^4$. For the last chain we use $L^{j_q-q} \geq L^3 M R_{K'}$ and, by Lemma 3.4(ii) applied $K'-k$ times, $R_k \leq L^{K'-k} R_{K'}$. *q.e.d.*

The inequality $a_k \geq L^2 M R_k$ says that every segment of a loop, and every gap between two of its segments or between two loops, is at least L^2 cubes of the step k long, at every step up to the stopping step. On the stopping lattice itself a loop at level q has $P_{K'}(C) = P_q(C) L^3 M R_{K'-1}$ bonds, a number which does not depend on K.

The steps $k \leq K'$ are those at which the loops will be carried, and their couplings are bounded by a number which does not depend on K. Indeed, for a history $(x_0, \dots, x_k)$ of the tube with $k \leq K'$ we have $x_k \geq \ell_k = t + b(K-k) - C_W \geq t + bj_q - C_W$ by (3.3), because $K - k \geq K - K' = j_q$, and $\hat{g}_k \leq g_k$ by Lemma 3.2(i). With

$$\gamma_q \coloneqq (\bar{g}^{-2} + bj_q - C_W)^{-1/2} \tag{3.46}$$

we therefore have $\hat{g}_k \leq g_k \leq \gamma_q$ for $0 \leq k \leq K'$. The number γ_q is small if $\bar{g}$ is small and also if q is large: since $\bar{g}^{-2} - C_W \geq \gamma^{-2}$ by (3.2) and $j_q \geq q$ by (3.44),

$$\gamma_q \leq (\bar{g}^{-2} + bq - C_W)^{-1/2} \leq \min\Big((\bar{g}^{-2} - C_W)^{-1/2},\ (\gamma^{-2} + bq)^{-1/2} \Big)\,, \tag{3.47}$$

and $(\bar{g}^{-2} - C_W)^{-1/2} \leq (1 - \beta_0/4)^{-1/2} \bar{g} \leq (1 + \beta_0)\bar{g}$, because $C_W \leq \beta_0 \bar{g}^{-2}/4$ by (3.2) and $(1+\beta_0)^2(1-\beta_0/4) \geq 1$ for $0 < \beta_0 \leq 1$. Every requirement of the paper which involves the representations of an ensemble concerns the steps $k \leq K'$ only, and it is imposed as an upper bound on γ_q; by (3.47) such a bound holds at every coupling $\bar{g} \leq \gamma$ once the level q is large enough.

4 The Coupling Flow

The construction of Section 3 takes a bare coupling and returns the running couplings. What the lattice theory of Section 1.1 needs is the converse: for a prescribed value $\bar{g}$ of the coupling at the unit scale and for every number K of steps, a bare coupling g_0 with $g_K = \bar{g}$, together with bounds on the couplings in between which do not depend on K. Bałaban states this as a theorem for $G = SU(2)$ and writes that "a proof of this theorem, based on perturbative calculations, will be given in a separate paper" [8, Theorem 2, p. 259]; in [9] the bounds on the couplings are a hypothesis, as Theorem 3.6 shows. In this section we prove the statement, for a simple group and for the construction of this paper.

Theorem 4.1 (Coupling Flow)**.** *There are $\gamma' > 0$ and constants $0 < b \leq b'$ and $C_W \geq 0$ such that, for every integer $K \geq 1$ and every $\bar{g} \in (0, \gamma']$, there is a bare coupling $g_0 = g_0(K, \bar{g})$ for which the running couplings $g_0, g_1, \dots, g_K$ of the construction for the target $(K, \bar{g})$ satisfy $g_K = \bar{g}$ and, for $0 \leq k \leq l \leq K$,*

$$\bar{g}^{-2} + b(K-k) - C_W \leq g_k^{-2} \leq \bar{g}^{-2} + b'(K-k) + C_W \qquad \text{and} \qquad g_k^{-2} \geq g_l^{-2} - C_W\,. \tag{4.1}$$

The constants are $b = 2c \ln L - \delta'$ and $b' = 2c \ln L + \delta'$, where c is the number (4.3), the first coefficient of the β-function, and $0 \leq \delta' \leq (2/5)c \ln L$.

The *target* $(K, \bar{g})$ stands for the tube of Definition 3.1 with these b, b', and C_W. The proof has three inputs of different kinds. The first is a computation at one loop: at zero

coupling the inverse square of the coupling decreases by $2c \ln L$ per step, up to an error which stays bounded when it is summed over any number of steps (Theorem 4.4). The second is perturbative as well, but concerns Bałaban's expansions: the true decrease differs from the one at zero coupling by a multiple of g_k^2, uniformly in the step and in the earlier couplings (Proposition 4.3(i), (ii)). With these two facts every bare coupling would produce a sequence that satisfies (4.1) up to its last step, but the last coupling would be whatever it turns out to be. To prescribe it we need the third input, the continuity of the couplings as functions of the bare coupling (Proposition 4.3(iii)), and it is for this that the parameters of the construction were detached from the running couplings in Section 3.1. Given the three inputs, the bare coupling is found by the intermediate value theorem.

4.1 The Coefficients of the Flow

We first say what the coefficient β_{k+1} of Definition 3.11 is. Let $\mathfrak{g} \subset \mathfrak{u}(N)$ be the Lie algebra of G, a space of anti-Hermitian matrices, with the inner product $\langle X, Y\rangle \coloneqq -\operatorname{tr}(XY)$ and the Killing form $\mathbf{K}(X, Y) \coloneqq \operatorname{Tr}(\operatorname{ad} X \operatorname{ad} Y)$.

Lemma 4.2 (Invariant Forms)**.** *Every bilinear form on $\mathfrak{g}$ which is invariant under $\mathrm{Ad}(G)$ is a real multiple of the inner product. In particular there is a number $\kappa > 0$, depending on G and on its realisation, with*

$$\mathbf{K}(X, Y) = \kappa \operatorname{tr}(XY) \,, \tag{4.2}$$

and $\kappa = 2N^2$ for $G = SU(N)$ in its defining representation.

Proof. An invariant bilinear form can be written as $(X, Y) \mapsto \langle SX, Y\rangle$ with a linear map S of $\mathfrak{g}$ which commutes with $\mathrm{Ad}(G)$, hence with every $\operatorname{ad} X$, and we treat its symmetric and antisymmetric parts separately, each of which commutes with ad as well, because the maps $\operatorname{ad} X$ are antisymmetric for the invariant inner product. A symmetric S has a real eigenvalue λ, and $\ker(S - \lambda)$ is a nonzero ideal of $\mathfrak{g}$, since $S[X, Y] = [X, SY]$. As $\mathfrak{g}$ is simple, $S = \lambda$. For an antisymmetric S the square S^2 is symmetric and commutes with ad, so that $S^2 = -\mu^2$ with $\mu \geq 0$ by what has been shown and $\langle S^2 X, X\rangle = -|SX|^2$. If μ were positive, $J \coloneqq S/\mu$ would be a complex structure on $\mathfrak{g}$ with $J[X, Y] = [X, JY]$, so that $\operatorname{ad}(JX) = J \operatorname{ad} X$ and $\mathbf{K}(JX, JX) = \operatorname{Tr}(J \operatorname{ad} X \, J \operatorname{ad} X) = \operatorname{Tr}(J^2 (\operatorname{ad} X)^2) = -\mathbf{K}(X, X)$. This is impossible, because $\mathbf{K}(X, X) = -\operatorname{Tr}((\operatorname{ad} X)^T \operatorname{ad} X)$ is negative for every $X \neq 0$, the centre of a simple Lie algebra being trivial. Hence $S = 0$. The Killing form is invariant and negative definite, which gives (4.2) with $\kappa > 0$, and for $\mathfrak{su}(N)$ one has $\mathbf{K}(X, Y) = 2N \operatorname{Tr}(XY) = 2N^2 \operatorname{tr}(XY)$. *q.e.d.*

The number

$$c \coloneqq \frac{11}{48\pi^2} \kappa \tag{4.3}$$

is the first coefficient of the β-function in the normalisation of the action (1.2): for $SU(N)$ and the usual coupling g_{s}, with $g^{-2} = 2N g_{\mathrm{s}}^{-2}$, the statement that g^{-2} decreases by $2c \ln L$ when the lattice spacing is multiplied by L is $\mathrm{d} g_{\mathrm{s}}^{-2} / \mathrm{d} \ln \varepsilon = -11N/(24\pi^2)$. From now on the numbers b_* and b^* of Section 3.1 are

$$b_* \coloneqq c \ln L \,, \qquad b^* \coloneqq 3c \ln L \,, \tag{4.4}$$

and a tube is always one that satisfies (3.2) with them.

The polarisation coefficient. We use the units of the step $k + 1$, in which the block field of that step lives on the unit lattice. Let F be a function of the background field of the step $k + 1$ on the lattice $\mathbb{Z}^4$, which is invariant under gauge transformations and under the

symmetries of the lattice and is a sum of local terms of the kind (3.27), so that its derivatives at the trivial field are summable. For a block field $V = \exp iB$, with B a field on the bonds with values in $i\mathfrak{g}$ and components $B^a_\mu(x)$ in an orthonormal basis, the second derivative of $B \mapsto F(U_{k+1}(\exp iB))$ at $B = 0$ with respect to $B^a_\mu(x)$ and $B^{a'}_\nu(x')$ is, for fixed μ, x, ν, x', an invariant bilinear form on $\mathfrak{g}$, because F is invariant under constant gauge transformations. By Lemma 4.2 and translation invariance it equals $\delta^{aa'}\Pi_{\mu\nu}(F; x - x')$, and we put

$$\Lambda(F) \coloneqq \sum_{x\in\mathbb{Z}^4} \Pi_{\mu\nu}(F; x)\, x_\mu x_\nu \tag{4.5}$$

for a pair $\mu \neq \nu$, on which the value does not depend. This is Bałaban's definition of the β-function from the vacuum polarisation [8, (1.20)–(1.22), p. 264], where the independence of the pair is part of the statement and the limit of the infinite lattice is taken with the help of the local representation. The coefficient of the step $k+1$ is Λ of the regular part which that step generates on the whole lattice,

$$\beta_{k+1} \coloneqq \Lambda(\mathbf{E}^{(k+1)}) \,, \tag{4.6}$$

and [9, (3.57)–(3.64), pp. 281–283] shows that this is the number for which the counterterm $-\beta_{k+1}A(\phi_{k+1}, U)$ of (3.27) removes the marginal part of $\mathbf{E}^{(k+1)}$. The regular part contains the logarithm of the normalisation factor of the Gaussian measure of the fluctuation field of the step,

$$Z^{(k)}(U) \coloneqq \int \delta(\tilde{Q}B) \exp[-\tfrac{1}{2}\langle B, \Delta^{(k)}(U)B\rangle]\, \mathrm{d}B \tag{4.7}$$

[8, (1.3), (1.4), p. 260], which is its part of order zero in the couplings, and the rest is the fluctuation integral proper, $\mathbf{E}^{(k+1)} = \log Z^{(k)} + \mathbf{E}^{(k+1)}_{\mathrm{fl}}$ up to a constant. Since Λ is linear,

$$\beta_{k+1} = b_{k+1} + \Lambda(\mathbf{E}^{(k+1)}_{\mathrm{fl}}) \,, \qquad b_{k+1} \coloneqq \Lambda(\log Z^{(k)}) \,. \tag{4.8}$$

The numbers $b_1, b_2, \ldots$ are the *coefficients at zero coupling*. They are determined by the quadratic forms $\Delta^{(k)}(U)$ of [8, (1.5), p. 261], that is, by L, G, the averaging operations, and k, and they involve no coupling, threshold, or tube.

Histories which are not flows. For the argument that will produce the bare coupling we need β_{k+1} as a function on the whole set H_k of Definition 3.1, whereas the construction started at g_0 only visits the points of H_k with $x_{j+1} = x_j - \beta_{j+1}$ for all $j < k$. We therefore extend the construction. For $h = (x_0, \ldots, x_k) \in H_k$ the densities $\varrho^h_0, \ldots, \varrho^h_k$ are defined inductively: $\varrho^h_0 \coloneqq \exp[-x_0 A - E]$, and ϱ^h_{j+1} is obtained by applying the operations $\mathbf{R}T$ of Definition 3.11 to ϱ^h_j, with $g_j = x_j^{-1/2}$ as the coupling of the scale j, and by multiplying each term of the resulting expansion (3.23) by

$$\exp[-(x_{j+1} - x_j + \beta_{j+1}(x_0, \ldots, x_j))A(U_{j+1})] \,, \tag{4.9}$$

where $A(U_{j+1})$ is the Wilson action of the background field of the term and $\beta_{j+1}(x_0, \ldots, x_j)$ is the coefficient (4.6) of the step just performed. The effect is that the Wilson term of every term of ϱ^h_{j+1} has the coefficient function

$$\frac{1}{g^2_{j+1}(x)} = x_{j+1} + \sum_{i=1}^{j+1} \beta_i(x_0, \ldots, x_{i-1})(1 - \phi_i(x)) \,, \tag{4.10}$$

which is x_{j+1} on the domain $\Lambda^{\sim -1}_{j+1}$ where all the ϕ_i equal one. Indeed, by (3.26) the step turns the coefficient function y of ϱ^h_j into $y - \beta_{j+1}\phi_{j+1}$, and (4.9) adds the constant $x_{j+1} - x_j + \beta_{j+1}$.

If h is a flow, the factor (4.9) is one at every step and ϱ_j^h is the density of the construction started at $g_0 = x_0^{-1/2}$. In all cases $\beta_{k+1}(h)$ denotes the coefficient (4.6) of the step $k+1$ applied to ϱ_k^h.

Proposition 4.3 (The Coefficients on a Tube)**.** *There are constants C_Λ, C_3, C_{rem}, C_R, and $\gamma_1 > 0$ such that, for every $\gamma \in (0, \gamma_1]$, every tube, every $k < K$, and every history $h = (x_0, \dots, x_k) \in H_k$, with $g_k = x_k^{-1/2}$, the following holds. The densities $\varrho_0^h, \dots, \varrho_k^h$ are defined and have the properties stated in Proposition 3.12, and:*

(i) $|\beta_{k+1}(h) - b_{k+1}| \le \delta'$, *where* $\delta' := C_\Lambda C_3 \gamma p_0(\gamma) + C_R \gamma^{\kappa_0 - 4}$;

(ii) $|\beta_{k+1}(h) - b_{k+1}| \le (C_{\mathrm{rem}} + C_R \gamma^{\kappa_0 - 6})\, g_k^2$;

(iii) the function β_{k+1} is continuous on H_k.

The constants depend on those of the construction and not on γ, the tube, k, or h.

Proof. Supplement, Propositions S.1, S.4, and S.5. Let us say where the difficulty lies in each part. That the densities of a history which is not a flow have Bałaban's properties is part of Proposition S.1, since the coefficient function (4.10) differs from that of a flow only through coefficients which obey the same bounds. By (4.8) the difference in (i) and (ii) is Λ of the fluctuation integral, whose local terms Bałaban bounds by a multiple of the threshold ε_1 of the fluctuation field [10, (2.41), p. 21]. In the construction of this paper that threshold is ϑ_k, which in the units of the step is at most $\gamma p_0(\gamma)$, and since Λ is bounded on local terms with exponential decay this gives the first term of δ'. The second term accounts for the **R**-terms of the earlier steps, which enter the regular part through the fluctuation integral and which the bound (3.29) makes small. The improvement to g_k^2 in (ii) rests on a symmetry: when g_k is replaced by a complex number z in the exponent of the fluctuation integral and the threshold is held fixed, the integral is an even analytic function of z which vanishes at $z = 0$ up to a Gaussian tail. For (iii) one has to follow every term of the density through the step and to check that it is continuous in the history at each configuration, which is where the construction of Section 3 is used: no set on which a characteristic function or a test function is supported moves with the history. *q.e.d.*

The exponent κ_0 is that of (3.29), and Bałaban takes $\kappa_0 \ge 7$ [9, (2.46), p. 263], so that $\delta' \to 0$ as $\gamma \to 0$. The first input of the proof, on the other hand, concerns the numbers b_k alone.

Theorem 4.4 (The Coefficients at Zero Coupling)**.** *There is a constant C_W, depending on L, G, and the averaging operations only, such that for all integers $0 \le m < n$*

$$\Big| \sum_{j=m+1}^{n} b_j - 2c\,(n-m) \ln L \Big| \le C_W \ . \tag{4.11}$$

Proof. Supplement, Proposition S.6. The statement is the content of Bałaban's Theorem 2 of [8] at zero coupling, and we describe the structure of its proof, since the number c is computed there. It suffices to show that $\Phi_k := \sum_{j=1}^{k} b_j$ satisfies $|\Phi_k - 2ck \ln L| \le C_W/2$ for every k. By (4.8) and the local representation of the terms of order zero [8, (0.28)–(0.30), p. 258], Φ_k is, up to a bounded error, the coefficient of the polarisation of the Gaussian factor $\sum_{j<k} \log Z^{(j)}(U_k)$ of k steps, measured against the polarisation of the action of the

background field, whose symbol is bounded above and below at a fixed nonzero momentum [2, (1.67), p. 29]. The Gaussian factor of k steps is then written exactly as a difference $F_k - F_0$ of two functionals, each of which is one half of the logarithm of the determinant of a gauge-invariant quadratic form, of the action at the level k for F_k and of the action at the level 0 for F_0, divided by the determinant of the corresponding gauge operator, plus a sum of terms which come from the current J of [8, (1.5), (1.8), p. 261]. The identity is linear algebra (a Schur complement and the matrix determinant lemma), and it telescopes over the steps. Three estimates of symbols then give the theorem: the polarisation of the terms of the current is bounded uniformly in k, because gauge covariance makes it vanish to second order at zero momentum; the polarisation of F_k is bounded uniformly in k at a fixed momentum, because it is the one-loop polarisation of a theory on a lattice of unit spacing with a cutoff at unit scale; and the polarisation of $-F_0$, the one-loop polarisation of the bare lattice theory seen from the coarse lattice, is $c\ln(1/\eta_k^2)$ times the action of the background up to a bounded error, with $\eta_k = L^{-k}$. The last estimate is the computation of the one-loop coefficient of the lattice theory in a background field, in which the logarithm comes from the region of momenta between the two cutoffs, and the coefficient $11/48\pi^2$ is the one of the continuum computation, κ entering through Lemma 4.2. *q.e.d.*

We now show how the coupling flow follows.

4.2 Proof of the Coupling Flow

Proof of Theorem 4.1. The constants. Let C_W be the constant of Theorem 4.4. We fix $\gamma \in (0, \gamma_1]$ so small that the conditions of Section 3 hold with (4.4) and that

$$\delta' \le \tfrac{2}{5} c \ln L \qquad \text{and} \qquad \gamma^{-2} \ge \Big(\frac{4}{\beta_0} - 1\Big) C_W \,, \tag{4.12}$$

and we put $b := 2c\ln L - \delta'$, $b' := 2c\ln L + \delta'$, and $\gamma' := (\gamma^{-2} + C_W)^{-1/2}$. Then $b \ge (8/5)c\ln L \ge b_*$, $b' \le (12/5)c\ln L \le b^*$, and $b'/b \le 3/2$, and for $\bar g \le \gamma'$ the number $t = \bar g^{-2}$ satisfies $t \ge \gamma^{-2} + C_W \ge 4C_W/\beta_0$. So $(K, b, b', C_W, \bar g)$ is a tube in the sense of Definition 3.1, for every $K \ge 1$ and $\bar g \in (0, \gamma']$. The choice is not circular, since the constants of Proposition 4.3, and hence δ', are the same for all tubes.

The window of the coefficients. Let $0 \le m < n \le K$ and let $h_j \in H_j$ be arbitrary histories, $m \le j < n$. By Proposition 4.3(i) each $\beta_{j+1}(h_j)$ differs from b_{j+1} by at most δ', and by (4.11) the sum of the b_{j+1} differs from $2c(n-m)\ln L$ by at most C_W. Hence

$$b(n-m) - C_W \le \sum_{j=m}^{n-1} \beta_{j+1}(h_j) \le b'(n-m) + C_W \,. \tag{4.13}$$

The projected flow. The set H_k is closed, convex, and nonempty, since it contains $(u_0, \dots, u_k)$. Let P_k be the Euclidean projection of $\mathbb{R}^{k+1}$ onto it, which does not increase distances and is therefore continuous. For $x_0 \in \mathbb{R}$ we define $y_0 := x_0$ and

$$y_{k+1} := y_k - \beta_{k+1}(P_k(y_0, \dots, y_k)) \tag{4.14}$$

for $0 \le k < K$. This sequence is defined for every x_0, whether or not it stays in the tube, and each y_k is a continuous function of x_0, by induction on k and Proposition 4.3(iii). Moreover the arguments of the coefficients in (4.14) are histories, so that (4.13) applies and gives, for $0 \le m < n \le K$,

$$b(n-m) - C_W \le y_m - y_n \le b'(n-m) + C_W \,. \tag{4.15}$$

The bare coupling. With $m = 0$ and $n = K$, (4.15) says that y_K lies between $x_0 - b'K - C_W$ and $x_0 - bK + C_W$. At $x_0 = \ell_0 = t + bK - C_W$ this gives $y_K \le t$, and at $x_0 = u_0 = t + b'K + C_W$ it gives $y_K \ge t$. By the intermediate value theorem the continuous function $x_0 \mapsto y_K$ takes the value t on $[\ell_0, u_0]$; its set of solutions there is closed, and we let $g_0(K, \bar g)^{-2}$ be the smallest of them.

The projection is inactive. For this x_0, (4.15) with $n = K$ and $y_K = t$ gives $\ell_m \le y_m \le u_m$ for every $m < K$, by the definition (3.3) of ℓ_m and u_m, and (4.15) itself is the condition (3.4). So $(y_0, \dots, y_k) \in H_k$ for every $k \le K$, the projection P_k leaves it fixed, and (4.14) reads $y_{k+1} = y_k - \beta_{k+1}(y_0, \dots, y_k)$: the sequence is a flow, the densities ϱ_k^h of its histories are those of the construction for the tube started at g_0, and its running couplings are $g_k = y_k^{-1/2}$, with $g_K = t^{-1/2} = \bar g$. The first part of (4.1) is $\ell_k \le y_k \le u_k$, and the second follows from the lower bound of (4.15), since $y_k - y_l \ge b(l-k) - C_W \ge -C_W$ for $k < l$. *q.e.d.*

Three remarks are in order. First, the construction depends on the target: the parameters, the densities, the coefficients, and the bare coupling are those of the tube of $(K, \bar g)$, and objects of different targets are not identified. What is uniform are the constants. Secondly, by Lemma 3.2 the running couplings of the theorem satisfy Bałaban's inequalities (2.6), which he derives from the renormalisation group equations and uses in their place [9, p. 255]; the window (4.1) is the inequality (0.31) of [8] up to the additive constant C_W, with $\beta = b/\ln L$ and $\beta' = b'/\ln L$. Thirdly, the bare coupling window (1.3) of the introduction is (4.1) at $k = 0$.

5 Dressed Loops and the First Step

Bałaban's construction acts on densities, and an observable has to be carried along with them. This section says in what form the loops of an ensemble are carried, and proves that the first step puts them into that form.

5.1 Two Integrals

Let $\mathcal{C} = ((C_1, \chi_1), \dots, (C_n, \chi_n))$ be a loop ensemble at level q, let $K > j_q$, and let $g_0 = g_0(K, \bar g)$ be the bare coupling of Theorem 4.1. With $\varrho_0 := \exp[-g_0^{-2} A - E]$ the bare density of Section 3.2, the expectation of the observable (1.4) is the quotient

$$\mathbb{E}_K[W_{\mathcal{C}}] = \frac{Z_K(\mathcal{C})}{Z_K}\,, \qquad Z_K(\mathcal{C}) := \int W_{\mathcal{C}}(U)\varrho_0(U)\,\mathrm{d}U\,, \qquad Z_K := \int \varrho_0(U)\,\mathrm{d}U\,, \tag{5.1}$$

in which the constant E cancels. We apply the construction to the numerator and to the denominator separately. Each operation of the construction preserves the integral of a density: the block averaging T does, because $\int \delta(\bar U(c) V(c)^{-1})\,\mathrm{d}V(c) = 1$ for the Haar measure, and the **R**-operation is defined so that "integrals of the densities are unchanged" [11, p. 176]; the density it produces "may not be equal to ϱ_k, but is equivalent to it, in the sense that they have equal integrals" [11, p. 200]. Hence $Z_K = \int \varrho_k(V_k)\,\mathrm{d}V_k$ for every $k \le K$, and for the numerator we shall construct densities N_k with

$$Z_K(\mathcal{C}) = \int N_k(V_k)\,\mathrm{d}V_k\,, \qquad N_0 := W_{\mathcal{C}}\,\varrho_0\,, \tag{5.2}$$

for $0 \le k \le K' = K - j_q$. No law of the block variables enters: the quotient of the two integrals at the step K' is the expectation at the cutoff K. The point of the construction is the form of N_k, which is that of ϱ_k with one more factor in each term.

5.2 The First Step of the Construction

The first step integrates the fine field with its block averages V on the lattice of the step 1 held fixed. Bałaban decomposes the integrand by test functions into terms labelled by two domains, the domain Ω_1 on which the test functions of the plaquettes have passed and the domain $\Lambda_1 \subset \Omega_1$ on which the fluctuation field is small as well, and after the fluctuation integral on Λ_1 and the renormalisation of the coupling he obtains

$$\varrho_1(V) = \sum_{\{\Omega_1,\Lambda_1\}} \chi_1(\Omega_1)\, \mathbf{T}_1(\{\Omega_1,\Lambda_1\}) \exp A_1\Big(\frac{1}{g_1^2}, U_1\Big) \tag{5.3}$$

[9, (1.28), p. 253]. Here $\mathbf{T}_1$ is the operation which holds the integrals over the bare field V_0 on Ω_1^c and over the fluctuation field A on $\Omega_1 \cap \Lambda_1^c$, with the Gaussian weight of the step and the characteristic functions $\chi(\Omega_1 \cap \Lambda_1^c, S_1)$ of the large fluctuations [9, (1.29), p. 254], and the effective action is

$$\begin{aligned} A_1\Big(\frac{1}{g_1^2}, U_1\Big) = -A\Big(\frac{1}{g_1^2}, U_1\Big) + \{\mathbf{E}^{(1)}(\Lambda_1, g_0, U_1) - \mathbf{E}^{(1)}(\Lambda_1, g_0, 1) - \beta_1(g_0) A(\phi_1, U_1)\} \\ + \mathbf{B}^{(1)}(U_1, A, S_1) - E_1 \end{aligned} \tag{5.4}$$

[9, (1.30), p. 254], with the coupling of [9, (1.27), p. 253]. These integrals stay pending inside ϱ_1: Bałaban leaves the operations $\mathbf{T}_k$ of the later steps "undefined, only their basic general properties are formulated" [9, p. 254]. A *labelled term* of ϱ_1 is a term of (5.3) with given labels. For the numerator we use the refinement of the labels of [1, Section 9], in which the domain Λ_1 is further restricted near the loop by a retreat of the carrier of the loop's integral; the refined labels are again labels of Bałaban's form, and on each of them the operation $\mathbf{T}_1$ and the action A_1 are those of (5.3) (Supplement, Proposition S.7).

In the construction of this paper the thresholds of the first step are those of the tube at $\hat{g}_0$, and the test functions of the plaquettes are soft (Definition 3.11). The first step with the loop inserted is the subject of the companion manuscript [1], for one square loop, Bałaban's thresholds at g_0, and characteristic functions as test functions, and for two disjoint squares in [1, Section 10]. We use its results for a general ensemble and for the construction of this paper, and the changes this requires are proved in Section 5.4.

5.3 Dressed Loops

Let us describe the form in which the ensemble is carried, beginning with a heuristic. After k steps the fine field has been integrated out down to the scale k, and a loop C_i is a path of $P_k(C_i)$ bonds of that scale. If the block field were exactly flat, the loop would contribute to each term of the density a number, the product $\kappa^{(k)}(C_i, \chi_i)$ of the numbers which the steps $1, \dots, k$ extract from it; since it is not, the loop also contributes the Wilson loop of the background field, corrected by insertions along the loop which are small where the background is close to flat and decay away from the regions of large fields. The insertions interact with each other through the excluded volume of the polymer expansions which produced them, and that interaction is resummed into a factor attached to each family of compatible insertions. The precise form is as follows.

Fix $k \geq 1$ and a labelled term of ϱ_k, with the labels of the construction at every scale. A *traversal* at the step k is an oriented bond of the lattice of the scale k which is contained in one of the loops C_i, traversed in the orientation of C_i; by Lemma 3.14 each loop is a union of $P_k(C_i) = P_q(C_i)a_k$ traversals. For a field U on the fine lattice and an arc J of C_i, that

is, a set of consecutive traversals, $U(J)$ denotes the product of the transports along the fine bonds of J. The objects which make up a dressed loop are the following.

(a) *Groups.* A group W is made at a step $j \le k$. It consists of a nonempty finite set of unions of cubes of π_j, its *localisation domains*, and of a set $\mathcal{I}(W)$ of pairwise disjoint arcs of the loops, its *intervals*; the set $\mathbb{L}(W)$ of the loops which carry an interval of W is nonempty. Its *domain* X_W is the union of its localisation domains and of the cubes of π_j which meet a fine bond of its intervals. Its *size* $\sigma(W) \ge 1$ is a number which is fixed at the step j, and we write $\sigma_j(W)$ when the step is to be shown: for $j \ge 2$ it is $\sigma_j(W) := \#_j(X_W) + n_j(W)/M$, where $\#_j(X_W)$ is the number of the cubes of π_j in X_W and $n_j(W)$ the number of the traversals of the step j which meet its intervals, and for $j = 1$ it is the size which [1] gives to a group. A group is a *cross group* if $|\mathbb{L}(W)| \ge 2$. A group which is not a cross group has one interval, its *arc*; a cross group may have several intervals on one loop, and its intervals are also called its arcs. The groups of a term carry a symmetric relation, their *compatibility*, and a *family* is a finite set $\mathfrak{B}$ of pairwise compatible groups whose intervals are pairwise disjoint. Two groups whose intervals share a bond are incompatible, with one exception. A *link group* is a group made at the first step whose only interval is a single fine bond (Section 5.6); it is incompatible with the link groups of the same bond and compatible with every other group. A link group and a group whose interval contains its bond thus cannot belong to one family, although they are compatible, and no cluster of a reference gas is incompatible with a link group.

(b) *Insertions.* Each group W carries an *insertion* Y_W, an element of $\bigotimes_{I\in\mathcal{I}(W)} \mathrm{End}(V_{\pi_{i(I)}})$, where $C_{i(I)}$ is the loop which contains the interval I and V_{π_i} is the space of the representation π_i; for a group with one interval it is a linear map of V_{π_i}. It is a function of the block field V_k and of the outer variables of the term, the variables which $\mathbf{T}_k$ holds pending. Its norm is the projective norm

$$\|Y\| := \inf\Big\{ \sum_{\alpha} \prod_{I\in\mathcal{I}(W)} \|A_{\alpha,I}\| \;\Big|\; Y = \sum_{\alpha} \bigotimes_{I\in\mathcal{I}(W)} A_{\alpha,I} \Big\}\,, \tag{5.5}$$

the infimum over the finite sums of elementary tensors, with the operator norms of the factors. For a group with one interval this is the operator norm of Y.

(c) *Words.* Let $\mathfrak{B}$ be a family whose insertions are elementary tensors, $Y_W = \bigotimes_I A_{W,I}$. Its word $w_k(\mathfrak{B})$ is the product, over the loops C_i, of the normalised traces on V_{π_i} of the ordered products along C_i in which the transport $\pi_i(U_k(J))$ along every arc J outside the intervals of $\mathfrak{B}$ alternates with the operators $A_{W,I}$ at the intervals $I \subset C_i$. This expression is linear in each factor $A_{W,I}$, and the word of a family with arbitrary insertions is defined from it by multilinearity. Here U_k is the background field of the term, and on a large-field region crossed by a loop the transports also contain outer variables. For the empty family $w_k(\emptyset) = \prod_i f_k^{(i)}$ with the *leading words* $f_k^{(i)} := \chi_i(U_k(C_i))$.

(d) *Interactions.* $Q_k(\mathfrak{B}) = \exp(-\sum_\Gamma \varphi_k(\Gamma))$, the sum over the clusters Γ of a hard-core gas of groups, the *reference gas* of the step, which are incompatible with a member of $\mathfrak{B}$, with numbers $\varphi_k(\Gamma)$.

(e) *The scalar exponent.* $\mathcal{E}_k = \sum_{j\le k} \sum_X e_X^{(j)} + E_{\mathrm{const},k}$, a sum of local terms $e_X^{(j)}$ indexed by connected unions X of boxes of the lattice integrated at the step j, and a number $E_{\mathrm{const},k}$.

The norm (5.5) is the one in which the words are bounded, whatever the order in which the loops meet the intervals of the cross groups.

Lemma 5.1 (Words and Norms)**.** *Let $\mathfrak{B}$ be a family, and let the transports along the arcs outside its intervals be unitary. Then $|w_k(\mathfrak{B})| \le \prod_{W\in\mathfrak{B}} \|Y_W\|$.*

Proof. If the insertions are elementary tensors, the word is a product over the loops of normalised traces of ordered products of operators on the spaces V_{π_i}. Each of these traces is at most the product of the operator norms of its factors in modulus, and the transports have the norm one, so that $|w_k(\mathfrak{B})| \le \prod_{W\in\mathfrak{B}} \prod_{I\in\mathcal{I}(W)} \|A_{W,I}\|$. In general we write each Y_W as a finite sum of elementary tensors, expand the word by multilinearity, apply this bound to every term, and take the infimum over the representations in (5.5). *q.e.d.*

The operator norm of Y_W on the tensor product of the spaces of its loops would not serve here. Two loops need not meet the intervals of several cross groups in the same cyclic order, and the word is then not the trace of a product of operators on a tensor product: for three transpositions of the two factors of $V \otimes V$, met by the two loops in opposite cyclic orders, it equals $\dim V$, although a transposition has the operator norm one. The bounds on the insertions of cross groups in the sequel are bounds on sums and integrals of elementary tensors by the sums and integrals of the products of the norms of their factors, and therefore bounds in the norm (5.5); a product of insertions and transports along the intervals is bounded by the product of the norms, since $\|AB\| \le \|A\|\|B\|$ on each factor of an elementary tensor.

The bounds of the class are stated with the following quantities. The small parameters of the step k are, with the radius $r_k := A_1(\log g_k^{-2})^{p_0}$ of the fluctuation cutoff in units of the coupling,

$$\tau_k := C_\tau[g_k(1+r_k)^3 + e^{-c_\tau r_k^2}] \,, \qquad \rho_k := C_\rho g_k^2(1+r_k)^4 \,, \qquad \rho_{t,k} := C_t g_k(1+2r_k) \,, \quad (5.6)$$

with constants fixed in Appendix A. For a traversal $\mathfrak{n}$ and $t \ge 1$ let $\mathcal{P}_t(\mathfrak{n})$ be the set of the fine plaquettes of Ω_k within distance $C_{\mathrm{box}}M(t+1)$ of $\mathfrak{n}$, the distance measured in units of the scale k. The *deviation* of the background near $\mathfrak{n}$ is

$$\delta^*_{\mathfrak{n}}(U) := L^{2k} \sum_{t\ge 1} (t+1)^3 e^{-\eta_w t} \max\{|U(\partial p) - 1| \,|\, p \in \mathcal{P}_t(\mathfrak{n})\} \,, \quad (5.7)$$

the maximum over the empty set being 0; the factor L^{2k} converts the deviation of a fine plaquette into that of a plaquette of the scale k. Finally $d_k(\mathfrak{n})$ is the distance, in units of the scale k, from $\mathfrak{n}$ to the large-field data of the scale k (the large-field region Z_k of the term), and $N^{(k)}_{\min}$ is the smallest period of the torus in units of the scale k.

The deviation δ^* bounds a sum of insertions, and this is how the class enters most estimates. The inductive step, however, has to evaluate an insertion at a background other than the one at which it was bounded, and to sort the insertions according to the region which their bounds consult; a bound on a sum cannot be used for either, and the class therefore also records a bound for each insertion of the deviation species, with the region to which it refers. The quantities are the following; for a group W they are formed at the step j at which W was made and in the units of the scale j, and they are not changed afterwards. The *dependence set* X^+_W of W is the closed ℓ^∞-neighbourhood of its domain X_W of radius $8M$ in the units of the fine lattice, which is less than the side of a cube of any of the partitions π_j, $j \ge 1$; and $\ell_W := \#_j(X_W)$, the number of the cubes of π_j in the domain, for $j \ge 2$, while $\ell_W := 1$ for $j = 1$. Put $R_t := C_{\mathrm{box}}M(t+1)$ for $t \ge 0$. For a traversal $\mathfrak{n}$ of the step j in

an interval of W let $t_{\mathfrak{n}}(W)$ be the least $t \geq 1$ for which X_W lies in the ℓ^∞-ball $B_\infty(\mathfrak{n}, R_t)$ of radius R_t about $\mathfrak{n}$. The *comparison region* of W at $\mathfrak{n}$ is the ball $B_\infty(\mathfrak{n}, R_{t_{\mathfrak{n}}(W)+1})$, which contains the dependence set X_W^+, and the *deviation of the background at* W is

$$\mathrm{Dev}_{\mathfrak{n}}(W; U) \coloneqq \ell_W \, L^{2j} \max_p |U(\partial p) - 1| \,, \tag{5.8}$$

the maximum over the fine plaquettes p in the comparison region of W at $\mathfrak{n}$. The *extent* $\mathrm{ext}(W)$ is the least $R \geq 0$ such that X_W and the comparison regions of W at all the traversals of its intervals lie in the ball of radius RM about each of these traversals. For a traversal $\mathfrak{n}$ of the step $k \geq j$ in an interval of W, let $t_{\mathfrak{n}}^{(k)}(W)$ be the least $t \geq 1$ such that the comparison regions of W at the traversals of the step j contained in $\mathfrak{n}$ lie within the distance R_t of $\mathfrak{n}$ in the units of the scale k; since the units grow with the step, $t_{\mathfrak{n}'}^{(k+1)}(W) \leq t_{\mathfrak{n}}^{(k)}(W)$ for $\mathfrak{n} \subset \mathfrak{n}'$. A group W with one interval, made at the step j, *embeds* in the torus if $2R_{t_{\mathfrak{n}}(W)+1} + 1 < N_{\min}^{(j)}$ for every traversal $\mathfrak{n}$ of the step j in its interval, that is, if each of its comparison regions, a box whose longest side is $2R_{t_{\mathfrak{n}}(W)+1} + 1$, is the injective image of a box of the universal cover of the torus. Finally the weights of the tails are

$$w_t \coloneqq (t+1)^3 e^{-\eta_w t} \,, \qquad W_t \coloneqq \sum_{t' \geq t} w_{t'} \,, \tag{5.9}$$

so that $\delta_{\mathfrak{n}}^*(U) = L^{2k} \sum_{t \geq 1} w_t \max\{|U(\partial p) - 1| \,|\, p \in \mathcal{P}_t(\mathfrak{n})\}$ in (5.7).

Definition 5.2 (Dressed Loops)**.** Let $k \geq 1$. A *dressed loop of the step* k for the ensemble $\mathcal{C}$ on a labelled term of ϱ_k is a function of V_k and of the outer variables of the term of the form

$$\mathcal{W}_k = \prod_{i=1}^n \kappa^{(k)}(C_i, \chi_i) \cdot e^{\mathcal{E}_k} \Big[\prod_{i=1}^n f_k^{(i)} + \sum_{\mathfrak{B} \neq \emptyset} Q_k(\mathfrak{B}) \, w_k(\mathfrak{B}) \Big] \,, \qquad \kappa^{(k)}(C, \chi) \coloneqq \prod_{j=1}^k \kappa_j(C, \chi) \,, \tag{5.10}$$

the sum over the nonempty families, absolutely convergent, with the objects (a)–(e) and the step factors κ_j of Definition 7.8, such that the following hold.

(i) *Interactions.* $|Q_k(\mathfrak{B})| \leq \prod_{W \in \mathfrak{B}} e^{\theta(W)}$ with $0 \leq \theta(W) \leq \nu\sigma(W)/16$, where $\sigma(W)$ is the size of W at the step at which it was made; more precisely, the sum of $|\varphi_k(\Gamma)|$ over the clusters Γ of the reference gas which are incompatible with a group W is at most $\theta(W)$.

(ii) *Insertions.* For every traversal $\mathfrak{n}$, the sum $\nu_{\mathfrak{n}}^{(k)} \coloneqq \sum_{W \ni \mathfrak{n}} e^{\theta(W)} \|Y_W\|$ over the groups of the term made at the step k, and over the groups of earlier steps which are not attached to a large-field region of their step, the groups of the cross species defined below being excepted in both cases, satisfies

$$\nu_{\mathfrak{n}}^{(k)} \leq C_\chi \tilde{\rho}_k \delta_{\mathfrak{n}}^*(U_k) + C\hat{\rho}_k e^{-c_{\mathrm{att}} d_k(\mathfrak{n})/M} + C\rho_{t,k} \, \mathfrak{m}_k(\mathfrak{n}) + \sum_{j=1}^k 2C_{S'} L^{k-j} \rho_j e^{-c_{\mathrm{att}} N_{\min}^{(j)}/M} \,, \tag{5.11}$$

where $\tilde{\rho}_k \coloneqq \rho_k / \min(\alpha_{0,k}, \alpha_{1,k})$ for $k \geq 2$ and $\tilde{\rho}_1 \coloneqq \rho_1$, $\hat{\rho}_k \coloneqq C\rho_k$, and $\mathfrak{m}_k(\mathfrak{n})$ is the sum of the marks of the large-field data of the step k near the boundary of Z_k, weighted by $e^{-c_*' \, \mathrm{dist}(\cdot, \mathfrak{n})/M}$. The *marks* are the mark weights of [1, Lemma 9.24] with the parameters of the step k: a bond b of the region of integration carries the mark $\mathfrak{m}_b$, which bounds the effect of the conditional mean on the cutoff of b and satisfies $\mathfrak{m}_b \leq 2C_\varpi r_k^{\dim G} \sum_e e^{-\mu_H d(b,e)}$, the sum over the bonds e of the boundary of that region, and a *host* Y of the step with an input outside the region carries a mark $\mathfrak{m}_Y$, which is bounded by a constant of the construction. The hosts are not the localisation domains

of the expansions, which are connected unions of arbitrarily many cubes: a host is the elementary box in which one local primitive of the step is evaluated, a box of cubes of side M with at most four cubes along each side [1, (3.36), (3.75), Lemma B.10], and only hosts carry marks. A mark of a bond is assigned to the cube Δ of side LM of the scale k which contains the bond, and a mark of a host to the cube Δ which contains the lowest cube of the host; $\hat{\mathfrak{m}}_\Delta$ is the sum of the marks assigned to Δ, and

$$\mathfrak{m}_k(\mathfrak{n}) \coloneqq \sum_\Delta \hat{\mathfrak{m}}_\Delta e^{-c'_* \operatorname{dist}(\Delta,\mathfrak{n})/M} , \qquad \hat{\mathfrak{m}}_\Delta \le C_{\mathfrak{m}}(1+r_k)^{\dim G} . \tag{5.12}$$

The second holds because a cube of side LM has at most $8(LM+1)^4$ bonds, each with a mark of at most $2C_\varpi r_k^{\dim G}$ times a convergent sum, and because the hosts assigned to it are at most 4^4L^4 in number, a host being determined by its lowest cube, one of the L^4 cubes of side M of Δ, and by its four side lengths. For general localisation domains no such count holds, and none is used: the localisation domains of the step enter the bounds of the insertions through the sums of their activities, with their decay, and not through marks. This bound per cube is the one used when the attached parts are paid (Section 6.15). These are the marks of the first step. At a step $k \ge 2$ the marks are $\hat{\mathfrak{m}}_\Delta \coloneqq 1$ for the cubes Δ of π_k which lie in Λ_k^c or have a point in common with a cube of Λ_k^c, and 0 for the other cubes (Section 6.13). The groups attached at an earlier step $j < k$ to a component Z of the large-field region of the step j satisfy, summed over all traversals, the bound $C\hat{\rho}_j \sum_{\mathfrak{n}_j} e^{-c_{\text{att}} d_j^Z(\mathfrak{n}_j)/M}$, the sum over the traversals of the step j, and the corresponding bound with $\rho_{t,j}$ for those attached to the boundary layer of Z. The groups with one interval which are not attached to a large-field region are of two species, and the species of a group is fixed when the group is made: the *deviation species* consists of those which embed in the torus, and the *wrapping species* of the others. The groups of the wrapping species satisfy the last term of (5.11) with weights: $\sum_{W \ni \mathfrak{n}} e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)} \|Y_W\|$, the sum over these groups, is at most that term. The groups of the deviation species, whose insertions vanish at pure-gauge data on their arcs, satisfy the first term of (5.11) with weights: $\sum_{W \ni \mathfrak{n}} e^{\theta(W)+\nu\sigma(W)} \|Y_W\| \le C_\chi \tilde{\rho}_k \delta^*_{\mathfrak{n}}(U_k)$, the sum over these groups, with $\sigma(W)$ the size of W at the step at which it was made. They satisfy it in the following form, of which it is a consequence (Lemma 5.3). The comparison regions of a group W of the deviation species lie in Ω_k. For each such group there is a number $b_W \ge 0$, fixed at the step j at which W was made, such that for every traversal $\mathfrak{n}$ of the step j in an interval of W

$$e^{\theta(W)} \|Y_W\| \le b_W \operatorname{Dev}_{\mathfrak{n}}(W; U_k) , \tag{5.13}$$

and, with $\beta_W \coloneqq b_W \ell_W L^{2j}$, for every traversal $\mathfrak{n}$ of the step k and every $t \ge 1$,

$$\sum_{W \ni \mathfrak{n},\ t_{\mathfrak{n}}^{(k)}(W) \ge t} e^{\nu\sigma(W)+\nu_e \operatorname{ext}(W)} \beta_W \le C_\chi \tilde{\rho}_k L^{2k} W_t , \tag{5.14}$$

the sum over the groups of the deviation species with $\mathfrak{n}$ in an interval; $\sigma(W)$ and $\operatorname{ext}(W)$ are those of the step at which W was made. The *cross species* consists of the cross groups which are not attached to a large-field region and contain no part of an insertion attached at an earlier step; on the term without large-field regions these are all the cross groups. Its insertions carry no deviation bound, they need not vanish at the reference data, and they are not contracted. For every $j \le k$, the groups of the cross species present at the step k which descend from those made at the step j satisfy

$$\begin{gathered} \sum_W e^{\theta(W)+\lambda_k(W)\nu\sigma(W)} \|Y_W\| \le 2^{k-j+1} \mathsf{A}_j^\times , \\ \mathsf{A}_j^\times \coloneqq P_j(\mathcal{C}) \Big[C_{\text{cr}} \mathfrak{a}_j e^{-\nu_\times d_j/M} + \mathbf{1}_{\mathbf{R}}(j)\, C_{\text{cr}}^{\mathbf{R}} e^{-p_0(g_j)/4} e^{-\kappa d_j/(6MR_j)} \Big] , \end{gathered} \tag{5.15}$$

with rates $\lambda_k(W) \in [\frac{15}{16}, 1]$ and the decay rate κ of Bałaban's polymers at the **R**-steps (Supplement, Proposition S.10), where d_j is the least ℓ^∞-distance between two loops of the ensemble in units of the scale j, so that $d_j \geq a_j$ by Lemma 3.14, $\mathfrak{a}_j$ is the amplitude of Definition 7.2, and $\mathbf{1}_{\mathbf{R}}(j)$ is 1 at the steps at which the **R**-operation acts and 0 at the others; for a single loop the cross species is empty. On the term without large-field regions the groups in question are the groups of the cross species made at the step j, and the bound holds there without the factor 2^{k-j+1} (Lemma 6.37). A cross group one of whose constituents is attached belongs to the species of the first such constituent and satisfies the bounds of that species.

(iii) *Analyticity.* The insertion of a group W depends on the background field only through its restriction to the dependence set X_W^+. For real U_k in the support of the indicators of the term, the factors which depend on the background extend analytically to $\mathbf{U} = e^{iL^{-k}A'}U_k$, for complex A' in the polydisc of the radii $(1 - \beta(1 - 2^{-(k-m)}))\alpha_{1,m}L^{-m}$ per fine bond on $\Omega_m \setminus \Omega_{m+1}$ (and on Ω_k for $m = k$), and for $k \geq 2$ with the gradient condition of [9, (2.39), p. 261] and in the variable $\mathbf{J}$ of [9, (2.36)]; they are covariant under G-valued gauge transformations, and (5.11) holds there with the deviation of $\mathbf{U}$ from U_k added to $\delta^*_{\mathfrak{n}}$. The same factors are defined, and analytic in the same polydiscs, at the *reference data* of the term, the trivial field 1 in place of U_k, with $\mathbf{J} = 0$ and the outer variables trivial, whether or not the reference data lie in the support of the indicators. On the term without large-field regions the domain and the functions are the following, in the units of the fine lattice. For a configuration U with values in G and a field α on the fine bonds with values in the complexified Lie algebra $\mathfrak{g}^c$ we write $e^{i\alpha}U$ for the configuration $b \mapsto e^{i\alpha(b)}U(b)$. For a bond $b = \langle x, x + e_\mu\rangle$ and a unit vector e, one of the $\pm e_\nu$, let $b + e := \langle x + e, x + e + e_\mu\rangle$, and let the *covariant difference* of α be

$$(\delta_e\alpha)(b) := U(x, x + e)\,\alpha(b + e)\,U(x, x + e)^{-1} - \alpha(b)\,, \tag{5.16}$$

where $U(x, x + e)$ is the value of U on the bond from x to $x + e$. For $n \geq 2$ let $\mathcal{S}_n(U)$ be the set of the pairs $(e^{i\alpha}U, \mathbf{J})$, with fields α and $\mathbf{J}$ on the fine bonds with values in $\mathfrak{g}^c$, such that for all bonds b, unit vectors e, and fine plaquettes p

$$\begin{aligned} |\alpha(b)| < \alpha_{1,n}L^{-n}\,, \qquad |(\delta_e\alpha)(b)| < \alpha_{1,n}L^{-2n}\,, \\ |(e^{i\alpha}U)(\partial p) - 1| < c_F\,\alpha_{0,n}L^{-2n}\,, \qquad |\mathbf{J}(b)| < \alpha_{0,n}L^{-3n}\,, \end{aligned} \tag{5.17}$$

with a constant $c_F \in (0, 1]$; $\mathcal{S}_1(U)$ is the set of the configurations $e^{i\alpha}U$ with $|\alpha(b)| < \alpha_{1,1}L^{-1}$, without a variable $\mathbf{J}$. In the units of the scale n the fine lattice has the spacing $\xi = L^{-n}$ and the complex part is written $e^{i\xi A'}$, so that $\alpha = \xi A'$, and (5.17) are the conditions $|A'|, |\nabla^\xi_U A'| < \alpha_{1,n}$, $|\partial\mathbf{U} - 1| < c_F\alpha_{0,n}\xi^2$, and $|\mathbf{J}| < \alpha_{0,n}$ of [9, (2.39), (2.34), (2.36), p. 261] on the top domain, the second with the constant c_F; a current has the dimension of the third power of an inverse length [8, (1.8), p. 261], whence the factor L^{-3n}. The constant c_F is so small that these conditions imply those on the averaged configurations, [9, (2.35), (2.37)]: Bałaban states that the conditions (i)–(iii) of his spaces with smaller constants imply the condition (iv), by the bounds [8, (1.17), p. 263], which he derives in one line from [7, Proposition 9]. The *space of the step* k is the union of the sets $\mathcal{S}_k(U)$ over the configurations U which are the trivial configuration, the background $U_k(V_k)$ of a block field V_k in the support of the test functions of the term, or a transform of one of these by a G-valued gauge transformation. For a set X of bonds let $\mathcal{S}_n(U)|_X$ be the set defined by the conditions (5.17) on the bonds of X and on the plaquettes and differences which they contain. The insertion of a group W made at a step $j \leq k$ is a holomorphic function of the restriction of $(\mathbf{U}, \mathbf{J})$ to X_W^+, of $\mathbf{U}$

alone for $j = 1$, on the open set

$$\mathcal{D}_W := \bigcup_U \mathcal{S}_j(U)|_{X_W^+} , \tag{5.18}$$

the union over the same configurations U at the step j, and at the later steps the insertion of W is the same function of the configuration and of the current, both in fine units. For a group with one interval, from the site x_W to the site x'_W, and for G-valued gauge transformations v,

$$Y_W(\mathbf{U}^v, R(v)\mathbf{J}) = \pi(v(x_W))\, Y_W(\mathbf{U}, \mathbf{J})\, \pi(v(x'_W))^{-1} , \tag{5.19}$$

where $\mathbf{U}^v(b) = v(b_-)\mathbf{U}(b)v(b_+)^{-1}$ and $(R(v)\mathbf{J})(b) = v(b_-)\mathbf{J}(b)v(b_-)^{-1}$ [8, (1.10), p. 262], and π is the representation of the loop of W; for a group with several intervals each factor of the tensor is transformed in this way.

(iv) *Scalar parts.* $Q_k(\mathfrak{B})$ and $\mathcal{E}_k$ are numbers on each labelled term, independent of V_k and of the outer variables, and

$$\sum_{X \ni x} |e_X^{(j)}| e^{\kappa_e |X|} \le C \tau_j^2 e^{-c_{\text{th}} d^{(j)}(x)} , \qquad |E_{\text{const},k}| \le \sum_{j \le k} \epsilon_j(\mathcal{C}, T) , \tag{5.20}$$

where $d^{(j)}(x)$ is the distance of the box x of the scale $j-1$ from the complement of the carrier of the loop's integral at the step j, and $\epsilon_j(\mathcal{C}, T)$ is the remainder bound of the step j defined in (7.31).

(v) *Uniformity.* The constants depend on the constants of the construction, the characters, and the shape constants $\Lambda(C_i)$, and not on k, K, the torus, the labelled term, or the number of loops; for an ensemble with admissible loops they may be taken to be nondecreasing functions of $c_{\mathcal{C}}$ which depend on Λ_* in place of the $\Lambda(C_i)$, because $\Lambda(C_i)$ enters only through the counts of Lemma 5.5, which increase with it, and a character only through the number c_{π_i} of (1.7) (Section 1.3).

(vi) *Torus consistency.* On the trivial term and at its reference data, let $T \subset T'$ be two tori of the family $\mathfrak{T}$ of Section 7.1. For every insertion and every cluster weight $\varphi_k(\Gamma)$ of an object whose dependence set has diameter less than $\iota_{T,k}/4$, the values on T and on T', at fields which coincide under the covering on the $\iota_{T,k}/4$-neighbourhood of a lift of that set, differ by amounts whose sums rooted at a traversal are at most $C_{\text{rel}} e^{-\iota_{T,k}/(32M)}$ times the right side of (5.11) for the insertions, and at most $C_{\text{rel}} e^{-\iota_{T,k}/(32M)} \mathfrak{a}_k$ for the cluster weights.

(vii) *Geometry and sizes.* For every group W of the term, made at a step $j \le k$, the domain X_W is connected, in the sense that any two of its cubes are joined by a chain of its cubes in which consecutive ones have a point in common, and $\#_j(X_W) + n_j(W)/M \le u_\star \sigma(W)$ with a constant $u_\star \ge 1$; and the domains of two incompatible groups have a point in common. On the term without large-field regions the insertions satisfy in addition two bounds in which their sizes on the whole domain of analyticity enter. Let $\|Y_W\|_{(k)}$ be the supremum of $\|Y_W\|$ over the *space of the step* k, the set of the configurations $(\mathbf{U}, \mathbf{J})$ of (iii) around the real backgrounds in the support of the indicators of the term and around its reference data, and say that a traversal of the step k is *in* an interval of a group if it contains a fine bond of that interval. Then, for every traversal $\mathfrak{n}$ of the step k,

$$\sum_{W \ni \mathfrak{n}} e^{\theta(W) + \lambda_c \nu \sigma(W) + \nu_e \operatorname{ext}(W)} \|Y_W\|_{(k)} \le C_c \rho_k , \qquad \lambda_c := \frac{15}{16} - \frac{1}{8192} , \tag{5.21}$$

the sum over the groups of the deviation species with $\mathfrak{n}$ in an interval. Moreover, for a group W made at the step k let c_W be the number subtracted from its insertion at its production (Section 6.7), which is 0 if nothing was subtracted, and $S_W := \|Y_W\|_{(k)} + |c_W|$; then

$$\sum_{W \text{ at } p} e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)} S_W \, \ell_W \le 2C_{S'}\rho_k \tag{5.22}$$

for every traversal p of the step k and every cube p of π_k, the sum over the groups made at the step k which are *at* p: those with p in an interval, or with $p \subset X_W$. The groups of the cross species and of the wrapping species have domains which are long, and their parts of this sum are small accordingly: for the same p,

$$\begin{aligned} \sum_{W \text{ at } p} e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)} S_W \, \ell_W &\le 2C_{S'}\rho_k \, e^{-\nu_\times d_k/M} \ , \\ \sum_{W \text{ at } p} e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)} S_W \, \ell_W &\le 2C_{S'}\rho_k \, e^{-c_{\text{att}} N^{(k)}_{\min}/M} \ , \end{aligned} \tag{5.23}$$

the first sum over the groups of the cross species and the second over those of the wrapping species which are made at the step k and are at p.

The ensemble is *dressed at the step* k if $N_k = \sum \chi_k(\Omega_k)\mathbf{T}_k[\mathcal{W}_k \exp A_k]$, the sum over the labelled terms of ϱ_k, with a dressed loop $\mathcal{W}_k$ of the step k on each of them.

Lemma 5.3 (From the Tails to the Sum)**.** *Let $\mathfrak{n}$ be a traversal of the step k, and let U be a configuration at which the groups W of the deviation species with $\mathfrak{n}$ in an interval satisfy* (5.13)*, with U in the place of U_k, and* (5.14)*. Then*

$$\sum_{W \ni \mathfrak{n}} e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)} \|Y_W\| \le C_\chi \tilde\rho_k \delta^*_{\mathfrak{n}}(U) \ . \tag{5.24}$$

Proof. Let m_t be the maximum of $|U(\partial p) - 1|$ over $p \in \mathcal{P}_t(\mathfrak{n})$ for $t \ge 1$, and $m_0 := 0$; the sets $\mathcal{P}_t(\mathfrak{n})$ increase with t, so that m_t is nondecreasing. Let W be one of the groups, made at the step j, and $\mathfrak{n}_j \subset \mathfrak{n}$ a traversal of the step j in its interval. Its comparison region at $\mathfrak{n}_j$ lies in Ω_k and within the distance R_t of $\mathfrak{n}$ in the units of the scale k, for $t = t^{(k)}_{\mathfrak{n}}(W)$, so that its fine plaquettes belong to $\mathcal{P}_t(\mathfrak{n})$, and by (5.13) and (5.8)

$$e^{\theta(W)}\|Y_W\| \le b_W \ell_W L^{2j} \, m_t = \beta_W \, m_{t^{(k)}_{\mathfrak{n}}(W)} \ .$$

We write $m_t = \sum_{t' \le t}(m_{t'} - m_{t'-1})$, sum over the groups, and use (5.14) for each t':

$$\sum_{W \ni \mathfrak{n}} e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)} \|Y_W\| \le \sum_{t' \ge 1} (m_{t'} - m_{t'-1}) \, C_\chi \tilde\rho_k L^{2k} W_{t'} = C_\chi \tilde\rho_k L^{2k} \sum_{t \ge 1} m_t w_t \ ,$$

where the last step is a summation by parts, with $W_t - W_{t+1} = w_t$ and $m_t W_{t+1} \to 0$, the numbers m_t being bounded. The right side is $C_\chi \tilde\rho_k \delta^*_{\mathfrak{n}}(U)$ by (5.7) and (5.9). *q.e.d.*

Three comments explain the definition. First, the bound (5.11) separates two kinds of insertions. The first term, the *deviation part*, is small because the background is close to flat; it vanishes when U_k is a pure gauge, by Schur's lemma, which is why it is proportional to δ^*. The other terms are *attached* parts, which carry no deviation and are instead small through the decay away from the large-field regions, the boundary layers, and the wrapping of the torus. For several loops there is a third kind, the insertions of the cross species: the value of

such an insertion at flat data is invariant under the simultaneous conjugation of its factors by G only, and it is not a scalar in general, so that nothing can be subtracted from it, and it is small because a cross group joins two loops which are far apart in the units of every step, which is the content of (5.15) and (5.23). Secondly, no smallness of $\sum_W \|Y_W\|$ over a whole loop is asserted: the sum over families is controlled by the product $\prod_W (1 + e^{\theta(W)}\|Y_W\|)$, whose logarithm is a sum over the traversals of (5.11), so that it costs a factor which is exponential in the number of traversals with a small rate. Thirdly, the step factors κ_j are defined in Section 7 by an explicit formula, and since the normalisation in (5.10) is required to be exactly $\prod_j \kappa_j$, the class is an induction for the normalisation of Definition 1.3.

5.4 The First Step with a Loop Ensemble

Theorem 5.4 (The First Step)**.** *Let L and Λ_* be as in Theorem 1.5, let $\mathcal{C}$ be a loop ensemble at level q whose loops are admissible, and let $\bar{g}$ and q be as in Theorem 1.5 for $\mathcal{C}$. For every K larger than an index $K_{\text{tor}}(\mathcal{C})$, which depends on $\mathcal{C}$ only through $\max_i P_q(C_i)$ and on q and T/u, the density N_1 obtained from $N_0 = W_{\mathcal{C}}\varrho_0$ by the first step of the construction for the target $(K, \bar{g})$ satisfies* (5.2)*, and the ensemble is dressed at the step* 1*.*

The proof uses the first-step manuscript [1] for the analysis of one square loop and of two squares, and consists in checking that each of its statements holds for a general ensemble and for the construction of this paper. We first collect the properties of a general loop that replace those of a square in that analysis; they are consequences of Section 2. Throughout, C is a loop of $\mathcal{C}$, read in fine units on the lattice $\mathbb{T}_K$, with $P_0(C)$ bonds, and ℓ is a length which is at most a quarter of the separation $a_0 \geq L^3 M R_{K'} L^{K'}$ of Lemma 3.14.

Lemma 5.5 (Square Properties of a General Loop)**.** *Let C be a loop with chord–arc constant $\Lambda(C)$.*

(i) The bonds of C which meet a closed ℓ^∞-cube of side s are at most $\Lambda(C) \cdot 2(s+1)$, and the sites of C in a set of ℓ^∞-diameter D are at most $2\Lambda(C)(D+1)+1$.

(ii) For a set S of ℓ^∞-diameter D which meets C, the smallest arc of C which contains $S \cap C$ has at most $4\Lambda(C)D + 2$ bonds, or else contains more than half of C.

(iii) A set of ℓ^∞-diameter less than $r_0(C)$ which meets C meets at most two consecutive segments of C, and on two consecutive segments the chord–arc constant is 2*.*

Proof. (i) Two sites x, y of C in the cube have $|x - y|_\infty \leq s$, hence, by Definition 2.4, arc distance at most $\Lambda(C)s$. The sites of C in the cube therefore lie on an arc of at most $\Lambda(C)s$ bonds measured from any one of them in either direction, that is, on at most $2\Lambda(C)s + 1$ sites, and a bond which meets the cube has an end among them or is one of the two bonds at each end; this gives at most $2\Lambda(C)(s+1)$ bonds for $s \geq 1$, and the count of sites in a set of diameter D is the same argument. (ii) is Lemma 2.6, and (iii) is Lemma 2.10 together with Lemma 2.5 applied to the path formed by two consecutive segments, which is a bent path with two perpendicular arms. *q.e.d.*

Proof of Theorem 5.4. The identity (5.2) at $k = 1$ holds because the first step preserves integrals (Section 5.1). The rest of the proof goes through the analysis of [1] in the order of that paper and states, for each part, the change which a general ensemble or the construction of this paper requires.

The insertion at a fixed cutoff and the extraction. The sourced effective interaction of [1, Sections 2–3] is built from the fine bonds of the loop and the local operators of the step, whose ranges are fixed and do not depend on the loop. The loop enters through three counts: the hull of a cluster on the loop, the number of labels which meet a bond, and the number of traversals within a given distance of a point. For a square these are bounded by the elementary geometry of the square; for C they are bounded by Lemma 5.5: the hull constant $32(1+R_*)$ of [1, (5.24)] becomes $16\Lambda(C)(1+R_*)$ by (ii), and the number of source labels of one traversal which meet a bond, which is at most a constant q^{src} for each of the traversals whose initial sites lie in a set of diameter $2R_*+2$, is at most $q^{\mathrm{src}}(4\Lambda(C)(R_*+1)+1)$ by (i). The constants which depend on these counts are fixed before the coupling, and the conditions on the coupling which contain them depend on $\Lambda(C) \le \Lambda_*$. Every condition on the coupling in this proof which involves the loops is a condition on g_0 or on g_1, both of which are at most γ_q by (3.46), so that it is a requirement on γ_q, depending on $c_{\mathcal{C}}$.

The coefficients. The perimeter and corner coefficients of [1, Sections 5–6] are defined on the straight line and on the bent path of two perpendicular rays through the coarse sites, in the orientations of the square. We define them for every orientation $v \in \mathcal{V} \coloneqq \{\pm e_0, \dots, \pm e_3\}$ on the line, and for every ordered pair (v, w) of perpendicular orientations on the bent path (2.6); the proofs of their convergence and of their tail bounds use only that the path is a line, or two perpendicular rays, through coarse sites, and not which coordinate directions it uses, so they hold with the same constants. The extraction of the regional scalar of the loop is then the following: allocate each cluster of the reference gas to the coarse segment of its source, give each corner c the clusters allocated to the second half of its incoming segment and to the first half of its outgoing segment, and compare the clusters rooted there with those of the bent path B_c. By Lemma 2.10 a cluster which distinguishes C from B_c near c reaches a distance at least $r_0(C) - L/2 - 2R_*$ from its root, and by the tail bounds of the cluster expansion such clusters cost at most $C\tau_0^2 e^{-cr_0(C)/L}$ per root. With κ_C the exponential of the sum of the line coefficients over the segments and of the bent-path coefficients over the corners, this gives

$$\Big| F_{C,\infty} - \log \kappa_1(C,\chi) \Big| \le C\tau_0^2 \Big(\frac{P_0(C)}{L} + |\mathcal{K}(C)| \Big) e^{-cr_0(C)/L} \,, \tag{5.25}$$

in place of the square's bound, where $F_{C,\infty}$ is the bulk limit of the logarithm of the reference partition function of the loop and κ_1 is the step factor of Definition 7.8 at $j=1$. The boundary and periodic comparisons of [1, Section 7 and Appendix F] are made per coarse segment and use the torus only through its smallest period; they hold for C with the sums over the $4n$ coarse segments of a square replaced by the sums over the $P_0(C)/L$ coarse segments of C. The bounds of the leading term then hold with $e^{4C\tau_0^2 P_0(C)/L}$ in place of $e^{C\tau^2(9n+1)}$, and the constant $1+1/c$, which bounds $(n+1)e^{-cn}$ for a square, is replaced by $1+2P(C)/(e\,c\,r(C))$, since $(P_0(C)/L + |\mathcal{K}(C)|)e^{-cr_0(C)/L} \le 2(P(C)/r(C))\,(r_0(C)/L)e^{-cr_0(C)/L}$ by Lemma 2.5.

The retreated carrier and the labelled terms. The analysis of [1, Section 9] on the labelled terms uses the loop through the envelopes of the groups, which are arcs of fewer than half of the traversals or the whole loop read from a corner, through the pins of the groups, and through the lifting of the groups to the universal cover. By Lemma 5.5(ii) the number of cubes of side LM which meet an envelope $\mathcal{H}(W)$ is at most $4 + (1+2/L)|\mathcal{H}(W)|/(LM)$, in place of $8 + |\mathcal{H}(W)|/(LM)$, so that the constants of the envelopes are unchanged. In the pins, the hull of a component γ of a group has at most $(8\Lambda(C)+1)LMn_\gamma$ traversals, in place of $17LMn_\gamma$, and the condition on the contraction rate which V4 imposes becomes $1000\,\eta\,(8\Lambda(C)+2) < 1/128$, which holds for $\eta \le \eta(\mathcal{C}) \coloneqq 1/(1\,200\,000\,\Lambda(\mathcal{C}))$. A small interior group has its contacts within a set of diameter less than $r_0(C)$, so that by Lemma 5.5(iii) they lie on two consecutive segments, and its lifting is the one of the square. The torus conditions of [1] involve the loop

through its diameter and through $4ne^{-c\iota_T} \leq 1$, which reads $(P_0(C)/L)e^{-c\iota_T} \leq 1$; both hold for $K \geq K_{\text{tor}}(\mathcal{C})$, because $3\,\text{diam}(C) < T$ and $\log(P_0(C)/L) \leq \log P_q(C) + K \log L$ while ι_T grows like L^K.

Several loops: the groups. For two squares, [1, Section 10] forms the groups of the joint expansion, which lie on one loop or are cross groups, proves Schur's lemma for each loop separately, and bounds the cross terms by the distance between the squares. We use its definitions and statements for an ensemble of n loops, reading "every other loop" where it has the second square, and we go through the places at which the number of loops enters. By Lemma 3.14 two loops are at distance at least $a_0 = L^{K-q}$ on the fine lattice, so that a set of diameter less than a_0 meets at most one loop. The groups are those of [1, Definition 10.1], with its condition (A2) imposed for every loop. A component γ of a term has a hull H^i_γ on every loop C_i on which it has a contact; a group which is long on a loop contains every component with a contact on that loop; and the interval set $\mathcal{H}_i(W)$ of a group W on C_i is the whole loop if W is long on C_i, and the union of the hulls H^i_γ of its components otherwise. The proofs of [1, Lemmata 10.2–10.5] consider one square at a time and hold for each loop of the ensemble: the source is the product over the loops of the normalised traces of the words of the loops, which is the normalised trace of the tensor product of these words, and every matrix factor belongs to one loop [1, Lemma 10.4]. A group on one loop has one interval [1, Lemma 10.2(d)], while a cross group may have several intervals on a loop. Let $n(W)$ be the number of the cubes of π_1 in the roots of W and in the domains of its cluster occurrences, counted with multiplicity, and n_γ the same number for a component, so that $n(W) = \sum_\gamma n_\gamma$ [1, Section 10.5]. Then W has at most $n(W)$ maximal intervals. Indeed, a maximal interval on a loop C_i is the whole loop, in which case W has a component with a contact on C_i, or it contains the hull H^i_γ of a component; the number of the maximal intervals is therefore at most the number of the pairs of a component of W and a loop on which that component has a contact. The contacts of γ lie within the distance $2L$ of its domain [1, proof of Lemma 10.6(b)], which is a union of at most n_γ cubes of π_1, and a cube of side LM is within the distance $2L$ of at most one loop, because $LM + 4L < a_0$. Hence γ has contacts on at most n_γ loops, which proves the claim.

Several loops: the pins. In the proof of the pins [1, Lemma 10.6] the number of loops enters at two places. The first is the connectedness of the union of the cubes of the components of a group W and of the cubes along its intervals, when W is long on the loops of a set T. Every overlap class of W has a contact on a loop of T, since otherwise it could be split off, as in that proof. Join two loops of T if a class of W has contacts on both of them. If the graph so defined on T were disconnected, let T_1 be one of its connected components, W_1 the union of the classes with a contact on a loop of T_1, and W_2 the union of the others. A class of W_1 has no contact on a loop of $T \setminus T_1$, because it has one on a loop of T_1 and no loop of $T \setminus T_1$ is joined to a loop of T_1, and a class of W_2 has none on a loop of T_1, by its definition. Both parts are nonempty, since W is long on a loop of T_1 and on a loop of $T \setminus T_1$ and therefore has components with contacts on each of them. So W_1 contains every component with a contact on a loop of T_1 and is long on no other loop, and W_2 contains every component with a contact on a loop of $T \setminus T_1$ and is long on no other loop. Replacing W by W_1 and W_2 would therefore give an admissible partition, in contradiction to the definition of the groups as the blocks of the finest one. So the graph is connected. Each class is connected together with the cubes along its hulls, it meets the cubes along a loop of T on which it has a contact, and the cubes along a whole loop form a connected set [1, proof of Lemma 10.6(a)]; hence the union is connected. The second place is the total mass of the groups which are long on some loop. Let $M^{(\rho)}(W)$ be the majorant of a group, the product of the weights $\rho\, m(D)$ of its roots, with the number ρ of [1, (E.11)], of the weights of its cluster occurrences, and

of $1/k(W)!$ for the number $k(W)$ of these occurrences [1, Section 10.4], and let $P_0(C_i) = P_q(C_i)L^{K-q} \geq 4L^{K-q}$ be the number of the fine traversals of C_i. For the groups which are long on exactly the loops of T and contain a class long on C_{i_0}, the computation of [1, proof of Lemma 10.6(c)] for a group long on both squares applies with the loops of T in the place of the two squares: the hulls of the components cover half of each loop of $T \setminus \{i_0\}$, which gives the factor $\exp(-15\nu \sum_{i\in T\setminus\{i_0\}} P_0(C_i)/(LM))$ against the weights $e^{30\nu\mathfrak{s}(\gamma)}$ of the components, and the classes other than the long one are anchored at the traversals of the loops of T. It bounds the weighted mass of these groups by $C\rho\, P_0(C_{i_0}) \exp(-10\nu \sum_{i\in T} P_0(C_i)/(LM))$. Put $x_K \coloneqq 4L^{K-q}$, and suppose that $x_K \geq LM/(10\nu)$, so that $x \mapsto xe^{-10\nu x/(LM)}$ decreases on $[x_K, \infty)$, and that $n\, e^{-10\nu x_K/(LM)} \leq 1$. Summing over the sets $T \ni i_0$ and then over i_0, we obtain

$$\begin{aligned}\sum_{W \text{ long}} M^{(\rho)}(W) e^{4\nu\mathfrak{s}(W)} &\leq C\rho \sum_{i_0} P_0(C_{i_0}) e^{-10\nu P_0(C_{i_0})/(LM)} \prod_{i\neq i_0} (1 + e^{-10\nu P_0(C_i)/(LM)}) \\ &\leq e\, n\, C\rho\, x_K e^{-10\nu x_K/(LM)} ,\end{aligned}$$

the product being at most $\exp(n\, e^{-10\nu x_K/(LM)}) \leq e$. If moreover $e\, n\, x_K e^{-10\nu x_K/(LM)} \leq 2LM/(5e\nu)$, the right side is at most the bound $C'\rho$ which that lemma has for the pair. The three conditions are lower bounds on K in terms of n, q, L, M, and ν, and n is at most the number $(2L^{m_T+q})^4$ of the sites of $\mathbb{T}_q$; we include them in the definition of $K_{\text{tor}}(\mathcal{C})$. The remaining counts of that proof are counts per traversal or per cube, and the traversals whose stencils meet a given cube of π_1 lie within a fixed multiple of LM of it, hence on one loop. The pins therefore hold for the groups of the ensemble, the cross groups included, with a constant C_{pin} which does not depend on the number of loops: for every fine bond and for every cube of π_1,

$$\sum_{W} M^{(\rho)}(W) e^{4\nu\mathfrak{s}(W)} \leq C_{\text{pin}}\, \rho , \tag{5.26}$$

the sum over the groups with that bond in an interval in the first case, and over the groups with that cube in the footprint in the second, where $\mathfrak{s}(W)$ is the size defined below [1, Lemma 10.6(d) and, on the labelled terms, Lemmata 9.27(a) and 10.9].

Several loops: the insertions of the cross groups. The insertion of a cross group W is the tensor

$$Y_W \coloneqq w(W) \int \bigotimes_{I\in\mathcal{I}(W)} \pi_{i(I)}(T_{W,I}(a))\, \mathrm{d}\mu_W(a) , \tag{5.27}$$

where a stands for the variables of the roots of W, μ_W is the product of the measures of these roots, $w(W)$ is the weight of the cluster occurrences of W, and $T_{W,I}(a)$ is the ordered product of the reconstructed links along the interval I; the word of a family with these insertions, in the sense of Section 5.3, is the expression which [1, proof of Lemma 10.5] integrates. That paper bounds the integral of a product of traces of such words by the majorant $M^{(\rho)}(W)$ [1, Lemmata 10.4 and E.3], and its proof gives the same bound for Y_W in the norm (5.5). The proof has three steps: a finite difference of the source in the selected variables is bounded by its supremum times the total variation of the measures; for a root with an odd number of selected variables and no selected potential, the part of the difference which is homogeneous of that degree integrates to zero, the kernel being even; and for such a root with selected potentials a Cauchy difference in a parameter of the potentials is taken [1, (E.6), (E.60), (E.61), and the proof of Lemma E.3]. All three are linear in the source, so that they apply to the function $a \mapsto \bigotimes_I \pi_{i(I)}(T_{W,I}(a))$, whose values are tensors. The suprema are those of the differences and of their remainders, which by the product rule are sums of elementary tensors in which some of the links are replaced by derivatives. The number of these tensors is that of

the scalar case, the norm (5.5) of an elementary tensor is at most the product of the operator norms of its factors, and these are bounded as in [1, proof of Lemma 10.4], the links which are not differentiated being unitary. Since the norm of an integral is at most the integral of the norm against the total variation, we conclude that $\|Y_W\| \le M^{(\rho)}(W)$. The cross groups have the reference scalar 0, and their flat values need not vanish [1, Section 10.3]. The cross exponent is the sum over the clusters which meet at least two loops; a cluster may meet three or more, and it is in the bound of the exponent, not in the exponent itself, that such a cluster is charged to one pair of the loops which it meets. On every labelled term the total of the cross groups is at most $C_\times \rho \Delta_\times$, with $\Delta_\times$ the sum of $e^{-c_\times \delta_a/(LM)}$ over the fine traversals a and δ_a the distance of a from the other loops [1, Lemmata 10.7(a) and 10.9]; since a traversal of the scale 1 consists of L fine traversals at L times its distance in fine units, and $\nu_\times \le c_\times$, this is (5.15) at $k = j = 1$. The bounds of the class on the term without large-field regions are proved in Corollary 5.8, for the cross groups together with the others.

The construction of this paper. Two features of the construction differ from those assumed in [1]. The thresholds are those of the tube at $\hat{g}_0$: by Lemma 3.5 the radius of the fluctuation cutoff is $t_0 \coloneqq \vartheta_0/g_0 \in [\rho^{-1/2} r_0, r_0]$, and every statement of [1] which uses the radius as an upper bound on a passed field holds a fortiori, while each one which uses it as a lower bound holds with the constant in its exponent divided by $\rho \le 3/2$; the only condition which changes is the one behind $e^{-c_{\mathrm{ph}} r^2} \le g_0$ in [1, (9.29)(iii)], with $c_{\mathrm{ph}} = 9 c_n \alpha_G/128$, which at the radius t_0 reads $(9 c_n \alpha_G/(64\rho)) A_1^2 \log g_0^{-2} \ge 1$ and is a smallness condition on the coupling (Supplement, Proposition S.7). The test functions of the plaquettes are soft: by Lemma 3.8 a term of the decomposition is supported where the passed plaquettes are below $(1 + \kappa_s)$ times the threshold, and every bound of [1] drawn from a passed test function holds at the threshold multiplied by $1 + \kappa_s$ (Supplement, Proposition S.2).

The groups of the first step: domains, sizes, and interactions. We verify the clauses of Definition 5.2 which concern the geometry of the groups and their interactions, on a labelled term. A group W of [1, Definition 3.24], formed for the carrier of the term [1, Lemma 9.21(a)], consists of its roots, of the occurrences of clusters attached to them, and of its intervals, whose union is $\mathcal{H}(W)$: one interval if W lies on one loop, and at most $n(W)$ maximal intervals in general, as was shown above. Its *constituent domains*, the roots and the domains of the occurrences, are connected unions of cubes of π_1, and they are the localisation domains of W in the sense of (a). With $n(W)$ the number of the cubes of the constituent domains, counted with multiplicity, as above, and $|\mathcal{H}(W)|$ the number of the fine bonds of the intervals, the size of the group is $\mathfrak{s}(W) = n(W) + |\mathcal{H}(W)|/(LM)$ [1, (E.17)], and we put $\sigma(W) \coloneqq \mathfrak{s}(W)$, so that $\sigma(W) \ge 1$. The domain X_W consists of the cubes of the constituent domains and of the cubes of π_1 which contain a site of an interval, each site of the fine lattice being counted to one cube of the partition [1, Lemma E.8]. It is connected in the sense of clause (vii): the proof of [1, Lemma E.8(b)] shows this for one loop, since consecutive bonds of the interval lie in equal or adjacent cubes, a constituent domain is connected, the domains of an occurrence are linked by incompatibilities, an occurrence is incompatible with a root, and every root has a contact on the interval within the distance $2L$; for a cross group which is long on no loop the same proof applies [1, proof of Lemma 10.6(a)], and for the others the connectedness was shown in the paragraph on the pins. Two groups are incompatible only if their intervals have a bond in common or if two of their constituent domains have a cube or a wall in common [1, proof of Lemma E.8(d)], and in both cases their domains have a common point. For the count, a straight piece of n_i fine bonds of a loop has $n_i + 1$ collinear sites, which lie in at most $n_i/(LM) + 2$ cubes of π_1, and by Lemma 2.8 an arc of n fine bonds consists of at most $n/s_0 + 2$ straight pieces, where $s_0 \ge L^3 M$ is the least number of fine bonds of a segment (Lemma 3.14); an interval of n fine bonds therefore contributes at most $(1 + 2/L) n/(LM) + 4$

cubes to X_W, and it meets at most $n/L + 2$ traversals of the step 1. Since W has at most $n(W)$ maximal intervals,

$$\#_1(X_W) + \frac{n_1(W)}{M} \le n(W) + \Big(2 + \frac{2}{L}\Big)\frac{|\mathcal{H}(W)|}{LM} + \Big(4 + \frac{2}{M}\Big)n(W) \le 11\,\sigma(W)\,, \qquad (5.28)$$

which is the bound of clause (vii) for the groups of the first step, with 11 in the place of $u_\star$. The constant of the class is the larger number $u_\star := (2 \cdot 3^d + 2^{d+4} + 17)13^d\Lambda_*$, which the groups made by the **R**-operation require (Supplement, Proposition S.10(viii)); the groups made at a step $j \ge 2$ by the expansion of the step satisfy the bound with 1 in its place, by the definition of their size (Section 6.6). The interactions are those of the reference gas of the term, $Q_1(\mathfrak{B}) = \exp(-\sum_\Gamma \Phi^\circ(\Gamma))$, the sum over the clusters which contain a group incompatible with a member of $\mathfrak{B}$, and with $\theta(W) := \sum|\Phi^\circ(\Gamma)|$, the sum over the clusters which contain a group incompatible with W, we have $|Q_1(\mathfrak{B})| \le \prod_{W\in\mathfrak{B}} e^{\theta(W)}$ and $\theta(W) \le \nu\sigma(W)/16$ [1, Lemma 9.21(d)], with the constant ν which is called η there; this is clause (i). Finally the insertion of W depends on the background only through its restriction to the dependence set X_W^+. Indeed, every bond of the background which the insertion consults lies in the host of one of its primitives, in the stencil of an entry of the Hessian at a bond of a constituent domain, or in the stencil of a bond of an interval [1, Section E.10.4]. A host is a rectangle of at most four cubes of side M of the fine lattice in each direction which contains its base and, in its interior, the bonds of the background which its primitive consults [1, (3.36), (3.75)]; hence the largest distance $R_{\rm h}$ from the base of a host to a point of it is at most $4M$, and since the hosts of the group have their bases within $R_{\rm h}$ of its constituent domains, they lie within $2R_{\rm h} \le 8M$ of these domains [1, Lemma E.8]. The stencil of an entry of the Hessian lies within $3L$ of its bond [1, Lemma B.3], and the stencil of a bond of an interval within $2L$ of its initial site [1, Lemma E.8]. All these bonds are therefore within $\max(8M, 3L) = 8M$ fine units of X_W, because $M \ge 8L^2$ [1, Section 3.7].

The deviation of a single insertion. On the term without large-field regions we now prove the bound (5.13) and the tail clause (5.14) at $k = 1$, with $\ell_W = 1$. On this term the carrier of the integral of the loops is the whole region of integration and there are no exterior data, so that the two operators Υ_W^0 and Υ_W^1 of [1, Lemma 9.26] coincide; the insertion of a group W on one loop is this operator Υ_W, and $\|\Upsilon_W\| \le K_\chi \mathfrak{d}_W$ with $\mathfrak{d}_W := \sup_{h\in G}|c_W(U_1; h) - a_W\chi(h)|$, the supremum of the difference between the closed contraction of the group at the background U_1 and at the reference data [1, Lemma 9.26(a)]. Let Q_W be the comparison rectangle of the group, the smallest coordinate rectangle of the fine lattice which contains its constituent domains, its interval, and every bond of the background which the insertion consults [1, Sections E.7.4 and E.10.4], and $\delta_{Q_W}(U) := \max_{p\subset Q_W}|U(\partial p) - 1|_\sigma$. Let W be a group of the deviation species, that is, a group on one loop which embeds in the torus (Section 5.3). The set which Q_W is to contain lies in the dependence set X_W^+, and this lies in the comparison regions of W, which are boxes that lift injectively to the universal cover of the torus; hence Q_W is defined, it lies in each comparison region, and it lifts injectively. For a group whose rectangle lifts, $\mathrm{diam}(Q_W) \le C_{\rm box}LMn(W) \le C_{\rm box}LM\mathfrak{s}(W)$ [1, Section E.7.4]. For such a group we claim

$$\mathfrak{d}_W \le C'_\chi\,\mathfrak{s}(W)^3\,\delta_{Q_W}(U_1)\,M^{(\rho)}(W)\,, \qquad (5.29)$$

where $M^{(\rho)}(W)$ is the majorant of the group with the weight ρ of [1, (E.11)] for each root [1, proof of Proposition 9.40] and C'_χ is a constant of that proof and of L, M. The first-step manuscript proves this bound with $\delta_{Q_W}(U_1)$ replaced by its bound $C\varepsilon_1$, for the groups with $n(W) < b/(4C_{\rm box}LM)$ on the histories on which the loop is far from the large-field region [1, (E.74) and Proposition 9.40]. Its proof uses the restriction on $n(W)$ and on the history for three purposes: to place Q_W in the region Λ_1 and outside S_1, which here is no condition; to

lift Q_W injectively, which holds because W embeds; and to bound $\delta_{Q_W}(U_1)$ by $C\varepsilon_1$, which we do not do [1, Sections E.7.4 and E.10.3]. The argument is then the following. Suppose first that $C_{\mathrm{tg}} \operatorname{diam}(Q_W)^2 \delta_{Q_W}(U_1) \le \tau_{\mathrm{b}}/2$, the phase test of [1, Section E.8.1]. In the tree gauge of the lifted rectangle Q_W rooted at the initial site of the interval the background is e^{iB_W} with $\|B_W\|_\infty \le C_{\mathrm{tg}} \operatorname{diam}(Q_W)^2 \delta_{Q_W}(U_1)$ [1, Lemma E.21 and (E.65)], a lemma which requires of the rectangle only that it lifts injectively. The contraction is defined at the backgrounds $U_{W,t} := e^{itB_W}$, $0 \le t \le 1$, because only the bonds of Q_W are consulted and the charts of the rows hold there by the phase test [1, Section E.8.1; proof of Proposition 9.40]; it equals $c_W(U_1; h)$ at $t = 1$, by this locality and by gauge covariance, and $a_W \chi(h)$ at $t = 0$ [1, (E.71)]; and its derivative along the path is at most $C_\chi \|B_W\|_\infty (n(W) + |\mathcal{H}(W)|) M^{(\rho)}(W)$ [1, (E.46), (E.73); proof of Proposition 9.40]. Integrating over t and using $n(W) + |\mathcal{H}(W)| \le CLM\mathfrak{s}(W)$ [1, Section E.11.2] and the bound of the diameter, we obtain (5.29) with the constant $C_\chi C_{\mathrm{tg}} C_{\mathrm{box}}^2 C (LM)^3$. If the phase test fails, then $1 < (2C_{\mathrm{tg}}/\tau_{\mathrm{b}})(C_{\mathrm{box}} LM\mathfrak{s}(W))^2 \delta_{Q_W}(U_1)$, and the bounds $|c_W(U_1; h)| \le M^{(\rho)}(W)$ and $|a_W| \le M^{(\rho)}(W)$ of the two contractions [1, (E.73); proof of Proposition 9.40] give $\mathfrak{d}_W \le 2M^{(\rho)}(W)$, which is at most the right side of (5.29) with the constant $(4C_{\mathrm{tg}}/\tau_{\mathrm{b}})(C_{\mathrm{box}} LM)^2$, since $\mathfrak{s}(W) \ge 1$. This proves the claim.

We translate (5.29) into the quantities of the class. Let $\mathfrak{n}$ be a traversal of the step 1 in the interval of W; it contains a fine bond of the interval, which lies in Q_W. Every point of X_W is within the distance LM of Q_W, hence within $C_{\mathrm{box}} LM\mathfrak{s}(W) + LM$ fine units of $\mathfrak{n}$. The radius $R_t = C_{\mathrm{box}} M(t+1)$, in the units of the scale 1, is $C_{\mathrm{box}} LM(t+1)$ fine units, which is at least that distance for $t = \lceil \mathfrak{s}(W) \rceil$; so $t_{\mathfrak{n}}(W) \le \lceil \mathfrak{s}(W) \rceil$. The neighbourhood $B_\infty(\mathfrak{n}, R)$ of a straight traversal is a coordinate rectangle, and Q_W is the smallest coordinate rectangle which contains a set within $8M$ fine units of X_W, by the last sentences of the preceding paragraph; since $X_W \subset B_\infty(\mathfrak{n}, R_{t_{\mathfrak{n}}(W)})$ and one step $C_{\mathrm{box}} LM$ of the radii exceeds $8M$, the rectangle Q_W lies in the comparison region $B_\infty(\mathfrak{n}, R_{t_{\mathfrak{n}}(W)+1})$. Therefore, by (5.8) with $\ell_W = 1$ and $j = 1$, and with a constant c_σ which compares the two norms of matrices,

$$e^{\theta(W)} \|Y_W\| \le b_W \operatorname{Dev}_{\mathfrak{n}}(W; U_1) \,, \qquad b_W := c_\sigma K_\chi C'_\chi \, e^{\theta(W)} \mathfrak{s}(W)^3 M^{(\rho)}(W) \, L^{-2} \,,$$

which is (5.13), with $\beta_W = b_W L^2$. For the tail clause we note first that $t^{(1)}_{\mathfrak{n}}(W) = t_{\mathfrak{n}}(W) + 1 \le \lceil \mathfrak{s}(W) \rceil + 1$, so that $t^{(1)}_{\mathfrak{n}}(W) \ge t$ implies $\mathfrak{s}(W) > t - 2$; secondly that the comparison regions of W at two traversals of its interval lie within $C_{\mathrm{box}} M(\mathfrak{s}(W) + 3)$ of one of them and the two traversals within $C_{\mathrm{box}} M\mathfrak{s}(W) + 2$ of each other, in the units of the scale 1, so that $\operatorname{ext}(W) \le C_{\mathrm{box}}(2\mathfrak{s}(W) + 4)$ and $\nu_e \operatorname{ext}(W) \le \nu\mathfrak{s}(W) + 2\nu$, because $\nu_e \le \nu/(2C_{\mathrm{box}})$ (Appendix A); and thirdly that a group with $\mathfrak{n}$ in its interval has one of the L fine bonds of $\mathfrak{n}$ in its interval. The groups whose interval contains a given fine bond satisfy $\sum_W M^{(\rho)}(W) e^{4\nu\mathfrak{s}(W)} \le C_{\mathrm{pin}} \rho$ [1, Lemma 9.27(a), for the weight ρ as in the proof of Proposition 9.40]. With $\theta(W) \le \nu\mathfrak{s}(W)/16$ the exponent of a group in the tail clause is at most $2\nu + (2 + \frac{1}{16})\nu\mathfrak{s}(W)$, which leaves the rate $\frac{31}{16}\nu$ of the weight $e^{4\nu\mathfrak{s}}$, and for $s \ge \max(1, t-2)$

$$s^3 e^{-\frac{31}{16}\nu s} \le \Big(\sup_{s>0} s^3 e^{-\frac{13}{8}\nu s} \Big) e^{-\frac{5}{16}\nu(t-2)} = \Big(\frac{24}{13e\nu} \Big)^3 e^{\frac{5}{8}\nu} \, e^{-\frac{5}{16}\nu t} \le \Big(\frac{24}{13e\nu} \Big)^3 e^{\frac{5}{8}\nu} \, e^{-\eta_w t} \,,$$

since $\eta_w \le \frac{5}{16}\nu$ (Appendix A). Hence, for every traversal $\mathfrak{n}$ of the step 1 and every $t \ge 1$,

$$\sum_{W \ni \mathfrak{n},\ t^{(1)}_{\mathfrak{n}}(W) \ge t} e^{\nu\sigma(W) + \nu_e \operatorname{ext}(W)} \beta_W \le L \, c_\sigma K_\chi C'_\chi C_{\mathrm{pin}} \Big(\frac{24}{13e\nu} \Big)^3 e^{\frac{21}{8}\nu} \rho \, e^{-\eta_w t} \,,$$

and $e^{-\eta_w t} \le w_t \le W_t$ by (5.9). Since $\rho \le (1 + \beta_0)^6 \rho_1$, by the conversion of the parameters at g_0 into those at g_1 described below, this is (5.14) at $k = 1$, where $\tilde{\rho}_1 = \rho_1$, for every $C_\chi \ge L^{-1}(1 + \beta_0)^6 c_\sigma K_\chi C'_\chi C_{\mathrm{pin}} (24/(13e\nu))^3 e^{21\nu/8}$.

Assembly. With these changes, the main theorem of [1] and its form on the labelled terms give, on every labelled term, the contribution of the ensemble as $\prod_i \kappa_1(C_i, \chi_i) e^{\mathcal{E}_1} [\prod_i f_1^{(i)} + \sum_{\mathfrak{B} \neq \emptyset} Q_1(\mathfrak{B}) w_1(\mathfrak{B})]$, with the insertions bounded per fine traversal by the deviation of the background and the attached parts, the interactions resummed by the cluster expansion against the reference gas, and the scalar exponent the sum of the carrier part, which is a sum of local terms, and of the torus and corner part, which is a constant. The bounds (i)–(iv) of Definition 5.2 at $k = 1$ follow by summing the bounds of [1] over the L fine traversals of each traversal of the scale 1, which multiplies the constants by L and converts the decay lengths LM in fine units into M in units of the scale 1, and by the inequality $g_0 \leq (1 + \beta_0) g_1$ of Lemma 3.2, which converts the parameters at g_0 into those at g_1. Clause (i) and the geometry of clause (vii) were verified in the paragraph on the groups of the first step; on the term without large-field regions, the bound (5.13) and the tail clause (5.14) of the deviation species were proved in the preceding paragraph, and clause (iii) with the bounds (5.21), (5.22), and (5.23) is Corollary 5.8 below, for all the groups of that term. Clause (vi) holds at $k = 1$ with vanishing differences: on the trivial term every insertion and every cluster weight of the first step is a finite sum of absolutely convergent walk series and parameter integrals in local data, each term of which is the same expression on a lift [1, Sections E.9.2 and E.10.4]. The constants depend on the loops only through $\Lambda(C_i)$, $P(C_i)/r(C_i)$, and the characters, and on the number of loops not at all, because every count above is a count per traversal or per cube, and a cube of side less than a_0 meets one loop. *q.e.d.*

5.5 The First Step at Complex Backgrounds

The clauses (iii) and (vii) of Definition 5.2 concern the insertions as functions on the space of the step, and at $k = 1$ this is more than the first-step manuscript contains. All its estimates are at real backgrounds, and several of its proofs use that the Hessian of the step is a positive real matrix, that the Gaussian measures are positive, and that the links of the background are unitary. We now extend them to the configurations $\mathbf{U} = e^{i\alpha} U$ of the space $\mathcal{S}_1(U)$ of (5.17), on the term without large-field regions. Let h be the invariant norm of [1] on $\mathfrak{g}$, $h_{\mathbb{C}}(x + iy)^2 := h(x)^2 + h(y)^2$ its Hermitian extension to $\mathfrak{g}^c$ [1, Lemma 9.25], and $c_h \geq 1$ a constant with $h_{\mathbb{C}}(X) \leq c_h |X|$ on $\mathfrak{g}^c$; the radius of the space in this norm is

$$\varrho_1 := c_h \, \alpha_{1,1} L^{-1} \, . \tag{5.30}$$

The extension costs each group a factor which is exponential in its size, with a rate proportional to ϱ_1, and since ϱ_1 tends to zero with the coupling the factor is absorbed by the weights of the groups. Two of the choices of constants of [1] leave no room for it, however, and we begin with the linear algebra and with these two choices.

Lemma 5.6 (Complex Symmetric Matrices)**.** *Let $H = R + iS$ be a complex symmetric matrix on $\mathbb{C}^n$, with R and S real symmetric, $R \geq \gamma' > 0$ as a quadratic form, and $\|H\| \leq m$ in the operator norm of the Hermitian product $\langle \cdot, \cdot \rangle$.*

(i) For every principal submatrix H_E of H and every $x \in \mathbb{C}^E$, $\mathrm{Re}\langle x, H_E x \rangle \geq \gamma' \|x\|^2$; in particular H_E is invertible and $\|H_E^{-1}\| \leq 1/\gamma'$.

(ii) With $q := (1 - \gamma'^2/m^2)^{1/2}$ we have $\|1 - (\gamma'/m^2) H\| \leq q$.

(iii) $1 \leq |\det H| / \det R \leq \exp(n \|S\|^2 / (2\gamma'^2))$.

(iv) $\int_{\mathbb{R}^n} e^{-\frac{1}{2} x^{\mathsf{T}} H x} \, \mathrm{d}x = (2\pi)^{n/2} (\det H)^{-1/2}$, with the branch of the root which is continuous on the convex set of the complex symmetric matrices with pos-

itive real part and positive on the real ones; and for the normalised integral $E_H[F] \coloneqq (2\pi)^{-n/2}(\det H)^{1/2}\int F(x)e^{-\frac12 x^{\mathsf T}Hx}\,\mathrm{d}x$ *and every measurable* F, $|E_H[F]| \le e^{n\|S\|^2/(4\gamma'^2)}E_R[|F|]$.

(v) If H_0 *is real symmetric with* $\gamma \le H_0 \le m_0$ *and* $\|H-H_0\| \le s < \gamma$, *then* $R \ge \gamma - s$, $\|S\| \le s$, *and* $\|H\| \le m_0 + s$.

Proof. (i) For $x \in \mathbb{C}^n$ the numbers $\langle x, Rx\rangle$ and $\langle x, Sx\rangle$ are real, because R and S are Hermitian, so that $\mathrm{Re}\langle x, Hx\rangle = \langle x, Rx\rangle \ge \gamma'\|x\|^2$; for a principal submatrix we apply this to the vectors supported in E. Then $\gamma'\|x\|^2 \le \|x\|\|H_E x\|$ gives the inverse. (ii) With $m' \coloneqq m^2/\gamma'$,

$$\|(1-H/m')x\|^2 = \|x\|^2 - \frac{2}{m'}\,\mathrm{Re}\langle x, Hx\rangle + \frac{\|Hx\|^2}{m'^2} \le \Big(1 - \frac{2\gamma'^2}{m^2} + \frac{\gamma'^2}{m^2}\Big)\|x\|^2\,,$$

by (i) and $\|Hx\| \le m\|x\|$. (iii) We have $H = R^{1/2}(1+iK)R^{1/2}$ with $K \coloneqq R^{-1/2}SR^{-1/2}$ real symmetric and $\|K\| \le \|S\|/\gamma'$, and $|\det(1+iK)| = \prod_k (1+\lambda_k^2)^{1/2}$ over the eigenvalues λ_k of K, which lies between 1 and $\exp(\frac12\sum_k \lambda_k^2)$. (iv) Both sides of the identity are holomorphic in the entries of S on the set where $R > 0$, on which $\det H \ne 0$ by (i), and they agree for $S = 0$; the set is convex, so they agree on it. For real x, $|e^{-\frac12 x^{\mathsf T}Hx}| = e^{-\frac12 x^{\mathsf T}Rx}$, and the bound follows from (iii). (v) The entrywise real and imaginary parts of a matrix A are $(A \pm \bar A)/2$ and $(A - \bar A)/(2i)$, and $\|\bar A\| = \|A\|$; applied to $A = H - H_0$, whose real part is $R - H_0$ and whose imaginary part is S, this gives the three bounds. *q.e.d.*

Two choices of constants. The constants of [1] are fixed in a chain, each from the earlier ones and all of them before the coupling [1, Section B.8]. We keep the chain and its order and change two of its links.

(K1) Let $\gamma \le H \le h_+$ be the bounds of the Hessian of the step [1, (3.56)]. Put $\gamma_c \coloneqq \gamma/2$ and $h_c \coloneqq h_+ + \gamma/2$, and compute from (γ_c, h_c) every constant which the chain computes from (γ, h_+): $s_H = 2h_c$, $q_H = 1 - \gamma_c/s_H$, and with them n_*, $C_{\mathrm w}$, δ_M, and $A_* \coloneqq h_c(\gamma_c^{-1/2}+1)$ [1, Section 3.5]. With $C_{\mathrm d}^K \coloneqq d_G \max(1/\gamma_c + 2A_*^2/\gamma_c^2,\ 2h_c^2/\gamma_c)$, where $d_G = \dim G$, require

$$4C_{\mathrm d}^K \delta_M^2 (LM+1)^4 \le \frac14\,, \qquad \alpha_G \le \min\Big(\frac{\gamma_c}{2},\ \frac{\gamma_c^2}{16A_*^2},\ \frac{1}{8C_{\mathrm d}^K(LM+1)^4}\Big)\,, \tag{5.31}$$

in the place of the conditions $4C_{\mathrm d}\delta_M^2(LM+1)^4 \le \frac12$ and $\alpha_G \le (8C_{\mathrm d}(LM+1)^4)^{-1}$ of [1, (3.1) and Lemma 3.16], which together bound the exponent of the Gaussian payment by $N_1(Z)$ with no room to spare.

(K2) The local maps of [1, Lemmata B.7 and B.9], namely the row solution, the reconstruction, the relative means, the logarithms of the Haar density and of the Jacobian of the constraint, and the primitives, are compositions of exponentials, block averages, a fibre inverse, and a chart logarithm. They are stated for the backgrounds whose links on a stencil are $e^{i\phi(b)}$ with $\|\phi\|_\infty \le \tau_L/2$ in a suitable gauge, a compact set $\mathcal{K}_L$ of families of links, and their constants are suprema over $\mathcal{K}_L$. These maps are holomorphic in the links on an open set which contains $\mathcal{K}_L$; hence there is $\rho_0 \in (0,1]$, depending on L and G only, such that they are holomorphic on the set $\mathcal{V}_{\rho_0}$ of the families of complex matrices within the operator distance ρ_0 of $\mathcal{K}_L$, bond by bond, and bounded on its closure. We take the constants of the two lemmata as suprema over the closure of $\mathcal{V}_{\rho_0}$, and every later constant by the formulae of [1].

Both are choices made before the coupling. The first makes α_G smaller and M larger, the second replaces constants by larger ones, and the conditions of [1] on these constants are monotone in these directions: those on M are lower bounds or conditions of divisibility, α_G enters through upper bounds and through conditions on the coupling, and the radius ϑ_* of the charts, which is chosen after the constants of (K2), through upper bounds [1, Sections 3.5 and B.8, Lemmata 3.16 and B.7]. The statements of [1] therefore hold with the changed constants, for a smaller coupling.

Proposition 5.7 (The Insertions of the First Step at Complex Backgrounds)**.** *Consider the term without large-field regions, with the constants of [1] chosen with (K1) and (K2), and let γ be sufficiently small. Let U be the background U_1 of a block field in the support of the term, or the trivial configuration, and let W be a group on one loop. Then the insertion Υ_W of W extends to a holomorphic function of the restriction of $\mathbf{U} = e^{i\alpha}U$ to the dependence set X^+_W, on the set $\mathcal{S}_1(U)|_{X^+_W}$; the extension is given by a formula in $\mathbf{U}$ which does not involve U, it satisfies the covariance law* (5.19)*, and*

$$\|\Upsilon_W(\mathbf{U})\| \le 2K_\chi\, e^{C'\mathfrak{s}(W)\varrho_1} M^{(\rho)}(W)\,, \qquad |a_W| \le M^{(\rho)}(W)\,, \qquad C'\varrho_1 \le \frac{\nu}{8}\,, \tag{5.32}$$

where $C' \coloneqq 8C_\pi LM$ with the constant C_π of [1, Lemma 9.25], a_W is the reference scalar of the group, and $M^{(\rho)}(W)$ is the majorant of (5.26)*. For a cross group W the insertion Y_W of* (5.27) *extends in the same way to a holomorphic function on $\mathcal{S}_1(U)|_{X^+_W}$, each factor of which satisfies the covariance law* (5.19) *at the ends of its interval, and $\|Y_W(\mathbf{U})\| \le e^{C'\mathfrak{s}(W)\varrho_1} M^{(\rho)}(W)$ in the norm* (5.5)*.*

Proof. Step 1: what has to be shown. The insertion is $\Upsilon_W = X_W - a_W$ with an operator X_W whose normalised traces against $\pi(h)$, $h \in G$, are the closed contractions $c_W(U;h)$ of the group [1, Lemma 9.26], and $\|X\| \le K_\chi \sup_{h\in G} |\dim(\pi)^{-1} \operatorname{tr}(X\pi(h))|$ for every operator X on the space of π [1, Lemma 9.26(a)], an inequality which does not involve the background. The contraction is built from three kinds of objects: maps of the background on bounded stencils, which are the entries of the Hessian [1, Lemma 3.3] and the maps of (K2); integrals over real fluctuation variables, whose cutoffs are functions of these variables only; and sums over selections, walks, and clusters. An integral of a holomorphic integrand with an integrable majorant which is locally uniform in the parameter is holomorphic, and so is a locally uniformly convergent series of holomorphic functions. It therefore suffices to define each object at $\mathbf{U}$ by its formula and to bound it by the majorant which [1] has for it at U, up to the factors of (5.32), uniformly on $\mathcal{S}_1(U)$.

Step 2: the charts. The local maps are evaluated at U in a site gauge in which the phases of the links on the stencil are at most $\tau_L/2$ [1, Lemmata B.7 and 9.13(c)]. In this gauge $\mathbf{U}(b) = e^{i\alpha'(b)}U(b)$ with $\alpha'(b)$ a conjugate of $\alpha(b)$ by an element of G, so that $h_{\mathbb{C}}(\alpha'(b)) < \varrho_1$. Since the operator norm of $X \in \mathfrak{g}^c$ is at most $\sqrt{2N}h_{\mathbb{C}}(X)$ for $G \subset U(N)$, and $\|e^{iX}-1\| \le 2\|X\|$ for $\|X\| \le \log 2$, every link of $\mathbf{U}$ on the stencil is within $\rho_0/2$ of the corresponding link of U once $\varrho_1 \le r_e \coloneqq \rho_0/(4\sqrt{2N})$. For $\varrho_1 \le r_e/2$ the stencils of $\mathbf{U}$ therefore lie in $\mathcal{V}_{\rho_0}$, with room for discs of radius $r_e/2$ in each variable, and by (K2) the local maps and the primitives are holomorphic at $\mathbf{U}$ and satisfy the bounds of [1, Lemmata B.7 and B.9] with the changed constants. In particular the potentials of the hosts satisfy the bound $C_0 g_0 (1+2r_0)^3 e^{-\kappa d_0(Y)}$ of [1, Lemma 9.13(d)] at $\mathbf{U}$, with the radius r_0 of the cutoff. The terms of the gauge-fixing functional need an argument of their own, because their bounds at a real background use that each summand is nonnegative [1, Lemma B.7(iv)]. For small $x, y \in \mathfrak{g}^c$ let $F(x,y) \coloneqq -i\log(e^{ix}e^{iy})$, which is holomorphic with $F(x,0) = x$ and $F(0,y) = y$. On a stencil, in the original gauge, a phase q at the background $\mathbf{U}$ is the phase $\tilde{q} \coloneqq F(q,\alpha)$ at the background

U, since $e^{iq}\mathbf{U} = e^{i\tilde{q}}U$, and we define the relative mean at $\mathbf{U}$ by $M_x^{\mathbf{U}}(q) \coloneqq M_x(\tilde{q})M_x(\alpha)^{-1}$ with the relative mean M_x at U of that lemma, which is holomorphic on its ball of radius ϑ_*; there is $r_{\text{iv}} > 0$, depending on G, L, and ϑ_*, such that for $h_{\mathbb{C}}(\alpha) < r_{\text{iv}}$ and q in the ball of radius $\vartheta_*/2$ both $\tilde{q}$ and α lie in that ball. The gauge term $\sum_x(1 - \frac{1}{2}\operatorname{tr}(M_x^{\mathbf{U}}(q) + M_x^{\mathbf{U}}(q)^{-1}))$ is then holomorphic in (q, α) and bounded by a constant, it is the term of [1] for real α, and it vanishes at $q = 0$ together with its first derivative in q: indeed $M_x^{\mathbf{U}}(0) = 1$ because $F(0, \alpha) = \alpha$, and the derivative of $\frac{1}{2}\operatorname{tr}(M + M^{-1})$ at $M = 1$ is zero. Cauchy's estimates bound its derivatives in q, and the constants of (K2) are taken at least as large as these bounds. We put $r_H \coloneqq \min(r_e, r_{\text{iv}})$ and require $\varrho_1 \le r_H/2$.

Step 3: the Hessian. Let H and H_U be the Hessians of the step at $\mathbf{U}$ and at U, compressed to the region of integration. An entry $H(b, c)$ depends on the links within the range $3L$ of b [1, Lemma B.3], of which there are n_e, a number of L; by Step 2 it is holomorphic in α on the polydisc of radius r_H and bounded there by a constant C_{ent}. By Cauchy's estimate on discs of radius $r_H/2$ each entry differs at $\mathbf{U}$ and at U by at most $(2n_eC_{\text{ent}}/r_H)\varrho_1$, and since a row or a column has at most $V_1 = 4(2R+3)^4$ nonzero entries, with the range R of [1, (3.56)], Schur's test gives

$$s \coloneqq \|H - H_U\| \le C_H\varrho_1 , \qquad C_H \coloneqq \frac{2V_1n_eC_{\text{ent}}}{r_H} . \tag{5.33}$$

The matrix H is complex symmetric, because $H^{\mathsf{T}} - H$ is holomorphic in α on a polydisc and vanishes for real α, where $e^{i\alpha}U$ is a real background. By Lemma 5.6(v), for $s \le \gamma/4$ its real part R and imaginary part S satisfy $\gamma_c \le \gamma - s \le R \le h_+ + s \le h_c$ and $\|S\| \le s$, the range is unchanged, and by Lemma 5.6(i) the same lower bound holds for every principal submatrix, which replaces the appeal to the minimum–maximum principle in [1, Lemma 9.13(a)].

Step 4: the walk series and the conditioning. The expansion of [1, Section 3.5] uses of the matrix $\mathsf{A}_H = 1 - H/s_H$ only that a block of A_H^n has norm at most q_H^n and vanishes beyond the distance nR [1, Lemma 3.13]. At $\mathbf{U}$ we have $\mathsf{A}_H = (1 - R/s_H) - iS/s_H$ with $0 \le 1 - R/s_H \le 1 - (\gamma - s)/s_H$, hence $\|\mathsf{A}_H\| \le 1 - (\gamma - 2s)/s_H \le q_H$ for the q_H of (K1), because $\gamma - 2s \ge \gamma_c$. The walk series, its bounds, and its holomorphy in the weakening parameters σ therefore hold at $\mathbf{U}$ with the constants of (K1), and the binomial series sums to a symmetric square root of H^{-1}, of norm at most $(\gamma - 2s)^{-1/2}$. The identity of the conditioning [1, Lemma 3.14] is proved there for a positive matrix by the conditional law of a Gaussian measure. Both of its sides are holomorphic in H on the set $\mathcal{N}$ of the complex symmetric matrices within the distance s_*, defined in (5.34), of the convex set of the real symmetric matrices between γ and h_+: the left side by Lemma 5.6(iv), the right side by the bounds of Step 5, which are uniform on $\mathcal{N}$. The set $\mathcal{N}$ is convex and open, and the two sides agree on its real points, which form an open subset of the real symmetric matrices; hence they agree on $\mathcal{N}$, which contains H.

Step 5: the Gaussian payment. Let E_0 be the bonds of a seed, $H_E = R_E + iS_E$ the compression of H to E_0, $C \coloneqq H_E^{-1} = C' + iC''$ with real symmetric C', C'', and $\Gamma = \Gamma_1 + i\Gamma_2$ the matrix $\Gamma_{E_0}(\sigma)$ of [1, (3.64)] with its real and imaginary parts, for σ in the polydisc of that section; put $\gamma' \coloneqq \gamma - s$. We bound the integral of the modulus of the density $e^{-\frac{1}{2}\langle \Gamma X, C\Gamma X\rangle}e^{-\langle B, \Gamma X\rangle}\,\mathrm{d}\mu_C(B)\,\mathrm{d}\mu_0(X)$ of [1, (3.66)] against $e^{\alpha_G\|B\|^2/2}$, the pairings being bilinear. The modulus of the Gaussian density of B is $(2\pi)^{-n/2}|\det H_E|^{1/2}e^{-\frac{1}{2}\langle B, R_EB\rangle}$, the modulus of $e^{-\langle B, \Gamma X\rangle}$ is $e^{-\langle B, \Gamma_1 X\rangle}$, and the integral over B equals

$$\Big(\frac{|\det H_E|}{\det R_E}\Big)^{1/2} \det(1 - \alpha_G R_E^{-1})^{-1/2} \exp(\tfrac{1}{2}\langle \Gamma_1 X, (R_E - \alpha_G)^{-1}\Gamma_1 X\rangle) .$$

With the real part of $\langle \Gamma X, C\Gamma X\rangle$ this leaves the exponent $\frac{1}{2}\langle X, T^c X\rangle$ in X, where

$$T^c := \Gamma_1^{\mathsf{T}}[(R_E-\alpha_G)^{-1}-C']\Gamma_1+\Gamma_2^{\mathsf{T}}C'\Gamma_2+\Gamma_1^{\mathsf{T}}C''\Gamma_2+\Gamma_2^{\mathsf{T}}C''\Gamma_1$$

is real symmetric and supported on the bonds J_Z of the final domain [1, Lemma 3.15]. We bound its norm. From $C-R_E^{-1}=-iCS_ER_E^{-1}$ and Lemma 5.6(i), (v) we get $\|C'-R_E^{-1}\|, \|C''\| \le s/\gamma'^2$, and $\|(R_E-\alpha_G)^{-1}-R_E^{-1}\| \le \alpha_G/(\gamma'(\gamma'-\alpha_G))$. The tail bound of the walk series gives $\|\Gamma(\sigma)-\Gamma(1)\| \le h_c\delta_M$ as in [1, Lemma 3.16(i)], and $\|\Gamma_1\| \le \|\Gamma\| \le h_c((\gamma-2s)^{-1/2}+1) \le A_*$. The matrix $\Gamma(1)$ at the real Hessian H_U is real, and along the segment $H_t := H_U + t(H-H_U)$ the square root of H_t^{-1} has the derivative of norm at most $\frac{1}{2}(\gamma-2s)^{-3/2}s$, by the binomial series with its positive coefficients; hence $\|\Gamma(1)-\Gamma(1)|_{H_U}\| \le C_\Gamma s$ with $C_\Gamma := \gamma_c^{-1/2}+\frac{1}{2}h_c\gamma_c^{-3/2}$, and $\|\Gamma_2\| \le h_c\delta_M + C_\Gamma s$. Collecting the terms and using (5.31), which bounds $2\alpha_G A_*^2/\gamma_c^2$ and $2h_c^2\delta_M^2/\gamma_c$ by $\frac{1}{8}$ each,

$$\|T^c\| \le \frac{2\alpha_G A_*^2}{\gamma_c^2}+\frac{2h_c^2\delta_M^2}{\gamma_c}+C_3 s \le \frac{1}{4}+C_3 s\,, \qquad C_3 := \frac{A_*^2}{\gamma_c^2}+\frac{2A_*(h_c+C_\Gamma\gamma/4)}{\gamma_c^2}+\frac{C_\Gamma^2\gamma}{2\gamma_c}\,,$$

so that $\|T^c\| \le \frac{1}{3}$ for $s \le 1/(12C_3)$, and the integral over X is $\det_{J_Z}(1-T^c)^{-1/2}$. For the three determinants, Lemma 5.6(iii) gives $\frac{1}{2}\log(|\det H_E|/\det R_E) \le d_G|E_0|s^2/(4\gamma'^2)$, and $-\frac{1}{2}\log(1-t) \le |t|$ for $|t| \le \frac{1}{2}$ gives $\log\det(1-\alpha_G R_E^{-1})^{-1/2} \le \alpha_G d_G|E_0|/\gamma'$ and $\log\det_{J_Z}(1-T^c)^{-1/2} \le d_G|J_Z|\|T^c\|$. Put $C_4 := d_G(C_3+\gamma/(16\gamma_c^2))$ and

$$s_* := \min\Big(\frac{\gamma}{4},\ \frac{1}{12C_3},\ \frac{1}{16C_4(LM+1)^4}\Big)\,. \tag{5.34}$$

Since $E_0 \subset J_Z$ and $|J_Z| \le 4(LM+1)^4 N_1(Z)$ [1, Section B.4], for $s \le s_*$ the logarithm of the integral is at most

$$(C_{\mathrm{d}}^K(\alpha_G+\delta_M^2)+C_4 s)|J_Z| \le N_1(Z)\,,$$

the three parts contributing at most $\frac{1}{2}$, $\frac{1}{4}$, and $\frac{1}{4}$ of $N_1(Z)$ by (5.31) and the choice of s_*. This is the bound $e^{N_1(Z)}$ of [1, Lemma 3.16(ii)] at $\mathbf{U}$, and it factorises over the components of the final domain, since T^c, C, and Γ are block diagonal [1, Lemma 3.15]. The third part of that lemma is an inequality for real B and is unchanged.

Step 6: the selected integrals and the cluster series. The proofs of the polymer representation, of the selected majorant, of the bounds of the activities and of the vacuum clusters, of the probe measures, and of the joint bound of the selected integrals [1, Lemmata 9.14, 9.15, 9.20, E.3, E.13, E.14; Corollary 9.16] use the background only through the bounds and the range of the Hessian, the charts and the bounds of the primitives, and the hosts [1, proof of Proposition 9.40]; Steps 2–5 supply the first two at $\mathbf{U}$, and the hosts do not depend on the background. Their other steps are the following. The cutoffs are indicators of the real fluctuation variables, with values in $\{0,1\}$, and are not changed. The modulus of every Gaussian density is paid by Step 5. The evenness of the kernel under $(B,X) \mapsto (-B,-X)$, on which the cancellation of the odd terms rests, is an identity of the formulae and holds for the complex Gaussian. The comparison of the hosts uses $|e^{\zeta\mathbf{V}}-1| \le e^{|\zeta\mathbf{V}|}-1$, which holds for complex $\mathbf{V}$, with the bound of the potentials of Step 2. The statements on normalisers, namely the positivity of the kernel and of the partition functions and the identities which divide by them, are not used: at $\mathbf{U}$ the weights of the groups are defined by the cluster series [1, Corollary 9.16 and Lemma 9.21(a)], whose convergence by the criterion of Kotecký and Preiss uses the activities in absolute value. The reference data do not depend on α.

Step 7: the words. At a real background the links of the loop are unitary and their words have the norm one; at $\mathbf{U}$ they do not. For $X = x+iy$ with $x, y \in \mathfrak{g}$ and $V \in G$ the Hermitian

parts of $\pm i\pi(X)$ are $\mp\pi(y)$, so that $\|\pi(e^{iX}V)\|$ and $\|\pi(e^{iX}V)^{-1}\|$ are at most $e^{\|\pi(y)\|} \le e^{C_\pi h(y)}$. The link of a traversal is $e^{iq_{\mathbf{U}}(x)}\mathbf{U}(r)$ at its bond r, with the reconstruction phase q at a real input x which does not depend on the background [1, Lemma 9.25(a)]. In the gauge of the stencil the phase is holomorphic in the background with the derivative at most $Cg_0(1+r_0)$ on the joint domain of the background and the phases [1, (E.52)], and for ϱ_1 small the segment from U to $\mathbf{U}$ stays in that domain; since the phase is real at U, its imaginary part at $\mathbf{U}$ has $h(\operatorname{Im} q_{\mathbf{U}}(x)) \le \varrho_1$ for γ small. Hence every link and every inverse link has norm at most $e^{2C_\pi\varrho_1}$ in the representation. The contraction of W contains at most $4|\mathcal{H}(W)|$ of them [1, Lemmata 9.25(b) and 9.26(d)], and $|\mathcal{H}(W)| \le LM\mathfrak{s}(W)$ by the definition of the size, so that the words of the group have norm at most $e^{8C_\pi LM\mathfrak{s}(W)\varrho_1} = e^{C'\mathfrak{s}(W)\varrho_1}$ in the place of one. The derivatives of the links are bounded in [1, Lemma 9.25(a)] by Cauchy's estimate for the factor which contains the phase, on the complex ball of the chart, and they gain the same factor.

Step 8: the bound and the covariance. With Steps 2–7 the proof of the majorant of the contraction [1, (E.73) and proof of Proposition 9.40] applies at $\mathbf{U}$ and gives $|c_W(\mathbf{U};h)| \le e^{C'\mathfrak{s}(W)\varrho_1}M^{(\rho)}(W)$ for every $h \in G$, locally uniformly, and at the reference data $|a_W| \le M^{(\rho)}(W)$. By Step 1 the insertion is holomorphic and bounded by $K_\chi(e^{C'\mathfrak{s}(W)\varrho_1}+1)M^{(\rho)}(W)$, which is (5.32). The formula is that of [1] with $\mathbf{U}$ in the place of U_1, the region, the cutoffs, and the index W being fixed, and at $\alpha = 0$ it is the insertion at U. Its covariance under a G-valued gauge transformation g, with α conjugated, is proved as at real backgrounds [1, Section E.10.2]: the formulae of the rows intertwine with the action R_g of g on the phases, the small solution of the row equation is unique in the complex ball of Step 2, the plaquette actions and the Haar density are invariant, and the change of variables $B \mapsto R_gB$, $X \mapsto R_gX$ with the real orthogonal R_g maps the complex Gaussian measure of covariance C to that of covariance $R_gCR_g^{-1}$. The dependence on the background through X_W^+ alone was shown in the proof of Theorem 5.4. The conditions on the coupling are $C_H\varrho_1 < s_*$, $\varrho_1 \le r_H/2$, $C'\varrho_1 \le \nu/8$, the two conditions of Step 7 on the phase, and the thresholds of [1] with the constants of (K1) and (K2); each holds for γ small, because ϱ_1 and $g_0(1+r_0)$ tend to zero with the coupling.

Step 9: the cross groups. Let W be a cross group. Its insertion (5.27) is the integral, against the measures of its roots, of the tensor of the words of its intervals, multiplied by the weight of its cluster occurrences. Steps 2–6 concern these measures and weights and do not involve the loops, so that they hold for W as they stand. By Step 7 each of the $|\mathcal{H}(W)|$ links of the words has norm at most $e^{2C_\pi\varrho_1}$ at $\mathbf{U}$, and the derivatives of the links gain the same factor. In the bound of the integral in the norm (5.5), which was derived in the proof of Theorem 5.4, the product of the norms of the factors of every elementary tensor is therefore at most $e^{2C_\pi|\mathcal{H}(W)|\varrho_1} \le e^{C'\mathfrak{s}(W)\varrho_1}$ times its bound at a real background, because $|\mathcal{H}(W)| \le LM\mathfrak{s}(W)$. Hence $\|Y_W(\mathbf{U})\| \le e^{C'\mathfrak{s}(W)\varrho_1}M^{(\rho)}(W)$, locally uniformly, and Y_W is holomorphic by the criterion of Step 1. Under a gauge transformation the measures and the weights are transformed as in Step 8, and the word of an interval is multiplied at its two ends, which is the covariance law for each factor of the tensor. *q.e.d.*

Corollary 5.8 (The Supremum Clauses at the First Step)**.** *On the term without large-field regions, let $\nu_\times \le \nu_e$ and $c_{\text{att}} \le \nu_e/8$. Then clause (iii) of Definition 5.2 holds at $k = 1$ in the form* (5.18), (5.19)*, and the bounds* (5.21), (5.22)*, and* (5.23) *hold at $k = 1$ for every pair of constants with $C_c \ge 2C_{S'} \ge L(2K_\chi+1)e^{2\nu}C_{\text{pin}}(1+\beta_0)^6$.*

Proof. We first bound the sizes S_W, then the exponents of the groups in the three clauses, and sum with the pins (5.26); the small factors of the cross and of the wrapping species come from the part of the weight of the pins which the first two clauses leave unused.

The space of the step 1 is the union of the sets $\mathcal{S}_1(U)$ over the configurations U of Proposition 5.7 and their gauge transforms, and a point of it has $h_{\mathbb{C}}(\alpha(b)) < \varrho_1$ by (5.30); the proposition, applied in a gauge and transported by the covariance, gives clause (iii) and, for a group on one loop, $\|Y_W\|_{(1)} \le 2K_\chi e^{\nu\mathfrak{s}(W)/8} M^{(\rho)}(W)$. The number subtracted from the insertion of such a group at its production is a_W, so that $S_W \le (2K_\chi+1)e^{\nu\mathfrak{s}(W)/8}M^{(\rho)}(W)$; from the insertion of a cross group nothing is subtracted, and $S_W = \|Y_W\|_{(1)} \le e^{\nu\mathfrak{s}(W)/8}M^{(\rho)}(W)$ by the last statement of the proposition. In both cases $\ell_W = 1$. The domain X_W is a connected set of at most $11\mathfrak{s}(W)$ cubes by (5.28), so that it lies within $11M\mathfrak{s}(W)$ of every traversal of its intervals in the units of the scale 1; as in the proof of Theorem 5.4 this gives $t_{\mathfrak{n}}(W) \le \lceil \mathfrak{s}(W) \rceil$, since $C_{\text{box}} \ge 11$, then $\text{ext}(W) \le C_{\text{box}}(2\mathfrak{s}(W)+4)$ and $\nu_e \,\text{ext}(W) \le \nu\mathfrak{s}(W) + 2\nu$. With $\theta(W) \le \nu\mathfrak{s}(W)/16$ the exponent of a group in each of the clauses is at most $2\nu + (2+\frac{3}{16})\nu\mathfrak{s}(W)$, the factor $e^{\nu\mathfrak{s}(W)/8}$ included, which is less than $2\nu + 4\nu\mathfrak{s}(W)$ by $\frac{29}{16}\nu\mathfrak{s}(W)$. A group with a traversal p of the step 1 in an interval has one of the L fine bonds of p in that interval, and a group with a cube $p \subset X_W$ has p in its footprint, because p is a cube of a constituent domain or contains a site of an interval. By the pins (5.26), the clauses (5.21) and (5.22) therefore hold with the right side $L(2K_\chi+1)e^{2\nu}C_{\text{pin}}\rho$, and $\rho \le (1+\beta_0)^6\rho_1$.

For (5.23) we keep the factor $e^{-\frac{29}{16}\nu\mathfrak{s}(W)}$ which this computation leaves unused. The domain of a cross group contains sites of two loops, and it is a connected union of at most $11\mathfrak{s}(W)$ cubes of side M in the units of the scale 1; hence $11M\mathfrak{s}(W) \ge d_1$, and the factor is at most $e^{-\frac{29}{176}\nu d_1/M} \le e^{-\nu_\times d_1/M}$, because $\nu_\times \le \nu_e \le \frac{29}{176}\nu$. A group of the wrapping species has a traversal $\mathfrak{n}$ in its interval with $N^{(1)}_{\min} \le 2R_{t_{\mathfrak{n}}(W)+1} + 1$, and $R_{t_{\mathfrak{n}}(W)+1} = C_{\text{box}}M(t_{\mathfrak{n}}(W)+2) \le C_{\text{box}}M(\mathfrak{s}(W)+3)$, so that $\mathfrak{s}(W) \ge N^{(1)}_{\min}/(2C_{\text{box}}M) - 4$. The factor is then at most

$$e^{\frac{29}{4}\nu}\exp\Big(-\frac{29}{32}\,\frac{\nu N^{(1)}_{\min}}{C_{\text{box}}M}\Big) \le \exp\Big(-\frac{1}{16}\,\frac{\nu N^{(1)}_{\min}}{C_{\text{box}}M}\Big) \le e^{-c_{\text{att}}N^{(1)}_{\min}/M}\,,$$

where the first inequality holds because $\frac{27}{32}N^{(1)}_{\min}/(C_{\text{box}}M) \ge \frac{29}{4}$, the periods of the torus being $2L^{m_T+q}a_1 \ge 2a_1 \ge 2L^2M$ in the units of the scale 1 (proof of Lemma 3.14), and the second because $c_{\text{att}} \le \nu_e/8 \le \nu/(16C_{\text{box}})$. *q.e.d.*

5.6 The First Step on a Term with Large-Field Regions

The first-step paper makes its expansion on the labelled terms on which the loops are *covered*: every bond of a loop lies in Ω_1, and every free input of its links lies in the carrier of the term [1, Sections 9.4.2 and 9.6.1]. On the other terms it bounds the conditional expectation of the loop by 1 and the total weight of these terms by the probability of a large field near the loop [1, Proposition 9.39]. For a single step this is sufficient. For the induction it is not, because the later steps act on each labelled term separately, and Definition 5.2 requires the form (5.10) on every one of them. Three things stand in the way. The mean of the conditional Gaussian of a term depends on its whole region of integration, so that every factor of the expansion depends on the labels of the term far from its own support. The expansion uses the hypothesis that the loops are covered. And the links of the loops near the complement of the region of integration are not the transports of the background. We begin with the first.

We use the notation of Supplement, Section 5: lengths are measured in fine units, $d(b,b')$ is the ℓ^∞-distance of the initial sites of two bonds, $\ell = LMR_1$, and a labelled term has the retreated region Λ, the carrier $D := D(\Lambda)$, the exterior $J(\Lambda)$, and the collar $J_{\text{col}}(\Lambda)$. Let H be the restriction to D of the Hessian of the first step, and write $\gamma := \gamma_L/2$ and $h_+ := m_L$. Every principal restriction of H satisfies $\gamma \le H \le h_+$, its entries vanish between bonds at a distance greater than $\rho_L = 4L$, and the sum of the norms of the entries of a row of the

Hessian which belongs to D is at most m_L [1, Section 9.1, (I1)–(I3)]; consequently the entries of the inverse of a principal restriction H_{AA} satisfy

$$\|(H_{AA})^{-1}(b,b')\| \le \gamma^{-1} e^{-\mu_H d(b,b')} , \qquad \mu_H := \frac{\gamma}{h_+ \rho_L} , \tag{5.35}$$

by [1, Lemma 9.1], whose proof uses the two-sided bound and the range alone. Here the norm of an entry is its operator norm for the norm h on $\mathfrak{g}$, and $h \le |\cdot|_\sigma \le c_{\mathrm{h}} h$ for the norm of the cutoff [1, Remark 9.3]. Let A^Λ be the exterior variable of the term on $J(\Lambda)$, let $r := t_0$ be the radius of the cutoff of the fluctuation field (Section 5.4), and let $\vartheta_V = 1/(4c_{\mathrm{h}} C_H^{\mathrm{near}})$, with $C_H^{\mathrm{near}} := \gamma^{-1} m_L N_{\mathrm{lat}}(\mu_H)$ and $N_{\mathrm{lat}}(\mu) := \sum_{s\ge 0} 4(2s+1)^4 e^{-\mu s}$, be the constant of the collar test [1, Lemma 9.2 and Remark 9.3]. The linear term of the conditional density is $\ell_\Lambda := H_{DJ(\Lambda)} A^\Lambda \in \mathfrak{g}^D$, and it involves the exterior variable only on the set $J_1(\Lambda) \subset J_{\mathrm{col}}(\Lambda)$ of the exterior bonds which have a nonzero entry of the Hessian with a bond of D. On the support of the term $|A^\Lambda(e)|_\sigma < \vartheta_V r$ for $e \in J_{\mathrm{col}}(\Lambda)$ (Supplement, Lemma 5.2(iii)), so that

$$\sup_{c\in D} h(\ell_\Lambda(c)) \le m_L \vartheta_V r , \tag{5.36}$$

by the bound of the rows. The first-step paper centres the conditional Gaussian at its mean $m := -H^{-1}\ell_\Lambda$, for which $|m_b|_\sigma \le r/4$ and

$$h(m_b) \le C_m \vartheta_V r \sum_{e\in J_1(\Lambda)} e^{-\mu_H d(b,e)} , \qquad C_m := \gamma^{-1} m_L \cdot 4(2\rho_L+1)^4 e^{\mu_H \rho_L} , \tag{5.37}$$

by [1, (9.11) and Lemma 9.19]. The entries of H^{-1} decay, but they do not vanish at any distance, so that m_b depends on the carrier and on the exterior variable everywhere. The following lemma replaces m by a vector which is determined within a distance of the order of $\log g_0^{-2}$, at the price of a linear term which is small in the coupling.

Lemma 5.9 (The Local Shift)**.** *There are integers $\ell_s \ge \rho_L + 1$ and N_0, which depend on L and G, with the following property. Put $\mu_x := \mu_H/20$, let N be the least integer $N \ge N_0$ with $e^{-\mu_H \ell_s N/4} \le g_0$, and let*

$$R_t := 5\ell_s(N+2) + 2\rho_L \le \frac{10}{\mu_H} \log g_0^{-2} + 5\ell_s(N_0+3) + 2\rho_L . \tag{5.38}$$

On every labelled term of the first step, and for the exterior variable in the support of the term, there is a real vector $\hat m \in \mathfrak{g}^D$ such that, with $\hat\delta := H\hat m + \ell_\Lambda$:

(i) Locality. *For every $b \in D$, the vectors $\hat m_b$ and $\hat\delta(b)$ are functions of the set of the bonds of D within the distance R_t of b, of the entries of the Hessian between bonds within that distance, and of the restriction of A^Λ to the bonds of $J_1(\Lambda)$ within that distance. Moreover $\hat m_b = 0$ unless $d(b, J_1(\Lambda)) \le R_t$, and $\hat\delta(b) = 0$ unless $d(b, J_1(\Lambda)) \le R_t + \rho_L$.*

(ii) Sizes. *For every $b \in D$,*

$$\begin{aligned} |\hat m_b|_\sigma \le \frac{r}{4} , \qquad h(\hat m_b) &\le 2C_m \vartheta_V r \sum_{e\in J_1(\Lambda)} e^{-\mu_x d(b,e)} , \\ h(\hat\delta(b)) &\le C_\delta \vartheta_V r\, g_0 \sum_{e\in J_1(\Lambda)} e^{-\mu_x d(b,e)} , \end{aligned} \tag{5.39}$$

with $C_\delta := 2m_L e^{3\mu_H\rho_L/20}$, and $\sup_b h(\hat\delta(b)) \le 2m_L \vartheta_V r\, g_0^2$.

(iii) The identity. *For every t and every $B \in \mathfrak{g}^D$, with $a = B + t\hat{m}$,*

$$-\tfrac{1}{2}\langle a, Ha\rangle - t\langle a, \ell_\Lambda\rangle = -\tfrac{1}{2}\langle B, HB\rangle - t\langle B, \hat{\delta}\rangle - t^2(\tfrac{1}{2}\langle \hat{m}, H\hat{m}\rangle + \langle \hat{m}, \ell_\Lambda\rangle) \; . \tag{5.40}$$

Proof. We construct an approximate inverse of H from the inverses of its restrictions to boxes, and we take for $\hat{m}$ a finite part of the resulting series for m.

The boxes. Let $\ell_s \geq \rho_L + 1$ be an integer, to be fixed below. We partition every coordinate circle of the fine torus into intervals whose lengths lie in $[\ell_s, 2\ell_s)$, which is possible because the periods of the torus exceed ℓ_s, and we call the products of these intervals *boxes*. A bond belongs to the box of its initial site. For a box Δ let $\tilde{\Delta}$ be the set of the bonds within the distance ℓ_s of Δ, and $\hat{\Delta}$ the set of those within the distance ρ_L of $\tilde{\Delta}$. A bond lies in $\hat{\Delta}$ for at most $N_{\mathrm{nb}} \coloneqq 7^4$ boxes, since such a box lies within the distance $2\ell_s$ of the bond and an interval of length $4\ell_s + 1$ meets at most seven intervals of length at least ℓ_s. The partition does not depend on the term.

The approximate inverse. For a box Δ let G_Δ be the inverse of the restriction of H to $\tilde{\Delta} \cap D$, extended by zero, let χ_Δ be the projection onto $\mathfrak{g}^{\Delta \cap D}$, and put

$$P \coloneqq \sum_\Delta G_\Delta \chi_\Delta \; , \qquad K_\Delta \coloneqq -\mathbf{1}_{D \setminus \tilde{\Delta}} H G_\Delta \chi_\Delta \; , \qquad K \coloneqq \sum_\Delta K_\Delta \; . \tag{5.41}$$

For a vector u supported in $\tilde{\Delta} \cap D$, the vector Hu is $H_{\tilde{\Delta} \cap D} u$ on $\tilde{\Delta} \cap D$ and $\mathbf{1}_{D \setminus \tilde{\Delta}} Hu$ outside it. With $u = G_\Delta \chi_\Delta f$ this gives $HG_\Delta \chi_\Delta f = \chi_\Delta f - K_\Delta f$, and since $\sum_\Delta \chi_\Delta$ is the identity of $\mathfrak{g}^D$,

$$HP = 1 - K \; . \tag{5.42}$$

We measure a matrix A indexed by bonds by $\|A\|_\infty \coloneqq \sup_b \sum_{b'} \|A(b, b')\|$, which bounds its action on the vectors with the norm $\sup_b h(v_b)$. An entry $K_\Delta(y, z)$ is the sum of $-H(y, x) G_\Delta(x, z)$ over $x \in \tilde{\Delta} \cap D$, for $y \in D \setminus \tilde{\Delta}$ and $z \in \Delta$. It vanishes unless $d(y, x) \leq \rho_L$ for some such x; then $y \in \hat{\Delta}$, and $d(x, z) > \ell_s - \rho_L$, because $d(y, z) > \ell_s$. At most $32(2s + 1)^3$ bonds are at the distance s from a given bond, four for each of the at most $(2s + 1)^4 - (2s - 1)^4 \leq 8(2s + 1)^3$ sites. By (5.35) for the restriction to $\tilde{\Delta} \cap D$ and by the bound of the rows of the Hessian, the sum of the norms of the entries of a row of K_Δ is therefore at most

$$\varepsilon_s \coloneqq m_L \gamma^{-1} \sum_{s > \ell_s - \rho_L} 32(2s + 1)^3 e^{-\mu_H s} \; , \tag{5.43}$$

and the rows of K_Δ lie in $\hat{\Delta}$, so that $\|K\|_\infty \leq q_s \coloneqq N_{\mathrm{nb}} \varepsilon_s$. We fix ℓ_s, depending on L and G, so large that $q_s \leq e^{-\mu_H \ell_s / 2}$; this is possible because ε_s is at most a polynomial in ℓ_s times $e^{-\mu_H(\ell_s - \rho_L)}$. By (5.35) for H itself, and since at most $4(2s+1)^4$ bonds lie within the distance s of a bond, $\|H^{-1}\|_\infty \leq \gamma^{-1} N_{\mathrm{lat}}(\mu_H)$.

The shift. For an integer $N \geq 0$ put $\tilde{m} \coloneqq -P \sum_{n=0}^N K^n \ell_\Lambda$. By (5.42), $H\tilde{m} + \ell_\Lambda = -(1 - K) \sum_{n \leq N} K^n \ell_\Lambda + \ell_\Lambda = K^{N+1} \ell_\Lambda =: \delta_N$, and since $Hm = -\ell_\Lambda$ we have $\tilde{m} - m = H^{-1} \delta_N$. With (5.36),

$$\begin{gathered} \sup_b h(\delta_N(b)) \leq q_s^{N+1} m_L \vartheta_V r \; , \qquad \sup_b |\tilde{m}_b - m_b|_\sigma \leq \tfrac{1}{2} \varepsilon_N \; , \\ \varepsilon_N \coloneqq 2 c_{\mathrm{h}} C_H^{\mathrm{near}} \vartheta_V r \, q_s^{N+1} = \tfrac{1}{2} r \, q_s^{N+1} \; . \end{gathered} \tag{5.44}$$

Since $|m_b|_\sigma \leq r/4$, the vector

$$\hat{m} \coloneqq \lambda \tilde{m} \; , \qquad \lambda \coloneqq \frac{r/4}{r/4 + \varepsilon_N/2} = \frac{1}{1 + q_s^{N+1}} \; , \tag{5.45}$$

satisfies $|\hat m_b|_\sigma \le \lambda(r/4+\varepsilon_N/2) = r/4$. The factor λ is a number which depends on N and on the constants only. For the vector of the statement, $\hat\delta = \lambda(H\tilde m + \ell_\Lambda) + (1-\lambda)\ell_\Lambda = \lambda\delta_N + (1-\lambda)\ell_\Lambda$, and $1-\lambda \le q_s^{N+1}$, so that

$$\sup_b h(\hat\delta(b)) \le 2m_L\vartheta_V r\, q_s^{N+1}\ . \tag{5.46}$$

The locality. We expand PK^n into the terms $G_{\Delta_0}\chi_{\Delta_0}K_{\Delta_1}\cdots K_{\Delta_n}$. In the entry (b,c) of such a term, $b\in\tilde\Delta_0$ lies within $\ell_s+2\ell_s-1$ of every bond of Δ_0; a bond of a row of K_{Δ_i} lies in $\hat\Delta_i$, hence within $\ell_s+\rho_L+2\ell_s-1<4\ell_s$ of every bond of Δ_i; and $c\in\Delta_n$. So the entry vanishes unless $d(b,c)\le 3\ell_s+4\ell_s n\le 5\ell_s(n+1)$. The term is built from the entries of H between the bonds of D in the sets $\hat\Delta_0,\dots,\hat\Delta_n$, which lie within $5\ell_s(n+1)+\ell_s+\rho_L$ of b when the entry does not vanish. Finally $\ell_\Lambda(c)$ involves A^Λ on the bonds of $J_1(\Lambda)$ within ρ_L of c and vanishes if there are none. Hence $\tilde m_b$, $\hat m_b$, and $\delta_N(b)$ are functions of the data named in (i) within the distance $5\ell_s(N+2)+2\rho_L$ of b, and $\tilde m_b=0$ unless $d(b,J_1(\Lambda))\le 5\ell_s(N+1)+\rho_L$; the vector $\hat\delta(b)$ vanishes unless $\delta_N(b)$ or $\ell_\Lambda(b)$ does not. With the N of the statement these are the assertions of (i).

The sizes. By the choice of N and of ℓ_s, $q_s^{N+1}\le e^{-\mu_H\ell_s N/2}\le g_0^2$, and (5.46) gives the last bound of (ii). For the decay we note that, for $R\ge 0$,

$$\mathbf{1}\{d(b,J_1(\Lambda))\le R\} \le e^{\mu_x R}\sum_{e\in J_1(\Lambda)} e^{-\mu_x d(b,e)}\ , \qquad q_s^{N+1}e^{\mu_x(R_t+\rho_L)} \le e^{3\mu_H\rho_L/20}e^{-\mu_H\ell_s N/4}\ , \tag{5.47}$$

the second because $q_s\le e^{-\mu_H\ell_s/2}$ and $\mu_x(R_t+\rho_L)=\frac14\mu_H\ell_s(N+2)+\frac{3}{20}\mu_H\rho_L$. The third bound of (5.39) follows from (5.46), from the support of $\hat\delta$, and from $e^{-\mu_H\ell_s N/4}\le g_0$. For the second, $h(\hat m_b)\le h(\tilde m_b)\le h(m_b)+\frac12\varepsilon_N$ where $\hat m_b\ne 0$, and $\frac12\varepsilon_N = c_{\rm h}C_H^{\rm near}\vartheta_V r\, q_s^{N+1}$; by (5.37), by (5.47), and by $e^{-\mu_H d}\le e^{-\mu_x d}$,

$$h(\hat m_b) \le (C_m + c_{\rm h}C_H^{\rm near}e^{3\mu_H\rho_L/20}e^{-\mu_H\ell_s N/4})\vartheta_V r\sum_{e\in J_1(\Lambda)} e^{-\mu_x d(b,e)}\ .$$

We let N_0 be the least integer with $c_{\rm h}C_H^{\rm near}e^{3\mu_H\rho_L/20}e^{-\mu_H\ell_s N_0/4}\le C_m$, which depends on L and G. The bound of R_t in (5.38) holds because $N\le N_0+1+2\log g_0^{-2}/(\mu_H\ell_s)$.

The identity. Since H is symmetric, $\frac12\langle a,Ha\rangle = \frac12\langle B,HB\rangle + t\langle B,H\hat m\rangle + \frac12 t^2\langle\hat m,H\hat m\rangle$ and $t\langle a,\ell_\Lambda\rangle = t\langle B,\ell_\Lambda\rangle + t^2\langle\hat m,\ell_\Lambda\rangle$; adding the two and using $H\hat m+\ell_\Lambda=\hat\delta$ gives (5.40). *q.e.d.*

The expansion of the first-step paper is made in the variable $B=a-tm$, in which the conditional density of the term is a Gaussian with the covariance H^{-1}, restricted by the cutoffs and perturbed by local potentials. We now make it in the variable $B=a-t\hat m$. By (5.40) the density then keeps the linear term $-t\langle B,\hat\delta\rangle$, and we add this term to the potentials. Let us recall the objects. On a labelled term, and for the exterior data in its support, the conditional density of $a\in\mathfrak{g}^D$ is $K_t^\Lambda(a)=\mathbf{1}\{|a_b|_\sigma<r \text{ for all } b\in D\}\exp[-\frac12\langle a,Ha\rangle - t\langle a,\ell_\Lambda\rangle + P_t^\Lambda(a)]$, for $t\in[0,1]$, where $P_t^\Lambda(a)=\sum_Y \mathbf{V}_t^\Lambda(Y,a)$ is a sum over the *hosts* $Y\in\mathbf{D}_0(\Lambda)$, boxes of cubes of side M with at most four cubes along each side; the potential $\mathbf{V}_t^\Lambda(Y,\cdot)$ depends on a on the set $E(Y)$ of the bonds of D which meet Y, and $|\mathbf{V}_t^\Lambda(Y,a)|\le\omega_t e^{-\kappa d_0(Y)}$ with $\omega_t := C_0 g_0(1+2r)^3$ when $|a_b|_\sigma<r$ on $E(Y)$, where $d_0(Y)$ is the tree length of Y in units of M [1, Section 9.5 and (9.12)]. Every bond $b\in D$ has a *probe host* $Y_b\in\mathbf{D}_0(\Lambda)$, with $b\in E(Y_b)$ [1, Lemma 9.13(d)]. The conditional measure is $\mathrm{d}\nu_t^\Lambda := K_t^\Lambda\,\mathrm{d}a/\int K_t^\Lambda\,\mathrm{d}a$.

Lemma 5.10 (The Conditional Measure at the Local Shift)**.** *On a labelled term of the first step, with the vectors $\hat m$ and $\hat\delta$ of Lemma 5.9, put for $b\in D$, $Y\in\mathbf{D}_0(\Lambda)$, $t\in[0,1]$, and*

$B \in \mathfrak{g}^D$

$$\begin{aligned} \hat\chi^t_b(B) &\coloneqq \mathbf{1}\{|B_b + t\hat m_b|_\sigma < r\}\ , \qquad \hat\Theta^t_E \coloneqq \prod_{b\in E} \hat\chi^t_b\ , \\ \tilde W_t(Y,B) &\coloneqq \mathbf{V}^\Lambda_t(Y, B + t\hat m) - t\,\hat\Theta^t_{E(Y)}(B) \sum_{b\,:\,Y_b = Y} \langle \hat\delta(b), B_b\rangle\ , \end{aligned} \tag{5.48}$$

where $\mathbf{V}^\Lambda_t(Y,\cdot)$ *is zero unless* $\hat\Theta^t_{E(Y)} = 1$.

(i) For every bounded Borel function F *on* $\mathfrak{g}^D$,

$$\int F\,\mathrm{d}\nu^\Lambda_t = \frac{\int F(B + t\hat m)\,\hat\Theta^t_D(B) \exp[-\frac12\langle B, HB\rangle + \sum_Y \tilde W_t(Y,B)]\,\mathrm{d}B}{\int \hat\Theta^t_D(B) \exp[-\frac12\langle B, HB\rangle + \sum_Y \tilde W_t(Y,B)]\,\mathrm{d}B}\ . \tag{5.49}$$

(ii) The function $\tilde W_t(Y,\cdot)$ *depends on* B *on* $E(Y)$ *only, it vanishes unless* $\hat\Theta^t_{E(Y)}(B) = 1$, *and*

$$\begin{aligned} |\tilde W_t(Y,B)| &\le \hat\omega_t\, e^{-\kappa d_0(Y)}\ , \\ \hat\omega_t &\coloneqq (C_0 + C_\tau)\, g_0 (1+2r)^3\ , \qquad C_\tau \coloneqq 5120\, M^4 m_L \vartheta_V e^{255\kappa}\ . \end{aligned} \tag{5.50}$$

(iii) The linear part of $\tilde W_t(Y,\cdot)$ *vanishes unless* Y *contains a point within the distance* $R_t + \rho_L + 1$ *of* $J_1(\Lambda)$, *and both parts are functions of the labels of the term and of its exterior variable within the distance* R_t *of* Y.

Proof. (i) Put $a = B + t\hat m$. The cutoff of K^Λ_t becomes $\hat\Theta^t_D(B)$, and by (5.40) the quadratic and the linear term become $-\frac12\langle B, HB\rangle - t\langle B, \hat\delta\rangle$ plus a number which does not depend on B. Where $\hat\Theta^t_D(B) = 1$, the first parts of the functions $\tilde W_t(Y,B)$ sum to $P^\Lambda_t(B + t\hat m)$. Their second parts sum to $-t\langle B, \hat\delta\rangle$ there, because $\hat\delta$ is supported in D, every bond of D has exactly one probe host, and $\hat\Theta^t_{E(Y)} = 1$ for every Y. By the invariance of the Lebesgue measure under translations, the integral of FK^Λ_t is therefore the numerator of (5.49) times that number, and with $F = 1$ the same holds for the denominator.

(ii) The first two statements hold for both parts of $\tilde W_t$, because $Y_b = Y$ implies $b \in E(Y)$. Where $\hat\Theta^t_{E(Y)}(B) = 1$, the argument $B + t\hat m$ satisfies the condition of the bound of $\mathbf{V}^\Lambda_t$, which gives $\omega_t e^{-\kappa d_0(Y)}$ for the first part. For the second, $|B_b|_\sigma < r + |\hat m_b|_\sigma \le \frac54 r$ on $E(Y)$ by (5.39), and $|\langle \hat\delta(b), B_b\rangle| \le h(\hat\delta(b)) h(B_b) \le 2 m_L \vartheta_V r\, g_0^2 \cdot \frac54 r$, by the Cauchy–Schwarz inequality for the inner product of $\mathfrak{g}$, whose norm is h, by $h \le |\cdot|_\sigma$, and by Lemma 5.9(ii). A host has at most 4^4 cubes of side M, each of which meets at most $8M^4$ bonds [1, Section B.1], so that the second part is at most $2048 M^4 \cdot \frac52 m_L \vartheta_V r^2 g_0^2$ in modulus. The tree length of a host is at most the number of its cubes minus one [1, Lemma B.1(1)], hence at most 255. Since $r^2 g_0^2 \le g_0 (1+2r)^3$ for $g_0 \le 1$, the second part is at most $C_\tau g_0 (1+2r)^3 e^{-\kappa d_0(Y)}$.

(iii) This is Lemma 5.9(i), together with the fact that a potential $\mathbf{V}^\Lambda_t(Y,\cdot)$ reads the exterior variable on the inputs of its primitives, which lie in Y [1, Lemma 9.13(d)]. *q.e.d.*

The polymer expansion at the local shift. The first-step paper writes the normaliser of the conditional measure as a polymer gas and bounds its activities [1, Lemmata 9.14 and 9.15, Corollary 9.16]. The proofs use the shift and the potentials through three properties only. Each cutoff is a function of one bond variable with values in $\{0,1\}$. Each potential depends on B on $E(Y)$, and where the cutoffs of $E(Y)$ do not vanish it is at most $\omega_* e^{-\kappa d_0(Y)}$ in modulus, with the constant $\omega_* \le \frac12$ of that paper. And $|m_b|_\sigma \le r/4$, which enters through the inequality $\mathbf{1}\{|B_b + t m_b|_\sigma \ge r\} \le \mathbf{1}\{|B_b|_\sigma \ge \frac34 r\}$ for the bonds at which a cutoff fails [1,

proofs of Lemmata 9.14 and 9.15]. By Lemma 5.9(ii) and Lemma 5.10(ii), the cutoffs $\hat\chi_b^t$ and the potentials $\tilde W_t$ have these properties as soon as

$$\hat\omega_t = (C_0 + C_\tau)\, g_0 (1+2r)^3 \le \tfrac12 \omega_* \,, \tag{5.51}$$

a smallness condition on the coupling, which takes the place of the conditions [1, (9.15) and (9.20)(ii)] on ω_t. Under it the three statements hold on every labelled term with $\hat\chi_b^t$ and $\tilde W_t$ in the places of the cutoffs and the potentials of that paper, with the same constants: the denominator of (5.49) is a Gaussian normalisation times a polymer gas of final domains, whose activities satisfy the bounds of [1, Corollary 9.16] uniformly in t, in the exterior data, and in the labels. A probe host which carries no potential in that paper carries here the linear part alone, with the same bound.

The same holds for the measures through which the first-step paper expands the conditional expectation of a function of the variable of integration [1, Lemma 9.20]. Its part (c), the comparison which bounds the total variation of these measures by the weights of the final domains, is stated there for all potentials which are at most $\frac12 \omega_* e^{-\kappa d_0(Y)}$ in modulus on the support of the cutoffs of their host, and this is (5.50) with (5.51). Its parts (a) and (b) rest on the polymer representation and on the bound of the activities, with the three properties above, and the supports of the measures lie in the box $|B_b|_\sigma \le \frac54 r$ because $|t\hat m_b|_\sigma \le r/4$. Its part (d), the expansion itself, is an identity between measures with the same Laplace transform, for which the family of the potentials with a further linear term has to have the same three properties; the functions $\tilde W_t$ have them with any further linear term, since they differ from the potentials of that paper by a linear term of the same kind [1, Section 9.6.3]. Hence, for every bounded Borel function F on $\mathfrak{g}^D$ and on every labelled term,

$$\int F \,\mathrm{d}\nu_t^\Lambda = F(t\hat m) + \sum_{\mathcal{A}\neq\emptyset} \hat q_\Lambda^t(\mathcal{A})\, \hat{\mathfrak{Q}}_\Lambda^t(\mathcal{A}; F(\cdot + t\hat m)) \,, \tag{5.52}$$

the sum over the nonempty compatible families $\mathcal{A}$ of final domains, where $\hat q_\Lambda^t$ and $\hat{\mathfrak{Q}}_\Lambda^t$ are the ratios of polymer gases and the integrals against the centred measures of [1, Lemma 9.20(d)], formed with $\hat\chi_b^t$ and $\tilde W_t$. This identity does not involve the loops.

The loops on an arbitrary term. We turn to the second obstacle. The function which the first-step paper integrates is the product of the characters of the loops in the reconstructed fine field [1, (3.26)]. For $a \in \mathfrak{g}^D$ let $U^\Lambda(a)$ be that field on the term: its value on a bond $r \notin \Omega_1$ is $V_0(r)$, and on a bond $r \in \Omega_1$ it is $e^{iq(r)} U_1(r)$, where the phase $q(r)$ is a function of the free variables on the *free-input set* $\mathcal{F}_r$ of r, a set of free bonds within the distance $2L+1$ of r [1, Lemma 3.5 and Section 9.6.1]. The free variables are a on the bonds of D and the exterior variable of the term on the others: the variable of Bałaban's conditional integral on $I \setminus D$, where it is no longer integrated, and his variable outside Λ_1 on $\Omega_1^* \setminus I$. So the link of a fine traversal depends on a through its restriction to $\mathcal{F}_r \cap D$ alone, and not at all if $r \notin \Omega_1$ or $\mathcal{F}_r \cap D = \emptyset$; we put $\mathcal{F}_r := \emptyset$ for $r \notin \Omega_1$. Let $f(a) := \prod_i \chi_i(U^\Lambda(a)(C_i))$ and

$$F^\Lambda := \int f \,\mathrm{d}\nu_1^\Lambda \,, \tag{5.53}$$

the conditional expectation of the loops on the term. On a covered term every traversal lies in Ω_1 and has $\mathcal{F}_r \subset D$, and this is the hypothesis under which [1, Section 9.6] works. It is used there at one place only where it cannot simply be dropped, the bounds of the derivatives of the links, and for these a weaker property suffices, which every term has.

Lemma 5.11 (The Links of a Loop on an Arbitrary Term)**.** *Let the standing conditions of [1, Lemma 3.11] and the threshold $4C_{\mathsf{C}} g_0(1+2r) \le \vartheta_*/2$ of [1, Lemma 9.25] hold. On every*

labelled term of the first step, for the exterior data in the support of the term, and for every fine traversal with bond $r \in \Omega_1$ and $\mathcal{F}_r \cap D \neq \emptyset$:

(i) the link $L_r(a) \coloneqq U^\Lambda(a)(r)$ is defined for every real a with $|a_b|_\sigma \leq 2r$ on D, and it is unitary;

(ii) for such a, every k with $1 \leq k \leq 2m_*$, bonds $b_1, \dots, b_k \in \mathcal{F}_r \cap D$, and unit vectors $e_1, \dots, e_k$ of $\mathfrak{g}$ for the norm h,

$$\| \partial_{e_1,b_1} \cdots \partial_{e_k,b_k} \pi(L_r)(a) \| \leq (2C_{\text{link}} g_0)^k \,, \tag{5.54}$$

and for $k \leq m_*$ the same with C_{link} in the place of $2C_{\text{link}}$, where m_* and C_{link} are the constants of [1, Lemma 9.25], which depend on L, G, and the representation.

Consequently the bounds of the mixed finite differences [1, Lemmata 9.25(b) and 9.31(b)] hold on every labelled term, for the words whose factors are links of the loops, their inverses, and fixed matrices, with the differences taken in the variables of the bonds of D and with the constants of these lemmata.

Proof. Let c be the row of r and $\mathcal{E}_c$ its two-block set, so that $\mathcal{F}_r \subset \mathcal{E}_c$ and the link is a function of the free variables on $\mathcal{E}_c$ [1, Lemma 3.5]. A bond of $\mathcal{F}_r \cap D$ belongs to I, so that c is one of the rows "within $2R_{\text{c}} + 2L$ of I" for which [1, Lemma 3.6] constructs the row solution, on every term of Bałaban's decomposition: by the remark which precedes that lemma, a row is within this distance of I as soon as one bond of $\mathcal{E}_c$ is within $2R_{\text{c}} + 1$ of I. The lemma states that every point $a' + \Psi(A_{\text{e}})$, with $a' \in \mathfrak{g}^I$ in the box $|a'_b| \leq 2r$ and A_{e} the variable outside Λ_1 on the support of its cutoffs, lies in the polydisc of the row, because $\Psi(A_{\text{e}}) = A_{\text{e}}$ and $|A_{\text{e}}(b)| < r$ on the bonds of $\mathcal{E}_c$ [1, Lemma 3.11(i), (iii)]. On our term the free variable on I is $a' = (a, a_{\text{old}})$, with a on D and the exterior variable a_{old} on $I \setminus D$, and $|a_{\text{old},b}|_\sigma < r$ on the support of the term, since the density of Bałaban's conditional integral contains the cutoff of every bond of I [1, Lemma 9.36]. So a' lies in the box whenever $|a_b|_\sigma \leq 2r$ on D, the row solution is defined at the input of the row, which gives (i), and every component of that input is at most $2r$ in norm. The latter is what the preamble of [1, Lemma 9.25] derives from the hypothesis that the loops are covered and uses for the bound $\|A\| \leq \vartheta_*/4$ of the phase A of the row. In the proof of part (a) of that lemma the link is then $\Pi(A)\pi(U_1(r))$ with a function Π which is analytic and bounded by C_Π on the ball of radius $\vartheta_*/2$, and a derivative in a direction e_b, $b \in \mathcal{F}_r \cap D$, is $g_0 D\Pi(A)[\mathsf{C}_U e_b]$; the estimate of Cauchy on the ball of radius $\vartheta_*/4$ about A involves neither the other components of the input nor the hypothesis. This gives (ii), for $k \leq m_*$ as in [1, Lemma 9.25(a)] and for $k \leq 2m_*$ as in [1, Lemma 9.31(a)]; for a free bond r the link is $e^{ig_0\pi(a_r)}\pi(U_1(r))$ and the bound is that of Duhamel's formula, as there. The parts (b) of the two lemmata are deduced from the parts (a) by the product rule inside the normalised trace, with the undifferentiated links unitary and with the number N_{tr} of the traversals which have a given bond in their free-input sets; a link which does not depend on a, because $r \notin \Omega_1$ or $\mathcal{F}_r \cap D = \emptyset$, is a fixed unitary factor, which these parts admit. *q.e.d.*

With the lemma and the expansion (5.52) the analysis of [1, Sections 9.6 and 9.7] can be repeated on every term, at the local shift. We state the outcome for one loop C with the character χ of the representation π; for an ensemble the groups, the tensors, and the norms are those of the paragraphs on several loops in the proof of Theorem 5.4, and nothing else changes. The objects are those of [1, Lemmata 9.21 and 9.26]. For a connected final domain Z let L_Z be the set of the bonds $b \in D$ with $Y_b \subset Z$, and call Z *marked* if some fine traversal has a bond of L_Z in its free-input set; the *groups* W of the term are those of [1,

Definition 3.24] for the carrier D, each with a set $\mathcal{A}(W)$ of marked final domains, its roots, a list of clusters of final domains, and an interval $\mathcal{H}(W)$ of fine traversals. They are defined by the domains and the loop alone. For a group W and $t \in \{0,1\}$ let $T^t_W(B)$ be the ordered product of the links of $U^\Lambda(\mathbf{1}_{\mathcal{A}(W)}B + t\hat{m})$ along $\mathcal{H}(W)$, where $\mathbf{1}_{\mathcal{A}(W)}B$ is B on the bonds of the sets L_Z, $Z \in \mathcal{A}(W)$, and 0 elsewhere, let $P^t_W \coloneqq T^t_W(0)$, and

$$\Upsilon^t_W \coloneqq \hat{w}^t(W) \int \pi(T^t_W(B)(P^t_W)^{-1})\, \mathrm{d}\Big(\bigotimes_{Z\in\mathcal{A}(W)} \hat{\nu}^t_Z \Big)(B) - a^\Lambda_W \,\mathrm{id}\,, \tag{5.55}$$

with the centred measures $\hat{\nu}^t_Z$ of the roots and the weight $\hat{w}^t(W)$ of the clusters, formed as in [1, Lemmata 9.20 and 9.21] from the expansion (5.52), and with the number a^Λ_W of (ii) below. Let $M_\Lambda(W)$ and $M^{\mathrm{ph}}_\Lambda(W)$ be the majorants of a group with the weights ρ_t and ρ_{ph} for each root [1, Lemma 9.26(d) and Section 9.7]; they are defined by the domains of W alone, and ρ_t is of the order of $g_0(1+2r)$, $\rho_{\mathrm{ph}} \le C_{\mathrm{ph}} g_0^2 (1+2r)^4$.

Proposition 5.12 (The Expansion of a Loop on an Arbitrary Term)**.** *Let γ be so small that* (5.51) *holds together with the thresholds [1, (9.15), (9.20), (9.29)] in which the constant C_0 is replaced by $C_0 + C_\tau$, and let the hypotheses of Lemma 5.11 hold. On every labelled term of the first step and for the exterior data in its support the following hold, with constants which do not depend on the term.*

(i) The identity. *There are numbers $S^\circ_\Lambda \neq 0$ and $q^\circ_\Lambda(\mathfrak{B})$ such that*

$$F^\Lambda = S^\circ_\Lambda \Big[f(\hat{m}) + \sum_{\mathfrak{B}\neq\emptyset} q^\circ_\Lambda(\mathfrak{B})\, \mathrm{tr}(\Upsilon^1_{W_1}\mathsf{U}_1 \cdots \Upsilon^1_{W_n}\mathsf{U}_n) \Big]\,, \tag{5.56}$$

the sum over the finite sets $\mathfrak{B} = \{W_1, \dots, W_n\}$ of pairwise compatible groups, in the cyclic order of their intervals, absolutely convergent, where U_j is π of the ordered product of the links of $U^\Lambda(\hat{m})$ along $\mathcal{H}(W_j)$ and along the traversals between $\mathcal{H}(W_j)$ and $\mathcal{H}(W_{j+1})$.

(ii) The reference gas. *The reference data of the term are the trivial background, the exterior variable zero, and the links outside Ω_1 equal to 1. At these data the first term of* (5.55)*, which does not depend on t there, is a multiple $a^\Lambda_W\,\mathrm{id}$ of the identity; S°_Λ is the partition function of the hard-core gas of the groups with the activities a^Λ_W, $q^\circ_\Lambda(\mathfrak{B}) = \exp(-\sum_\Gamma \Phi^\circ_\Lambda(\Gamma))$ with the sum over the clusters of that gas which contain a group incompatible with a member of $\mathfrak{B}$, and $\sum_\Gamma |\Phi^\circ_\Lambda(\Gamma)| \le \nu\sigma(W)/16$ for the clusters which contain W or a group incompatible with W.*

(iii) The sizes. *For every group W and $t \in \{0,1\}$,*

$$|a^\Lambda_W| \le M^{\mathrm{ph}}_\Lambda(W)\,, \qquad \|\Upsilon^t_W\| \le 2K_\chi M^{\mathrm{ph}}_\Lambda(W)\,, \tag{5.57}$$

and, for every fine traversal $\mathfrak{n}$ and every cube Δ of π_1,

$$\sum_{W\ni\mathfrak{n}} e^{4\nu\sigma(W)} M^{\mathrm{ph}}_\Lambda(W) + \sum_{W\,:\,\Delta\in F(W)} e^{4\nu\sigma(W)} M^{\mathrm{ph}}_\Lambda(W) \le C_{\mathrm{pin}}\rho_{\mathrm{ph}}\,, \tag{5.58}$$

the first sum over the groups with $\mathfrak{n}$ in the interval and the second over those with Δ in the footprint $F(W)$, the set of the cubes of π_1 which meet a constituent domain of W, a host of one of its primitives, or the free-input set of a traversal of its interval [1, Section E.5.2]; the same holds with M_Λ and ρ_t.

Proof. We go through the statements of [1, Sections 9.6 and 9.7] in their order and say, for each, where its proof uses the mean m and the hypothesis that the loops are covered, and why the use is available here. The expansion of the measure at the local shift, (5.52), has been established on every term.

The groups and the identity. By [1, Lemma 9.21(a)] the term of a family $\mathcal{A}$ of final domains in the expansion of $\int f \, \mathrm{d}\nu_t^\Lambda$ vanishes if a domain of $\mathcal{A}$ is not marked, and the remaining terms are regrouped into the sum over the compatible families of groups. Its proof uses of the loop that the link of a traversal depends on the variable of integration through its restriction to $\mathcal{F}_r \cap D$, "the de-younged bonds [being] constants"; for an unmarked domain the function $B \mapsto f(\mathbf{1}_K B + t\hat{m})$ then does not depend on the variables of L_Z and the centred measure of Z integrates it to zero. This property holds on every term, with the convention $\mathcal{F}_r = \emptyset$ outside Ω_1, as we saw. The regrouping is combinatorial, and its convergence rests on the total variations of the centred measures and on the sum of the cluster weights of the vacuum gas which are incompatible with a family, that is, on [1, Lemma 9.20(a) and Corollary 9.16], which hold at the local shift. The reference data are defined on every term, and constant gauge transformations fix them: the exterior variable is zero and the links outside Ω_1 are 1. So the proof of Schur's lemma for the groups [1, Lemma 9.21(b)], which uses the covariance of the reconstruction, of the Hessian, of the cutoffs, and of the hosts under constant gauge transformations, applies as it stands. The identity

$$\int f \, \mathrm{d}\nu_t^\Lambda = S_\Lambda^\circ f(t\hat{m}) + \sum_{\mathfrak{B} \neq \emptyset} S_\Lambda^{\circ[\mathfrak{B}]} R_\Lambda^t(\mathfrak{B})$$

of [1, Lemma 9.21(c)] is the inversion of a finite Boolean sum and an interchange of absolutely convergent sums, and the form $R_\Lambda^t(\mathfrak{B}) = \mathrm{tr}(\Upsilon_{W_1}^t \mathsf{U}_1^t \cdots \Upsilon_{W_n}^t \mathsf{U}_n^t)$ of [1, Lemma 9.26(b)] uses only that a traversal of $\mathcal{H}(W_j)$ has no bond of a root of another group in its free-input set and that the link of a traversal outside all the intervals is taken at $t\hat{m}$, which are the definitions. Dividing by S_Λ° at $t = 1$ we obtain (i), once (ii) is proved.

The reference gas and the first-order sizes. By Lemma 5.11, which holds at the reference data as [1, Lemma 9.25] does, the mixed difference of order $|\mathsf{J}|$ of a closed word of links is at most $u_t^{|\mathsf{J}|}$, with $u_t = C'_\chi g_0(1 + 2r)$, at a base point a_0 with $|a_{0,b}|_\sigma \leq r/4$ and for increments with $|B_b|_\sigma \leq \frac{5}{4} r$; the base points $t\hat{m}$ and the supports of the measures satisfy these conditions by Lemma 5.9(ii). With the comparison [1, Lemma 9.20(c)] this gives $|a_W^\Lambda| \leq M_\Lambda(W)$ and $\sup_h |c_W^t(h)| \leq M_\Lambda(W)$ for the functions $c_W^t(h) := \mathrm{tr}(\Upsilon_W^t \pi(h)) + a_W^\Lambda \chi(h)$, as in [1, Lemmata 9.21(d) and 9.26(e)]. The pins [1, Lemma 9.27(a)] are sums of the majorants over the groups, "the field-independent domain/cluster/interval geometry", and involve neither the carrier nor the loop beyond the counts of the paragraph on the retreated carrier in the proof of Theorem 5.4; for an ensemble they are (5.26). The criterion of Kotecký and Preiss for the reference gas, and with it (ii), follows from the pins and from the bound of a_W^Λ as in [1, Lemma 9.21(d)].

The second-order sizes. The bound $|c_W^t(h)| \leq M_\Lambda^{\mathrm{ph}}(W)$ of [1, Lemma 9.33] improves the weight of a root from ρ_t to ρ_{ph} by three mechanisms [1, Section 9.7]. A root with two or more selected bonds has the factor u_t^2 from the finite differences. A root with one selected bond and a selected host has the factor $2\hat{\omega}_t/\omega_*$ from a complex parameter which multiplies the potentials of its selected hosts [1, Lemma 9.32(c)]; the proof uses the potentials through their bound on the support of the cutoffs, which is (5.50), with the radius $\omega_*/(2\hat{\omega}_t) \geq 1$ of the parameter. A root with one selected bond and no selected host has a measure which is even at $t = 0$ and at the reference data, so that the part of the word which is linear in its variable integrates to zero, and which at $t = 1$ differs from the measure at $t = 0$ by differences of the

cutoffs, paid by the small factor of the large values of the field [1, Lemma 9.32(a), (b)]; no potential occurs in such a measure, hence no linear term, and of the shift only $|\hat{m}_b|_\sigma \le r/4$ is used. The Taylor remainders are bounded by [1, Lemma 9.31(b)], which holds on every term by Lemma 5.11. The proof of [1, Lemma 9.33] combines these with the statements already discussed, and it is here that the first-step paper bounds the functions c_W^t for all groups without the hypothesis on the loops entering otherwise than through the links. This gives (5.57), by Burnside's theorem in the form [1, Lemma 9.26(a)], $\|\Upsilon\| \le K_\chi \sup_h |\mathrm{tr}(\Upsilon\pi(h))|$, and (5.58), by the pins with the weight ρ_{ph}. In the bound $\rho_{\mathrm{ph}} \le C_{\mathrm{ph}} g_0^2 (1+2r)^4$ the constant C_{ph} of [1, Section 9.7] is taken with $C_0 + C_\tau$ in the place of C_0, because $\hat{\omega}_t$ has taken the place of ω_t. *q.e.d.*

The proposition gives the form of a dressed loop on the term, up to three points which the next paragraphs settle. The insertions Υ_W^1 have to be divided into those which vanish with the curvature of the background and the others, and the others have to be attached to the large-field region. The leading word $f(\hat{m})$ and the words U_j are formed with the links of $U^\Lambda(\hat{m})$, which near the complement of the region of integration are not the transports of the background. And the number S_Λ° has to be identified with the step factors times the exponential of a sum of local terms. The first-step paper compares the insertions at $t = 1$ and at $t = 0$ group by group, through marks which decay from the complement of the region [1, Lemmata 9.24 and 9.26(d)], because its shift does not vanish at any distance. With the local shift no such comparison is needed: beyond the range of the shift the two insertions of a group are equal, and the groups within that range are bounded by their sizes and paid by the large-field region as a whole.

Interior groups and atoms. Put $R_{\mathrm{at}} \coloneqq R_t + \rho_L + 1$, with the range R_t of (5.38), and recall that $\ell = LMR_1$ is the side of the cubes of which Λ is a union; for γ small $R_{\mathrm{at}} + 8M + 2L \le \ell$, since R_t grows like $\log g_0^{-2}$ and $R_1 \ge (\log \hat{g}_1^{-2})^r$ with $r = 2$. A group W of a labelled term of the first step is *interior* if every cube of its footprint is at a distance greater than R_{at} from Λ^c. An interior group is *regular* if it is a cross group, or does not embed in the torus, or embeds and none of its comparison regions contains a point of Λ^c. Every other group is an *atom* of the first step, and we assign it to a connected component Z of Λ^c, its *species* being $(1, Z)$: the first component, in a fixed order, which is within the distance R_{at} of a cube of its footprint or, for an interior group, which one of its comparison regions meets. On the term without large-field regions every group is regular.

Lemma 5.13 (The Interior Groups of the First Step)**.** *Under the hypotheses of Proposition 5.12, let W be an interior group of a labelled term of the first step.*

(i) Every free bond with an endpoint in a cube of $F(W)$ belongs to D, the vectors $\hat{m}_b$ and $\hat{\delta}(b)$ vanish for these bonds, and no primitive whose host meets a cube of $F(W)$ has an input outside D.

(ii) $\Upsilon_W^1 = \Upsilon_W^0$.

(iii) The domains and the interval of W define a group of the term without large-field regions, with the same size, and Υ_W^1 and a_W^Λ are the functions of the restriction of the background U_1 to X_W^+ which the insertion and the reference scalar of that group are on that term.

(iv) If W is regular and lies on one loop, the bound (5.29) *holds for it, and with it* (5.13) *at $k = 1$ with the number b_W of the term without large-field regions.*

Proof. (i) A cube of $F(W)$ lies in Λ, being at a positive distance from Λ^c. The carrier D consists of the bonds of I which have an endpoint in Λ, and a free bond which is not in I

has both endpoints outside $\Lambda_1 \supset \Lambda$ [1, Corollary 9.29]; so every free bond with an endpoint in Λ belongs to D, and every bond of $J(\Lambda)$ has both endpoints in Λ^c. A bond b with an endpoint in a cube of $F(W)$ has its initial site within the distance 1 of that cube, so that $d(b,e) > R_{\mathrm{at}} - 1 = R_t + \rho_L$ for every $e \in J_1(\Lambda) \subset J(\Lambda)$, and $\hat{m}_b = 0$, $\hat{\delta}(b) = 0$ by Lemma 5.9(i). The inputs of a primitive lie in its host [1, Lemma 9.13(d)], and a host which meets a cube of $F(W)$ lies within $4M + 1 \le R_{\mathrm{at}}$ of it, hence in Λ, so that its inputs belong to D.

(ii) The insertions (5.55) at $t = 1$ and at $t = 0$ are built from the measures of the selections whose final domains are the roots and the domains of the clusters of W, and from the links along $\mathcal{H}(W)$. In such a measure the dependence on t is in the cutoffs $\hat{\chi}_b^t$ of the bonds which meet the final domain and in the potentials $\tilde{W}_t$ of the hosts in it. By (i) the cutoffs do not depend on t, the linear part of $\tilde{W}_t$ vanishes, and its other part is the sum of the primitives of the host at the argument $B + t\hat{m} = B$ with the exterior inputs multiplied by t [1, Lemma 9.24, proof], of which there are none. The links along $\mathcal{H}(W)$ are functions of the variables on their free-input sets, which lie in cubes of $F(W)$, at the argument $\mathbf{1}_{\mathcal{A}(W)}B + t\hat{m} = \mathbf{1}_{\mathcal{A}(W)}B$. So both insertions are the same integral.

(iii) By [1, Lemma 9.37] the selected integrals at $t = 0$ with a final domain Z are the same for the carrier D and for the carrier I of Bałaban's conditional integral, provided that every bond of I with an endpoint in Z lies in D, and a group all of whose domains have this property is a group for both carriers, with the same weight and the same reference scalar; the proof shows that such an integral involves the entries of the Hessian between the bonds which meet Z, the primitives of the hosts in Z, and nothing else of the term. By (i) this applies to W. Every stencil which meets a bond of D lies in the region $\Omega_1^{\sim -1} \setminus R_1'$ [1, Lemma 3.11(i) and Section 7], in which the entries of the Hessian, the primitives, and the reconstruction of the links are given by the formulae of the term without large-field regions as functions of the background on their stencils: these formulae are those of finitely many types of rows and blocks modulo translations [1, Sections 3.2 and B], and they see the term only through the set of the free bonds in the stencil. The size is defined by the domains and the interval, and the dependence set was determined in the proof of Theorem 5.4.

(iv) The proof of (5.29) in the proof of Theorem 5.4 uses the term in three ways: the rectangle Q_W lies in Λ_1 and outside S_1, it lifts injectively, and the contraction of the group consults the background on the bonds of Q_W only. Here Q_W lies in a comparison region of W, as was shown there, hence in $\Lambda \subset \Lambda_1$, which does not meet S_1 [9, p. 251]; it lifts because W embeds; and the contraction is that of the term without large-field regions, by (iii). *q.e.d.*

Consequently the regular groups of a term are among the groups of the term without large-field regions, with the same insertions as functions of the background, the same sizes, and the same majorants; the sums of the clauses (5.14), (5.21), (5.22), and (5.23) over the regular groups of the term are therefore parts of the sums which were bounded on that term in the proof of Theorem 5.4 and in Corollary 5.8, and the suprema over the space of the step are those of Proposition 5.7, whose proof uses the background on the dependence set of the group only, through the charts and the bounds of the Hessian there.

The links near the complement of the region. The words of (5.56) are formed with the links $\hat{T}_r := U^{\Lambda}(\hat{m})(r)$, while a dressed loop has the transports $T_r := U_1(r)$ of the background. We compare the two bond by bond. Put

$$Y_r := \pi(\hat{T}_r)\pi(T_r)^{-1} - 1 \,, \qquad \pi(\hat{T}_r) = (1 + Y_r)\,\pi(T_r) \,, \tag{5.59}$$

for a fine bond r of a loop with the representation π. There are three cases. If $r \notin \Omega_1$, then $\hat{T}_r = V_0(r) = U_1(r)$, because the background is the critical point among the configurations

which equal V_0 on Ω_1^c [9, (1.12), p. 248], and $Y_r = 0$. If $r \in \Omega_1$, then $\hat{T}_r = e^{iq(r)}U_1(r)$ with the phase $q(r)$ of the reconstruction at the free variables on $\mathcal{F}_r$, which are $\hat{m}$ on D and the exterior variable on the other bonds, and $q(r) = 0$ if all of these vanish [1, Lemma 3.5]. Now $\hat{m}_b \neq 0$ only if $d(b, J_1(\Lambda)) \leq R_t$, the bonds on which the exterior variable lives have both endpoints in Λ^c, and $\mathcal{F}_r$ lies within $2L+1$ of r. Hence

$$Y_r \neq 0 \quad \text{only if} \quad \operatorname{dist}(r, \Lambda^c) \leq R_t + 2L + 2 \leq R_{\text{at}} \,, \tag{5.60}$$

because $\rho_L = 4L$. Among these bonds we distinguish those whose row is within $2R_{\text{c}} + 2L$ of I, the *near* bonds, from the others. For a near bond the input of the row lies in the polydisc of the row, by the proof of Lemma 5.11, and the phase is small with the coupling. A bond which is not near has no bond of I in its free-input set, so that its phase does not involve $\hat{m}$ or the labels of the retreat, and it lies within $2L$ of Λ_1^c, because a free bond outside I has both endpoints outside Λ_1; its phase is real, so that $\|Y_r\| \leq 2$, and nothing better holds, since the exterior variable is not small there.

The phase of a near bond depends on the term through $\hat{m}$, hence on the labels and on the exterior variable within the distance R_t, which is of the order of $\log g_0^{-2}$. An insertion of the class has to be localised in a set whose size is comparable with its own, and we therefore decompose Y_r according to the boxes of the proof of Lemma 5.9 through which $\hat{m}$ reaches the bonds of $\mathcal{F}_r$. For a family $s = (s_\Delta)$ of complex numbers indexed by the boxes put

$$K(s) \coloneqq \sum_\Delta s_\Delta K_\Delta \,, \qquad \hat{m}(s) \coloneqq -\lambda P \sum_{n=0}^{N} K(s)^n \ell_\Lambda \,, \tag{5.61}$$

so that $\hat{m}(1) = \hat{m}$, and let $Y_r(s)$ be the expression (5.59) with $\hat{m}(s)$ in the place of $\hat{m}$ in the free variables. Call two boxes *adjacent* if their distance is at most $5\ell_s$; a box has at most $N_{\text{adj}} \coloneqq 13^4$ boxes adjacent to it or equal to it, since an interval of length less than $12\ell_s$ meets at most thirteen intervals of length at least ℓ_s. Let $\mathrm{St}(r)$ be the set of the boxes Δ with $\tilde{\Delta} \cap \mathcal{F}_r \neq \emptyset$; they are pairwise adjacent, because $\mathcal{F}_r$ has diameter at most $4L + 2 \leq \ell_s + 1$. Put

$$R_0 \coloneqq 1 + 2N_{\text{adj}}^2 e^{32\nu} \,, \qquad C_{\text{lk}} \coloneqq \frac{4\sqrt{2}\, C_\pi Q_* C_{\mathsf{C}}}{\vartheta_*} \,, \qquad \bar{Y} \coloneqq C_{\text{lk}} m_* \Big(\frac{N_{\text{nb}}}{2c_{\text{h}}} + 1 \Big) g_0 (1+r) \,, \tag{5.62}$$

with the constants C_π, Q_*, C_{C}, ϑ_*, m_* of [1, Lemma 9.25], and suppose that the integer ℓ_s of Lemma 5.9, which may be taken larger, satisfies $R_0 e^{-\mu_H \ell_s/2} \leq \frac{1}{2}$, that $LM \geq 4\ell_s + 4\rho_L$, and that γ is so small that $\bar{Y} \leq 1$ and $C_{\mathsf{C}} m_* (N_{\text{nb}}/(2c_{\text{h}}) + 1) g_0 (1+r) \leq \vartheta_*/2$.

Lemma 5.14 (The Pieces of a Link)**.** *On a labelled term of the first step and for the exterior data in its support, let r be a near bond of a loop.*

(i) The function $Y_r(s)$ is holomorphic on the polydisc $|s_\Delta| \leq R_0$ and $\|Y_r(s)\| \leq \bar{Y}$ there.

(ii) $Y_r = \sum_X Y_{r,X}$, the sum over the finite sets X of boxes, with

$$Y_{r,X} \coloneqq \prod_{\Delta \in X} \int_0^1 \mathrm{d}s_\Delta \, \partial_{s_\Delta} Y_r(s) \Big|_{s_{\Delta'}=0,\ \Delta' \notin X} \,, \qquad \|Y_{r,X}\| \leq \bar{Y} (R_0 - 1)^{-|X|} \,, \tag{5.63}$$

and $Y_{r,X} = 0$ unless every box of X is joined to a box of $\mathrm{St}(r)$ by a chain of adjacent boxes of X.

(iii) $Y_{r,X}$ *depends on the background, on the labels of the term, and on the exterior variable only through their restrictions to the set*

$$D_r(X) \coloneqq B_\infty(r, 2L+2) \cup \bigcup_{\Delta \in X \cup \mathrm{St}(r)} B_\infty(\Delta, \ell_s + 2\rho_L) \,.$$

The cubes of π_1 *which meet* $D_r(X)$ *form a set which is connected in the sense of Definition 5.2(vii), and their number* $\sigma_{r,X}$ *satisfies* $\sigma_{r,X} \le 16(1 + N_{\mathrm{adj}} + |X|)$*; every one of them lies within* $2(|X| + N_{\mathrm{adj}} + 1)$ *sides of a cube which contains a site of* r*.*

(iv) For every near bond r*,*

$$\sum_X e^{32\nu(1+N_{\mathrm{adj}}+|X|)} \|Y_{r,X}\| \le W_{\mathrm{lk}} \bar{Y} \,, \qquad W_{\mathrm{lk}} \coloneqq 2^{N_{\mathrm{adj}}+2} N_{\mathrm{adj}}^{2N_{\mathrm{adj}}} e^{32\nu(N_{\mathrm{adj}}+1)} \,, \tag{5.64}$$

and the exponent on the left is at least $2\nu\sigma_{r,X}$*.*

Proof. (i) By the proof of Lemma 5.9, $\|K_\Delta\|_\infty \le \varepsilon_s$ with rows in $\hat{\Delta}$, so that $\|K(s)\|_\infty \le R_0 q_s \le R_0 e^{-\mu_H \ell_s/2} \le \frac{1}{2}$ on the polydisc, and $\sum_{n \le N} \|K(s)\|_\infty^n \le 2$. The entries of G_Δ satisfy (5.35) and its rows lie in $\tilde{\Delta}$, which contains a given bond for at most N_{nb} boxes; hence $\|P\|_\infty \le N_{\mathrm{nb}} \gamma^{-1} N_{\mathrm{lat}}(\mu_H)$, and with (5.36) and $\lambda \le 1$

$$\sup_b h_{\mathbb{C}}(\hat{m}(s)_b) \le 2N_{\mathrm{nb}} \gamma^{-1} N_{\mathrm{lat}}(\mu_H)\, m_L \vartheta_V r = \frac{N_{\mathrm{nb}}}{2c_{\mathrm{h}}}\, r \,,$$

by the definitions of C_H^{near} and ϑ_V. The other free variables on $\mathcal{F}_r$ are real and less than r in norm on the support of the term, as in the proof of Lemma 5.11. The phase of the row is $A = g_0 \mathsf{C}_U B$ with the free variables B on the at most $m_* - 1$ bonds of $\mathcal{F}_r$, and $\|\mathsf{C}_U e_b\| \le C_{\mathsf{C}}$, so that $\|A\| \le g_0 C_{\mathsf{C}} \sum_b h_{\mathbb{C}}(B_b) \le \vartheta_*/2$ by the last condition before the lemma. On the closed ball of this radius the phase $q(r)$ is a holomorphic function $\Phi(A)$ with $h_{\mathbb{C}}(\Phi) \le Q_*$ [1, Lemma 9.25], and $\Phi(0) = 0$, so that $h_{\mathbb{C}}(\Phi(A)) \le (2Q_*/\vartheta_*)\|A\|$ by Schwarz's lemma; for a free bond r the phase is $g_0 B_r$. Since $Y_r(s) = e^{i\pi(q(r))} - 1$ with $\|\pi(X)\| \le \sqrt{2} C_\pi h_{\mathbb{C}}(X)$, and $\|e^{iZ} - 1\| \le 2\|Z\|$ for $\|Z\| \le 1$, we obtain $\|Y_r(s)\| \le C_{\mathrm{lk}} g_0 \sum_{b \in \mathcal{F}_r} h_{\mathbb{C}}(B_b) \le \bar{Y}$. The function is holomorphic in s because $\hat{m}(s)$ is a polynomial in s.

(ii) The function $Y_r(s)$ depends on finitely many of the variables s_Δ, and the fundamental theorem of calculus in each of them gives $Y_r(1) = \sum_X Y_{r,X}$, as in (6.7). Cauchy's estimate on the circles of radius $R_0 - 1$ about the points of $[0,1]$, which lie in the polydisc, gives the bound. With $s_{\Delta'} = 0$ outside X, the vector $\hat{m}(s)_b$ for $b \in \mathcal{F}_r$ is a sum of the terms $G_{\Delta_0} \chi_{\Delta_0} s_{\Delta_1} K_{\Delta_1} \cdots s_{\Delta_n} K_{\Delta_n} \ell_\Lambda$ with $\Delta_0 \in \mathrm{St}(r)$ and $\Delta_1, \dots, \Delta_n \in X$, and such a term vanishes unless consecutive boxes are at a distance of at most $\ell_s + \rho_L \le 5\ell_s$: the columns of K_{Δ_i} and of χ_{Δ_0} lie in the box and the rows of $K_{\Delta_{i+1}}$ in $\hat{\Delta}_{i+1}$, by the proof of Lemma 5.9. If a box of X is not joined to $\mathrm{St}(r)$ through adjacent boxes of X, the function does not depend on its variable, and the derivative vanishes.

(iii) The link consults the background on the stencil of its row, within $2L + 2$ of r [1, Lemma 3.11], and the exterior variable on $\mathcal{F}_r$. In the terms of (ii), G_Δ involves the entries of the Hessian between the bonds of D in $\tilde{\Delta}$, K_Δ those between $\hat{\Delta}$ and $\tilde{\Delta}$, and $\ell_\Lambda(c)$ the exterior variable and the entries within ρ_L of $c \in \Delta_n$; an entry of the Hessian consults the background within ρ_L of its bonds. All of this lies in $D_r(X)$. Each of the at most $1 + N_{\mathrm{adj}} + |X|$ sets of which $D_r(X)$ is the union has a diameter less than $4\ell_s + 4\rho_L \le LM$, so that it meets at most 2^4 cubes of π_1, and these have pairwise a point in common. The sets form a chain in which consecutive ones are at a distance less than LM, by (ii), so that their cubes are equal or have a point in common; this gives the connectedness, the count, and the distance.

(iv) A set X with k boxes for which $X \cup \mathrm{St}(r)$ is connected under adjacency is determined by a connected set of at most $k+N_{\mathrm{adj}}$ boxes through a fixed box of $\mathrm{St}(r)$ and by its intersection with $\mathrm{St}(r)$. A connected set of n boxes through a given box is the set of the vertices of a closed walk of $2(n-1)$ steps, each to one of fewer than N_{adj} adjacent boxes, so that there are at most $N_{\mathrm{adj}}^{2(n-1)}$ of them, and at most $2N_{\mathrm{adj}}^{2(k+N_{\mathrm{adj}})}$ with at most $k+N_{\mathrm{adj}}$ boxes; hence the sets X with k boxes are at most $2^{N_{\mathrm{adj}}+1}N_{\mathrm{adj}}^{2N_{\mathrm{adj}}}(N_{\mathrm{adj}}^2)^k$ in number. With (ii),

$$\sum_X e^{32\nu(1+N_{\mathrm{adj}}+|X|)}\|Y_{r,X}\| \le 2^{N_{\mathrm{adj}}+1}N_{\mathrm{adj}}^{2N_{\mathrm{adj}}}e^{32\nu(N_{\mathrm{adj}}+1)}\,\bar{Y}\sum_{k\ge 0}\Big(\frac{N_{\mathrm{adj}}^2 e^{32\nu}}{R_0-1}\Big)^k ,$$

and the ratio of the series is $\frac{1}{2}$ by the choice of R_0. The last statement is the count of (iii). *q.e.d.*

The form of a dressed loop on a term. We insert (5.59) into the leading word and into the words U_j of (5.56), for the bonds outside the intervals of the groups, and expand the products: every such bond contributes the transport $\pi(T_r)$ of the background, or one of the pieces $Y_{r,X}\pi(T_r)$ if it is a near bond, or $Y_r\pi(T_r)$ if it is a bond with $Y_r \neq 0$ which is not near. The links inside the interval of a group W of the family are not expanded: they are collected with Υ^1_W into

$$Y_W := \Upsilon^1_W\,\pi(P^1_W)\,, \tag{5.65}$$

the insertion of the group at its interval $\mathcal{H}(W)$. A piece of a link is treated as a group of its own, a *link group*: its only interval is the bond r, its localisation domain is the set of the cubes of π_1 which meet $D_r(X)$, or $B_\infty(r, 2L+2)$ for a bond which is not near, its size is the number of these cubes, its insertion is $Y_{r,X}\pi(T_r)$, respectively $Y_r\pi(T_r)$, and it is an atom of the species $(1, Z)$ for the first component Z of Λ^c within the distance R_{at} of r, which exists by (5.60). A link group is incompatible with the groups whose intervals contain its bond and with the other link groups of the same bond, and with no others; in particular it is compatible with every group of the reference gas whose interval does not contain r, whatever their domains, and no cluster of the reference gas is counted as incompatible with it, so that $\theta = 0$ for a link group. With these conventions (5.56) reads

$$F^\Lambda = S^\circ_\Lambda\Big[\chi(U_1(C)) + \sum_{\mathfrak{B}\neq\emptyset} Q_1(\mathfrak{B})\,w_1(\mathfrak{B})\Big]\,, \qquad Q_1(\mathfrak{B}) := q^\circ_\Lambda(\mathfrak{B}')\,, \tag{5.66}$$

the sum over the families $\mathfrak{B}$ of pairwise compatible groups and link groups with pairwise disjoint intervals, where $\mathfrak{B}'$ is the set of the groups of $\mathfrak{B}$ which are not link groups and $w_1(\mathfrak{B})$ is the word of Section 5.3, formed with the insertions (5.65), those of the link groups, and the transports of U_1. The sum converges absolutely, by Proposition 5.12 and because the loop has finitely many bonds, each with pieces whose norms are summable by (5.64).

6 The Inductive Step

We now show that a renormalisation step maps a dressed loop of the step k to one of the step $k+1$, with the same constants. The difficulty is one of counting. A loop has $P_k(C)$ traversals at the step k, and $P_k(C) = P_q(C)a_k$ grows like L^{K-k}, so that every quantity which a step contributes in proportion to the perimeter is, summed over the loop, unbounded as $K \to \infty$, even when it is of order g_k^2 per traversal. Such quantities can therefore be neither estimated nor neglected; each of them has to be either absorbed exactly into the step factor κ_{k+1}, or carried along without ever being summed in absolute value over the loop. The deviation parts

of the insertions are of the second kind, and what keeps them under control is a contraction: a traversal of the step $k+1$ consists of L traversals of the step k, while the deviation of the background, measured in units of the scale $k+1$, is L^2 times the same deviation measured in units of the scale k, so that with the normalisation $\rho_k \delta^*$ of (5.11) an old insertion contributes to the new bound with a factor of order $L \cdot L^{-2} = L^{-1}$.

Theorem 6.1 (The Inductive Step)**.** *Let L, Λ_*, $\bar{g}$, $\mathcal{C}$, and $K \geq K_{\text{tor}}(\mathcal{C})$ be as in Theorem 5.4, and let $1 \leq k < K'$. If the ensemble is dressed at the step k, then it is dressed at the step $k+1$, with the same constants in Definition 5.2, and* (5.2) *holds at $k+1$.*

Corollary 6.2 (Dressed Loops at Every Step)**.** *Under the hypotheses of Theorem 5.4, the ensemble is dressed at every step $1 \leq k \leq K'$, with constants independent of k, K, the torus, and the number of loops.*

Proof. Induction on k, with Theorem 5.4 as its base and Theorem 6.1 as its step. *q.e.d.*

The proof of Theorem 6.1 occupies the rest of this section. It follows Bałaban's step [9, Sect. 3] on the terms with small fields, with the dressed loop carried through each of its operations, his treatment of large fields [12, Sect. 1] for the terms whose large-field regions grow, and the **R**-operation of [11, 12]. We fix a labelled term of ϱ_k and write U_k for its background field, U_{k+1} for the background field of the step $k+1$, and $V^{(k)} := M^k(U_{k+1})$ for the block field of the step k determined by U_{k+1}.

6.1 The Evaluation Point

The fluctuation field of the step $k+1$ is the deviation of V_k from $V^{(k)}$, and every factor of the dressed loop has to be expanded around the configuration at which this deviation vanishes. Two facts make this possible. First, $V^{(k)} = M^k(U_{k+1})$ [8, (2.3), p. 265], and since the minimal orbit with given averages is unique [9, p. 256], $U_k(V^{(k)}) = U_{k+1}$ in the gauge of the step. Secondly, $V^{(k)}$ lies in the support of the test functions of the step k on $\Omega^{\sim}_{k+1}$: by [9, (3.19), p. 268] the block field $V^{(k)}$ is within ϑ_k of the local reference fields on the cubes of [9, (3.16)], so that it lies in the support of the test function [9, (3.3), p. 265] with threshold $2\vartheta_k$, and Bałaban concludes that "$\chi_k(\square) = 1$ for $\square \subset \Omega^{\sim}_{k+1}$" [9, p. 266]. For the soft test functions the same holds with the margin of Lemma 3.8. On the cubes of the band $\Omega^{\sim}_{k+1} \setminus \Omega^{\sim -1}_{k+1}$, which lie in no cube of [9, (3.16)], the localisation of the minimiser with constraints of two scales is needed; it is not printed, and it is proved in Supplement, Proposition S.8, from the uniqueness of the minimal orbit [7, Theorem 1].

The regularity of the new background is [9, p. 268]: $|\partial U_{k+1} - 1| < 2B_3 \varepsilon_{k+1} (L^{-1}\eta)^2$ on Ω_{k+1}. By the lasso identity, which writes the holonomy around a plaquette of the scale j as an ordered product of conjugates of the holonomies of the L^{2j} fine plaquettes inside it, and by $|\prod_i A_i - 1| \leq \sum_i |A_i - 1|$ for unitary A_i, a plaquette of the scale k in Ω_{k+1} has deviation at most $2B_3 \varepsilon_{k+1} L^{-2}$ under U_{k+1}. Consequently, for the deviation (5.7),

$$\delta^{*,(k+1)}_{\mathfrak{n}'}(U) = L^2\, \delta^{*,(k)}_{\mathfrak{n}'}(U) \qquad \text{and} \qquad \delta^{*,(k+1)}_{\mathfrak{n}'}(U_{k+1}) \leq 2c_\sigma B_3 \varepsilon_{k+1} \sum_{t \geq 1} (t+1)^3 e^{-\eta_w t}\,, \quad (6.1)$$

where the superscript indicates the scale whose units are used, the first identity holds for the same plaquettes and radii, and c_σ converts Bałaban's matrix norm into the norm of the insertions.

6.2 The Identity of the Step

The first thing to establish is that the step can be made with the dressed loop inside the integrand. Bałaban passes from the density ϱ_k to the expansion of $T\varrho_k$ by a sequence of operations on the integrand of a fixed term [9, pp. 264–270], and for the density only the integral of the result matters. The dressed loop is a further factor, which depends on the integration variables, and each operation has to be an identity for the product. Most of them are identities for every integrand. Two kinds are not: the gauge fixing requires that the integrand be gauge invariant, and the changes of variables have to be changes of variables in the proper sense, bijections between the domains of the old and of the new variables, with which the further factor can be composed.

We fix a labelled term of ϱ_k and sets Ω_{k+1}, Λ_{k+1}, S_{k+1} of the step. On the bonds of Ω_{k+1} the step writes the block field as $V_k = e^{iB'}V^{(k)}$ with $B' = g_k C A_k - h\tilde{D}(g_k C A_k)$, where A_k is the fluctuation field [9, (3.11), p. 266; (3.15), pp. 267–268], and on the bonds of S_{k+1} it changes the variables by the map Ψ_{k+1} of [9, p. 269] (Supplement, Lemma 17.23). Let

$$F \coloneqq \mathcal{W}_k(e^{iB'}V^{(k)}) \circ \Psi_{k+1} \tag{6.2}$$

be the dressed loop of the term at this block field, as a function of the new fluctuation variables, of V_{k+1}, and of the outer variables. The field A_k is integrated on Λ_{k+1}, where we write A for it, and it is an outer variable of the step on $\Omega_{k+1} \cap \Lambda_{k+1}^c$ [9, p. 270]. Let $\mathcal{Z}_{k+1}$ be the last integral of [9, (3.25), p. 270], the integral over A with its factor $z^{(k)}$, and let $\langle\,\cdot\,\rangle$ be the expectation in the probability measure [9, (3.27), p. 271] at $s = t = 1$, which is the integrand of that integral divided by $\mathcal{Z}_{k+1}$.

Proposition 6.3 (The Identity of the Step)**.** *Let $1 \le k < K'$, and let the ensemble be dressed at the step k. Then*

$$(TN_k)(V_{k+1}) = \sum \chi_{k+1}(\Omega_{k+1})\, \mathbf{T}_{k+1}\Big[\exp\Big(A_k\Big(\frac{1}{g_k^2}, U_{k+1}\Big) + E_0^{(k)}\Big)\, \mathcal{Z}_{k+1}\,\langle F\rangle\Big]\,, \tag{6.3}$$

the sum over the sequences of domains of [9, (3.25)]; this is Bałaban's expansion of $T\varrho_k$ with the factor $\langle F\rangle$ in each term. The factor $\langle F\rangle$ is a function of V_{k+1} and of the outer variables of $\mathbf{T}_{k+1}$, which are those of $\mathbf{T}_k$, the field A_k on $\Omega_{k+1} \cap \Lambda_{k+1}^c$, and the field V_k on Ω_{k+1}^c.

Proof. By Definition 5.2, $N_k = \sum \chi_k(\Omega_k)\mathbf{T}_k[\mathcal{W}_k \exp A_k]$, and TN_k is the expression [9, (3.1), p. 264] with the factor $\mathcal{W}_k$ in each term. We follow the operations by which Bałaban transforms a term of (3.1) into the terms of (3.25), and we show that each of them is an identity when the integrand is multiplied by a bounded Borel function of the integration variables, composed with the changes of variables where there are any. The function $\mathcal{W}_k$ is of this kind. Indeed, the transports of a word are unitary at real fields, so that by Lemma 5.1 and Definition 5.2(i)

$$\Big|\sum_{\mathfrak{B}\neq\emptyset} Q_k(\mathfrak{B})\, w_k(\mathfrak{B})\Big| \le \prod_W (1 + e^{\theta(W)}\|Y_W\|) \le \exp\Big(\sum_W e^{\theta(W)}\|Y_W\|\Big)\,,$$

and the last sum is at most the sum, over the finitely many traversals of the torus, of the bounds of Definition 5.2(ii), which are bounded on the support of the test functions of the term.

Decompositions of unity. The decompositions [9, (3.2), (3.3), p. 265; (3.16), p. 268] multiply the integrand by a sum which equals one, for the soft test functions by Lemma 3.8(i). They hold for every integrand.

Test functions which equal one. On $\Omega^{\sim}_{k+1}$ the test functions χ_k and χ_{Ax} of the step k are removed, because they equal one on the support of the new ones [9, (3.8), (3.9), p. 266], and on the cubes of R^c_{k+1} the test function of [9, (3.3)] is removed for the same reason [9, (3.17)–(3.19), pp. 268–269]; for the soft test functions the removals hold on the plateaus (Supplement, Proposition S.2(a)). These are identities between factors of the integrand at each point, and they hold for every integrand.

Regrouping. At fixed Ω_{k+1} the terms are summed over the sets P_{k+1} and Q_{k+1} which determine it, and "the resummation applies only to the characteristic functions and the gauge fixing terms localized in $\Omega_k \backslash \Omega_{k+1}$" [9, p. 267]. The sets R_{k+1} are then replaced by the pairs (Λ_{k+1}, S_{k+1}) of [9, (3.20), p. 269]. The dressed loop is a function of V_k and of the outer variables which does not depend on these sets, so that it is a common factor of the terms which are regrouped.

The gauge fixing. In the blocks of a set P^1_{k+1} of points of the scale $k+1$ Bałaban introduces "the axial gauge fixing as in (1.5), with V_k, ε_k, g_k instead of U, ε_0, g_0", and obtains "an expansion of the form (1.6)" [9, p. 265]. The identity [9, (1.5), p. 247] is an integral over gauge transformations $u(x)$ at the points $x \neq y$ of the blocks $B(y)$, $y \in P^1_{k+1}$, and the Faddeev–Popov procedure which follows it replaces V_k by a gauge transform under the integral. It uses that the other factors of the integrand are invariant under the gauge transformations u of V_k which are supported at these points, in Bałaban's words "the invariance of all the expressions determining the density with respect to the gauge transformations of the new field" [9, p. 250]. We show that $\mathcal{W}_k(V_k^u) = \mathcal{W}_k(V_k)$ for these u. Let $\tilde{u}(x') := u(x)$ for the fine points x' of the block $B^k(x)$ of a point x of the scale k, which is the extension that [9, p. 250] uses at the first step. The averages are covariant, $M(\{u\mathbf{U}_j v\}) = uM(\{\mathbf{U}_j\})v$ [8, (0.6), p. 253], and the action is invariant, so that $U \mapsto U^{\tilde{u}}$ maps the minimal orbit of V_k onto that of V_k^u. The representative $U_k(V)$ is singled out in the minimal orbit by axial conditions inside the blocks, at every level of their hierarchy: conditions on the averaged contour variables $\mathbf{U}(y,x)$ of [8, (0.11), p. 253], or on the transports along fixed contours [6, (1.15), p. 78]. In a block $B^k(x)$ the transformation $\tilde{u}$ is constant, so that it conjugates these variables by $u(x)$ and preserves the conditions, and outside these blocks it equals one. Since the conditions "determine uniquely an element in each orbit" [6, p. 78], it follows that $U_k(V_k^u) = U_k(V_k)^{\tilde{u}}$. By Definition 5.2(iii) the factors of the words are covariant, so that a closed word is invariant under a gauge transformation of the background which leaves the outer variables fixed; and $\tilde{u}$ leaves them fixed, because the blocks of P^1_{k+1} lie in $(Z_k'^{\sim 5})^c$ [9, p. 265], whereas the outer variables of the term are fields on bonds of the large-field region $Z_k \subset Z_k'$. This proves the invariance.

The fluctuation field. On $B(\Gamma_{k+1})$ the block field is written $V_k = V_k' V^{(k)}$ and $A_k' = (1/i)\log V_k'$; the δ-functions of the averaging are linearised and removed "using the operator C", and the field is scaled, $A_k' = g_k A_k$ [9, (3.11), p. 266; p. 267]. Here "all the applied formulas are the same as in Sect. 2 [I]", where the δ-functions serve to "eliminate the variables $B'(b_0(c))$" [8, p. 268]. These operations express the integration variables on the support of the δ-functions through the variables A_k, with a Jacobian which Bałaban includes in the action, and every other factor of the integrand is composed with this parametrisation. For the dressed loop the result is the function $\mathcal{W}_k(e^{iB'}V^{(k)})$ of (6.2).

The change of variables on S_{k+1}. By Supplement, Lemma 17.23, whose two hypotheses hold when γ is small, the map Ψ_{k+1} is a diffeomorphism between open sets which contain the supports of the test functions in the new and in the old variables. Therefore the integral of Bałaban's integrand times a bounded Borel function of the old variables is the integral of his integrand in the new variables times the composition of that function with Ψ_{k+1}, which

for the dressed loop is F.

The conditional integral. Finally the integral over A_k is split into the integral over its restriction A to Λ_{k+1} and the integral over its restriction to $\Omega_{k+1} \cap \Lambda_{k+1}^c$, which becomes part of the operation $\mathbf{T}^{(k+1)}$, the quadratic form being completed as in [9, (3.23)–(3.25), p. 270]. This is Fubini's theorem on the finite torus for an integrable function, and it holds with the bounded factor F.

After these operations the term of (3.1) with the factor $\mathcal{W}_k$ is the sum of the terms of (3.25) with the factor F under the last integral. We have $\mathcal{Z}_{k+1} \neq 0$, since Bałaban takes the logarithm of this integral in [9, (3.26), p. 271], and by the definition of the expectation the last integral with the factor F equals $\mathcal{Z}_{k+1}\langle F\rangle$. This is (6.3). The variables on which $\langle F\rangle$ depends are those of F which are not integrated. *q.e.d.*

Since the averaging preserves integrals (Section 5.1), the proposition gives $\int (TN_k)\,\mathrm{d}V_{k+1} = \int N_k\,\mathrm{d}V_k$, and the task of the step is to write $\langle F\rangle$ in the form of a dressed loop of the step $k+1$. At the steps at which the **R**-operation acts it is applied afterwards (Section 6.15).

6.3 The Expansion of the Words in the Localisation Parameters

To write $\langle F\rangle$ as a dressed loop we first have to see on which part of the fluctuation field each word of F depends, and the difficulty is that this dependence is not local. By [8, p. 265 and (2.6), p. 266], whose formulae [9, p. 267] transfers to the operators of a sequence of domains, the background of the step k at the block field $e^{iB'}V^{(k)}$ is a gauge transform of $e^{i\eta \mathbf{H}_k(B')}U_{k+1}$, where $\mathbf{H}_k$ is the function [10, (1.17), p. 6] of the fluctuation variable B' of [10, (1.19), p. 6], built from the propagators H, $\tilde{G}$, and H_0 of the step. These propagators have kernels which decay exponentially and do not vanish at any distance, so that the transport of the new configuration along a single bond of a loop depends on the fluctuation field on the whole lattice. Bałaban meets the same difficulty for the terms of the effective action, and he resolves it by parameters which switch off the propagators across the walls of a partition [10, pp. 3–7]. We apply his device to the words of the dressed loop.

The parameters. Let σ_k be the partition of the torus into the cubes Δ of side M of the scale k, the family which [10, p. 3] takes in its example, here with no cube exempt. To each cube we assign a parameter $s(\Delta)$, and we define the propagators $H(s)$, $\tilde{G}(s)$, $H_0(s)$ by the rule of [10, p. 3]: the term of a walk ω in the expansion [10, (1.6)] of a propagator is multiplied by $s(\Delta_1)\cdots s(\Delta_m)$, where $\Delta_1, \ldots, \Delta_m$ are the cubes of σ_k which meet the localisation domain of ω. Let $\mathbf{H}(s, B')$ be the function [10, (1.17)] with these propagators; by [10, (1.5), p. 3] the solutions which enter it are functionals of the propagators with the properties of the original ones. For a set $\sigma \subset \sigma_k$ let $Y(\sigma)$ be the union of its cubes, and call a union of cubes *connected* if any two of its cubes are joined by a chain of its cubes in which consecutive ones share a wall of dimension $d-1$ [10, p. 4]; the *components* of $Y(\sigma)$ are its maximal connected parts. We use three properties.

(a) $\mathbf{H}(1, B') = \mathbf{H}_k(B')$, since the propagators at $s = 1$ are the original ones [10, p. 3], and $\mathbf{H}(0, B') = 0$, since the localisation domain of every walk meets a cube, so that all three propagators vanish at $s = 0$.

(b) If $s(\Delta) = 0$ for every $\Delta \notin \sigma$, then $\mathbf{H}(s, B')$ vanishes on a neighbourhood of $Y(\sigma)^c$, and on a component Y of $Y(\sigma)$ it depends only on the restrictions of s, of B', and of the configuration to Y; we write $\mathbf{H}_Y(s, B')$ for it. This is the argument of [10, p. 4]: the kernels of the propagators vanish unless both arguments lie in the interior of one

component, the equations [10, (1.3), (1.4)] separate into equations in the components, and B' is a local function of the fluctuation field.

(c) For complex parameters with $|s(\Delta)| \le e^{\kappa_1}$, where κ_1 is the constant of [10, p. 5] with $\delta_1 M \ge \kappa_1$, and for $|B'| < \varepsilon_3$, the function $\mathbf{H}_Y(s, B')$ is analytic in s and B' and

$$|\mathbf{H}_Y(s, B')| \le 4B_0 e^{16\kappa_1} |B'|_Y \ , \tag{6.4}$$

with $|B'|_Y$ the supremum over Y, by [10, (1.18), p. 6] applied to B' restricted to Y; the constants B_0, ε_2, ε_3 are those of [10, pp. 3, 5, 6], with $4C_2 B_0 e^{16\kappa_1}\varepsilon_2 \le 1$, $4C_4 B_0^2 e^{32\kappa_1}\varepsilon_3 \le 1$, and $2B_0 e^{16\kappa_1}\varepsilon_3 \le \varepsilon_2$, and "all the bounds above are uniform on the domain" of the configurations [10, p. 6]. None of the estimates [10, (1.11)–(1.18)] uses the cube which Bałaban exempts, so that they hold with every cube carrying a parameter.

The term and the domain of a member. In this subsection the labelled term of ϱ_k is the term without large-field regions, for which $\Omega_j = \Lambda_j$ is the whole torus for every $j \le k$ and there are no outer variables; the sets Ω_{k+1}, Λ_{k+1}, S_{k+1} of the step are arbitrary. On this term the operators of the step k are those of one domain, to which [10] applies as it stands. The terms of ϱ_k with large-field regions, on which the propagators are those of a sequence of domains and the factors of the words depend on outer variables, are the subject of Section 6.15. For a group W of a family of the step k, made at a step $j \le k$, let X_W be its domain and X_W^+ its dependence set (Section 5.3); the domain contains the cubes of π_j that meet a fine bond of its intervals, and the insertion Y_W depends on the background field only through its restriction to X_W^+ (Definition 5.2(iii)).

The domain of the factors. The words have to be evaluated at the configurations $e^{i\eta\mathbf{H}(s,B')}U_{k+1}$, which for $s \ne 1$ are not backgrounds of block fields, and we have to make sure that their factors are defined there. On the support of the test functions of the step, $|g_k A_k| \le r_y \vartheta_k$ on every bond; this is the bound [9, (3.14), p. 267] in the form which the soft test functions give it, with the constant r_y of Supplement, Lemma 17.23, in whose proof this form is derived. For γ small the right side is below the constant ε_1 of [10, p. 7], and therefore, by [10, (1.21), p. 7],

$$|\mathbf{H}(s, B')| \le h_k \coloneqq 4B_0 C_1 e^{16\kappa_1} r_y \vartheta_k \tag{6.5}$$

for $|s(\Delta)| \le e^{\kappa_1}$, "and the same for all admissible norms", with the absolute constant C_1 of [10, (1.20), p. 6]. Among these norms are those of the condition [8, (3.14), p. 272], under which Bałaban evaluates the terms of the action of the earlier steps at a configuration $e^{i\xi\mathbf{A}}\mathbf{U}$, with the argument [8, (3.13)] in the place of $\mathbf{J}$: he requires of $\mathbf{H}_k(s, B')$ the bounds of that condition [10, p. 7], and he obtains them from [10, (1.21)]. The factors of the dressed loop are analytic on the same spaces as those terms (Definition 5.2(iii)). In the units of the step k the radii of the space of the step $k+1$ are smaller than those of the step k by at least $\frac{1}{2}\min(\beta, 1)\min(\alpha_{0,k}, \alpha_{1,k})$: on $\Omega_k \setminus \Omega_{k+1}$ by the factor $1 - \frac{1}{2}\beta$ of [9, (2.34)–(2.39), p. 261], and on Ω_{k+1} because $\alpha_{i,k+1}/L \le \frac{1}{2}\alpha_{i,k}$, by (3.18) and $(1+\beta_0)^2 \le L/2$. Let $c_{\mathbf{J}} \ge 1$ be an absolute constant such that, for a point $(\mathbf{U}, \mathbf{J})$ of the space of the step $k+1$ and a field $\mathbf{A}$ whose norms in [8, (3.14)] are at most $a \le 1$, the configuration $e^{i\xi\mathbf{A}}\mathbf{U}$ with the argument [8, (3.13)] lies in the space of the step k whenever $c_{\mathbf{J}} a \le \frac{1}{2}\min(\beta, 1)\min(\alpha_{0,k}, \alpha_{1,k})$; it exists because that argument differs from the current of $\mathbf{U}$ by $D_{\mathbf{U}}^{\xi *} D_{\mathbf{U}}^{\xi} \mathbf{A} + \mathbf{F}(\mathbf{U}, \mathbf{A})$ with a local operator $\mathbf{F}$ of the second order in $\mathbf{A}$ and $\nabla_{\mathbf{U}}^{\xi}\mathbf{A}$ [8, (3.11), p. 272]. We assume that

$$c_{\mathbf{J}} h_k \le \tfrac{1}{4}\min(\beta, 1)\min(\alpha_{0,k}, \alpha_{1,k}) \ , \tag{6.6}$$

which holds for γ small, since $\vartheta_k/\alpha_{i,k} = (A_1/C_i^{\alpha})(\log \hat{g}_k^{-2})^{p_0-q_i}$ and $q_0 = q_1 > p_0$ (Appendix A). Then, for every s with $|s(\Delta)| \le e^{\kappa_1}$, the factors of the words of the step k are defined and analytic at $e^{i\eta \mathbf{H}(s,B')}U_{k+1}$, with half of the margin between the spaces of the two steps left.

The expansion of a word. By the covariance of its factors (Definition 5.2(iii)) the word of a family $\mathfrak{B}$ is invariant under gauge transformations of the background, so that at the block field $e^{iB'}V^{(k)}$ it is the same word evaluated at the configuration $e^{i\eta \mathbf{H}_k(B')}U_{k+1}$. For a family $\mathfrak{B}$, the empty family included, let $\Phi_{\mathfrak{B}}(s)$ be its word evaluated at $e^{i\eta \mathbf{H}(s,B')}U_{k+1}$. For $k \ge 2$ the factors are functions of a configuration and of the variable $\mathbf{J}$ of [9, (2.36), p. 261], and they are evaluated at the argument which [8, (3.10)–(3.13), p. 272] substitutes for $\mathbf{J}$ when the configuration is $e^{i\xi \mathbf{A}}U_{k+1}$: by [8, (3.11), (3.12)] the current of that configuration is $(L^{j-1}\eta)^3 J_{k+1} + D^{\xi *}_{U_{k+1}} D^{\xi}_{U_{k+1}} \mathbf{A} + \mathbf{F}(U_{k+1}, \mathbf{A})$, in the normalisation of the step j at which the factor was made, and we take this expression at $\mathbf{A} = \mathbf{H}(s, B')$. By (a), $\Phi_{\mathfrak{B}}(1)$ is the word at the configuration of integration and $\Phi_{\mathfrak{B}}(0)$ is the word at U_{k+1}. The torus has finitely many cubes, and the fundamental theorem of calculus in each parameter gives, as in [10, (1.9), p. 4],

$$\Phi_{\mathfrak{B}}(1) = \sum_{\sigma \subset \sigma_k} \prod_{\Delta \in \sigma} \int_0^1 \partial_{s(\Delta)} \Phi_{\mathfrak{B}}(s)\big|_{s(\sigma^c)=0} \, \mathrm{d}s(\Delta) \, . \tag{6.7}$$

The term $\sigma = \emptyset$ is $\Phi_{\mathfrak{B}}(0)$. In the term of σ, by (b), the configuration on a fine bond is $e^{i\eta \mathbf{H}_Y}U_{k+1}$ if the bond lies in a component Y of $Y(\sigma)$, and U_{k+1} otherwise. A factor of the word therefore has its value at U_{k+1} unless the interior of a component meets the set on which it depends: for a transport, the bonds of its arc, and for the insertion of a member W of $\mathfrak{B}$, the dependence set X^+_W. We say that a component *touches* a member W if its interior meets X^+_W, and we call a component *active* for $\mathfrak{B}$ if it contains a fine bond of a loop or touches a member of $\mathfrak{B}$. If a component of $Y(\sigma)$ is not active, the word does not depend on its parameters and the term vanishes, so that only the sets σ all of whose components are active remain.

Arcs and groups. Let Y be an active component and C_i a loop. Let $S'_i(Y)$ be the set of the fine bonds of C_i in Y together with the intervals on C_i of the members of $\mathfrak{B}$ which Y touches. An interval J of a member W of $\mathfrak{B}$ on C_i which contains a bond b of $S'_i(Y)$ lies in $S'_i(Y)$. Indeed, if b lies in an interval of a member touched by Y, that interval is J, since the intervals of a family are pairwise disjoint. Otherwise b is a bond of C_i in Y; the cube of the partition π_j of the step at which W was made which contains an end site of b belongs to X_W, and this site is an interior point of Y, the sites being interior points of the cubes of the partitions (Section 7.1), so that Y touches W and again $J \subset S'_i(Y)$. We put no arcs between the parts of $S'_i(Y)$: to join them is left to the regrouping of Section 6.5, where the arcs which are added are bounded by the sizes of the objects which they join. Two active components are *linked* if they touch a common member of $\mathfrak{B}$, and the *groups* G of the term are the classes of the equivalence relation which this generates. The *arc set* of a group is $I(G) \coloneqq \bigcup_i \bigcup_{Y \in G} S'_i(Y)$, a union of arcs of the loops. The arc sets of distinct groups are disjoint. Indeed, a fine bond of a loop lies in at most one component, two components having no cube in common; a bond of an interval of a member lies in the sets $S'_i(Y)$ of the components which touch that member, and these are linked; and if a bond of a component Y lies in an interval of a member, then Y touches that member, by what was shown. A group whose arc set meets two loops contains a component with bonds or touched members on both, because the intervals of a member which two linked components touch belong to the sets S'_i of both.

A member touched by a component of G has all its intervals in $I(G)$, and it is touched by no component of another group, by the definition of the groups. A member of $\mathfrak{B}$ which no

component touches is *untouched*; its insertion has its value at U_{k+1}, by (b), and its intervals are disjoint from the arc sets of all the groups, by what was shown. Let $\mathfrak{B}(G)$ be the set of the members touched by the components of G, and call the maximal arcs of $I(G)$ the *pieces* of G. We define the tensor Y_G by applying the operations $\prod_{Y\in G}\prod_{\Delta\subset Y}\int_0^1 \mathrm{d}s(\Delta)\,\partial_{s(\Delta)}$ to the ordered products, along the pieces of G, of the insertions of the members of $\mathfrak{B}(G)$ and of the transports of $e^{i\eta\mathbf{H}(s,B')}U_{k+1}$ along the bonds of the pieces between them, at $s(\Delta)=0$ for every cube outside the components of G; for insertions which are not elementary tensors the ordered products are defined by multilinearity, as in Section 5.3. So Y_G has one factor for each of its pieces, a linear map between the fibres at the two ends of the piece. By what was said, the parameters of the components of another group do not enter Y_G.

Lemma 6.4 (The Expansion of a Word)**.** *Let $\mathfrak{B}$ be a family of the step k on the term without large-field regions, and suppose* (6.6)*. For real fluctuation fields in the support of the test functions of the step, the word of $\mathfrak{B}$ at the configuration of integration equals*

$$\sum_{\mathfrak{Y}} w(\mathfrak{B},\mathfrak{Y})\;, \tag{6.8}$$

where the sum runs over the finite sets $\mathfrak{Y}$ of connected unions of cubes of σ_k which are active for $\mathfrak{B}$ and no two of which have a cube or a wall in common, the empty set included, and $w(\mathfrak{B},\mathfrak{Y})$ is the word formed, in the cyclic order of each loop, from the tensors Y_G of the groups G of $\mathfrak{Y}$, the insertions at U_{k+1} of the untouched members of $\mathfrak{B}$, and the transports of U_{k+1} between them. The term $\mathfrak{Y}=\emptyset$ is the word of $\mathfrak{B}$ at U_{k+1}; for $\mathfrak{B}=\emptyset$ it is the product of the leading words of the step $k+1$. Each Y_G depends on the fluctuation field and on the parameters only on the components of G, and on U_{k+1} only along the pieces of G, on its components, and on the dependence sets of the members of $\mathfrak{B}(G)$.

Proof. By (6.7) and the remarks which follow it, the word is the sum over the sets σ all of whose components are active. The map which sends σ to the set $\mathfrak{Y}$ of the components of $Y(\sigma)$ is a bijection onto the sets of the statement: the components of $Y(\sigma)$ are connected, active, and no two of them share a cube or a wall, and conversely the components of the union of the members of such a set $\mathfrak{Y}$ are these members. In the term of σ the integrand is the word in which the factors on the pieces of the groups are evaluated at the configurations $e^{i\eta\mathbf{H}_Y}U_{k+1}$, and all other factors, the insertions of the untouched members among them, at U_{k+1}. The operations $\int_0^1 \mathrm{d}s(\Delta)\,\partial_{s(\Delta)}$ of the cubes of a component act only on the factors of its group, by (b) and the two remarks before the lemma, and the word is multilinear in its factors, so that the term of σ is $w(\mathfrak{B},\mathfrak{Y})$. The locality of Y_G is (b) with the locality of the factors of the word. *q.e.d.*

The expansion is useful because its terms are small: each component of a group contributes a factor which is proportional to the fluctuation field on it and decays exponentially in the number of its cubes, at the rate κ_1 of the parameters. We prove this here for the groups which touch no member, whose tensors are built from transports alone; the groups with members are estimated in Lemma 6.7 and Proposition 6.9 from the bounds of the class. For complex X in the complexified Lie algebra, $\|\pi(e^{iX})\|\le e^{c'_\pi|X|}$ with $c'_\pi := 2\sqrt{N}c_\pi$: writing $X=X_1+iX_2$ with X_1,X_2 in the Lie algebra, we have $|X_1|,|X_2|\le|X|$ for the norm of the test functions, which is invariant under the adjoint, and $\|\mathrm{d}\pi(X)\|\le c_\pi(\|X_1\|+\|X_2\|)$ by (1.7), with $\|X_j\|\le\sqrt{N}|X_j|$. Let $\bar\Lambda := \max_i \Lambda(C_i)$, write $\#(Y)$ for the number of the cubes of a component and $\#(G) := \sum_{Y\in G}\#(Y)$, and fix a constant $\varepsilon_H<\varepsilon_3$ with

$$16\bar\Lambda c'_\pi M B_0 e^{16\kappa_1}\varepsilon_H \le 1\;, \tag{6.9}$$

where c'_π is taken for the representation with the largest c_π.

Lemma 6.5 (The Size of a Tensor without Members)**.** *In the setting of Lemma 6.4 let* $\kappa_1 \geq 1$*, and let* G *be a group of a set* $\mathfrak{Y}$ *with* $\mathfrak{B}(G) = \emptyset$*. Let the background be* $\mathbf{U} = e^{i\eta a'} U_{k+1}$ *with a field* a' *with values in the complexified Lie algebra for which* $\mathbf{U}$ *lies in the space of the step* $k+1$*, and let* $\|a'\|_\infty$ *be the supremum of* $|a'(b)|$ *over the fine bonds of* $I(G)$*. If* $|B'|_Y \leq \varepsilon_H$ *for every component* Y *of* G*, then the tensor* Y_G *formed with* $\mathbf{U}$ *in the place of* U_{k+1} *satisfies*

$$\|Y_G\| \leq \exp(c'_\pi \|a'\|_\infty |I(G)|) \prod_{Y \in G} \frac{|B'|_Y}{\varepsilon_H} \, e^{-(\kappa_1 - 2)\#(Y)} \,, \tag{6.10}$$

where $|I(G)|$ *is the length of the arc set in the units of the step* k *and the norm is the projective norm* (5.5) *over the pieces of* G*. Moreover* $|I(G)| \leq 4\bar{\Lambda} M \#(G)$*.*

Proof. For a family $t = (t_Y)_{Y \in G}$ of complex numbers let $\Phi(s,t)$ be the tensor product, over the pieces of G, of the ordered products of the transports $\pi_i(e^{i\eta \mathbf{H}_Y(s, t_Y B')(b)} e^{i\eta a'(b)} U_{k+1}(b))$ along the fine bonds b of the piece, where Y is the component which contains b and the first factor is absent if there is none. By property (c) of the parameters, with the uniformity of Bałaban's bounds on the space of the configurations, Φ is analytic in (s,t) for $|s(\Delta)| \leq e^{\kappa_1}$ and $|t_Y| \leq T_Y := \varepsilon_H / |B'|_Y$, since $|t_Y B'|_Y \leq \varepsilon_H < \varepsilon_3$ there; and $T_Y \geq 1$. By its definition,

$$Y_G = \prod_{Y \in G} \prod_{\Delta \subset Y} \int_0^1 \partial_{s(\Delta)} \Phi(s,t) \, \mathrm{d}s(\Delta) \Big|_{t=1} \,,$$

at $s(\Delta) = 0$ for the cubes outside the components of G. We bound the integrand $g(s,t) := \prod_Y \prod_{\Delta \subset Y} \partial_{s(\Delta)} \Phi(s,t)$ at $t = 1$ and real s in three steps.

First, g vanishes whenever one t_Y vanishes: then $\mathbf{H}_Y(s, t_Y B') = 0$ for every s, by (6.4), so that Φ does not depend on the parameters of the cubes of Y and its derivatives in them vanish. Schwarz's lemma, applied in each variable t_Y on the disc of radius T_Y, therefore gives $\|g(s,1)\| \leq \prod_Y T_Y^{-1} \sup_t \|g(s,t)\|$, the supremum over the polydisc. Secondly, each derivative in a parameter $s(\Delta)$ at a point of $[0,1]$ is bounded by Cauchy's estimate on the circle of radius $e^{\kappa_1} - 1$ about that point, which lies in the disc $|s(\Delta)| \leq e^{\kappa_1}$; since $e^{\kappa_1} - 1 \geq e^{\kappa_1 - 1}$ for $\kappa_1 \geq 1$, this gives $\|g(s,t)\| \leq e^{-(\kappa_1 - 1)\#(G)} \sup \|\Phi\|$, the supremum over the polydisc in s and t. Thirdly, Φ is an elementary tensor, and its norm is the product of the norms of its factors, each of which is a product of transports along single bonds. The transport along a bond b has norm at most $\exp(c'_\pi \eta(|\mathbf{H}_Y(s, t_Y B')(b)| + |a'(b)|))$, because $U_{k+1}(b)$ is unitary, and $|\mathbf{H}_Y(s, t_Y B')| \leq 4 B_0 e^{16\kappa_1} \varepsilon_H$ on the polydisc by (6.4). The fine bonds of the loops in a cube of σ_k have total length at most $4\bar{\Lambda} M$ in the units of the step k, by Lemma 5.5(i) at the scale k and because a cube of side $M < a_k$ meets one loop (Lemma 3.14). Hence

$$\sup \|\Phi\| \leq \exp(c'_\pi \cdot 4\bar{\Lambda} M \#(G) \cdot 4 B_0 e^{16\kappa_1} \varepsilon_H) \exp(c'_\pi \|a'\|_\infty |I(G)|) \leq e^{\#(G)} \exp(c'_\pi \|a'\|_\infty |I(G)|) \,,$$

by (6.9). Multiplying the three bounds, and integrating over the parameters in $[0,1]$, we obtain (6.10); the projective norm of an integral of elementary tensors is at most the integral of the products of the norms of their factors.

For the length, a group which touches no member consists of one component, by the definition of the groups, and its arc set is the set of the fine bonds of the loops in that component, of total length at most $4\bar{\Lambda} M \#(G)$ by the count of the bonds in a cube which was used above. *q.e.d.*

For a group of one component the bound (6.10) is of the first order in the fluctuation field, hence of the order of g_k up to logarithms, whereas the insertions of the class are of the

order of g_k^2. The first order is there: it is the linear response of the transport of a loop to the fluctuation field. It is harmless because it is odd in the fluctuation field and the measure of the step is even up to its potentials (Supplement, Lemma 17.28), and what has to be shown here is that the even part of the tensor is of the second order. On the support of the test functions, $|g_k A_k| \le r_y \vartheta_k$, as was used for (6.5), and therefore $|B'| \le b_k := C_1 r_y \vartheta_k$ by [10, (1.20), p. 6]. The middle member of that display, $O(1) g_k |B| + 4C_2 (O(1) g_k |B|)^2$, bounds the two terms of $B' = g_k C A_k - h\tilde{D}(g_k C A_k)$ separately: the function $\tilde{D}$ "has an expansion beginning with quadratic terms" [8, p. 267], and $|h\tilde{D}(g_k C A_k)| \le 4C_2 (c_0 g_k |A_k|)^2$, where c_0 denotes the absolute constant $O(1)$ of that member.

Lemma 6.6 (The Even Part of a Tensor of One Component)**.** *In the setting of Lemma 6.5, let the group G consist of one component Y, let $b_k \le \frac{1}{2}\varepsilon_H$, and put $S_Y := \exp(c'_\pi \|a'\|_\infty |I(G)|)\, e^{-(\kappa_1 - 2)\#(Y)}$. Let $Y_G = Y_G^{\mathrm{ev}} + Y_G^{\mathrm{od}}$ be the decomposition of the tensor, as a function of the fluctuation field A_k, into its even and odd parts under $A_k \mapsto -A_k$. Then, on the support of the test functions of the step,*

$$\|Y_G^{\mathrm{ev}}\| \le c_{\mathrm{ev}} \vartheta_k^2 S_Y \,, \qquad \|Y_G^{\mathrm{od}}\| \le \Big(\frac{C_1 r_y}{\varepsilon_H} \vartheta_k + c_{\mathrm{ev}} \vartheta_k^2 \Big) S_Y \,, \qquad c_{\mathrm{ev}} := \frac{4C_2 c_0^2 r_y^2}{\varepsilon_H} + \frac{2C_1^2 r_y^2}{\varepsilon_H^2} \,. \tag{6.11}$$

Proof. By property (b) of the parameters, Y_G depends on the fluctuation field through the restriction X of B' to Y; we write $Y_G(X)$. The proof of Lemma 6.5, without its first step, shows that Y_G is analytic in X and $\|Y_G(X)\| \le S_Y$ for $|X|_Y \le \varepsilon_H$, and $Y_G(0) = 0$. Let Λ be its differential at 0, a linear map, and $R(X) := Y_G(X) - \Lambda(X)$. Cauchy's estimates for the function $t \mapsto Y_G(tX)$ on the disc of radius $\varepsilon_H / |X|_Y$ give

$$\|\Lambda(X)\| \le S_Y \frac{|X|_Y}{\varepsilon_H} \,, \qquad \|R(X)\| \le S_Y \sum_{n \ge 2} \Big(\frac{|X|_Y}{\varepsilon_H} \Big)^n \le 2 S_Y \Big(\frac{|X|_Y}{\varepsilon_H} \Big)^2$$

for $|X|_Y \le \frac{1}{2}\varepsilon_H$, the first for every X by linearity. Now $B' = g_k C A_k - h\tilde{D}(g_k C A_k)$, where C is linear [8, p. 268], so that the even part of B' under $A_k \mapsto -A_k$ is the even part of $-h\tilde{D}(g_k C A_k)$, and its restriction to Y is at most $4C_2 c_0^2 r_y^2 \vartheta_k^2$ on the support, by the bound recalled before the lemma and $|g_k A_k| \le r_y \vartheta_k$. The function Λ of the odd part of B' is odd. Hence Y_G^{ev} is the sum of Λ of the even part of B' and of the even part of $R(B')$, and with $|B'|_Y \le b_k = C_1 r_y \vartheta_k \le \frac{1}{2}\varepsilon_H$ this gives the first bound of (6.11). The second follows from $\|Y_G\| \le S_Y b_k / \varepsilon_H$, which is (6.10), and the first. *q.e.d.*

A group which touches members of the family is estimated in the same way, with one difference. The insertions of the members are not entire functions of the configuration, as the transports are, so that the fluctuation field on a component may be scaled only as far as the interpolated configurations stay in the domain on which these insertions are analytic. By the argument which gave (6.6), this is the case as long as $c_{\mathbf{J}} \cdot 4B_0 e^{16\kappa_1}$ times the scaled field is at most $\frac{1}{4}\min(\beta, 1)\min(\alpha_{0,k}, \alpha_{1,k})$; we put

$$r_\alpha := \frac{\min(\beta, 1)\min(\alpha_{0,k}, \alpha_{1,k})}{16 c_{\mathbf{J}} B_0 e^{16\kappa_1}} \,, \tag{6.12}$$

so that (6.6) reads $b_k \le r_\alpha$. The factor which a component gains is then the ratio of its fluctuation field to $\min(\varepsilon_H, r_\alpha)$, a negative power of $\log g_k^{-2}$ in place of a power of the coupling, and the smallness of the tensor comes from the insertions of its members.

Lemma 6.7 (The Size of a Tensor with Touched Members)**.** *In the setting of Lemma 6.5, with a field a' for which $e^{i\eta a'} U_{k+1}$ lies in the space of the step $k+1$, let G be a group of a*

set $\mathfrak{Y}$ with the set $\mathfrak{B}(G)$ of touched members, and suppose (6.6) *and $b_k \le \varepsilon_H$. Then, on the support of the test functions of the step,*

$$\|Y_G\| \le \exp(4\bar\Lambda M c'_\pi \|a'\|_\infty \#(G)) \prod_{Y\in G} \frac{|B'|_Y}{\min(\varepsilon_H, r_\alpha)}\, e^{-(\kappa_1-2)\#(Y)} \prod_{W\in\mathfrak{B}(G)} \sup\|Y_W\|\,, \qquad (6.13)$$

where $\|a'\|_\infty$ is the supremum of $|a'(b)|$ over the fine bonds of the loops in the components of G and each supremum is taken over the configurations $e^{i\eta\mathbf{H}(s,tB')}e^{i\eta a'}U_{k+1}$, with the corresponding argument in the place of $\mathbf{J}$, for $|s(\Delta)| \le e^{\kappa_1}$ and $|t_Y||B'|_Y \le \min(\varepsilon_H, r_\alpha)$, which lie in the domain of analyticity of the factors of the step k.

Proof. We repeat the proof of Lemma 6.5 with the function $\Phi(s,t)$ which is the tensor product, over the pieces of G, of the ordered products of the transports and of the insertions Y_W, $W \in \mathfrak{B}(G)$, at the configuration $e^{i\eta\mathbf{H}(s,tB')}e^{i\eta a'}U_{k+1}$, where tB' denotes the field $t_Y B'$ on each component Y of G; the products are formed by multilinearity. For $|t_Y||B'|_Y \le \min(\varepsilon_H, r_\alpha)$ the field $\mathbf{H}(s,tB')$ is at most $4B_0 e^{16\kappa_1} r_\alpha$ in the admissible norms, by (6.4), and by the definitions of $c_{\mathbf{J}}$ and r_α the configuration lies in the space of the step k, on which the insertions are analytic (Definition 5.2(iii)). So Φ is analytic on the polydisc $|s(\Delta)| \le e^{\kappa_1}$, $|t_Y| \le T_Y \coloneqq \min(\varepsilon_H, r_\alpha)/|B'|_Y$, and $T_Y \ge 1$ because $|B'|_Y \le b_k \le \min(\varepsilon_H, r_\alpha)$. The three steps are as before. If one t_Y vanishes, $\mathbf{H}_Y = 0$, no factor depends on the parameters of the cubes of Y, and the derivatives vanish, so that Schwarz's lemma gives $\prod_Y T_Y^{-1}$. Cauchy's estimate in the parameters gives $e^{-(\kappa_1-1)\#(G)}$. And the norm of Φ is at most the product of the norms of the transports, which run along the fine bonds of the loops in the components of G outside the intervals of the members and are bounded as before by $e^{\#(G)}\exp(4\bar\Lambda M c'_\pi\|a'\|_\infty\#(G))$, and of the projective norms of the insertions of the members, since the contraction of tensors along a piece does not increase the product of the norms (Section 5.3). *q.e.d.*

6.4 The Fluctuation Integral with Marked Polymers

By Lemma 6.4 the function F is, on the term without large-field regions, a finite sum of numbers times words, and in each word only the tensors Y_G depend on the field A which the step integrates, each of them through the fluctuation field on the components of its group. The composition with Ψ_{k+1} in (6.2) does not interfere, because the bonds of $S_{k+1}^{(k)*}$ are separated from Λ_{k+1} by a layer of cubes [9, (3.20), p. 269], so that Ψ_{k+1} changes outer variables of the step only. The expectation of a word is therefore a contraction of the expectation of the tensor product of its tensors Y_G, and what we need is an expansion of such an expectation into local pieces. Bałaban makes this expansion for one localised factor under the integral: the factor marks a domain Z_0 of the polymer expansion, whose activity $H'(Z_0)$ contains it, while the other polymers have the activities of the expansion of the normalisation, and the sum over these is exponentiated [9, (3.44)–(3.46), pp. 277–278]. We need the same with several marked domains and with factors which are tensors.

The setting. We fix a family $\mathfrak{B}$ and a set $\mathfrak{Y}$ as in Lemma 6.4, with the set $\mathfrak{G}$ of its groups, and put

$$\mathcal{X} \coloneqq \bigotimes_{G\in\mathfrak{G}} Y_G\,, \qquad (6.14)$$

a function of A and of the other variables with values in the tensor product of the spaces of the tensors Y_G. The word $w(\mathfrak{B},\mathfrak{Y})$ is multilinear in its factors, so that $\langle w(\mathfrak{B},\mathfrak{Y})\rangle$ is the contraction of $\langle\mathcal{X}\rangle$ with the factors which do not depend on A, the insertions of the untouched members and the transports of U_{k+1}; and $\langle\mathcal{X}\rangle = I(\mathcal{X})/I(1)$, where $I(f)$ denotes the last integral of [9, (3.25)] with the further factor f under it. For $G \in \mathfrak{G}$ let D_G be the

union of the components of G and of the cubes of π_k which share a point with them, and let $D^\# := \bigcup_{G\in\mathfrak{G}} D_G$. The bonds of the scale k on whose variables the tensor Y_G depends lie in the interior of D_G. Indeed Y_G depends on A through the restriction of B' to its components (Lemma 6.4), and the variable $B'(b)$ is $g_k A_k(b)$ unless b is a bond $b_0(c)$, for which it is a function of the variables in the blocks of the ends of c and between them, since the δ-function which eliminates it is that of the average [8, (0.4), p. 253] on the bond c [8, pp. 267–268]; these blocks have the side $L \le M$ and are contained in cubes of π_k at a distance less than M from the component, and two cubes of π_k at a distance less than M share a point.

The polymers. The expansion of [10, Sect. 2] writes the fluctuation integral as a sum over finite sets of *polymers*, localisation domains $Z \in \mathbf{D}_{k+1}$, with activities $H(Z)$ and the hard core ζ: $\zeta(Z,Z') = 0$ if $Z \cap Z'$ contains a cube or a wall of a cube, and $\zeta(Z,Z') = 1$ otherwise [10, (2.8)–(2.11), p. 14]. We call two polymers *compatible* if $\zeta = 1$, and write $Z \not\sim Z'$ if they are not. For the measure [9, (3.27)] of the step the expansion is that of [9, pp. 275–279], where "we repeat all the considerations of Sect. 3.7 [I]" with the cover and the localised action of [9, pp. 275–277]; the section of [I] which is meant is [10, Sect. 2], whose formulae (2.1)–(2.13) are cited there as (I.7.1)–(I.7.13) (Section 14). We call the polymers of the expansion of the normalisation $I(1)$ *vacuum polymers.* A *cluster* is a finite sequence $\mathcal{C} = (Z_1, \dots, Z_n)$ of vacuum polymers whose incompatibility graph is connected, with the weight

$$\varphi^T(\mathcal{C}) := \frac{1}{n!}\rho^T(Z_1,\dots,Z_n)H(Z_1)\cdots H(Z_n)\,, \tag{6.15}$$

where ρ^T is the function of [10, (2.12), p. 14], which vanishes unless that graph is connected; a cluster is incompatible with a union X of cubes, $\mathcal{C} \not\sim X$, if one of its polymers is. Bałaban exponentiates the expansion of the normalisation, "the factor is represented as $\exp \tilde{\mathbf{E}}^{(k+1)}(\Lambda_{k+1})$", and the sum over the polymers compatible with a marked domain as well [9, pp. 277–278], and "the other expectation values in (3.26), (3.28) are treated in exactly the same way" [9, p. 279], the first of which is the expectation in the measure (3.27). We use this in the following form: the series $\sum_{\mathcal{C}\not\sim X}|\varphi^T(\mathcal{C})|$ converges for every union X of cubes of the torus, and the sum of the products of the activities over the finite sets of pairwise compatible vacuum polymers which are compatible with X equals $\exp\sum_{\mathcal{C}}\varphi^T(\mathcal{C})$, the sum over the clusters none of whose polymers is incompatible with X. Both statements, with the bound $\sum_{\mathcal{C}\not\sim X}|\varphi^T(\mathcal{C})|e^{d(\mathcal{C})} \le cN(X)$, where $N(X)$ is the number of the cubes of π_{k+1} in X, $c = \frac{9}{64}e^{-4\kappa}$, and $d(\mathcal{C})$ is $9\log 2$ times the total number of the cubes of the polymers of $\mathcal{C}$, are derived from Bałaban's bound [10, (2.38), p. 20] of the activities in Supplement, Lemma 17.24, for $\kappa \ge 2^9(1+16\log 2)$.

Proposition 6.8 (Marked Polymers)**.** *In the setting above, call two connected components of a union Z of cubes of π_{k+1}* linked *if a set D_G, $G \in \mathfrak{G}$, meets both, and call Z linked if this relation connects all its components. Then*

$$\langle \mathcal{X} \rangle = \sum_{\mathfrak{M}} Q_{k+1}(\mathfrak{M}) \bigotimes_{M\in\mathfrak{M}} \hat{H}(M)\,, \tag{6.16}$$

where:

(i) the sum runs over the finite sets $\mathfrak{M}$ of pairwise compatible marked polymers*: linked unions M of cubes of π_{k+1}, each with a nonempty set $\mathfrak{G}(M) \subset \mathfrak{G}$ of groups whose sets D_G it contains, such that the sets $\mathfrak{G}(M)$, $M \in \mathfrak{M}$, form a partition of $\mathfrak{G}$;*

(ii) $\hat{H}(M) := e^{-\Psi(M)}\tilde{H}(M)$, where the marked activity *$\tilde{H}(M)$ is the activity of M in the expansion [10, (2.1)–(2.11)] of the integral with the factor $\bigotimes_{G\in\mathfrak{G}(M)} Y_G$ under it, a tensor with one factor for each piece of each group of $\mathfrak{G}(M)$, and $\Psi(M) := \sum_{\mathcal{C}\not\sim M}\varphi^T(\mathcal{C})$;*

(iii) with $\Psi_{\geq}(T) := \sum \varphi^T(\mathcal{C})$*, the sum over the clusters which are incompatible with every member of a set* T *of marked polymers,*

$$Q_{k+1}(\mathfrak{M}) := \exp\Big(\sum_{T \subset \mathfrak{M},\ |T| \geq 2} (-1)^{|T|} \Psi_{\geq}(T) \Big) \,. \tag{6.17}$$

The tensor $\hat{H}(M)$ *depends on* M *and on the groups of* $\mathfrak{G}(M)$ *alone, and* $\Psi_{\geq}(T)$ *on* T *alone.*

Proof. We go through the construction [10, (2.1)–(2.11), pp. 11–14] for the integral $I(\mathcal{X})$ and note at each step what the factor $\mathcal{X}$ changes.

The Mayer expansion [10, (2.1)] of the exponential of the localised action is an identity for the exponential alone, with $\mathcal{X}$ as a further factor of the integrand. In place of [10, (2.2)] we put $Y_0 := \bigcup_{Y \in \mathbf{D}} Y \cup D^{\#}$.

The test functions. The test function $\chi^{(k)}$ is a product of functions of single bond variables [9, (3.16), p. 268], and the decomposition [10, (2.3)] of the factors of the bonds outside Y_0 applies to it, with the thresholds of the construction; Z_0 is the smallest localisation domain which contains Y_0 and the set P of that decomposition, as in [10, p. 12].

The conditioning [10, (2.4)–(2.6)] on the variables outside Z_0 uses of the integrand only that it is localised in Z_0 with respect to the integration variable. Since $D^{\#} \subset Z_0$, the factor $\mathcal{X}$ is a function of the variable which is conditioned, so that [10, (2.5), (2.6)] hold with $\mathcal{X}$ inside the function G of [10, (2.6)].

The localisation [10, (2.8), (2.9)] introduces parameters in the operators $C^{(k)}(Z_0)$, $(C^{(k)})^{1/2}$, and Δ_k, none of which occurs in $\mathcal{X}$; it gives $I(\mathcal{X}) = \sum_Z H(Z)$ with activities which now contain $\mathcal{X}$.

The factorisation. Bałaban's activities factorise over the connected components of Z [10, (2.10)], because every factor of the integrand, the Mayer factors, the factors of the test functions, and the conditioned Gaussian integrals at vanishing parameters outside Z, splits over them. The factor $\mathcal{X}$ splits over the groups, and the tensor Y_G of a group depends on the integration variable in D_G only, which may meet several components. Hence the activity factorises over the classes of linked components: a class which contains no set D_G has the activity of a vacuum polymer, and a class M which contains the sets D_G of the groups $G \in \mathfrak{G}(M) \neq \emptyset$ has an activity $\tilde{H}(M)$ with values in the tensor product of the spaces of these Y_G. A set D_G which meets a class is contained in it, because $D_G \subset D^{\#} \subset Z_0 \subset Z$ and the components of Z which D_G meets are linked. As in [10, (2.11)] we obtain

$$I(\mathcal{X}) = \sum_{\mathfrak{M}} \Big(\bigotimes_{M \in \mathfrak{M}} \tilde{H}(M) \Big) \, \Xi(\mathfrak{M}) \,, \tag{6.18}$$

the sum over the sets $\mathfrak{M}$ of (i), where $\Xi(\mathfrak{M})$ is the sum of the products of the activities over the finite sets of pairwise compatible vacuum polymers which are compatible with every $M \in \mathfrak{M}$; and $I(1) = \Xi(\emptyset)$.

The ratio. By the exponentiation in the form stated before the proposition, applied to the union of the members of $\mathfrak{M}$ and to the empty set,

$$\frac{\Xi(\mathfrak{M})}{\Xi(\emptyset)} = \exp\Big(- \sum_{\mathcal{C} \not\sim \mathfrak{M}} \varphi^T(\mathcal{C}) \Big) \,, \tag{6.19}$$

the sum over the clusters which are incompatible with at least one member of $\mathfrak{M}$. For such a cluster let $s(\mathcal{C}) \geq 1$ be the number of the members of $\mathfrak{M}$ with which it is incompatible. It

occurs $s(\mathcal{C})$ times in $\sum_{M\in\mathfrak{M}}\Psi(M)$ and once in the exponent of (6.19), and

$$\sum_{T\subset\{M\,|\,\mathcal{C}\not\sim M\},\ |T|\geq 2}(-1)^{|T|}=(1-1)^{s(\mathcal{C})}-1+s(\mathcal{C})=s(\mathcal{C})-1\ ,$$

so that it occurs with the coefficient $s(\mathcal{C})-1$ in the exponent of (6.17). The rearrangement is permitted, because a cluster lies in at most $2^{s(\mathcal{C})}$ of the sums $\Psi_{\geq}(T)$, the torus has finitely many cubes, so that $s(\mathcal{C})\leq|\mathfrak{M}|$, and $\sum_{\mathcal{C}\not\sim\mathfrak{M}}|\varphi^T(\mathcal{C})|$ converges. Hence

$$\frac{\Xi(\mathfrak{M})}{\Xi(\emptyset)}=\prod_{M\in\mathfrak{M}}e^{-\Psi(M)}\cdot Q_{k+1}(\mathfrak{M})\ , \tag{6.20}$$

and dividing (6.18) by $I(1)=\Xi(\emptyset)$ gives (6.16). The last statement holds because the activity of a class of linked components is defined by the integral over the variables of that class. *q.e.d.*

A marked polymer may carry several groups, and a group several pieces, on one loop or on several, so that the tensors $\hat{H}(M)$ are the first objects of the step with several intervals. The marked polymers of a set $\mathfrak{M}$ have a hard core among themselves and interact softly through Q_{k+1}, which is a product of factors, one for each set T of at least two of them; they have no hard core with the domains of the untouched members of $\mathfrak{B}$, which took no part in the expansion.

The size of a marked activity. How small is $\tilde{H}(M)$? Bałaban's bound for the activities of the vacuum polymers cannot hold for it, since the factor $\mathcal{X}_M\coloneqq\bigotimes_{G\in\mathfrak{G}(M)}Y_G$ has a prescribed support in which nothing decays, and since one of the terms of the activity is the conditional expectation of this factor, which carries no small constant. What holds instead is a bound by the supremum of the factor, with a decay in the part of M outside its support and a price for each cube of the support (Supplement, Lemma 17.27). The supremum is small by the lemmata of Section 6.3, with one exception: a factor which consists of the tensor of a single component without members is of the first order in the fluctuation field. Its odd part enters the activity only through the odd part of the potentials of the fluctuation integral (Supplement, Lemma 17.28), and this gives the second order. For a marked polymer M let $\mathfrak{B}(M)\coloneqq\bigcup_{G\in\mathfrak{G}(M)}\mathfrak{B}(G)$ be the set of its touched members, $n(M)$ the number of the components of its groups, $\#(G_M)\coloneqq\sum_{G\in\mathfrak{G}(M)}\#(G)$ the number of their cubes, which are cubes of π_k, and $\#(M)$ the number of the cubes of π_{k+1} in M. With the constant A of Supplement, Lemma 17.27, and the number y_k of Supplement, Lemma 17.28, put

$$c_\kappa\coloneqq 1+81(5+\tfrac{3}{2}L)\kappa+\tfrac{6561}{16}(L+2)\kappa+1594323\,A\ ,\qquad a_k^{(1)}\coloneqq c_{\mathrm{ev}}\vartheta_k^2+\frac{L^4y_k}{2A}\Big(\frac{C_1r_y}{\varepsilon_H}\vartheta_k+c_{\mathrm{ev}}\vartheta_k^2\Big)\ . \tag{6.21}$$

Proposition 6.9 (The Size of a Marked Activity)**.** *On the terms without large-field regions of ϱ_k and of ϱ_{k+1}, suppose that the hypotheses of Supplement, Lemmata 17.27 and 17.28, hold, together with* (6.6)*, $b_k\leq\frac{1}{2}\min(\varepsilon_H,r_\alpha)$, and $4\bar{\Lambda}Mc'_\pi\alpha_{1,k+1}\leq L$. Then, for every marked polymer M and at every configuration of the space of the step $k+1$,*

$$\|\tilde{H}(M)\|\leq\mathfrak{a}(M)\,e^{-(\kappa_1-2-c_\kappa)\#(G_M)}\,e^{-(L+2)\kappa\#(M)/1296}\prod_{W\in\mathfrak{B}(M)}\|Y_W\|_{(k)}\ , \tag{6.22}$$

where $\mathfrak{a}(M)\coloneqq a_k^{(1)}$ if $\mathfrak{B}(M)=\emptyset$ and $n(M)=1$, $\mathfrak{a}(M)\coloneqq(b_k/\varepsilon_H)^{n(M)}$ if $\mathfrak{B}(M)=\emptyset$ and $n(M)\geq 2$, and $\mathfrak{a}(M)\coloneqq(b_k/\min(\varepsilon_H,r_\alpha))^{n(M)}$ if $\mathfrak{B}(M)\neq\emptyset$. Moreover $a_k^{(1)}\leq c_a\rho_k$ and $(b_k/\varepsilon_H)^2\leq c_a\rho_k$, with a constant c_a which depends on the constants of the construction and of (6.11) *and not on k.*

Proof. Let $D := \bigcup_{G\in\mathfrak{G}(M)} D_G$, with m connected components, and let D' be the smallest union of cubes of π_{k+1} which contains D and the cubes of π_k that touch it, as in Supplement, Section 17. We first count. The set D_G consists of cubes of the components of G and of cubes which share a point with them, so that D has at most $81\,\#(G_M)$ cubes of π_k, and at most as many connected components; the cubes of D and those which touch it are at most $81^2\,\#(G_M)$, and each cube of D' contains one of them. Hence

$$m \le 81\,\#(G_M)\,, \qquad \#(D') \le 6561\,\#(G_M)\,. \tag{6.23}$$

By Supplement, Lemma 17.27, applied to M with the factor $\mathcal{X}_M$, and by the remark which follows that lemma,

$$\|\tilde H(M)\| \le \|\mathcal{X}_M\|_{\sup}\, e^{81A\#(D')}\, e^{(5+\frac32 L)\kappa(m-1)}\, e^{-(L+2)\kappa(\#(M)-81\#(D'))/1296}\,,$$

where $\|\mathcal{X}_M\|_{\sup}$ is the supremum of the projective norm of $\mathcal{X}_M$ over the fluctuation fields of the support of the test functions and over the space of the step $k+1$. By (6.23) the three exponential factors other than $e^{-(L+2)\kappa\#(M)/1296}$ are together at most $\exp((c_\kappa-1)\#(G_M))$, where we have bounded $81A$ by $243A$ and used $81\cdot 6561/1296 = 6561/16$ and $243\cdot 6561 = 1594323$.

The norm of a tensor product is at most the product of the norms, so that $\|\mathcal{X}_M\|_{\sup} \le \prod_{G\in\mathfrak{G}(M)} \sup\|Y_G\|$. On the support of the test functions $|B'|_Y \le b_k$ for every component Y (Section 6.3), and a configuration $e^{i\eta a'}U_{k+1}$ of the space of the step $k+1$ has $|a'(b)| \le \alpha_{1,k+1}/L$ on the fine bonds, in the units of the step k, by the radii of Definition 5.2(iii), so that $\exp(4\bar\Lambda M c'_\pi \|a'\|_\infty \#(G)) \le e^{\#(G)}$ by the last hypothesis. For a group G without members, which is a single component, Lemma 6.5 therefore gives $\|Y_G\| \le (b_k/\varepsilon_H)e^{-(\kappa_1-3)\#(G)}$, and for a group with members Lemma 6.7 gives

$$\|Y_G\| \le \prod_{Y\in G} \frac{b_k}{\min(\varepsilon_H, r_\alpha)} \cdot e^{-(\kappa_1-3)\#(G)} \prod_{W\in\mathfrak{B}(G)} \|Y_W\|_{(k)}\,,$$

since the configurations over which that lemma takes the suprema lie in the space of the step k. Multiplying over the groups, and using $b_k/\varepsilon_H \le b_k/\min(\varepsilon_H, r_\alpha) \le 1$, we obtain (6.22) in the second and the third case.

In the first case $\mathcal{X}_M = Y_G$ for a group which consists of one component Y without members. By Supplement, Lemma 17.28, the bound of Lemma 17.27 holds with $\|\mathcal{X}^{\rm ev}\|_{\sup} + (L^4 y_k/(2A))\|\mathcal{X}^{\rm od}\|_{\sup}$ in the place of $\|\mathcal{X}_M\|_{\sup}$ and $243A$ in the place of $81A$, where $\mathcal{X}^{\rm ev}$ and $\mathcal{X}^{\rm od}$ are the even and the odd part of the factor under the reflection of the fluctuation field. By Lemma 6.6, whose hypothesis $b_k \le \frac12\varepsilon_H$ holds, these parts are at most $c_{\rm ev}\vartheta_k^2 S_Y$ and $(C_1 r_y \vartheta_k/\varepsilon_H + c_{\rm ev}\vartheta_k^2)S_Y$ with $S_Y \le e^{-(\kappa_1-3)\#(Y)}$, which gives (6.22) with $\mathfrak{a}(M) = a_k^{(1)}$.

For the last assertion, $y_k = 4\alpha_4(\vartheta_k/\varepsilon_1)(8M^4 t_k^2 + K_{\mathbf{D}})$ with $t_k = \vartheta_k/g_k$ (Supplement, Section 17), and $\vartheta_k \le 1$, so that $a_k^{(1)} \le c\,\vartheta_k^2(1+t_k^2)$ with $c := c_{\rm ev} + (2L^4\alpha_4/(A\varepsilon_1))\max(8M^4, K_{\mathbf{D}})(C_1 r_y/\varepsilon_H + c_{\rm ev})$. By (3.17), $\vartheta_k \le g_k r_k$, hence $t_k \le r_k$ and $\vartheta_k^2(1+t_k^2) \le g_k^2(1+r_k)^4 = \rho_k/C_\rho$ by (5.6). Likewise $(b_k/\varepsilon_H)^2 = (C_1 r_y/\varepsilon_H)^2\vartheta_k^2 \le (C_1 r_y/\varepsilon_H)^2\rho_k/C_\rho$, and we take $c_a := \max(c, (C_1 r_y/\varepsilon_H)^2)/C_\rho$. *q.e.d.*

The bound (6.22) shows where the smallness of a new group comes from. A marked polymer without members is of the order of ρ_k, the square of the coupling up to logarithms. One with members is small through the insertions of the members, which are of that order themselves, and gains for each of its components a factor $b_k/\min(\varepsilon_H, r_\alpha)$, which tends to zero

with the coupling as a power of its logarithm, because $q_0 > p_0$. The exponent c_κ is the price of the support, and the rate κ_1 of the localisation parameters, which is chosen after κ and L and before M [10, p. 5], is taken so large that it pays for it and for the counting of the next section.

6.5 The Groups of the Step

In this subsection the labelled term of ϱ_{k+1} is the term without large-field regions. By Lemma 6.4 and Proposition 6.8, applied to every word of F,

$$\langle F \rangle = \prod_{i=1}^{n} \kappa^{(k)}(C_i, \chi_i) \cdot e^{\mathcal{E}_k} \sum_{\mathfrak{B}} Q_k(\mathfrak{B}) \sum_{\mathfrak{Y}} \sum_{\mathfrak{M}} Q_{k+1}(\mathfrak{M})\, w(\mathfrak{B}, \mathfrak{Y}, \mathfrak{M}) \,, \tag{6.24}$$

where $Q_k(\emptyset) \coloneqq 1$, the sums run over the families of the step k, the sets $\mathfrak{Y}$ of Lemma 6.4, and the sets $\mathfrak{M}$ of Proposition 6.8, and $w(\mathfrak{B}, \mathfrak{Y}, \mathfrak{M})$ is the word in which the tensors $\hat{H}(M)$ are contracted, in the cyclic order of each loop, with the insertions at U_{k+1} of the untouched members of $\mathfrak{B}$ and the transports of U_{k+1}. This is not yet the form of a dressed loop, for three reasons. The tensors $\hat{H}(M)$ have several pieces, between which other factors of the word may lie. The terms are tied together by two soft interactions, Q_k and Q_{k+1}, the second of which depends on the fields. And the marked polymers have no hard core with the untouched members, whose domains they may meet. All three defects are removed by a regrouping of the terms of (6.24) which changes none of them: the soft interactions are expanded, and whatever an interaction, an overlap, or the order along a loop ties together is collected into one group of the step $k+1$. We call two unions of cubes of the partitions of the construction *compatible* if their intersection has dimension at most $d-2$, so that it contains neither a cube nor a wall of a cube; for two polymers this is the condition $\zeta = 1$.

The flat parts of the vacuum weights and the links. The activities of the vacuum polymers are analytic functions of a configuration $(\mathbf{U}, \mathbf{J})$ on the spaces of the step $k+1$, with bounds which are uniform on these spaces [10, pp. 6, 15], and the reference data $(1, 0)$ belong to them. Let $\varphi^T_\circ(\mathcal{C})$ be the weight (6.15) of a cluster at the reference data and $y_\mathcal{C} \coloneqq \varphi^T(\mathcal{C}) - \varphi^T_\circ(\mathcal{C})$, a function of the background field on the union $\operatorname{supp}\mathcal{C}$ of the polymers of $\mathcal{C}$, since each activity "is localized in the interior of Z with respect to the external gauge fields" [10, p. 14]; the statements on the cluster series used in Proposition 6.8 hold at the reference data as well. We write $\Psi_\circ$, $\Psi_{\geq,\circ}$, and $Q_\circ$ for the quantities of that proposition with $\varphi^T_\circ$ in the place of φ^T. For a set $\mathfrak{M}$ of marked polymers, a *link* is a pair $\lambda = (T, \mathcal{C})$ of a nonempty subset $T \subset \mathfrak{M}$ and a cluster $\mathcal{C}$ which is incompatible with every member of T; its *domain* is $\operatorname{supp}\mathcal{C}$, its *factor* is

$$u_\lambda \coloneqq \exp((-1)^{|T|} y_\mathcal{C}) - 1 \,, \tag{6.25}$$

which vanishes at the reference data, and it is *carried* by the first member of T in a fixed order of the unions of cubes. Since $\prod_M e^{-\Psi(M)} \cdot Q_{k+1}(\mathfrak{M})$ is the exponential of $\sum_{T \neq \emptyset} (-1)^{|T|} \Psi_\geq(T)$, where $\Psi_\geq(\{M\}) = \Psi(M)$, and $\Psi_\geq(T) - \Psi_{\geq,\circ}(T)$ is the sum of $y_\mathcal{C}$ over the clusters incompatible with every member of T,

$$\prod_{M \in \mathfrak{M}} e^{-\Psi(M)} \cdot Q_{k+1}(\mathfrak{M}) = \prod_{M \in \mathfrak{M}} e^{-\Psi_\circ(M)} \cdot Q_\circ(\mathfrak{M}) \prod_\lambda (1 + u_\lambda) = \prod_{M \in \mathfrak{M}} e^{-\Psi_\circ(M)} \cdot Q_\circ(\mathfrak{M}) \sum_{\mathcal{L}} \prod_{\lambda \in \mathcal{L}} u_\lambda \,, \tag{6.26}$$

the sum over the finite sets $\mathcal{L}$ of links. The product converges absolutely, because a cluster belongs to at most $2^{|\mathfrak{M}|}$ links, $|u_\lambda| \leq |y_\mathcal{C}| e^{|y_\mathcal{C}|}$, and $\sum_{\mathcal{C} \not\subset \mathfrak{M}} (|\varphi^T(\mathcal{C})| + |\varphi^T_\circ(\mathcal{C})|)$ converges. The flat factor is a number, and as in the proof of Proposition 6.8 it equals $\exp \sum_\mathcal{C} (s(\mathcal{C}) - 1) \varphi^T_\circ(\mathcal{C})$,

the sum over the clusters which are incompatible with $s(\mathcal{C}) \geq 2$ members of $\mathfrak{M}$; hence

$$Q_\circ(\mathfrak{M}) = \sum_{\mathcal{O}} \prod_{\mathcal{C}\in\mathcal{O}} v_{\mathcal{C}} \;, \qquad v_{\mathcal{C}} \coloneqq \exp((s(\mathcal{C})-1)\varphi_\circ^T(\mathcal{C})) - 1 \;, \tag{6.27}$$

the sum over the finite sets $\mathcal{O}$ of such clusters, the *vacuum occurrences*, again absolutely convergent.

The old occurrences. By Definition 5.2, $Q_k(\mathfrak{B})$ is the exponential of $-\sum_\Gamma \varphi_k(\Gamma)$ over the clusters Γ of the reference gas of the step k which are incompatible with a member of $\mathfrak{B}$, and

$$Q_k(\mathfrak{B}) = \sum_{m\geq 0} \frac{(-1)^m}{m!} \sum_{(\Gamma_1,\dots,\Gamma_m)} \prod_{j=1}^{m} \varphi_k(\Gamma_j) \;, \tag{6.28}$$

the inner sum over the sequences of such clusters, the *old occurrences*. The series converges absolutely, because the sum of $|\varphi_k(\Gamma)|$ over the clusters incompatible with a given group is finite (for $k \geq 2$ by Lemma 7.3 at the step k, and for $k = 1$ by the corresponding bound of the first step).

Inserting (6.26), (6.27), and (6.28) into (6.24), we obtain a sum over *terms* $t = (\mathfrak{B}, \mathfrak{Y}, \mathfrak{M}, \mathcal{L}, \mathcal{O}, (\Gamma_j))$. The *constituents* of a term are its marked polymers, the members of $\mathfrak{B}$, and the groups of its old occurrences. A vacuum occurrence, like a link, is *carried* by the first, in the fixed order, of the marked polymers with which it is incompatible. A constituent has a *domain*: a marked polymer M the set M together with the domains of the links which it carries, and a group W its dependence set X_W^+; two such sets are compatible, as before, if their intersection has dimension at most $d-2$. It also has *pieces* on the loops. The pieces of a group are its intervals. If the arc sets of the groups of $\mathfrak{G}(M)$ lie on one loop C_i, the marked polymer M has one piece, the arc $I_i(M)$ of C_i obtained by removing the longest maximal run of bonds which are not in these arc sets, the first such run from the base point of C_i if several are longest, and $I_i(M) \coloneqq C_i$ if they cover C_i; otherwise its pieces are the pieces of the groups of $\mathfrak{G}(M)$. A constituent *lies on the loop* C_i if all its pieces are arcs of C_i, and it then has a single piece, since a group which is not a cross group has one interval (Section 5.3).

Definition 6.10 (The Groups of the Step)**.** Two constituents of a term are *tied* in each of the following cases.

(i) A member of $\mathfrak{B}$ is touched by a component of a group of $\mathfrak{G}(M)$: it is tied to M.

(ii) A marked polymer belongs to the set T of a link of $\mathcal{L}$, or is incompatible with a vacuum occurrence of $\mathcal{O}$: it is tied to the marked polymer which carries the link or the occurrence.

(iii) The domains of a marked polymer and of a group, a member of $\mathfrak{B}$ or a group of an old occurrence, are incompatible.

(iv) The domain of a link carried by a marked polymer is incompatible with another marked polymer or with the domain of a link which that one carries.

(v) Two groups of one old occurrence are incompatible, or a group of an old occurrence is incompatible with a member of $\mathfrak{B}$; here the relation is the compatibility of the groups of the step k, and two constituents which are the same group of the step k count as incompatible.

The *arc set* of a set of constituents is the union of their pieces and, for each tied pair among them both members of which lie on the same loop C_i and have disjoint pieces, of the shorter of the two arcs of C_i between these pieces; if the two arcs have the same length, the one which follows the earlier piece in the orientation of C_i is taken. No arc is added for a tied pair one of whose members has pieces on several loops. The *groups of the step* $k+1$ in the term t are the classes of the smallest equivalence relation on its constituents which contains the tied pairs and for which, in addition, two classes whose arc sets share a bond coincide, and a member of $\mathfrak{B}$ one of whose intervals lies in the arc set of a class belongs to that class. For a group $\mathcal{G}$ we write $I(\mathcal{G})$ for its arc set, whose maximal arcs are its *intervals*; its localisation domains are the unions of the cubes of π_{k+1} which meet the domain of one of its constituents or a polymer of one of its vacuum occurrences; and its *insertion* $Y_{\mathcal{G}}$ is the contraction, along its intervals and in the order of each loop, of the tensors $e^{-\Psi_\circ(M)}\tilde{H}(M)$ of its marked polymers, the insertions at U_{k+1} of its untouched members, and the transports of U_{k+1} between them, multiplied by the factors u_λ of the links which its marked polymers carry, the factors $v_{\mathcal{C}}$ of its vacuum occurrences, and $((-1)^m/m!)\prod_j \varphi_k(\Gamma_j)$ for the sequence of its m old occurrences. Two groups, each given with its constituents, links, and occurrences, are *compatible* if their arc sets are disjoint; the domain of every marked polymer of the one is compatible with the domain of every constituent of the other; every member of $\mathfrak{B}$ of the one is compatible, as a group of the step k, with every member of $\mathfrak{B}$ of the other; no vacuum occurrence of the one is incompatible with a marked polymer of the other; and no group of an old occurrence of the one is incompatible, as a group of the step k, with a member of $\mathfrak{B}$ of the other. Two old occurrences of different groups are thus not required to be compatible with each other, as the series (6.28) imposes no condition between its clusters.

The equivalence relation exists and is unique: the relation in which all constituents are equivalent satisfies the conditions, and the conditions are preserved by intersections of equivalence relations, since the arc set of a set of constituents increases with the set. A link or an occurrence belongs to one group: the marked polymers of the set T of a link are tied to its carrier by (ii), and so are the marked polymers with which a vacuum occurrence is incompatible; and any two groups of an old occurrence are joined by a chain of groups of the occurrence in which consecutive ones are incompatible, the incompatibility graph of a cluster being connected, so that (v) ties them. An interval J of a member W of $\mathfrak{B}$ which is not in a group $\mathcal{G}$ is disjoint from $I(\mathcal{G})$. Indeed, J does not lie in $I(\mathcal{G})$, by the definition of the groups. If it met $I(\mathcal{G})$, it would contain an end bond of a maximal arc of $I(\mathcal{G})$ and a bond outside $I(\mathcal{G})$. That end bond belongs to a piece of a constituent of $\mathcal{G}$, because an arc which is added between two pieces has a piece at each of its ends; so J would contain a bond of a piece without lying in that piece. For a piece of a marked polymer M this is excluded by the first statement on the sets $S_i'(Y)$ in Section 6.3, by which an interval of a member that contains a bond of such a set lies in it: if M has pieces on several loops, a bond of a piece is a bond of a set $S_i'(Y)$, which is contained in that piece, and if M lies on one loop, J contains an end bond of $I_i(M)$, which is a bond of a set $S_i'(Y) \subset I_i(M)$. It is excluded for an interval of a member of $\mathfrak{B}$, because the intervals of a family are disjoint; and for an interval of a group of an old occurrence, because that group would then be incompatible with W, two groups whose intervals share a bond being incompatible when neither is a link group, and W would belong to $\mathcal{G}$ by (v); here W is not a link group, since J has more than one bond. So the arc sets of the groups of a term are pairwise disjoint, by the definition; every interval of a constituent lies in the arc set of its group; and no interval of a member meets the arc set of another group. Consequently the word of the term is the word, in the sense of Section 5.3, of the insertions $Y_{\mathcal{G}}$ at the intervals of the groups, with the transports of U_{k+1} between them; this uses only that the word is an ordered product along each loop which is multilinear in its

factors, and that the numerical factors of a term are distributed over the groups as in the definition, the factor $1/m!$ of (6.28) because the sequences (Γ_j) which induce given sequences in the groups are $m!/\prod_{\mathcal{G}} m_{\mathcal{G}}!$ in number.

Proposition 6.11 (The Regrouping)**.** *On the terms without large-field regions of ϱ_k and of ϱ_{k+1}, suppose that the sum over the terms t of the products of the absolute values of their numerical factors and of the norms of their tensors converges. Then*

$$\langle F \rangle = \prod_{i=1}^{n} \kappa^{(k)}(C_i, \chi_i) \cdot e^{\mathcal{E}_k} \Big[\prod_{i=1}^{n} f_{k+1}^{(i)} + \sum_{\mathfrak{F} \neq \emptyset} w_{k+1}(\mathfrak{F}) \Big] , \tag{6.29}$$

the sum over the finite sets $\mathfrak{F}$ of pairwise compatible groups of the step $k+1$, each with its constituents, links, and occurrences, where $w_{k+1}(\mathfrak{F})$ is the word of the insertions $Y_{\mathcal{G}}$, $\mathcal{G} \in \mathfrak{F}$. Each $Y_{\mathcal{G}}$ depends on the background field U_{k+1} only on the domains of the constituents of $\mathcal{G}$ and along its intervals.

Proof. By the convergence assumed, the series obtained by inserting (6.26), (6.27), and (6.28) into (6.24) may be summed in any order, and by the remarks after Definition 6.10 the contribution of a term t is $w_{k+1}(\mathfrak{F})$ for the set $\mathfrak{F}$ of its groups; the term without constituents gives the product of the leading words. It remains to see that $t \mapsto \mathfrak{F}$ is a bijection onto the sets of pairwise compatible groups, the sequences of old occurrences being recorded group by group. The groups of a term are pairwise compatible: their arc sets are disjoint, as was shown; the marked polymers of $\mathfrak{M}$ are pairwise compatible and the members of $\mathfrak{B}$ are pairwise compatible as groups of the step k, by the definitions of these sets; two constituents in different groups are not tied, which gives the compatibility of the domain of a marked polymer with the domain of a group by (iii), and with that of another marked polymer by (iv); and the conditions on the vacuum and on the old occurrences hold by (ii) and (v). Conversely, let $\mathfrak{F}$ be a set of pairwise compatible groups, and let t be the union of their data. Its marked polymers are pairwise compatible, so that their components, which lie in the sets D_G of their groups, form a set $\mathfrak{Y}$ of Lemma 6.4; the members form a family; and the number $s(\mathcal{C})$ of a vacuum occurrence, and with it the marked polymer which carries it, is the same in t as in its group, since it is incompatible with no marked polymer of another group. No constituent of one group is tied in t to one of another: for (i), a component which touches a member W lies in a marked polymer, and the intersection of that polymer with X_W^+ then contains an open set, so that the two domains are incompatible; for (ii), a link belongs to one group and a vacuum occurrence is compatible with the marked polymers of the other groups; for (iii) and (iv) by the compatibility of the domains of the marked polymers with those of the other constituents; for (v) by the condition on the old occurrences, each of which is contained in one group. Their arc sets are disjoint, so that no member has an interval in the arc set of another group, and no further identification occurs. Hence the groups of t are the members of $\mathfrak{F}$. The locality of $Y_{\mathcal{G}}$ follows from that of the tensors $\tilde{H}(M)$ (Proposition 6.8), of the factors u_λ, and of the insertions of the members, the other factors being numbers. *q.e.d.*

Distinct groups of a set $\mathfrak{F}$ interact only through the hard core of the compatibility: no soft interaction is left. The hypothesis of the proposition is verified in Section 6.6 together with the bounds of the insertions.

6.6 The Count of the Groups of the Step

Proposition 6.11 is so far an identity between a sum and a rearrangement of it, and what it asserts depends on the convergence of that sum. We now prove the convergence, and with

it the bound on the groups of the step $k+1$ from which their place in the class will follow. What makes this a problem of counting is that a group is not the sum of its parts. Its size σ_{k+1}, in which the bounds of the class are weighted, counts the cubes and the traversals of its domain, and the domain contains arcs which belong to none of the constituents: the arcs which the regrouping adds between them. If the weights of the constituents are to pay for the weight of the group, these arcs have to be charged to the constituents which they join, and each constituent may be charged only in proportion to its own size, however many others it is tied to. We arrange this by giving every constituent a set of cubes which is large enough to contain all that the constituent can add to a group, and whose number is bounded by its own size. Tied constituents then have a cube in common, a group becomes a connected set in the graph of this relation, and the connected sets of a graph are counted by a tree bound, which is the criterion of Kotecký and Preiss in the form we need.

Throughout this subsection the labelled terms of ϱ_k and of ϱ_{k+1} are the terms without large-field regions, lengths and distances are measured in the units of the scale $k+1$ unless the contrary is said, distances are ℓ^∞-distances, and a *cube* is a cube of π_{k+1}, of side M. By Lemma 3.14 at the step $k+1$ two loops are at distance at least $a_{k+1} \geq L^2 M$ and the segments of the loops have at least $L^2 M$ bonds, so that a cube meets at most one loop, and by Lemma 5.5(i) it meets at most $2\bar{\Lambda}(M+1) \leq 4\bar{\Lambda}M$ traversals of the step $k+1$.

Lemma 6.12 (Coarse Cubes)**.** *Let $j \leq k$, and let X be a union of n cubes of π_j which is connected in the sense of Definition 5.2(vii). Then* $\operatorname{diam}_\infty(X) \leq nM/L$*, and X meets at most* $625(1+n/L)$ *cubes.*

Proof. A cube of π_j has the side $ML^{-(k+1-j)} \leq M/L$ in the units of the scale $k+1$, and two points of X are joined by a chain of at most n of its cubes in which consecutive ones have a common point; this gives the diameter. For the count let $\mathcal{Q}$ be the set of the cubes which X meets, and let $\mathcal{Q}' \subset \mathcal{Q}$ be maximal among the subsets whose members have centres at distance at least $3M$ from each other. Every member of $\mathcal{Q}$ has its centre within $2M$ of the centre of a member of $\mathcal{Q}'$, so that $|\mathcal{Q}| \leq 5^4 |\mathcal{Q}'|$. For $Q \in \mathcal{Q}'$ let Q^* be the union of Q and of the 80 cubes which touch it. The sets Q^*, $Q \in \mathcal{Q}'$, have no cube in common, and every cube of π_j lies in one cube, the partitions being nested. If $\mathcal{Q}'$ has at least two members, then X contains, besides a point of Q, a point of another member of $\mathcal{Q}'$, whose distance from Q is at least $2M$. A chain of cubes of π_j in X from the first point to the second begins with a cube contained in Q^* and leaves Q^*, and its cubes up to the first one which is not contained in Q^* form a connected set which reaches from a point of Q to a point of the boundary of Q^*, at distance M from Q; since each of them has diameter at most M/L, at least L of them are contained in Q^*. Hence $n \geq L|\mathcal{Q}'|$ in this case, and $|\mathcal{Q}| \leq 625 \max(1, n/L)$ in both. *q.e.d.*

Bodies, windows, and cube sets. Fix a term t of Section 6.5. For a marked polymer M of t let $\mathcal{K}(M)$ be the set of the clusters of the links and of the vacuum occurrences which M carries. The *body* of a constituent, a group W or a marked polymer M, is

$$B(W) \coloneqq X_W^+ , \qquad B(M) \coloneqq M \cup \bigcup_{\mathcal{C} \in \mathcal{K}(M)} \operatorname{supp} \mathcal{C} \cup \bigcup_{W \in \mathfrak{B}(M)} X_W^+ , \tag{6.30}$$

and $d(c) \coloneqq \operatorname{diam}_\infty B(c)$. If the constituent c lies on a loop C_i, with the piece $p(c)$, its *window* $\operatorname{win}(c)$ is the arc of C_i which consists of $p(c)$ and of the points of C_i whose distance from an end of $p(c)$, measured along C_i, is at most $2\bar{\Lambda} d(c)$. The *cube set* $K(c)$ is the set of the cubes which meet $B(c)$ or, if c lies on a loop, $\operatorname{win}(c)$; and $n(c)$ is the number of the traversals of the step $k+1$ which meet $\operatorname{win}(c)$ if c lies on a loop, and which meet a piece of c otherwise.

Lemma 6.13 (The Covering of a Group)**.** *Let $\mathcal{G}$ be a group of the step $k+1$ in the term t.*

(i) The body of a constituent is connected, and it contains the domain of the constituent, the polymers of the vacuum occurrences which it carries, and the end sites of its pieces. A piece of a constituent which does not lie on a loop is contained in its body.

(ii) The bodies of two tied constituents c, c' have a point in common. If both lie on the loop C_i and their pieces are disjoint, the arc which Definition 6.10 adds between these pieces has length at most $\bar{\Lambda}(d(c)+d(c'))$, and it lies in win$(c)$ *if $d(c) \geq d(c')$ and in* win(c') *otherwise.*

(iii) The arc set $I(\mathcal{G})$ is contained in the union of the windows of the constituents of $\mathcal{G}$ which lie on a loop and of the pieces of the others, $X_{\mathcal{G}}$ is contained in the union of the cubes of the sets $K(c)$, and any two constituents of $\mathcal{G}$ are joined by a chain of constituents of $\mathcal{G}$ in which the cube sets of consecutive ones have a cube in common.

(iv) The domain $X_{\mathcal{G}}$ is connected in the sense of Definition 5.2(vii). Its number of cubes satisfies $\#(X_{\mathcal{G}}) \leq |\bigcup_c K(c)|$, the number of the traversals of the step $k+1$ which meet $I(\mathcal{G})$ satisfies $n_{k+1}(\mathcal{G}) \leq \sum_c n(c)$, and ext$(\mathcal{G}) \leq 2\,\#(X_{\mathcal{G}}) + 3C_{\text{box}} + 1$*; the union and the sum run over the constituents of $\mathcal{G}$.*

(v) The insertion $Y_{\mathcal{G}}$ depends on the background field only through its restriction to $X_{\mathcal{G}}$, and the domains of two incompatible groups of the step $k+1$ have a point in common.

Proof. (i) For a group W the body is a closed neighbourhood of the connected set X_W, which contains the intervals of W. Let M be a marked polymer. Each of its connected components, in the sense of the walls, is a connected set, and so is, for each component Y of a group of $\mathfrak{G}(M)$, the union $Y^\sim$ of Y and of the cubes of π_k which share a point with it; the sets $Y^\sim$ lie in M, because D_G is their union over the components of G. By the definition of a linked union in Proposition 6.8, any two connected components of M are joined by a chain of connected components in which consecutive ones meet one set D_G, hence meet the sets $Y^\sim$ and $Y'^\sim$ of two components of one group G; and two components of a group are joined by a chain of components of that group in which consecutive ones touch a common member $W \in \mathfrak{B}(M)$, whose dependence set is connected and meets both. So $M \cup \bigcup_{W\in\mathfrak{B}(M)} X_W^+$ is connected. The support of a cluster is connected, its polymers being connected unions of cubes with a connected graph of incompatibility, and it shares a cube or a wall with M if M carries a link or an occurrence of that cluster. Hence $B(M)$ is connected, and it contains the domain of M and the polymers of its vacuum occurrences by definition. The pieces of the groups of $\mathfrak{G}(M)$ consist of bonds of the loops in their components, which lie in M, and of intervals of members of $\mathfrak{B}(M)$, which lie in the domains of these members; if M lies on a loop, its piece $I_i(M)$ begins and ends with such bonds.

(ii) We go through the cases of Definition 6.10. In (i) the member belongs to $\mathfrak{B}(M)$, and its body is a part of $B(M)$. In (ii) the marked polymer is incompatible with a cluster of $\mathcal{K}(M)$, for the carrier M, so that it shares a cube or a wall with the support of that cluster. In (iii) and (iv) the two domains share a cube or a wall, and they are contained in the bodies. In (v) the two groups are the same group or incompatible groups of the step k, and their domains have a common point by Definition 5.2(vii). Now let c and c' lie on C_i with disjoint pieces p and p', and let x and x' be end sites of p and of p'. They lie in the two bodies, which have a common point, so that $|x-x'|_\infty \leq d(c)+d(c')$, and by Definition 2.4 the shorter of the two arcs of C_i between x and x' has length at most $\Lambda(C_i)(d(c)+d(c'))$. This arc begins at an end of p. It may first run through p; it then leaves p at an end and runs outside p until

it meets p', which it does at an end of p'. Its part between these two ends is one of the two arcs between the pieces, and the shorter of the two, which is the arc that the definition adds, is not longer. That arc begins at an end of p, ends at an end of p', and has length at most $2\bar{\Lambda}\max(d(c), d(c'))$, which gives the last statement.

(iii) By its definition $I(\mathcal{G})$ is the union of the pieces of the constituents of $\mathcal{G}$ and of the arcs of (ii), and the piece of a constituent which lies on a loop is a part of its window; this is the first statement. A localisation domain of $\mathcal{G}$ consists of cubes which meet the domain of a constituent or a polymer of a vacuum occurrence, hence a body, by (i); and a cube which meets a fine bond of $I(\mathcal{G})$ meets a window, or a piece which lies in a body by (i). So $X_{\mathcal{G}}$ consists of cubes of the sets $K(c)$. For the third statement, call two constituents of the term *joined* if they are connected by a chain of constituents in which the cube sets of consecutive ones have a cube in common. This is an equivalence relation, and it has the three properties which define the groups. Tied constituents are joined, since every cube which contains a common point of their bodies belongs to both cube sets. If the arc sets of two classes share a bond, that bond lies in a window or in a piece of a constituent of each of them, by (ii), and a cube which meets it belongs to the cube sets of both constituents, so that the classes coincide. And a member of $\mathfrak{B}$ one of whose intervals lies in the arc set of a class has the bonds of that interval in its own body and in a window or a piece of a constituent of the class. Since the groups are the classes of the smallest equivalence relation with these properties, the constituents of a group are joined.

(iv) Let F be the union of the bodies of the constituents of $\mathcal{G}$ and of $I(\mathcal{G})$. The bodies of the constituents of $\mathcal{G}$ cover the same set as their domains and the polymers of their vacuum occurrences, since the members of $\mathfrak{B}(M)$ and the clusters of the links are constituents and parts of domains of constituents of $\mathcal{G}$; so $X_{\mathcal{G}}$ is the union of the cubes which meet F. The set F is connected: the bodies are connected, those of tied constituents meet, every piece and every arc added between two pieces is a connected set which meets the bodies of the constituents concerned at their end sites, and the classes which the definition merges because their arc sets share a bond, or because an interval of a member lies in an arc set, have a common point there. A finite family of closed cubes each of which meets a connected set, and which together cover it, is connected in the sense of Definition 5.2(vii), since otherwise that set would be the union of two disjoint nonempty closed sets. The bounds on $\#(X_{\mathcal{G}})$ and on $n_{k+1}(\mathcal{G})$ follow from (iii). For the extent put $N \coloneqq \#(X_{\mathcal{G}})$, and let $\mathfrak{n}$ be a traversal of the step $k+1$ in an interval of $\mathcal{G}$. It meets a cube of $X_{\mathcal{G}}$, and $X_{\mathcal{G}}$ is a connected union of N cubes of side M, so that $X_{\mathcal{G}}$ lies in the ball of radius $NM+1$ about $\mathfrak{n}$. The radii $R_t = C_{\mathrm{box}}M(t+1)$ increase in steps of $C_{\mathrm{box}}M \geq 1$, so that $R_{t_{\mathfrak{n}}(\mathcal{G})} \leq \max(2C_{\mathrm{box}}M, NM+1+C_{\mathrm{box}}M)$, and the comparison region of $\mathcal{G}$ at $\mathfrak{n}$ is a ball of radius at most $(N+3C_{\mathrm{box}})M$ about $\mathfrak{n}$. Another traversal of the intervals of $\mathcal{G}$ lies within $NM+1$ of $\mathfrak{n}$, and therefore $X_{\mathcal{G}}$ and all the comparison regions lie in the ball of radius $(2N+3C_{\mathrm{box}}+1)M$ about each of these traversals.

(v) The first statement is the last statement of Proposition 6.11, the domains of the constituents and the intervals of $\mathcal{G}$ being contained in $X_{\mathcal{G}}$. For the second we go through the conditions of the compatibility in Definition 6.10. If the arc sets share a bond, the cubes which meet it lie in both domains. If the domain of a marked polymer of the one is incompatible with the domain of a constituent of the other, or a vacuum occurrence of the one with a marked polymer of the other, the two sets share a cube or a wall, and both are covered by the localisation domains of their groups. If a member of the one, or a group of an old occurrence of the one, is incompatible with a member of the other as a group of the step k, their domains have a common point by Definition 5.2(vii) at the step k, and these domains lie in the dependence sets, which the localisation domains cover. *q.e.d.*

The lemma reduces the geometry of a group to the cube sets of its constituents, and we now bound these sets. For a polymer the bound has to contain the members which it touches, since the arcs of the polymer contain their intervals.

Lemma 6.14 (The Sizes of the Cube Sets)**.** *Let* $L \geq 81\bar{\Lambda}$.

(i) For a group W *of the step* k *put* $s(W) \coloneqq |K(W)| + n(W)/M$. *Then*

$$d(W) \leq \frac{(u_\star \sigma(W) + 1)M}{L} , \qquad s(W) \leq 3000\,\bar{\Lambda} + 3 \cdot 10^4\,\bar{\Lambda} u_\star \frac{\sigma(W)}{L} , \tag{6.31}$$

and for every cube of $K(W)$ *there is a traversal of the step* k *in an interval of* W *whose distance from that cube, in the units of the scale* k*, is at most* $\max(\mathrm{ext}(W)M + 1,\, 2\bar{\Lambda}(u_\star\sigma(W) + 1)M)$.

(ii) For a marked polymer M *of the term let* $K(v) \coloneqq K(M) \cup \bigcup_{W \in \mathfrak{B}(M)} K(W)$, *let* $n(v)$ *be the number of the traversals of the step* $k+1$ *counted by* $n(M)$ *or by one of the numbers* $n(W)$, $W \in \mathfrak{B}(M)$, *and put* $s(v) \coloneqq |K(v)| + n(v)/M$ *and*

$$\hat{d}(v) \coloneqq \#(M) + \sum_{\mathcal{C} \in \mathcal{K}(M)} N(\mathcal{C}) + \sum_{W \in \mathfrak{B}(M)} \frac{u_\star \sigma(W) + 1}{L} , \tag{6.32}$$

with $N(\mathcal{C})$ *the number of the cubes of the polymers of* $\mathcal{C}$. *Then* $d(M) \leq M\hat{d}(v)$,

$$s(v) \leq 33 + 121\,\bar{\Lambda}\Big(\#(M) + \sum_{\mathcal{C} \in \mathcal{K}(M)} N(\mathcal{C})\Big) + \sum_{W \in \mathfrak{B}(M)} \Big(3001\,\bar{\Lambda} + 3.01 \cdot 10^4\,\bar{\Lambda} u_\star \frac{\sigma(W)}{L}\Big) , \tag{6.33}$$

and every cube of $K(v)$ *lies within the distance* $9\bar{\Lambda}M\hat{d}(v)$ *of every cube of* M.

Proof. (i) Let W be made at the step $j \leq k$, and put $N_W \coloneqq \#_j(X_W)$, so that $N_W \leq u_\star\sigma(W)$ and $n_j(W) \leq u_\star M\sigma(W)$ by Definition 5.2(vii). The dependence set lies in the union of X_W and of the cubes of π_j which touch it, a connected union of at most $81N_W$ cubes of π_j, since $8M$ fine bonds are less than the side of a cube of π_j. By Lemma 6.12, $B(W)$ meets at most $625(1 + 81N_W/L)$ cubes, X_W meets at most $625(1 + N_W/L)$, and $d(W) \leq N_W M/L + 16ML^{-(k+1)} \leq (N_W + 1)M/L$. If W is a cross group, its intervals lie in X_W, and each of the cubes which X_W meets is met by at most $4\bar{\Lambda}M$ traversals, so that $n(W)/M \leq 2500\bar{\Lambda}(1 + N_W/L)$. If W lies on a loop, its interval has the length $n_j(W)L^{-(k+1-j)} \leq u_\star M\sigma(W)/L$ and lies in X_W, and the window adds to it two arcs of length $2\bar{\Lambda}d(W)$. By Lemma 2.8, applied on the fine lattice, an arc of length ℓ meets at most $8\ell/M + 16\ell/(L^2M) + 32 \leq 9\ell/M + 32$ cubes, the segments having at least L^2M bonds of the scale $k+1$. Hence the two arcs meet at most $36\bar{\Lambda}(N_W + 1)/L + 64$ cubes, and $n(W) \leq u_\star M\sigma(W)/L + 4\bar{\Lambda}d(W) + 2$. Adding up, and using $\bar{\Lambda} \geq 2$ and $L \geq 81\bar{\Lambda}$, we obtain the bound on $s(W)$ in both cases, with room to spare. For the last statement, a cube of $K(W)$ meets $B(W)$ or one of the two arcs. In the first case its distance from every traversal of the step j in an interval of W is at most $\mathrm{ext}(W)M$ in the units of the scale j, by the definition of the extent, plus the $8M$ fine bonds of the dependence set; in the second its distance from the traversal at the corresponding end of the interval is at most $2\bar{\Lambda}d(W)$, which is $2\bar{\Lambda}(N_W + 1)M$ in the units of the scale k. A length in the units of the scale j is not larger in those of the scale k.

(ii) The body $B(M)$ is a connected union of the cubes of M, of the supports of the clusters of $\mathcal{K}(M)$, and of the bodies of the members of $\mathfrak{B}(M)$, by Lemma 6.13(i), and the diameter

of a connected finite union of closed sets is at most the sum of their diameters; with (i) this gives $d(M) \le M\hat{d}(v)$. The cubes which meet M or one of these supports are at most $81(\#(M) + \sum_{\mathcal{C}} N(\mathcal{C}))$, and those which meet the body of a member W belong to $K(W)$. If M lies on the loop C_i, the sites of the arc sets of its groups lie in $B(M)$, and $I_i(M)$ is the shortest arc of C_i which contains them, so that it has length at most $4\Lambda(C_i)d(M)$ by Lemma 2.6 and the remark which follows it. The window then has length at most $8\bar{\Lambda}d(M)$, it meets at most $72\bar{\Lambda}\hat{d}(v) + 32$ cubes, by the count of (i), and $n(M) \le 8\bar{\Lambda}M\hat{d}(v) + 2$. If M does not lie on a loop, its pieces consist of bonds of the loops in M, which are met by at most $4\bar{\Lambda}M\#(M)$ traversals, and of intervals of the members of $\mathfrak{B}(M)$, which are counted by the numbers $n(W)$. In both cases

$$s(v) \le 33 + 81\Big(\#(M) + \sum_{\mathcal{C}} N(\mathcal{C})\Big) + 80\,\bar{\Lambda}\hat{d}(v) + \sum_{W \in \mathfrak{B}(M)} s(W)\ ,$$

which is (6.33) by (i), $81 \le 41\bar{\Lambda}$, and $L \ge 81\bar{\Lambda}$. Finally, a point of $B(M)$ is within $d(M)$ of every cube of M; a point of the window of M is within $8\bar{\Lambda}d(M)$ of an end site of $I_i(M)$, which lies in $B(M)$; and a point of the window of a member $W \in \mathfrak{B}(M)$ is within $2\bar{\Lambda}d(W) \le 2\bar{\Lambda}M\hat{d}(v)$ of its interval, which lies in $B(M)$. So every cube of $K(v)$ is within $(8\bar{\Lambda}+1)M\hat{d}(v)$ of every cube of M. *q.e.d.*

The tree bound. The sum over the connected sets of a graph is controlled by one condition on the weights of the neighbours of a vertex. We state it with a second weight, which marks one vertex of the set; this is what will show that a group which contains an insertion of an earlier step is small in proportion to that insertion.

Lemma 6.15 (Connected Sets of a Graph)**.** *Let $\mathcal{V}$ be a countable set with a symmetric relation $\sim$, and let $w, b\colon \mathcal{V} \to [0,\infty)$ be such that, for every $v \in \mathcal{V}$,*

$$\sum_{v' \sim v} w(v')\, e^{2b(v')} \le \tfrac{1}{2} b(v)\ . \tag{6.34}$$

For $v_0 \in \mathcal{V}$ let $\mathcal{S}(v_0)$ be the set of the finite subsets of $\mathcal{V}$ which contain v_0 and are connected in the graph of the relation.

(i) $\sum_{S \in \mathcal{S}(v_0)} \prod_{v \in S \setminus \{v_0\}} w(v) \le e^{b(v_0)/2}$.

(ii) Let $m\colon \mathcal{V} \to [0,\infty)$ and $\epsilon \ge 0$ be such that $\sum_{v' \sim v} m(v') e^{2b(v')} \le \frac{1}{2}\epsilon\, b(v)$ for every v. Then

$$\sum_{S \in \mathcal{S}(v_0)} \sum_{v_1 \in S \setminus \{v_0\}} m(v_1) \prod_{v \in S \setminus \{v_0, v_1\}} w(v) \le \epsilon\, b(v_0)\, e^{b(v_0)/2}\ .$$

Proof. Put $T_0(v) \coloneqq 1$ and $F_0(v) \coloneqq 0$, and for $n \ge 0$

$$T_{n+1}(v) \coloneqq \sum_{A} \prod_{c \in A} w(c) T_n(c)\ ,$$

$$F_{n+1}(v) \coloneqq \sum_{A} \sum_{c \in A} [m(c) T_n(c) + w(c) F_n(c)] \prod_{c' \in A \setminus \{c\}} w(c') T_n(c')\ ,$$

the sums over the finite sets A of vertices $c \sim v$. Then $T_{n+1}(v) \le \exp \sum_{c \sim v} w(c) T_n(c)$, and by induction $T_n(v) \le e^{b(v)/2}$, because $\sum_{c \sim v} w(c) e^{b(c)/2} \le \frac{1}{2} b(v)$ by (6.34). Likewise $F_{n+1}(v) \le e^{b(v)/2} \sum_{c \sim v} [m(c) T_n(c) + w(c) F_n(c)]$, and by induction $F_n(v) \le \epsilon\, b(v) e^{b(v)/2}$: the

first terms sum to at most $\frac{1}{2}\epsilon\, b(v)$ by the hypothesis on m, and the second to at most $\epsilon \sum_{c\sim v} w(c)b(c)e^{b(c)/2} \le \frac{1}{2}\epsilon\, b(v)$, by (6.34) and $xe^{x/2} \le e^{2x}$.

It remains to compare the sums of the lemma with these numbers. For $S \in \mathcal{S}(v_0)$ let the *depth* of S be the largest distance from v_0 in the graph which the relation induces on S. We claim that the sum in (i) over the sets of depth at most n is at most $T_n(v_0)$, and the sum in (ii) at most $F_n(v_0)$; the lemma follows by letting n tend to infinity. For $n = 0$ the only set is $\{v_0\}$. Let S have depth at most $n+1$, and fix an enumeration of $\mathcal{V}$. Every $u \in S \setminus \{v_0\}$ has a parent, the first vertex in the enumeration among its neighbours in S which are closer to v_0 than u; this defines a tree on S with the root v_0. Let A be the set of the neighbours of v_0 in S, and for $c \in A$ let S_c be the set of c and of its descendants. Then $S_c \in \mathcal{S}(c)$ with depth at most n, the set S is the disjoint union of $\{v_0\}$ and of the sets S_c, and the product of the weights over $S \setminus \{v_0\}$ is the product over $c \in A$ of $w(c)$ times the product over $S_c \setminus \{c\}$. Since S is determined by A and the sets S_c, the sum over S is at most the sum over A of the product over $c \in A$ of $w(c)$ times the sum over S_c, which is the claim for (i) by induction. In (ii) the marked vertex v_1 belongs to exactly one of the sets S_c, where it is c itself or a marked vertex of $S_c \setminus \{c\}$, and the same argument gives the recursion of F_{n+1}. *q.e.d.*

The vertices. We describe a group of the step $k+1$ by a set of vertices of three kinds, each with a cube set K, a number s, and an activity. A *member vertex* is a group W of the step k, with $K(W)$ and $s(W)$ of Lemma 6.14 and the activity $\|Y_W\|_{(k)}$. An *occurrence vertex* is a pair (Γ, n) of a cluster Γ of the reference gas of the step k and an integer $n \ge 1$, with $K(\Gamma) := \bigcup_{W'\in\Gamma} K(W')$, $s(\Gamma) := \sum_{W'\in\Gamma} s(W')$, the sum over the distinct groups of Γ, and the activity $|\varphi_k(\Gamma)|^n/n!$. A *polymer vertex* is a quadruple $v = (M, \mathfrak{Y}_v, \mathfrak{B}_v, \mathcal{K}_v)$ of a union M of cubes, a nonempty finite set $\mathfrak{Y}_v$ of connected unions of cubes of π_k in M, a finite set $\mathfrak{B}_v$ of groups of the step k, and a finite set $\mathcal{K}_v$ of clusters of vacuum polymers which are incompatible with M, such that the dependence set of every $W \in \mathfrak{B}_v$ meets M. Its cube set, its number $s(v)$, and its reach $\hat{d}(v)$ are those of Lemma 6.14(ii), formed with $\mathfrak{B}_v$ and $\mathcal{K}_v$ in the places of $\mathfrak{B}(M)$ and $\mathcal{K}(M)$. To define its activity let $\bar{\varphi}(\mathcal{C})$ be the bound of $|\varphi^T(\mathcal{C})|$ on the space of the step $k+1$ of Supplement, Lemma 17.24(iv), let $d(\mathcal{C}) = 9\log 2 \cdot N(\mathcal{C})$, and put

$$\begin{gathered} \xi(\mathcal{C}) := \bar{\xi}(\mathcal{C})e^{\bar{\xi}(\mathcal{C})}\,, \qquad \bar{\xi}(\mathcal{C}) := 2e^{d(\mathcal{C})}\bar{\varphi}(\mathcal{C})\,, \\ \kappa' := \kappa_1 - 2 - c_\kappa\,, \qquad r_\kappa := \frac{(L+2)\kappa}{1296}\,, \qquad \varpi_k := \frac{b_k}{\min(\varepsilon_H, r_\alpha)}\,. \end{gathered} \tag{6.35}$$

The activity of the polymer vertex v is

$$\mathrm{act}(v) := \mathfrak{a}_v\, e^{c\#(M)}\, e^{-\kappa'\#(\mathfrak{Y}_v)}\, e^{-r_\kappa \#(M)} \prod_{W\in\mathfrak{B}_v} \|Y_W\|_{(k)} \prod_{\mathcal{C}\in\mathcal{K}_v} \xi(\mathcal{C})\,, \tag{6.36}$$

where $c = \frac{9}{64}e^{-4\kappa}$, $\#(\mathfrak{Y}_v)$ is the total number of the cubes of the members of $\mathfrak{Y}_v$, and $\mathfrak{a}_v := c_a\rho_k$ if $\mathfrak{B}_v = \emptyset$ and $\mathfrak{a}_v := \varpi_k$ otherwise, with the constant c_a of Proposition 6.9. For every vertex v we put

$$w(v) := \mathrm{act}(v)\, e^{4\nu s(v)}\,, \qquad b(v) := \nu|K(v)|\,, \tag{6.37}$$

and we write $v \sim v'$ if $K(v)$ and $K(v')$ have a cube in common. The *vertex set* $S(\mathcal{G})$ of a group $\mathcal{G}$ of the step $k+1$ consists of the polymer vertices $(M, \mathfrak{Y}_M, \mathfrak{B}(M), \mathcal{K}(M))$ of its marked polymers, where $\mathfrak{Y}_M$ is the set of the components of the groups of $\mathfrak{G}(M)$; of its members which no component touches; and of the pairs (Γ, n) of the distinct clusters Γ of its

old occurrences with their multiplicities n in the sequence of these occurrences. Let finally

$$P(\mathcal{G}) \coloneqq e^{c'_\pi \alpha_{1,k+1} n_{k+1}(\mathcal{G})} \prod_M \sup\|e^{-\Psi_\circ(M)} \tilde{H}(M)\| \prod_W \|Y_W\|_{(k)} \prod_\lambda \sup|u_\lambda| \prod_{\mathcal{C}} |v_{\mathcal{C}}| \cdot \frac{1}{m!} \prod_{j=1}^{m} |\varphi_k(\Gamma_j)| \tag{6.38}$$

be the product of the norms of the factors of $Y_{\mathcal{G}}$ in Definition 6.10, the suprema over the space of the step $k+1$ and the products over the marked polymers, the untouched members, the links, the vacuum occurrences, and the old occurrences of $\mathcal{G}$. The first factor bounds the transports: on the space of the step $k+1$ the transport along one traversal of the step $k+1$ has norm at most $e^{c'_\pi \alpha_{1,k+1}}$, by the radii of Definition 5.2(iii), and the insertion of an untouched member is evaluated at a configuration of that space, which belongs to the space of the step k (Section 6.3). Since a contraction of tensors does not increase the product of the norms (Section 5.3),

$$\sup\|Y_{\mathcal{G}}\| \le P(\mathcal{G}) \,, \tag{6.39}$$

the supremum over the space of the step $k+1$.

Lemma 6.16 (The Weight of a Group)**.** *Suppose that the hypotheses of Proposition 6.9 hold, that* $\kappa \ge 2^9(2+16\log 2)$*, and that* $4\bar{\Lambda} M c'_\pi \alpha_{1,k+1} \le \nu \le 1$*. Then the vertex set of every group of the step* $k+1$ *is connected in the graph of the relation* $\sim$*, and for every finite set* S *of vertices*

$$\sum_{\mathcal{G}\colon S(\mathcal{G})=S} e^{\frac{5}{2}\nu\sigma_{k+1}(\mathcal{G}) + \nu_e \operatorname{ext}(\mathcal{G})} P(\mathcal{G})\, \ell_{\mathcal{G}} \le \frac{e^{2\nu}}{\nu} \prod_{v\in S} w(v) \,, \tag{6.40}$$

the sum over the groups of the step $k+1$*, in all the terms* t*, with the vertex set* S*.*

Proof. The cube set of a polymer vertex of $S(\mathcal{G})$ contains those of the marked polymer and of the members which it touches, and the cube set of an occurrence vertex those of the groups of the occurrence, so that the vertex set is connected by Lemma 6.13(iii).

A group is determined by its vertex set up to the links and the vacuum occurrences which its marked polymers carry: the vertex set gives the marked polymers with their components and touched members, hence their groups, since two components are linked if they touch a common member; the untouched members; the old occurrences with their multiplicities, to which there correspond $m!/\prod n!$ sequences; and, for each marked polymer M, the set $\mathcal{K}(M)$ of the clusters of its links and vacuum occurrences. We bound the factors of (6.38) in turn.

The clusters. Fix M and $\mathcal{C} \in \mathcal{K}(M)$, and let s be the number of the marked polymers of the group which are incompatible with $\mathcal{C}$. They are pairwise compatible, and each of them contains a cube which belongs to a polymer of $\mathcal{C}$ or shares a wall with such a cube, so that $s \le 9N(\mathcal{C})$. The links of $\mathcal{C}$ which M carries are pairs $(T,\mathcal{C})$ with M the first member of T, at most 2^{s-1} in number, and at most one vacuum occurrence is $\mathcal{C}$. By (6.25) and (6.27), $|u_\lambda| \le 2\bar{\varphi} e^{2\bar{\varphi}}$ and $|v_{\mathcal{C}}| \le (s-1)\bar{\varphi} e^{(s-1)\bar{\varphi}}$ with $\bar{\varphi} = \bar{\varphi}(\mathcal{C})$. By Supplement, Lemma 17.24(iv), applied to a polymer of $\mathcal{C}$, $\bar{\varphi} e^{d(\mathcal{C})+N(\mathcal{C})} \le cN(\mathcal{C})$, whence $9N(\mathcal{C})\bar{\varphi} \le c$ and $\bar{\xi}(\mathcal{C}) \le 1$. The sum, over the nonempty sets of links and occurrences of $\mathcal{C}$ carried by M, of the products of these numbers is therefore at most

$$(1+2\bar{\varphi}e^{2\bar{\varphi}})^{2^{s-1}} (1+(s-1)\bar{\varphi}e^{(s-1)\bar{\varphi}}) - 1 \le x e^x \,, \qquad x \coloneqq 2^s \bar{\varphi} e^c + 2^{s-1}\bar{\varphi} e^c \le \bar{\xi}(\mathcal{C}) \,,$$

because $s-1 \le 2^{s-1}$, $\frac{3}{2}e^c \le 2$, and $2^s \le e^{d(\mathcal{C})}$. This is the factor $\xi(\mathcal{C})$ of (6.36).

The other factors. By Supplement, Lemma 17.24(ii), at the reference data, $|\Psi_\circ(M)| \le c\#(M)$. By Proposition 6.9, and since $\mathfrak{a}(M) \le \mathfrak{a}_v$ in each of its three cases, $\sup\|\tilde{H}(M)\|$

is at most the product of the factors of (6.36) other than $e^{c\#(M)}$ and those of the clusters. The factors of the untouched members are the activities of the member vertices, and the sum of $(1/m!)\prod_j|\varphi_k(\Gamma_j)|$ over the sequences with given multiplicities is the product of the activities of the occurrence vertices. So the sum of $P(\mathcal{G})e^{-c'_\pi\alpha_{1,k+1}n_{k+1}(\mathcal{G})}$ over the groups with the vertex set S is at most $\prod_{v\in S}\mathrm{act}(v)$.

The exponent. The arc set of a group depends on its links, and not on its vertex set alone, but the following bounds hold for every group $\mathcal{G}$ with the vertex set S. By Lemma 6.13(iv), with $N \coloneqq \#(X_\mathcal{G}) \le \sum_{v\in S}|K(v)|$ and $n_{k+1}(\mathcal{G}) \le \sum_{v\in S} n(v)$, we have $\sigma_{k+1}(\mathcal{G}) = N + n_{k+1}(\mathcal{G})/M \le \sum_{v\in S} s(v)$, $\nu_e\,\mathrm{ext}(\mathcal{G}) \le 2\nu_e N + 2\nu$ because $\nu_e \le \nu/(2C_{\mathrm{box}})$, and $\ell_\mathcal{G} = N \le \nu^{-1}e^{\nu N}$. The hypothesis on $\alpha_{1,k+1}$ gives $c'_\pi\alpha_{1,k+1}n_{k+1}(\mathcal{G}) \le \nu n_{k+1}(\mathcal{G})/M$. Altogether the exponent is at most $2\nu + (\frac{5}{2}+\frac{3}{2})\nu\,\sigma_{k+1}(\mathcal{G}) \le 2\nu + 4\nu\sum_{v\in S} s(v)$, which with (6.37) gives (6.40). *q.e.d.*

The rooted sums. By Lemma 6.15 the sum over the groups is now reduced to the sums of the weights of the vertices which contain a given cube, and these are what the class at the step k and the bounds of the polymers control. The class is stated per traversal, and the first thing to do is to pass from traversals to cubes; this is where the weight of the extent in Definition 5.2(vii) is spent. Let

$$\mathsf{E}_k \coloneqq \sup_{\mathfrak{n}} \sum_{W\ni\mathfrak{n}} e^{\theta(W)+\lambda_c\nu\sigma(W)+\nu_e\,\mathrm{ext}(W)}\|Y_W\|_{(k)} \tag{6.41}$$

be the supremum over the traversals of the step k of the sum over all the groups of the term of ϱ_k with $\mathfrak{n}$ in an interval, of whatever species; by (5.21) the groups of the deviation species contribute at most $C_c\rho_k$ to it. We use the constants

$$\begin{aligned} c_r &\coloneqq \frac{\nu}{11664\,\bar\Lambda u_\star}\,, \qquad K_{\mathrm{box}} \coloneqq 6\bar\Lambda LM\sum_{t\ge0}(t+3)^5 e^{-c_r(t-2\bar\Lambda)_+}\,, \\ c_p &\coloneqq \min\Big(\frac{1}{28800\,\bar\Lambda}, \frac{\nu L}{1152\,\bar\Lambda u_\star}\Big)\,, \qquad C_{\mathrm{root}} \coloneqq \sum_{t\ge0} 8(2t+1)^3 e^{-c_p(t-1)_+}\,, \end{aligned} \tag{6.42}$$

which depend on $\bar\Lambda$, $u_\star$, ν, L, and M only, and we say that a vertex v is *at* a cube Δ if $\Delta \in K(v)$.

Proposition 6.17 (The Rooted Sums)**.** *Suppose that the hypotheses of Lemma 6.16 hold, and moreover*

$$\nu \le \frac{1}{1.2\cdot10^6\,\bar\Lambda}\,, \qquad L \ge 6\cdot10^6\,\bar\Lambda u_\star\,, \qquad r_\kappa \ge 10\,, \qquad L^4e^{-\kappa'} \le \frac{\nu}{1.6\cdot10^5\,u_\star}\,. \tag{6.43}$$

Let Δ be a cube.

- *(i)* $\sum_{W\ at\ \Delta}\|Y_W\|_{(k)}e^{(\nu/16)\sigma(W)} \le K_{\mathrm{box}}\mathsf{E}_k$*, the sum over the groups of the term of* ϱ_k*, and* $\sum_{W\ at\ \Delta} S_W e^{(\nu/16)\sigma(W)} \le 2K_{\mathrm{box}}C_{S'}\rho_k$*, the sum over the groups made at the step* k*.*
- *(ii) The member vertices satisfy* $\sum_{W\ at\ \Delta} w(W)e^{2b(W)} \le e^{1/60}K_{\mathrm{box}}\mathsf{E}_k$*.*
- *(iii) If* $10368\,u_\star e^{1/60}C_{S'}\rho_k \le \nu$*, the occurrence vertices satisfy* $\sum_{(\Gamma,n)\ at\ \Delta} w(\Gamma,n)e^{2b(\Gamma)} \le 6K_{\mathrm{box}}C_{S'}\rho_k$*.*
- *(iv) The polymer vertices with* $\mathfrak{B}_v = \emptyset$ *and* $M \in \mathbf{D}_{k+1}$ *satisfy* $\sum_{v\ at\ \Delta} w(v)e^{2b(v)} \le C_\alpha\rho_k$ *with* $C_\alpha \coloneqq 2C_{\mathrm{root}}c_aL^4e^{-\kappa'}$*.*

(v) If $e^{1/50} K_{\text{box}} \mathsf{E}_k \le \frac{1}{2}$*, the polymer vertices of the vertex sets of the groups of the step* $k+1$ *which have* $\mathfrak{B}_v \neq \emptyset$ *satisfy* $\sum_{v \text{ at } \Delta} w(v) e^{2b(v)} \le C_\beta \varpi_k \mathsf{E}_k$ *with* $C_\beta := 3 C_{\text{root}} K_{\text{box}} L^4 e^{-\kappa'}$.

Proof. (i) Let W be at Δ. By Lemma 6.14(i) there is a traversal $\mathfrak{n}$ of the step k in an interval of W whose distance D from Δ, in the units of the scale k, satisfies $D \le \text{ext}(W)M + 1$ or $D \le 2\bar\Lambda(u_\star \sigma(W) + 1)M$. In the first case $\frac{1}{2}\nu_e \,\text{ext}(W) \ge \frac{1}{2}\nu_e (D-1)/M$, in the second $\frac{1}{64}\nu\sigma(W) \ge (\nu/(128\bar\Lambda u_\star))(D/M - 2\bar\Lambda)$, and in both

$$e^{\frac{1}{16}\nu\sigma(W)} \le e^{-c_r (D/M - 2\bar\Lambda)_+} \, e^{\lambda_c \nu\sigma(W) + \nu_e \,\text{ext}(W)} ,$$

since $\frac{1}{16} + \frac{1}{64} \le \lambda_c$ and $\frac{1}{2}\nu_e = \nu/(11664\Lambda_*) \ge c_r$, the latter because $\bar\Lambda u_\star \ge \Lambda_*$. The traversals of the step k at a distance less than $(t+1)M$ from Δ meet a cube of side at most $LM + 2(t+1)M + 2$ in the units of the scale k. By Lemma 3.14 the loops which meet that cube are at distance at least $L^2 M$ from each other, so that they are at most $(t+3)^4$ in number, and by Lemma 5.5(i) each of them has at most $2\bar\Lambda(LM + 2(t+1)M + 3) \le 6\bar\Lambda LM(t+3)$ traversals which meet it. Summing over $t \ge 0$ the traversals with $tM \le D < (t+1)M$, and for each of them the groups W with that traversal in an interval, we obtain the first bound from (6.41), as $\theta(W) \ge 0$. The second is the same computation with (5.22) at the traversals of the step k, where $\ell_W \ge 1$ and the weight $\nu\sigma(W)$ takes the place of $\lambda_c \nu\sigma(W)$.

(ii) By (6.31) and the first two conditions of (6.43),

$$4\nu s(W) + 2b(W) \le 6\nu s(W) \le \frac{1}{60} + \frac{\nu}{32}\sigma(W) , \tag{6.44}$$

because $18000\bar\Lambda\nu \le \frac{1}{60}$ and $1.8 \cdot 10^5 \bar\Lambda u_\star / L \le \frac{1}{32}$. So (ii) follows from (i).

(iii) The reference gas of the step k is the hard-core gas of the groups made at the step k with the activities c_W, $|c_W| \le S_W$, and the compatibility of the groups. Put $a(W) := (\nu/64)\sigma(W)$ and $d(W) := \frac{1}{60} + (\nu/32)\sigma(W)$. A group which is incompatible with W has a point of its domain in common with X_W, by Definition 5.2(vii), so that it is at one of the at most $81 u_\star \sigma(W)$ cubes of π_k which belong to X_W or touch it. By (5.22) at the cubes of π_k,

$$\sum_{W' \not\sim W} |c_{W'}| e^{a(W') + d(W')} \le 81 u_\star \sigma(W) \, e^{1/60} \cdot 2 C_{S'} \rho_k \le a(W) ,$$

by the hypothesis of (iii). This is the condition of [15, Theorem 5.4] with the functions a and d, in the form in which Lemma 7.3 uses that theorem: the sum of $|\varphi_k(\Gamma)| \prod_{W' \in \Gamma} e^{d(W')}$ over the clusters which contain a group incompatible with an external polymer is at most the sum of $|c_{W'}| e^{a(W') + d(W')}$ over the groups incompatible with it. With a cube of π_k in the domain of a group of Γ as the external polymer this gives $|\varphi_k(\Gamma)| \le 2 e^{1/60} C_{S'} \rho_k \le 1$, so that $\sum_{n \ge 1} |\varphi_k(\Gamma)|^n / n! \le e |\varphi_k(\Gamma)|$. With the cube Δ as the external polymer, incompatible with the groups which are at Δ, and with (6.44) for each group of an occurrence, it gives

$$\sum_{(\Gamma, n) \text{ at } \Delta} w(\Gamma, n) e^{2b(\Gamma)} \le e \sum_{W' \text{ at } \Delta} |c_{W'}| e^{\frac{1}{60} + \frac{3}{64}\nu\sigma(W')} \le 2 e^{1 + 1/60} K_{\text{box}} C_{S'} \rho_k ,$$

by (i).

(iv) and (v). Let v be a polymer vertex. By (6.33) and the first two conditions of (6.43), as in (ii),

$$4\nu s(v) + 2b(v) \le 198\nu + \frac{1}{1600}\Big(\#(M) + \sum_{\mathcal{C} \in \mathcal{K}_v} N(\mathcal{C})\Big) + \sum_{W \in \mathfrak{B}_v} \Big(\frac{1}{60} + \frac{\nu}{32}\sigma(W)\Big) .$$

We first reduce the sum over the vertices at Δ to a sum over the vertices whose polymer contains a given cube. By Lemma 6.14(ii) a vertex at Δ has $9\bar\Lambda\hat d(v) \ge t-1$ for every cube Δ_0 of M, where tM is the distance of the centres of Δ_0 and Δ; then $\#(M)+\sum_{\mathcal C} N(\mathcal C) \ge (t-1)/(18\bar\Lambda)$ or $\sum_{W\in\mathfrak B_v}(u_\star\sigma(W)+1)/L \ge (t-1)/(18\bar\Lambda)$, and in both cases the number

$$g(v) \coloneqq \frac{1}{1600}\Big(\#(M)+\sum_{\mathcal C\in\mathcal K_v} N(\mathcal C)\Big)+\frac{\nu}{64}\sum_{W\in\mathfrak B_v}(\sigma(W)+1)$$

is at least $c_p(t-1)_+$. There are at most $8(2t+1)^3$ cubes Δ_0 for each t, and therefore

$$\sum_{v \text{ at } \Delta} w(v)e^{2b(v)} \le C_{\text{root}} \sup_{\Delta_0} \sum_{v\colon \Delta_0\subset M} w(v)\, e^{2b(v)+g(v)}\,. \tag{6.45}$$

In the sum on the right we sum first over $\mathcal K_v$ and $\mathfrak Y_v$. By the two displays and (6.36),

$$w(v)e^{2b(v)+g(v)} \le e^{198\nu}\,\mathfrak a_v\, e^{-(r_\kappa-c-\frac{1}{800})\#(M)}\, e^{-\kappa'\#(\mathfrak Y_v)} \prod_{\mathcal C\in\mathcal K_v}\xi(\mathcal C)e^{N(\mathcal C)/800}\prod_{W\in\mathfrak B_v}\omega(W)\,,$$

where $\omega(W) \coloneqq \|Y_W\|_{(k)}\exp(\frac{1}{60}+\frac{1}{64}\nu+\frac{3}{64}\nu\sigma(W))$. The sum over the sets $\mathcal K_v$ of clusters incompatible with M of the product of the numbers $\xi(\mathcal C)e^{N(\mathcal C)/800} \le 2e\,\bar\varphi(\mathcal C)e^{d(\mathcal C)+N(\mathcal C)}$ is at most $\exp(2e\,c\,\#(M))$, by Supplement, Lemma 17.24(iv). The sets $\mathfrak Y_v$ are determined by the sets of their cubes, nonempty subsets of the $L^4\#(M)$ cubes of π_k in M, and

$$\sum_{\mathfrak Y_v} e^{-\kappa'\#(\mathfrak Y_v)} \le (1+e^{-\kappa'})^{L^4\#(M)}-1 \le L^4e^{-\kappa'}e^{2\#(M)}\,,$$

by the last condition of (6.43); if M has several connected components and each of them is to contain a member of $\mathfrak Y_v$, the same bound holds with one factor $L^4e^{-\kappa'}$ for each component. Since $7c+\frac{1}{800}\le 1$, each connected component P of M is thus left with the factor

$$\omega(P) \coloneqq L^4e^{-\kappa'}e^{-(r_\kappa-3)\#(P)}\,,\qquad \sum_{P\ni\Delta_0}\omega(P)e^{2\#(P)} \le L^4e^{-\kappa'}\sum_{n\ge1}64^{n-1}e^{-(r_\kappa-5)n} \le L^4e^{-\kappa'}\,,$$

the sum over the connected unions $P\in\mathbf D_{k+1}$ of cubes which contain Δ_0, of which there are at most 64^{n-1} with n cubes (Supplement, proof of Lemma 17.24); the last step uses $r_\kappa\ge10$.

For (iv), M is connected and $\mathfrak a_v = c_a\rho_k$, and the three bounds give the claim with $e^{198\nu}\le 2$.

For (v), let v belong to the vertex set of a group, so that M is a marked polymer with $\mathfrak B(M)\ne\emptyset$. Consider the graph whose vertices are the connected components P of M and the members $W\in\mathfrak B(M)$, a component and a member being adjacent if a cube of the component meets the dependence set of the member. It is connected, by the proof of Lemma 6.13(i), since a component of a group which touches a member lies in a connected component of M. So the sum over these vertices v with $\Delta_0\subset M$ is at most $e^{198\nu}\varpi_k$ times the sum, over the connected sets of this kind which contain the component $P_0\ni\Delta_0$ and at least one member, of the products of the numbers ω. We bound it by Lemma 6.15(ii), applied to the graph of all the connected unions P of cubes and all the groups W of the term, with the weights ω, the charges $b'(P)\coloneqq\#(P)$ and $b'(W)\coloneqq(\nu/128)(\sigma(W)+1)$, and the marks $m(W)\coloneqq\omega(W)$, $m(P)\coloneqq0$. The condition (6.34) holds at a member W, because its dependence set meets at most $625u_\star(\sigma(W)+1)$ cubes, by Lemma 6.12 and $L\ge81$, and

$$\sum_{P\sim W}\omega(P)e^{2\#(P)} \le 625\,u_\star(\sigma(W)+1)L^4e^{-\kappa'} \le \tfrac12 b'(W)$$

by the last condition of (6.43); and at a component P, with $\epsilon := 2e^{1/50}K_{\rm box}\mathsf{E}_k \le 1$, because by (i)

$$\sum_{W\sim P}\omega(W)e^{2b'(W)} \le \#(P)\sup_{\Delta'}\sum_{W \text{ at } \Delta'}\|Y_W\|_{(k)}e^{\frac{1}{50}+\frac{1}{16}\nu\sigma(W)} \le \tfrac{1}{2}\epsilon\, b'(P)\ ,$$

where we used $\omega(W)e^{2b'(W)} = \|Y_W\|_{(k)}e^{\frac{1}{60}+\frac{1}{32}\nu+\frac{1}{16}\nu\sigma(W)}$ and $\nu/32 \le \frac{1}{50}-\frac{1}{60}$, and that a member adjacent to P is at a cube of P. The same display is the hypothesis on the marks. Lemma 6.15(ii) therefore bounds the sum over the connected sets by $\epsilon\,\#(P_0)e^{\#(P_0)/2} \le \epsilon\, e^{2\#(P_0)}$, and the sum over $P_0 \ni \Delta_0$ gives $\epsilon L^4 e^{-\kappa'}$. With (6.45) and $2e^{198\nu+1/50} \le 3$ this is (v). *q.e.d.*

Theorem 6.18 (The Size Bound)**.** *Suppose that the hypotheses of Proposition 6.17 hold, with those of its parts (iii) and (v), and put*

$$E_\alpha := C_\alpha\rho_k\ ,\qquad E_\beta := C_\beta\varpi_k\mathsf{E}_k\ ,\qquad E_o := e^{1/60}K_{\rm box}\mathsf{E}_k + 6K_{\rm box}C_{S'}\rho_k\ . \tag{6.46}$$

If $E_\alpha + E_\beta + E_o \le \frac{1}{2}\nu$, then the hypothesis of Proposition 6.11 holds at every real background of the support of the term, and for every cube Δ

$$\sum_{\mathcal{G}\colon \Delta\subset X_{\mathcal{G}}} e^{\frac{5}{2}\nu\sigma_{k+1}(\mathcal{G})+\nu_e\,\mathrm{ext}(\mathcal{G})}P(\mathcal{G})\,\ell_{\mathcal{G}} \le \frac{e^{2\nu}}{\nu}\Big[E_\alpha+E_\beta+\frac{2}{\nu}(E_\alpha+E_\beta+E_o)E_o+\frac{2}{\nu}(E_\beta+E_o)E_\alpha\Big]\ , \tag{6.47}$$

the sum over the groups of the step $k+1$, in all the terms, other than those which consist of a single member of a family and nothing else.

Proof. Let $E := E_\alpha + E_\beta + E_o$. The vertex set of a group consists of member vertices, of occurrence vertices, and of polymer vertices which have $\mathfrak{B}_v = \emptyset$ and a connected polymer, as was shown in the proof of Proposition 6.9 for a marked polymer without members, or which have $\mathfrak{B}_v \ne \emptyset$. Let $\mathcal{V}$ be the set of these vertices, with the weights and the relation of (6.37). A vertex $v' \sim v$ is at one of the cubes of $K(v)$, so that by Proposition 6.17

$$\sum_{v'\sim v} w(v')e^{2b(v')} \le |K(v)|\,E \le \tfrac{1}{2}b(v)\ ,$$

which is (6.34). Now let $\mathcal{G}$ be a group with $\Delta \subset X_{\mathcal{G}}$. By Lemma 6.13(iii) one of its vertices is at Δ, and we choose a root v_0 among these: a polymer vertex with $\mathfrak{B}_v \ne \emptyset$ if there is one, otherwise a member or an occurrence vertex if there is one, and otherwise a polymer vertex without members. By Lemma 6.16 the left side of (6.47) is at most $e^{2\nu}/\nu$ times the sum over the roots v_0 at Δ and over the connected sets $S \ni v_0$ which occur, of the product of the weights, and we bound this sum for the three kinds of roots. For the roots of the first kind Lemma 6.15(i) gives at most $\sum_{v_0} w(v_0)e^{b(v_0)/2} \le E_\beta$. For a root of the second kind the set S has a further element, since a group which consists of a single member is excluded and an old occurrence is tied to a member; we apply Lemma 6.15(ii) with the marks $m := w$ and $\epsilon := 2E/\nu$ and obtain at most $\sum_{v_0} w(v_0)\,\epsilon\, b(v_0)e^{b(v_0)/2} \le (2E/\nu)E_o$. For a root of the third kind, either all the elements of S are polymer vertices without members, and Lemma 6.15(i) gives at most E_α, or S contains a vertex of another kind, and Lemma 6.15(ii), with the marks $m := w$ on the vertices of the other kinds and $m := 0$ on the polymer vertices without members, for which $\epsilon = 2(E_\beta+E_o)/\nu$, gives at most $(2(E_\beta+E_o)/\nu)E_\alpha$. In each case we used $b\,e^{b/2} \le e^{2b}$ and Proposition 6.17 at Δ.

For the convergence, the terms t correspond to the sets of pairwise compatible groups (Proposition 6.11), and the product of the norms of the tensors and of the numerical factors of a term is the product of the numbers $P(\mathcal{G})$ of its groups, without the factors of the

transports. The sum over the terms is therefore at most the exponential of the sum of $P(\mathcal{G})$ over all the groups. The groups which consist of a single member contribute the sum of $\|Y_W\|_{(k)}$ over the groups of the term of ϱ_k, which is finite by (6.41), the torus having finitely many traversals, and the others contribute at most the right side of (6.47) times the number of the cubes of the torus. *q.e.d.*

The four terms of (6.47) have different origins, and the difference matters for the induction. The first is the contribution of the groups which the step makes out of nothing, and it is of the order of ρ_k with a constant which does not involve the constants of the class. The others are proportional to the sums of the step k, with coefficients which tend to zero with the coupling: ϖ_k is a negative power of $\log g_k^{-2}$, and E_α, E_β, E_o are small. This is the form which a recursion with a fixed point requires.

6.7 The Small-Field Map and the Flat Parts

The groups and the interactions. By Proposition 6.11 the expectation $\langle F \rangle$ is, on the terms without large-field regions, the sum of the words of the families of pairwise compatible groups of Definition 6.10. A group whose intervals lie on one loop has one interval: each of its constituents then has a single piece, and every identification of Definition 6.10 joins two sets of constituents whose arc sets meet or adds the arc between two pieces, so that the arc set stays an arc. A group with intervals on several loops is a cross group. The polymers which belong to no group form Bałaban's expansion of the effective action. The families of compatible groups are resummed below against the gas of the groups which they exclude, which gives the interactions $Q_{k+1}(\mathfrak{B})$. The identity $N_{k+1} = \sum \chi_{k+1}(\Omega_{k+1}) \mathbf{T}_{k+1}[\mathcal{W}_{k+1} \exp A_{k+1}]$ is Proposition 6.3, with $\mathcal{W}_{k+1} = \langle F \rangle$ at the steps at which no **R**-operation acts, and (5.2) holds at $k+1$ because the step preserves integrals.

The insertions as functions on the space of the step. So far the insertions of the groups of the step $k+1$ are functions of the block field V_{k+1}, through the background U_{k+1} and its current J_{k+1}. The class requires them as holomorphic and covariant functions on the space of the step, and both properties are inherited from the ingredients. In his own construction Bałaban obtains the variables as follows: "we replace J_{k+1} by the variable **J**, and in the remaining expressions we replace U_{k+1} by the new variable **U**" [8, p. 272], and for the operators "we replace the configurations U_{k+1}, J_{k+1} occurring in operators there by the configurations **U**, **J**" [8, p. 273]. We do the same in the tensors Y_G, in the marked activities, and in the cluster weights.

Lemma 6.19 (The New Insertions on the Space of the Step)**.** *On the terms without large-field regions of ϱ_k and of ϱ_{k+1}, under the hypotheses of Theorem 6.18, the insertion $Y_\mathcal{G}$ of every group $\mathcal{G}$ of the step $k+1$ is the value at (U_{k+1}, J_{k+1}) of a function of the restriction of $(\mathbf{U}, \mathbf{J})$ to $X_\mathcal{G}$ which is holomorphic on the union of the sets $\mathcal{S}_{k+1}(U)|_{X_\mathcal{G}}$ over the configurations U of the space of the step $k+1$, and which satisfies the covariance law* (5.19) *for each of its intervals.*

Proof. Analyticity. We go through the ingredients of $Y_\mathcal{G}$ in Definition 6.10. The propagators with the localisation parameters, the function $\mathbf{H}(s, B')$, and the quadratic forms and covariances of the fluctuation integral are "analytic functions of the configurations **U**, **J** in a domain given by the conditions I.(i)–(iii) with some α_0', α_1', and all the bounds above are uniform on the domain" [10, p. 6], with constants "much bigger than α_0, α_1" [10, p. 15]; the conditions (i)–(iii) are the pointwise ones [8, (1.11)–(1.14), p. 262], so that this domain contains the sets $\mathcal{S}_{k+1}(U)$ of (5.17). The potentials $\mathbf{V}_k(Y, \cdot)$ and the activities of the vacuum polymers are

analytic on the spaces $\mathbf{U}^c_{k+1}(X, \alpha_0, \alpha_1)$ [10, Lemma 2, p. 11; p. 15], whose definition contains the conditions on the averaged configurations, and which contain the sets $\mathcal{S}_{k+1}(U)|_X$ by the choice of the constant c_F (Definition 5.2(iii)). For a point of $\mathcal{S}_{k+1}(U)$, a fluctuation field of the support of the test functions, and parameters with $|s(\Delta)| \le e^{\kappa_1}$, the interpolated configuration at which the insertions of the touched members are evaluated lies in the space of the step k, by (6.6), and these insertions are holomorphic there (Definition 5.2(iii) at the step k); the transports are entire functions. The tensors Y_G are therefore holomorphic in $(\mathbf{U}, \mathbf{J})$, with the bounds of Lemmata 6.5 and 6.7, which do not depend on the point. A marked activity $\tilde{H}(M)$ is a finite sum of terms [10, (2.14), p. 15], each an integral over real variables and over contours in the parameters of a product of these functions, which Bałaban considers "as an analytic function of $(\mathbf{U}, \mathbf{J})$ in the space $\mathbf{U}^c_{k+1}(X, \alpha_0, \alpha_1)$, and of the complex parameters" [10, p. 15]; the bounds of the integrands in the proof of Supplement, Lemma 17.27, are uniform on the space, so that $\tilde{H}(M)$ is holomorphic. So are the cluster weights $\varphi^T(\mathcal{C})$, sums of products of holomorphic activities which converge uniformly by Supplement, Lemma 17.24(iv), and with them the factors u_λ. The insertions of the untouched members are evaluated at $(\mathbf{U}, \mathbf{J})$ itself, a point of the space of the step k (Section 6.3). Finally $Y_{\mathcal{G}}$ is a contraction of finitely many of these functions with transports and numbers, and it depends on $(\mathbf{U}, \mathbf{J})$ only on $X_{\mathcal{G}}$ by Lemma 6.13(v).

Covariance. Let v be a G-valued gauge transformation. The operators of the step transform by conjugation with $R(v)$: $\Delta^\eta(U^v) = R(v)\Delta^\eta(U)R(v^{-1})$ and likewise for G', R, Δ_a, and G, and the averaging operators commute with $R(v)$ [5, (3.30)–(3.34), pp. 395–396]. The localisation parameters multiply the terms of the random walk expansions of the propagators [10, p. 3], which are products of such operators and of scalar functions, so that the propagators with parameters transform in the same way; this is what Bałaban uses when he states that, "by Lemma 2, and the transformation properties of the operators in (2.14) with respect to gauge transformations, e.g. see (3.28)–(3.34) [13], the expressions (2.14) are gauge invariant with respect to all G-valued transformations" [10, p. 21]. The function $\mathbf{H}(s, B')$ is the unique solution, in balls of norms which are invariant under $R(v)$, of the equations [10, (1.3), (1.4), p. 2], which are built from these propagators and from local functions that commute with the gauge transformations; hence $\mathbf{H}(s; \mathbf{U}^v, R(v)\mathbf{J}, R(v)B') = R(v)\mathbf{H}(s; \mathbf{U}, \mathbf{J}, B')$, and B' is transformed with the fluctuation field, as in the simultaneous transformations [8, (3.29), p. 276]. The interpolated configuration at the transformed data is therefore the gauge transform by v of the interpolated configuration, and the argument which takes the place of $\mathbf{J}$ in the insertions of the members, the current of that configuration [8, (3.10), p. 272], is transformed by $R(v)$. By (5.19) at the step k for the members and by the same law for the transports, each factor of a tensor Y_G at the data $(\mathbf{U}^v, R(v)\mathbf{J}, R(v)B)$ is the factor at $(\mathbf{U}, \mathbf{J}, B)$ conjugated by $\pi(v)$ at the two ends of its piece. In the integrals which define $\tilde{H}(M)$ at $(\mathbf{U}^v, R(v)\mathbf{J})$ we substitute $R(v)B$ and $R(v)X$ for the integration variables; $R(v)$ is orthogonal, the Gaussian densities and the potentials are invariant under the simultaneous transformations, the latter by [10, Lemma 2, p. 11], and the test functions depend on the numbers $|B(b)|$. Hence $\tilde{H}(M)$ transforms as its factor does, piece by piece. The activities of the vacuum polymers are invariant [10, p. 21], and so are the numbers $\Psi_\circ(M)$, the weights $y_{\mathcal{C}}$, and the factors u_λ. The untouched members and the transports of the background satisfy the law by hypothesis, and a contraction along an interval of factors which satisfy it at the ends of their pieces satisfies it at the ends of the interval. This is (5.19) for $Y_{\mathcal{G}}$. *q.e.d.*

The flat parts. Every group of the step has a *flat value*, its insertion at the reference data of the term, the trivial background with the labels of the term. By gauge covariance under constant gauge transformations, which fix the trivial background, the flat value of a

group on one loop commutes with $\pi_i(G)$, and by Schur's lemma it is a multiple $c_W\,\mathrm{id}$ of the identity. If the group is not attached to a large-field region and contains no part of an attached insertion, the insertion is split as $Y_W = c_W\,\pi_i(U_{k+1}(I_W)) + Y_W^{\mathrm{d}}$, where I_W is the interval of the group and $\pi_i(U_{k+1}(I_W))$ the transport of the background along it. A word with the first part at I_W is c_W times the word without the group, since that part restores the transport which the group had replaced; the scalars c_W form the reference gas of the step, and the second part Y_W^{d} vanishes at flat data and is a deviation. An attached group is not split: its insertion carries no deviation bound, with or without the subtraction, and it is kept whole (Section 7.1). Section 7 resums the reference gas along each loop and extracts from it, exactly, the step factor κ_{k+1} and the remainder E°_{k+1}, which joins $\mathcal{E}_{k+1}$. A cross group carries no flat value. Its insertion at the reference data is invariant under the simultaneous conjugation of its factors by $\pi_{i(I)}(u)$, $u \in G$, and not under the conjugation of a single factor, so that it is not a scalar in general and need not vanish; nothing is subtracted from it, and it is kept whole as an insertion of the cross species (Section 6.16).

The production bound. The class records, for the groups made at a step, a bound on their sizes on the whole space of that step, (5.22). For the step $k+1$ it is a consequence of Theorem 6.18, in which the old sums enter only with coefficients that vanish with the coupling; this is what allows the constant to be the same at every step.

Corollary 6.20 (The Production Bound)**.** *In the setting of Theorem 6.18, let $\mathcal{G}$ be a group made at the step $k+1$, that is, a group of the step $k+1$ which does not consist of a single member. Let $c_{\mathcal{G}}$ be its flat value if its insertion is split and 0 otherwise, $Y_{\mathcal{G}}^{\mathrm{d}}$ its insertion after the subtraction, and $S_{\mathcal{G}} := \|Y_{\mathcal{G}}^{\mathrm{d}}\|_{(k+1)} + |c_{\mathcal{G}}|$. Then $S_{\mathcal{G}} \le 3e^{\nu\sigma_{k+1}(\mathcal{G})}P(\mathcal{G})$, and for all numbers $0 \le \theta(\mathcal{G}) \le \nu\sigma_{k+1}(\mathcal{G})/16$, every cube p of π_{k+1}, and every traversal p of the step $k+1$,*

$$\sum_{\mathcal{G}\ at\ p} e^{\theta(\mathcal{G})+\nu\sigma_{k+1}(\mathcal{G})+\nu_e\,\mathrm{ext}(\mathcal{G})} S_{\mathcal{G}}\,\ell_{\mathcal{G}} \le \frac{3e^{2\nu}}{\nu}\Big[E_\alpha + E_\beta + \frac{2}{\nu}(E_\alpha + E_\beta + E_o)E_o + \frac{2}{\nu}(E_\beta + E_o)E_\alpha\Big]\,, \tag{6.48}$$

the sum over the groups made at the step $k+1$ which are at p: those with the cube p in $X_{\mathcal{G}}$, or with the traversal p in an interval; and the same bound holds with the further factor $e^{\frac{7}{16}\nu\sigma_{k+1}(\mathcal{G})}$ in each term of the sum. If moreover $\mathsf{E}_k \le C_{\mathsf{E}}\rho_k$ with a constant C_{E}, and

$$C_{S'} := \max\Big(\frac{3e^{2\nu}}{\nu}(1+\beta_0)^6 C_\alpha,\ \tfrac{1}{2}L(2K_\chi+1)e^{2\nu}C_{\mathrm{pin}}(1+\beta_0)^6\Big)\,, \tag{6.49}$$

the second number being the constant which the first step requires (Corollary 5.8), then the right side of (6.48) *is at most $2C_{S'}\rho_{k+1}$ for γ_q sufficiently small, depending on C_{E} and on the constants of Proposition 6.17.*

Proof. The number $c_{\mathcal{G}}$ is the value of $Y_{\mathcal{G}}$ at the reference data, which belong to the space of the step $k+1$, so that $|c_{\mathcal{G}}| \le P(\mathcal{G})$ by (6.39). The transport along the interval of $\mathcal{G}$ has norm at most $e^{c'_\pi\alpha_{1,k+1}n_{k+1}(\mathcal{G})} \le e^{\nu\sigma_{k+1}(\mathcal{G})}$ on that space, by the hypothesis $4\bar{\Lambda}Mc'_\pi\alpha_{1,k+1} \le \nu$ of Lemma 6.16 and $n_{k+1}(\mathcal{G}) \le M\sigma_{k+1}(\mathcal{G})$. Hence $\|Y_{\mathcal{G}}^{\mathrm{d}}\|_{(k+1)} \le (1+e^{\nu\sigma_{k+1}(\mathcal{G})})P(\mathcal{G})$ and $S_{\mathcal{G}} \le 3e^{\nu\sigma_{k+1}(\mathcal{G})}P(\mathcal{G})$. The exponent on the left of (6.48) is then at most $(\frac{1}{16}+2)\nu\sigma_{k+1}(\mathcal{G})+\nu_e\,\mathrm{ext}(\mathcal{G})$, which is less than the exponent of (6.47) by $\frac{7}{16}\nu\sigma_{k+1}(\mathcal{G})$, and a traversal in an interval of $\mathcal{G}$ meets a cube of $X_{\mathcal{G}}$; so (6.48), with or without the further factor, follows from Theorem 6.18.

For the last statement, $\rho_k \le (1+\beta_0)^6\rho_{k+1}$ by (6.55), so that the first term on the right of (6.48) is $(3e^{2\nu}/\nu)C_\alpha\rho_k \le C_{S'}\rho_{k+1}$. By (6.46), $E_\beta \le C_\beta C_{\mathsf{E}}\varpi_k\rho_k$ and $E_o \le K_{\mathrm{box}}(e^{1/60}C_{\mathsf{E}} + 6C_{S'})\rho_k$, so that each of the other terms is at most ρ_k times a constant times ϖ_k or ρ_k/ν. Both tend to zero with γ_q, uniformly in $k \le K'$: $\rho_k \le C_\rho\gamma_q^2(1+A_1(\log\gamma_q^{-2})^{p_0})^4$ because $g_k \le \gamma_q$ by

(3.46), and ϖ_k is at most a constant times $\vartheta_k/\min(\alpha_{0,k},\alpha_{1,k})$, a negative power of $\log \hat g_k^{-2}$ as in (6.6). The other terms together are therefore at most $C_{S'}\rho_{k+1}$ for γ_q small. *q.e.d.*

By (6.48) with $\theta = 0$ the activities $c_{\mathcal G}$ of the reference gas of the step $k+1$ satisfy the hypothesis of Lemma 7.3, whose proof verifies the condition of Kotecký and Preiss with the weight $a^\circ(\mathcal G) = (\nu/32)\sigma_{k+1}(\mathcal G)$; the theorem cited there bounds the sum of the absolute values of the weights of the clusters which are incompatible with a group $\mathcal G$ by $a^\circ(\mathcal G)$. With $\theta(\mathcal G)$ defined as that sum we have $\theta(\mathcal G) \le \nu\sigma_{k+1}(\mathcal G)/16$, which is clause (i) of Definition 5.2 at the step $k+1$ for the groups made there, and (6.48) with these numbers is the production bound (5.22) at the step $k+1$. The definition of θ uses the bound with $\theta = 0$ only, so that the argument is not circular.

6.8 The Deviation Bound of the New Groups

The production bound controls the size of a new group, and the class asks for more: at a real background the deviation part of an insertion has to be small in proportion to the curvature of the background near the group, as in (5.13). That it is so comes from the subtraction of the flat value. The deviation part $Y^{\mathrm d}_{\mathcal G}$ is a holomorphic function on the space of the step $k+1$ which vanishes at the trivial configuration, and near the group the background is, in a suitable gauge, the exponential of a field which is bounded by the local curvature times the linear size of the region (Supplement, Lemma 17.22). On the complex line through the trivial configuration and the background in that gauge, Schwarz's lemma converts the bound of the function on the space into a bound proportional to that field.

We keep the setting of Section 6.6, with the units of the scale $k+1$, in which the fine lattice has the spacing $\xi = L^{-(k+1)}$. Let $\mathcal G$ be a group made at the step $k+1$ with one interval, from the site $x_{\mathcal G}$ to the site $x'_{\mathcal G}$, let $\mathfrak n$ be a traversal of the step $k+1$ in that interval, $t = t_{\mathfrak n}(\mathcal G)$, and let $\mathcal Q = \mathcal Q_{\mathfrak n}(\mathcal G)$ be the smallest cube which is a union of cubes of π_{k+1} and contains the ball $B_\infty(\mathfrak n, R_t)$, hence $X_{\mathcal G}$. Let $\mathcal Q^{\sim 5}$ be the cube which is obtained from $\mathcal Q$ by adding five layers of cubes. The group $\mathcal G$ is of the deviation species if it embeds in the torus (Section 5.3); the cube $\mathcal Q^{\sim 5}$ then lies in the comparison region of $\mathcal G$ at $\mathfrak n$, as the proof below shows, and is therefore the injective image of a cube of the universal cover of the torus. The other groups with one interval are those of the wrapping species (Section 6.16). Put

$$C_g \coloneqq 3B_3(2C_{\mathrm{box}}+5)M\ , \qquad c_{\mathrm{an}} \coloneqq \min\Big(\frac34,\ \frac{c_F}{2c_G}\Big)\ , \qquad \bar\alpha_{k+1} \coloneqq \min(\alpha_{0,k+1},\alpha_{1,k+1})\ , \tag{6.50}$$

with the constant B_3 of [7, Theorem 1, p. 279], the constant c_F of (5.17), and the constant c_G of (6.63).

Proposition 6.21 (The Deviation Bound of a New Group)**.** *Let the standing hypotheses of Supplement, Lemma 17.22, and the hypotheses of Theorem 6.18 hold, let γ be sufficiently small, and let V_{k+1} be a block field in the support of the term. Then, for every group $\mathcal G$ of the deviation species made at the step $k+1$ and every traversal $\mathfrak n$ of the step $k+1$ in its interval,*

$$\|Y^{\mathrm d}_{\mathcal G}(U_{k+1}, J_{k+1})\| \le \frac{C_g}{c_{\mathrm{an}}\bar\alpha_{k+1}}\, S_{\mathcal G}\, \mathrm{Dev}_{\mathfrak n}(\mathcal G; U_{k+1})\ , \tag{6.51}$$

so that (5.13) *holds at the step $k+1$ with $b_{\mathcal G} \coloneqq (C_g/(c_{\mathrm{an}}\bar\alpha_{k+1}))e^{\theta(\mathcal G)}S_{\mathcal G}$. Moreover, under the hypotheses of Corollary 6.20, with $\beta_{\mathcal G} = b_{\mathcal G}\ell_{\mathcal G}L^{2(k+1)}$, for every traversal $\mathfrak n$ of the step $k+1$ and every $t \ge 1$,*

$$\sum_{\mathcal G\ni\mathfrak n,\ t_{\mathfrak n}(\mathcal G)\ge t} e^{\nu\sigma_{k+1}(\mathcal G)+\nu_e\,\mathrm{ext}(\mathcal G)}\beta_{\mathcal G} \le \frac{4C_gC_{S'}}{c_{\mathrm{an}}}\,\tilde\rho_{k+1}L^{2(k+1)}W_t\ , \tag{6.52}$$

the sum over the groups of the deviation species made at the step $k+1$ with $\mathfrak{n}$ in the interval.

Proof. The region. The cube $\mathcal{Q}$ has side at most $2R_t + 1 + 2M$, and $\mathcal{Q}^{\sim 5}$ lies in the ball $B_\infty(\mathfrak{n}, R_t + 6M)$, hence in the comparison region $B_\infty(\mathfrak{n}, R_{t+1})$ of $\mathcal{G}$ at $\mathfrak{n}$, because $R_{t+1} - R_t = C_{\rm box} M \geq 6M$. Let E be the largest value of $\xi^{-2}|U_{k+1}(\partial p) - 1|$ over the fine plaquettes p in $\mathcal{Q}^{\sim 5}$, as in Supplement, Lemma 17.22. Since $\xi^{-2} = L^{2(k+1)}$, the definition (5.8) gives $\ell_{\mathcal{G}} E \leq \mathrm{Dev}_{\mathfrak{n}}(\mathcal{G}; U_{k+1})$. The domain $X_{\mathcal{G}}$ is a connected union of $\ell_{\mathcal{G}}$ cubes of side M, one of which meets $\mathfrak{n}$ (Lemma 6.13(iv)), so that it lies in the ball of radius $M\ell_{\mathcal{G}} + 1$ about $\mathfrak{n}$, and therefore $R_t \leq \max(2C_{\rm box}M, M\ell_{\mathcal{G}} + 1 + C_{\rm box}M) \leq 2C_{\rm box}M\ell_{\mathcal{G}}$ by the definition of t. The number $M_{\mathcal{Q}}$ of that lemma is half the side of $\mathcal{Q}^{\sim 2}$, rounded up to a multiple of a number which is at most M; hence $M_{\mathcal{Q}} \leq R_t + 5M \leq (2C_{\rm box} + 5)M\ell_{\mathcal{G}}$, and

$$x := 3B_3 M_{\mathcal{Q}} E \leq C_g \,\mathrm{Dev}_{\mathfrak{n}}(\mathcal{G}; U_{k+1}) \,. \tag{6.53}$$

The trivial case. If $x \geq c_{\rm an}\bar{\alpha}_{k+1}$, the right side of (6.51) is at least $S_{\mathcal{G}}$, which bounds the left side, since (U_{k+1}, J_{k+1}) is a point of the space of the step $k+1$. We assume from now on that $x < c_{\rm an}\bar{\alpha}_{k+1}$. Then $2E \leq 2x/(3B_3M)$ and $2M_{\mathcal{Q}}E \leq 2x/(3B_3)$ are less than $\bar{\alpha}_{k+1}$, which tends to zero with γ, so that for γ small the two hypotheses of Supplement, Lemma 17.22, on E and on $M_{\mathcal{Q}}E$ hold, their right sides being constants; here we used $M_{\mathcal{Q}} \geq M$.

The gauge and the function. By Supplement, Lemma 17.22, applied to the cube $\mathcal{Q}$, which is embedded in the torus with its five layers because $\mathcal{Q}^{\sim 5}$ lies in the comparison region and $\mathcal{G}$ embeds, there is a G-valued gauge transformation u on a neighbourhood of $\mathcal{Q}^{\sim 2}$ such that $U^u_{k+1} = e^{i\xi A}$ on $\mathcal{Q}^{\sim 2}$ with $|A|, |\nabla^\xi A| \leq x$ and $|R(u)J_{k+1}| \leq \frac{4}{3}x$. For a complex number z with $|z|x < c_{\rm an}\bar{\alpha}_{k+1}$ the pair $(e^{iz\xi A}, zR(u)J_{k+1})$ satisfies the conditions (5.17) at $n = k+1$ around the trivial configuration, on the bonds of $\mathcal{Q}^{\sim 2}$. Indeed, in the units of the fine lattice its complex part is $\alpha = z\xi A$, of norm at most $|z|x\,\xi < \alpha_{1,k+1}L^{-(k+1)}$; its covariant differences at the trivial configuration are its differences, of norm at most $|z|x\,\xi^2 < \alpha_{1,k+1}L^{-2(k+1)}$; by (6.63) with $U = 1$, and since $|z|x \leq 1$, its curvature is at most $c_G\xi^2(|z|x + (|z|x)^2) \leq 2c_G|z|x\,\xi^2 < c_F\alpha_{0,k+1}L^{-2(k+1)}$; and the current has norm at most $\frac{4}{3}|z|x < \alpha_{0,k+1}$ in the units of the scale $k+1$. By Lemma 6.19 the function

$$f(z) := Y^{\rm d}_{\mathcal{G}}(e^{iz\xi A}, zR(u)J_{k+1}) = Y_{\mathcal{G}}(e^{iz\xi A}, zR(u)J_{k+1}) - c_{\mathcal{G}}\,\pi(e^{iz\xi A}(I_{\mathcal{G}}))$$

is therefore holomorphic on the disc of radius $R_X := c_{\rm an}\bar{\alpha}_{k+1}/x > 1$, where $\|f(z)\| \leq \|Y^{\rm d}_{\mathcal{G}}\|_{(k+1)} \leq S_{\mathcal{G}}$; here we used that $Y_{\mathcal{G}}$ and the transport along $I_{\mathcal{G}}$ depend on the configuration only on $X_{\mathcal{G}} \subset \mathcal{Q}$ (Lemma 6.13(v)). At $z = 0$ the configuration is the trivial one with the current 0, and $f(0) = 0$ by the definition of the flat value $c_{\mathcal{G}}$. At $z = 1$ it is the gauge transform of (U_{k+1}, J_{k+1}) by u on $X_{\mathcal{G}}$, and by (5.19), which the transport along $I_{\mathcal{G}}$ satisfies as well,

$$f(1) = \pi(u(x_{\mathcal{G}}))\, Y^{\rm d}_{\mathcal{G}}(U_{k+1}, J_{k+1})\, \pi(u(x'_{\mathcal{G}}))^{-1} \,,$$

with unitary factors. Schwarz's lemma, applied to the matrix elements of f, gives $\|Y^{\rm d}_{\mathcal{G}}(U_{k+1}, J_{k+1})\| = \|f(1)\| \leq S_{\mathcal{G}}/R_X = S_{\mathcal{G}}\, x/(c_{\rm an}\bar{\alpha}_{k+1})$, and (6.51) follows from (6.53).

The tails. By the definitions, $e^{\nu\sigma_{k+1}(\mathcal{G}) + \nu_e \,\mathrm{ext}(\mathcal{G})}\beta_{\mathcal{G}}$ is $(C_g/(c_{\rm an}\bar{\alpha}_{k+1}))L^{2(k+1)}$ times the term of $\mathcal{G}$ in (6.48). If $t_{\mathfrak{n}}(\mathcal{G}) \geq t \geq 2$, then $X_{\mathcal{G}}$ is not contained in the ball of radius $R_{t-1} = C_{\rm box}Mt$ about $\mathfrak{n}$, so that $M\ell_{\mathcal{G}} + 1 > C_{\rm box}Mt$ and $\sigma_{k+1}(\mathcal{G}) \geq \ell_{\mathcal{G}} \geq C_{\rm box}t - 1$. The further factor $e^{\frac{7}{16}\nu\sigma_{k+1}(\mathcal{G})}$ of Corollary 6.20 is then at least $e^{-\nu}e^{\eta_w t}$, because the rate of the weights (5.9) is $\eta_w \leq \frac{5}{16}\nu \leq \frac{7}{16}\nu C_{\rm box}$. So the sum of the terms of (6.48) over the groups with $t_{\mathfrak{n}}(\mathcal{G}) \geq t$ is at most $2e^{\nu}C_{S'}\rho_{k+1}e^{-\eta_w t} \leq 2e^{\nu}C_{S'}\rho_{k+1}W_t$, since $W_t \geq w_t \geq e^{-\eta_w t}$; and for $t = 1$ the sum over all the groups is at most $2C_{S'}\rho_{k+1} \leq 2C_{S'}\rho_{k+1}W_1$, since $W_1 \geq w_1 = 8e^{-\eta_w} \geq 1$. With $2e^{\nu} \leq 4$ and $\tilde{\rho}_{k+1} = \rho_{k+1}/\bar{\alpha}_{k+1}$ this is (6.52). *q.e.d.*

The factor $1/\bar{\alpha}_{k+1}$ in (6.51) is the price of Schwarz's lemma: the function is bounded on a space whose radii are $\alpha_{0,k+1}$ and $\alpha_{1,k+1}$, and it is evaluated at a point whose distance from the centre is of the order of the deviation. This is why the first term of (5.11) has the coefficient $\tilde{\rho}_{k+1} = \rho_{k+1}/\bar{\alpha}_{k+1}$, and it is the reason for the two radii having exponents $q_0 = q_1$ larger than that of the thresholds: the deviation of a background of the support is of the order of ε_{k+1}, and $\varepsilon_{k+1}/\bar{\alpha}_{k+1}$ tends to zero with the coupling.

6.9 The Contraction of the Deviation Parts

It remains to carry the groups of the step k which the step leaves alone. A member of a family which no component touches and which is tied to nothing forms a group of the step $k+1$ by itself, and its insertion is the old function, evaluated at the new background U_{k+1} (Proposition 6.11). Nothing has to be estimated anew for it. The new background is a background of the step k, so that the bounds of the class hold at it as they stand, and what changes is only the unit in which the traversals are counted: a traversal of the step $k+1$ consists of L traversals of the step k, each with its carried groups, while the deviation of the background in the units of the scale $k+1$ is L^2 times the same deviation in the units of the scale k. This is the contraction by $L \cdot L^{-2} = L^{-1}$ which was announced at the beginning of the section. The parts of an old insertion which the fluctuation of the step changes do not appear here: an insertion which a component touches lies in a marked polymer and thereby in a group made at the step $k+1$, whose deviation part was bounded in Proposition 6.21.

Proposition 6.22 (The Carried Groups at the Real Backgrounds)**.** *Let the ensemble be dressed at the step k, let γ be so small that Supplement, Proposition S.8, applies, and consider the terms without large-field regions of ϱ_k and of ϱ_{k+1}, with a block field V_{k+1} in the support of the second.*

(i) For every group W of the deviation species of the step k, made at a step $j \le k$, and every traversal $\mathfrak{n}$ of the step j in its interval, $e^{\theta(W)}\|Y_W(U_{k+1}, J_{k+1})\| \le b_W \operatorname{Dev}_{\mathfrak{n}}(W; U_{k+1})$, and Y_W vanishes at the reference data.

(ii) If (5.14) holds at the step k with the constant $C_\chi^{(k)}$, then for every traversal $\mathfrak{n}'$ of the step $k+1$ and every $t \ge 1$

$$\sum_{W \ni \mathfrak{n}',\ t_{\mathfrak{n}'}^{(k+1)}(W) \ge t} e^{\nu\sigma(W) + \nu_e \operatorname{ext}(W)} \beta_W \le L^{-1} \frac{\tilde{\rho}_k}{\tilde{\rho}_{k+1}} C_\chi^{(k)} \tilde{\rho}_{k+1} L^{2(k+1)} W_t \,, \tag{6.54}$$

the sum over the groups of the deviation species of the step k with $\mathfrak{n}'$ in an interval.

Proof. (i) The block field $V_k^* := M^k(U_{k+1})$ lies in the support of the term without large-field regions of ϱ_k, by Supplement, Proposition S.8, and $U_k(V_k^*) = U_{k+1}$ by the uniqueness of the minimal orbit (Section 6.1). The bound (5.13) of the step k holds at every block field of the support, in particular at V_k^*, where the background is U_{k+1} and the argument of the insertion in the place of $\mathbf{J}$ is the current of that background, which is J_{k+1}; the insertion is the same function at both steps (Definition 5.2(iii)). The insertions of the deviation species vanish at the reference data at the step at which they are made, by the subtraction of the flat value, and the reference data of the two steps are the same configuration.

(ii) A group W of the sum has a traversal $\mathfrak{n} \subset \mathfrak{n}'$ of the step k in its interval, one of the L traversals of the step k in $\mathfrak{n}'$, and $t_{\mathfrak{n}}^{(k)}(W) \ge t_{\mathfrak{n}'}^{(k+1)}(W)$ (Section 5.3). The left side is therefore at most the sum over these L traversals of the left side of (5.14) at the step k, that is, at most $L \cdot C_\chi^{(k)} \tilde{\rho}_k L^{2k} W_t$, which is the right side of (6.54). *q.e.d.*

The ratio of the amplitudes of the two steps is controlled by Lemma 3.2: $g_k \le (1+\beta_0) g_{k+1}$ by (3.9), $(\log g_k^{-2})^p \le (1+\beta_0)(\log g_{k+1}^{-2})^p$ by (3.10), and $\alpha_{i,k+1} \le (1+\beta_0)^2 \alpha_{i,k}$ by (3.18), so that

$$\frac{\rho_k}{\rho_{k+1}} \le (1+\beta_0)^6 , \qquad \frac{\tilde\rho_k}{\tilde\rho_{k+1}} \le (1+\beta_0)^8 . \tag{6.55}$$

For $k = 1$ the second ratio is $(\rho_1/\rho_2)\min(\alpha_{0,2}, \alpha_{1,2})$, since $\tilde\rho_1 = \rho_1$, and it is at most $(1+\beta_0)^6$.

The recursion. At the step $k+1$ the groups of the deviation species on the term without large-field regions are the carried ones and those made at the step, and both satisfy (5.13), by Propositions 6.22(i) and 6.21. Adding (6.54) and (6.52), we see that the tail clause (5.14) holds at the step $k+1$ with a constant which satisfies

$$C_\chi^{(k+1)} \le C_{\text{new}} + a C_\chi^{(k)} , \qquad C_{\text{new}} := \frac{4 C_g C_{S'}}{c_{\text{an}}} , \qquad a := L^{-1}(1+\beta_0)^8 , \tag{6.56}$$

by (6.55). No term of the second order in the old constants occurs: the contractions of the old insertions with each other and with the fluctuation field are parts of the groups made at the step, and they are contained in C_{new}, which by (6.49) involves no constant of the class. At the steps at which the **R**-operation acts, C_{new} also contains the constant $C^{\mathbf{R}}(g_k) \le C^{\mathbf{R}}(\gamma_q)$ of the tail clause of the groups of the deviation species which that operation makes (Section 6.15); the coefficient a is the same at these steps, because the operation bounds its groups as a whole, with the insertions which they take in, and leaves the other insertions as they are. Since $(1+\beta_0)^8 \le L/4$ (Appendix A), we have $a \le \frac12$ at every step. Let $C^* := \max(C_\chi^{(1)}, 2C_{\text{new}})$, with the constant $C_\chi^{(1)}$ of the first step. If $C_\chi^{(k)} \le C^*$, then

$$C_\chi^{(k+1)} \le C_{\text{new}} + a C^* \le \tfrac12 C^* + \tfrac12 C^* = C^* . \tag{6.57}$$

So the tail clause holds at every step with the constant $C_\chi := C^*$, and by Lemma 5.3 so does the first term of (5.11), with its weights. The factor $\tilde\rho_k = \rho_k / \min(\alpha_{0,k}, \alpha_{1,k})$ in these bounds is the price of Schwarz's lemma in Proposition 6.21. The supremum form (5.21) of the same groups at the step $k+1$ does not follow from a recursion of this kind, as we now explain.

6.10 The Carried Insertions at Complex Data

The clause (5.21) of Definition 5.2 bounds the insertions on the whole space of the step, and not only at its real backgrounds. For the groups which a step makes this is part of what the count of that step delivers, in the form (5.22). For a group W which was made at an earlier step j and is carried to the step $k+1$ the question is of another kind, since its insertion is the function which was bounded at the step j, and it has now to be bounded on the space of the step $k+1$ and summed over the groups at a traversal of that step. The obvious bound fails. In the units of the fine lattice the space of the step $k+1$ lies in that of the step j, so that the supremum $\|Y_W\|_{(j)}$ of the step j bounds the insertion; but a traversal of the step $k+1$ consists of L^m traversals of the step j, $m := k+1-j$, each with carried groups of its own, and the sum of these suprema is larger than the bound of the step j by the factor L^m. At the real backgrounds this factor is compensated by the contraction of the deviation (Section 6.9). At complex data there is no deviation to contract, and the compensation has to come from the size of the complex part: the radii of the space of the step $k+1$ are smaller than those of the step j by the factor L^{-m}, up to the ratio of the couplings, so that Schwarz's lemma on the larger space gains L^{-m} for each insertion. This cancels the number of the traversals and no more, and the sum over the generations $m = 1, \dots, k$ would still grow with k.

What removes the growth is gauge covariance at complex gauge transformations. The complex part of a configuration of the space of the step $k+1$ varies slowly, by the condition on its covariant differences, and on a set which is small in the units of the scale $k+1$ it is therefore, to first order, a complex gauge transformation with a gauge function linear in the position. The dependence set of a carried group is such a set, of diameter δ_W of the order of $ML^{-m}\sigma(W)$ in these units. Removing the linear part by a gauge transformation which is the identity at the first site of the interval of W leaves a complex part smaller by the factor δ_W, and Schwarz's lemma then gains $L^{-2m}\sigma(W)$, which is summable over the generations; the factor $\sigma(W)$, and the groups whose dependence sets are not small, are paid by a sixteenth of the weight $e^{\nu\sigma(W)}$. This is the reason for the rate $\lambda_c < 1$ in (5.21). The insertions are covariant under gauge transformations with values in the compact group only, and the first step is to see that the covariance extends.

Throughout this subsection we work on the labelled term without large-field regions, we fix steps $1 \le j \le k$ and put $m \coloneqq k+1-j$, and we measure lengths in the units of the fine lattice, so that a traversal of the step n consists of L^n fine bonds and a cube of π_n has the side ML^n. We write $\mathfrak{g}$ for the Lie algebra of G, $\mathfrak{g}^c \coloneqq \mathfrak{g} \otimes \mathbb{C}$, and G^c for the complexification of G, the connected complex Lie group $G\exp(i\mathfrak{g})$ of matrices with the Lie algebra $\mathfrak{g}^c$; a representation π of G extends to a holomorphic representation of G^c. The norm $|\cdot|$ is that of Definition 5.2, on the matrices of the size of G, and it is invariant under the conjugation by the elements of G. The sets $\mathcal{S}_n(U)$ are those of (5.17); each of them is open in the space of the pairs of a G^c-valued configuration and a $\mathfrak{g}^c$-valued field, because $\alpha \mapsto e^{i\alpha}U(b)$ maps a neighbourhood of 0 in $\mathfrak{g}^c$ biholomorphically onto a neighbourhood of $U(b)$ in G^c, and a gauge transformation v with values in G maps $\mathcal{S}_n(U)|_X$ onto $\mathcal{S}_n(U^v)|_X$, by the invariance of the norm.

What is used of the class. Let W be a group of the deviation species made at the step j; it has one interval I_W, from a site x_W to a site x'_W, which lies in its domain X_W. We use the following.

(a) By Definition 5.2(iii) the insertion Y_W is holomorphic on the open set $\mathcal{D}_W$ of (5.18), it is bounded by $\|Y_W\|_{(j)}$ there, it satisfies the covariance law (5.19), and at the later steps it is the same function of the configuration and of the current.

(b) For V_{k+1} in the support of the step $k+1$ the background U_{k+1} is a background of the step j: with $V_i^* \coloneqq M^i(U_{k+1})$ for $j \le i \le k$, the block field V_i^* lies in the support of the step i (Supplement, Proposition S.8, by descending induction on i), and $U_i(V_i^*) = U_{k+1}$ by the uniqueness of the minimal orbit (Section 6.1). Moreover $|U_{k+1}(\partial p) - 1| < 2B_3\varepsilon_{k+1}L^{-2(k+1)}$ for every fine plaquette p [9, p. 268], and for γ small the pair of U_{k+1} and its current J_{k+1} is a point of $\mathcal{S}_{k+1}(U_{k+1})$.

(c) The domain X_W is a connected union of at most $u_\star\sigma(W)$ cubes of π_j, and I_W has at most $u_\star M\sigma(W)L^j$ fine bonds (Definition 5.2(vii)).

(d) The radii of two steps $n' < n$ satisfy $\alpha_{i,n} \le \varkappa_{n-n'}\,\alpha_{i,n'}$ for $i = 0, 1$, with $\varkappa_m \coloneqq (1+\beta_0)^2 m^{1/2}$, by (3.18), and $\alpha_{1,n} = \lambda_{\mathrm{r}}\alpha_{0,n}$ with $\lambda_{\mathrm{r}} \coloneqq C_1^\alpha / C_0^\alpha$, since $q_1 = q_0$ (Appendix A).

We assume

$$L \ge 20(1+\beta_0)^2\,, \tag{6.58}$$

so that $\varkappa_m L^{-m} \le (1+\beta_0)^2 L^{-1} \le \frac{1}{20}$ for every $m \ge 1$, because $m^{1/2}L^{-(m-1)} \le 1$ for $L \ge 2$.

Lemma 6.23 (The Space of a Later Step Lies in That of an Earlier One)**.** *Let U be a G-valued configuration. Then $\mathcal{S}_{k+1}(U) \subset \mathcal{S}_j(U)$.*

Proof. By (d) and (6.58), $\alpha_{i,k+1}L^{-(k+1)} \le \varkappa_m L^{-m}\alpha_{i,j}L^{-j} \le \alpha_{i,j}L^{-j}$, and the same holds with the powers $L^{-2(k+1)}$ and $L^{-3(k+1)}$, which are smaller. Each of the four radii of (5.17) at $n = k+1$ is therefore at most the corresponding radius at $n = j$, and for $j = 1$ only the first condition is posed. *q.e.d.*

Lemma 6.24 (Complexified Covariance)**.** *Let X be a finite set of bonds with the set S of its sites, $x, x' \in S$, $\mathcal{D}$ an open set of pairs $q = (\mathbf{U}, \mathbf{J})$ of a G^c-valued configuration and a $\mathfrak{g}^c$-valued field on X which is invariant under the G-valued gauge transformations, and F a holomorphic function on $\mathcal{D}$ with values in the linear maps of the space of a representation π of G, such that $F(q^v) = \pi(v(x))F(q)\pi(v(x'))^{-1}$ for every $q \in \mathcal{D}$ and every $v \in G^S$, where $q^v \coloneqq (\mathbf{U}^v, R(v)\mathbf{J})$. Let $q \in \mathcal{D}$ and let $t \mapsto u_t$, $t \in [0,1]$, be a continuous path in $(G^c)^S$ with $u_0 = 1$ and $q^{u_t} \in \mathcal{D}$ for every t. Then*

$$F(q^{u_1}) = \pi(u_1(x))\, F(q)\, \pi(u_1(x'))^{-1} \,. \tag{6.59}$$

Proof. The map $u \mapsto q^u$ is holomorphic on $(G^c)^S$, so that $\mathcal{U} \coloneqq \{u \in (G^c)^S \mid q^u \in \mathcal{D}\}$ is open, and it contains G^S by the invariance of $\mathcal{D}$. Since π is holomorphic on G^c, the function

$$\Psi(u) \coloneqq \pi(u(x))^{-1} F(q^u) \pi(u(x')) - F(q)$$

is holomorphic on $\mathcal{U}$, and it vanishes on G^S by hypothesis. In the chart $Z \mapsto \exp(Z)$ of $(G^c)^S$ at the identity, with $Z \in (\mathfrak{g}^c)^S = \mathfrak{g}^S \oplus i\mathfrak{g}^S$, the function $\Psi \circ \exp$ is holomorphic near 0 and vanishes on the real subspace $\mathfrak{g}^S$, whose complex span is the whole space. All its derivatives at 0 in the real directions vanish, hence, by the complex linearity of the derivatives of a holomorphic function, all its derivatives at 0, and Ψ vanishes on a neighbourhood of the identity. By the identity theorem it vanishes on the connected component of $\mathcal{U}$ which contains the identity, and the path u_t lies in this component, because it lies in $\mathcal{U}$ and starts at the identity. Hence $\Psi(u_1) = 0$, which is (6.59). *q.e.d.*

We use the lemma for the insertion of W with the transport along its interval removed. Let $\mathbf{U}(I_W)$ be the ordered product of $\mathbf{U}$ along the fine bonds of I_W, from x_W to x'_W, an element of G^c, and

$$\hat{Y}_W(\mathbf{U}, \mathbf{J}) \coloneqq Y_W(\mathbf{U}, \mathbf{J})\, \pi(\mathbf{U}(I_W))^{-1} \,. \tag{6.60}$$

Since $\mathbf{U}^v(I_W) = v(x_W)\mathbf{U}(I_W)v(x'_W)^{-1}$, the function $\hat{Y}_W$ is holomorphic on $\mathcal{D}_W$ and satisfies (5.19) with x_W in the place of x'_W. By Lemma 6.24, for a path u_t as there with $u_1(x_W) = 1$,

$$\hat{Y}_W(\mathbf{U}^{u_1}, R(u_1)\mathbf{J}) = \hat{Y}_W(\mathbf{U}, \mathbf{J}) \,. \tag{6.61}$$

The gauge transformation which we shall use is built from the following function. Three properties of the norm enter, each with a constant which depends only on G and on the norm; we fix $\rho_0 \in (0,1]$ and $c_G \ge 1$ such that all three hold.

(N1) For $X, Q \in \mathfrak{g}^c$ with $|X| \le \rho_0$, $|e^{iX}Qe^{-iX} - Q| \le c_G|X||Q|$.

(N2) For $X, Y, Z, A \in \mathfrak{g}^c$ of norm less than ρ_0 the matrix $e^{iX}e^{iY}e^{iZ}e^{-i(X+A)}e^{-iZ}$ is within the radius of convergence of the logarithm about 1, and

$$\Phi(X, Y, Z, A) \coloneqq -i \log(e^{iX}e^{iY}e^{iZ}e^{-i(X+A)}e^{-iZ}) \tag{6.62}$$

is a holomorphic function with values in $\mathfrak{g}^c$; the function $\Psi \coloneqq \Phi - (Y - A)$ has all its first and second partial derivatives bounded by c_G in norm where the four arguments have norm at most $\rho_0/2$.

(N3) Let U be a G-valued configuration and α a field with $|\alpha(b)| \le a_\circ \le \rho_0$ and $|(\delta_e\alpha)(b)| \le d_\circ \le a_\circ$ on the bonds of a fine plaquette p. Then

$$|(e^{i\alpha}U)(\partial p) - 1| \le c_G(|U(\partial p) - 1| + d_\circ + a_\circ^2) \,. \tag{6.63}$$

Such constants exist. (N1) holds because $Q \mapsto e^{iX}Qe^{-iX} - Q$ is holomorphic in X and vanishes at $X = 0$. In (N2) the product is a holomorphic function of the arguments which equals 1 at the origin, the logarithm of a product of exponentials of small elements of $\mathfrak{g}^c$ is given by the series of Baker, Campbell, and Hausdorff and lies in $\mathfrak{g}^c$, and the derivatives are bounded by Cauchy's estimates. For (N3), let $b_1 = \langle x, x + e_\mu\rangle$, b_2, b_3, $b_4 = \langle x, x + e_\nu\rangle$ be the bonds of p in the order $(\partial p) = b_1b_2b_3^{-1}b_4^{-1}$, $U_i := U(b_i)$, $\alpha_i := \alpha(b_i)$, and $F := U(\partial p)$. Moving the factors U_i to the right we find

$$(e^{i\alpha}U)(\partial p) = e^{i\alpha_1}e^{i\beta_2}e^{-i\beta_3}e^{-i\beta_4}F \,,$$
$$\beta_2 := U_1\alpha_2U_1^{-1} \,, \qquad \beta_3 := FU_4\alpha_3U_4^{-1}F^{-1} \,, \qquad \beta_4 := F\alpha_4F^{-1} \,.$$

Here $U_1\alpha_2U_1^{-1} = \alpha_4 + (\delta_{e_\mu}\alpha)(b_4)$ and $U_4\alpha_3U_4^{-1} = \alpha_1 + (\delta_{e_\nu}\alpha)(b_1)$ by (5.16), and the conjugation by F changes an element Q by at most a constant times $|F-1||Q|$. The sum $\alpha_1 + \beta_2 - \beta_3 - \beta_4$ of the exponents is therefore at most $2d_\circ$ plus a constant times $|F-1|a_\circ$ in norm, the product of the four exponentials differs from $1 + i(\alpha_1 + \beta_2 - \beta_3 - \beta_4)$ by at most a constant times $a_\circ^2$, since its arguments are bounded by $2a_\circ$, and the multiplication by F changes norms by at most a constant factor. This gives (6.63).

Two consequences of the definition (6.62) will be used: $\Phi(X, 0, 0, 0) = 0$ for every X and $\Phi(0, 0, Z, 0) = 0$ for every Z, and the linear part of Φ at the origin is $Y - A$. Hence, where the four arguments have norm at most $\rho_0/2$,

$$|\Psi(X, Y, Z, A)| \le c_G[(|Y| + |A|)(|X| + |Y| + |Z| + |A|) + |X||Z|] \,,$$
$$|\partial_X\Psi| \le c_G(|Y| + |Z| + |A|) \,, \qquad |\partial_Z\Psi| \le c_G(|X| + |Y| + |A|) \,, \tag{6.64}$$
$$|\partial_Y\Psi| \le c_G(|X| + |Y| + |Z| + |A|) \,.$$

For the bounds of the derivatives we note that $\partial_X\Psi$ vanishes at $(X, 0, 0, 0)$, $\partial_Z\Psi$ at $(0, 0, Z, 0)$, and $\partial_Y\Psi$ at the origin, and we integrate the second derivatives along the segment from these points. For the first line, $\Psi(X, 0, Z, 0)$ vanishes if $X = 0$ or $Z = 0$, so that it equals the integral of its mixed second derivative over $[0, 1]^2$ along (sX, tZ) and is at most $c_G|X||Z|$; and $\Psi(X, Y, Z, A) - \Psi(X, 0, Z, 0)$ is the integral of the derivative in (Y, A) along (X, sY, Z, sA), which vanishes at the origin and is therefore at most $c_G(|X| + |Y| + |Z| + |A|)$.

Lemma 6.25 (The Gauge Reduction of a Slowly Varying Complex Part)**.** *Let Q be a cube of the fine lattice of side $D \ge 1$ and x_0 a site of Q with $x_0 + e_\mu \in Q$ for every μ. Let $U = e^{i\zeta}$ on the bonds of Q, with a field ζ with values in $\mathfrak{g}$, and let α be a field on the bonds of Q with values in $\mathfrak{g}^c$, such that for all bonds b and unit vectors e with $b, b + e$ in Q*

$$|\alpha(b)| \le a \,, \qquad |(\delta_e\alpha)(b)| \le a' \,, \qquad |\zeta(b)| \le z \,, \qquad |\zeta(b + e) - \zeta(b)| \le z' \,, \tag{6.65}$$

with $a' \le a$ and $4aD, a, z \le \rho_0/2$. Put

$$\omega(x) := \sum_\mu a_\mu\,(x - x_0)_\mu \,, \qquad a_\mu := \alpha(\langle x_0, x_0 + e_\mu\rangle) \,, \qquad u_t := e^{it\omega} \,, \tag{6.66}$$

for the sites x of Q and $t \in [0, 1]$. Then $u_t(x_0) = 1$, $|\omega| \le 4aD$, and $(e^{i\alpha}U)^{u_t} = e^{i\alpha_t}U$ on the bonds of Q, with fields α_t with values in $\mathfrak{g}^c$ which satisfy, with $a'' := a' + c_Gza$ and $r_\Phi := c_G[2a(4aD + 2a + z) + 4aDz]$,

$$|\alpha_1(b)| \le 4Da'' + r_\Phi \,, \qquad |\alpha_t(b)| \le 2a + r_\Phi \,, \tag{6.67}$$

$$|(\delta_e\alpha_t)(b)| \le a'' + c_Gz(2a + r_\Phi) + c_G[(2a + z)a + (4aD + 2a + z)a'' + (4aD + 2a)z'] \,. \tag{6.68}$$

Proof. The value $\omega(x_0) = 0$ and the bound on ω are immediate from (6.66), since $|a_\mu| \le a$ and $\sum_\mu |(x - x_0)_\mu| \le 4D$. For a bond $b = \langle x, x + e_\mu \rangle$ of Q we have $\omega(x + e_\mu) = \omega(x) + a_\mu$, and therefore

$$(e^{i\alpha} U)^{u_t}(b) = e^{it\omega(x)} e^{i\alpha(b)} e^{i\zeta(b)} e^{-i(t\omega(x) + t a_\mu)} = e^{i\alpha_t(b)} e^{i\zeta(b)} ,$$
$$\alpha_t(b) \coloneqq \Phi(t\omega(x), \alpha(b), \zeta(b), t a_\mu) ,$$

by the definition (6.62); the arguments have norm at most $\rho_0/2$ by hypothesis, and α_t has values in $\mathfrak{g}^c$ by (N2).

The plain differences of α. For $b, b + e$ in Q, with $b = \langle x, x + e_\mu \rangle$, the value of U on the bond from x to $x + e$ is $e^{\pm i\zeta}$ with the value of ζ on that bond, so that by (5.16), (N1), and (6.65)

$$|\alpha(b + e) - \alpha(b)| \le |(\delta_e \alpha)(b)| + c_G z \, |\alpha(b + e)| \le a' + c_G z a = a'' .$$

Two bonds of the direction μ in Q are joined by a chain of at most $4D$ translations by unit vectors which stays in Q, one coordinate after the other, because Q is a cube. Hence $|\alpha(b) - a_\mu| \le 4Da''$ for every bond b of the direction μ in Q.

The bounds (6.67). By (N2), $\alpha_t(b) = \alpha(b) - t a_\mu + \Psi(t\omega(x), \alpha(b), \zeta(b), t a_\mu)$, and by the first line of (6.64), with $|t\omega| \le 4aD$, the last term is at most r_Φ in norm. For $t = 1$ the first two terms are at most $4Da''$ together, and for general t at most $2a$.

The bound (6.68). By (5.16) and (N1), $|(\delta_e \alpha_t)(b)| \le |\alpha_t(b + e) - \alpha_t(b)| + c_G z(2a + r_\Phi)$. The arguments of Φ at $b + e$ are $t\omega(x) + t a_e$, with $a_e \coloneqq \pm a_\nu$ for $e = \pm e_\nu$, then $\alpha(b + e)$, $\zeta(b + e)$, and the same $t a_\mu$. The difference of the linear parts is $\alpha(b + e) - \alpha(b)$, at most a'', and by the second and third lines of (6.64) the difference of the values of Ψ is at most $c_G(2a + z)|a_e| + c_G(4aD + 2a + z)a'' + c_G(4aD + 2a)z'$, the three terms being the changes of the first, the second, and the third argument, each multiplied by the bound of the corresponding derivative. *q.e.d.*

One carried insertion. Let W be a group of the deviation species made at the step j. We fix a point of the space of the step $k + 1$,

$$(\mathbf{U}, \mathbf{J}) = (e^{i\alpha} U, \mathbf{J}) \in \mathcal{S}_{k+1}(U) , \tag{6.69}$$

around a real configuration U which is either U_{k+1}, for a block field V_{k+1} in the support, with the current $J \coloneqq J_{k+1}$, or the trivial configuration, with $J \coloneqq 0$; in both cases $(U, J) \in \mathcal{S}_{k+1}(U)$ and, by (b), U is one of the configurations of (5.18). By Lemma 6.23 and (a), both $(\mathbf{U}, \mathbf{J})$ and (U, J) lie in $\mathcal{D}_W$. We write $a_{(n)} \coloneqq \alpha_{1,n} L^{-n}$ for the radius of the complex part at the step n, and

$$a \coloneqq a_{(k+1)} , \qquad \mathsf{d}_W \coloneqq 2u_\star M \sigma(W) L^j , \qquad \delta_W \coloneqq \mathsf{d}_W L^{-(k+1)} = 2u_\star M L^{-m} \sigma(W) . \tag{6.70}$$

By (c) the ℓ^∞-diameter of X_W is at most $u_\star \sigma(W) M L^j$, and that of the dependence set X_W^+ exceeds it by at most $16M \le ML^j$; hence X_W^+ lies in a cube Q_W of the fine lattice of side d_W, and δ_W is its side in the units of the scale $k + 1$. The interval has at most $\mathsf{d}_W/2$ fine bonds, and $\delta_W \ge 2ML^{-m}$ because $\sigma(W) \ge 1$.

Proposition 6.26 (One Carried Insertion)**.** *Let γ be sufficiently small. With the notation above the following hold.*

(i) For every W,

$$\|Y_W(\mathbf{U}, \mathbf{J}) - Y_W(U, J)\| \le c_{\mathrm{pl}} \varkappa_m L^{-m} \|Y_W\|_{(j)} , \qquad c_{\mathrm{pl}} \coloneqq \max\Big(2, \frac{8 c_G \lambda_{\mathrm{r}}}{c_F L}\Big) . \tag{6.71}$$

(ii) Let $\mu_c \coloneqq \min(\frac{1}{2}, c_F M/(c_G\lambda_{\mathrm{r}}))$. *If* $\delta_W \le \mu_c$, *then*

$$\|Y_W(\mathbf{U},\mathbf{J})\| \le 2\|Y_W(U,J)\| + \varkappa_m L^{-m}\Big(\frac{40\,\delta_W}{\mu_c} + 14L^{-2m}\Big)e^{\nu\sigma(W)/8192}\,\|Y_W\|_{(j)}\,. \quad (6.72)$$

The first bound is the gain of the radii alone and holds for every group; the second has the further factor $\delta_W \le 2u_\star M L^{-m}\sigma(W)$ and is the one which is summable over the generations.

Proof. We write $\mathsf{r}_n \coloneqq \alpha_{0,n}L^{-3n}$ for the radius of the current in (5.17) and $a_2 \coloneqq \alpha_{1,k+1}L^{-2(k+1)} = aL^{-(k+1)}$ for that of the covariant differences at the step $k+1$. By (d) and (6.58),

$$\frac{a}{a_{(j)}} \le \varkappa_m L^{-m} \le \frac{1}{20}\,, \qquad \frac{a_2}{\alpha_{1,j}L^{-2j}} \le \varkappa_m L^{-2m}\,, \qquad \frac{\mathsf{r}_{k+1}}{\mathsf{r}_j} \le \varkappa_m L^{-3m} \le \frac{1}{20L^2}\,. \quad (6.73)$$

The curvature of the real configuration is $f_U \coloneqq \max_p|U(\partial p)-1| \le 2B_3\varepsilon_{k+1}L^{-2(k+1)}$ by (b), and $f_U = 0$ for the trivial configuration. Since $\varepsilon_{k+1}/\alpha_{0,k+1}$ and $\alpha_{1,n}$ tend to zero with γ, uniformly in the steps (Appendix A, $q_0 > p_0$), we may and do assume that

$$c_G f_U \le \tfrac{1}{4}c_F\,\alpha_{0,j}L^{-2j}\,, \qquad c_G\,\alpha_{1,n}^2 \le \tfrac{1}{4}c_F\,\alpha_{0,n}\,, \qquad \alpha_{1,n} \le \rho_0 \quad (6.74)$$

for all steps n; the first uses $\alpha_{0,k+1} \le \varkappa_m\alpha_{0,j}$ and $\varkappa_m L^{-2m} \le 1$.

(i). For $j \ge 2$ put

$$R' \coloneqq \min\Big(\frac{a_{(j)}}{2a},\ \frac{c_F\,\alpha_{0,j}L^{-2j}}{8c_G\,a_2},\ \frac{1}{4}\Big(\frac{\mathsf{r}_j}{\mathsf{r}_{k+1}} - 1\Big)\Big)\,,$$

and for $j = 1$ let R' be the first of the three numbers. If $2R' \le 1$, the left side of (6.71) is at most $2\|Y_W\|_{(j)} \le \|Y_W\|_{(j)}/R'$, because both points lie in $\mathcal{D}_W$. Let $2R' > 1$. For $|z| < 2R'$ the pair $(e^{iz\alpha}U, J + z(\mathbf{J}-J))$ lies in $\mathcal{S}_j(U)$: the complex part $z\alpha$ has norm less than $2R'a \le a_{(j)}$ and covariant differences less than $2R'a_2 \le a_{(j)}L^{-(k+1)} \le \alpha_{1,j}L^{-2j}$; by (6.63) with $a_\circ = a_{(j)} \le \rho_0$ and $d_\circ = 2R'a_2$, and by (6.74) and the second entry of R', its curvature is at most $c_G f_U + 2c_G R'a_2 + c_G a_{(j)}^2 \le \frac{3}{4}c_F\alpha_{0,j}L^{-2j}$; and the current has norm less than $\mathsf{r}_{k+1}+4R'\mathsf{r}_{k+1} \le \mathsf{r}_j$, by the third entry. The function $g(z) \coloneqq Y_W(e^{iz\alpha}U, J+z(\mathbf{J}-J)) - Y_W(U,J)$ is therefore holomorphic on the disc of radius $2R' > 1$, it is at most $2\|Y_W\|_{(j)}$ in norm there, and $g(0) = 0$, so that by Schwarz's lemma, applied to the matrix elements, $\|g(1)\| \le \|Y_W\|_{(j)}/R'$. In both cases the left side of (6.71) is at most $\|Y_W\|_{(j)}/R'$, and by (6.73) the reciprocals of the three entries of R' are at most $2\varkappa_m L^{-m}$, $(8c_G\lambda_{\mathrm{r}}/c_F)\varkappa_m L^{-2m}$, and $4\cdot\frac{20}{19}\varkappa_m L^{-3m}$, each of which is at most $c_{\mathrm{pl}}\varkappa_m L^{-m}$.

(ii), the gauge. The conditions (5.17) are invariant under the G-valued gauge transformations, which conjugate α and $\mathbf{J}$, and so are the norms in (6.72), by (5.19) and because π is unitary on G. For $U = U_{k+1}$ we apply Supplement, Lemma 17.22, to a cube $\mathcal{Q}$ of 2^4 cubes of π_{k+1} which contains Q_W; such a cube exists because $\delta_W \le \frac{1}{2}$ is less than the side M of these cubes. Its hypotheses hold for γ small, because the local curvature is $E \le 2B_3\varepsilon_{k+1}$ by (b) and $M_{\mathcal{Q}} \le 4M$. It gives a G-valued gauge transformation after which $U = e^{i\zeta}$ on Q_W with, in fine units,

$$|\zeta(b)| \le z \coloneqq \hat{b}L^{-(k+1)}\,, \qquad |\zeta(b+e)-\zeta(b)| \le z' \coloneqq \hat{b}L^{-2(k+1)}\,, \qquad \hat{b} \coloneqq 24B_3^2M\varepsilon_{k+1}\,,$$

and for the trivial configuration $\zeta = 0$. We work in this gauge and apply Lemma 6.25 to the cube Q_W, with $D = \mathsf{d}_W$, $x_0 = x_W$, the radii a and $a' = a_2$ of (6.69), and these z, z'; its

hypotheses hold because $4a\mathsf{d}_W = 4\alpha_{1,k+1}\delta_W \le 2\alpha_{1,k+1}$ and z are small. In the bounds of that lemma, $a_2\mathsf{d}_W = a\delta_W$, $za\mathsf{d}_W = \hat{b}\,a\delta_W$, $a^2 = \alpha_{1,k+1}L^{-(k+1)}a \le \alpha_{1,k+1}a\delta_W$, and $za = \hat{b}\,a_2$, so that every term is a multiple of $a\delta_W$, or of a, or of a_2, with a coefficient which is 4, respectively 2, respectively 1, up to terms which vanish with $\alpha_{1,k+1}$ and $\hat{b}$. For γ small we therefore have, for $t \in [0,1]$ and on the bonds of Q_W,

$$|\alpha_1| \le 5a\delta_W\ , \qquad |\alpha_t| \le 3a\ , \qquad |\delta_e\alpha_t| \le 2a_2\ , \qquad |\omega| \le 4\alpha_{1,k+1}\delta_W\ . \tag{6.75}$$

(ii), the path. We claim that $(\mathbf{U}^{u_t}, R(u_t)\mathbf{J}) \in \mathcal{S}_j(U)|_{X_W^+}$ for $t \in [0,1]$. By Lemma 6.25, $\mathbf{U}^{u_t} = e^{i\alpha_t}U$, and by (6.75) and (6.73) the complex part satisfies $|\alpha_t| \le 3a < a_{(j)}$ and $|\delta_e\alpha_t| \le 2a_2 \le 2\varkappa_m L^{-2m}\alpha_{1,j}L^{-2j} < \alpha_{1,j}L^{-2j}$. For the curvature we use that $\delta_W \le \mu_c$ and $\delta_W \ge 2ML^{-m}$ give $L^{-m} \le \mu_c/(2M)$, so that $2a_2 \le 2\varkappa_m L^{-m}\cdot L^{-m}\lambda_{\mathrm{r}}\alpha_{0,j}L^{-2j} \le (\mu_c\lambda_{\mathrm{r}}/(20M))\alpha_{0,j}L^{-2j} \le (c_F/(20c_G))\alpha_{0,j}L^{-2j}$ by the definition of μ_c; with (6.63) for $a_\circ = 3a \le a_{(j)}$ and (6.74) the curvature is less than $c_F\alpha_{0,j}L^{-2j}$. Finally $|R(u_t)\mathbf{J}| \le (1+c_G|\omega|)|\mathbf{J}| \le 2\mathsf{r}_{k+1} < \mathsf{r}_j$ by (N1). For $j=1$ only the first of these conditions is posed. Since $u_1(x_W)=1$, (6.61) and (6.60) give

$$Y_W(\mathbf{U},\mathbf{J}) = \hat{Y}_W(\mathbf{U}^{u_1}, R(u_1)\mathbf{J})\,\pi(\mathbf{U}(I_W))\ . \tag{6.76}$$

(ii), the disc. Put

$$R_A \coloneqq \frac{\mu_c\, a_{(j)}}{10\, a\delta_W}\ , \qquad R_J \coloneqq \frac{1}{3}\Big(\frac{\mathsf{r}_j}{\mathsf{r}_{k+1}} - 1\Big)\ , \qquad R \coloneqq \min(R_A, R_J)\ ,$$

with $R \coloneqq R_A$ for $j=1$, and $f(z) \coloneqq \hat{Y}_W(e^{iz\alpha_1}U, J + z(R(u_1)\mathbf{J} - J))$ for $|z| < R$, so that $f(1)$ is the first factor of (6.76) and $f(0) = \hat{Y}_W(U,J)$. Since $\delta_W \le \mu_c$, we have $R_A \ge a_{(j)}/(10a) \ge 2$ and $R_J \ge 2$ by (6.73). For $|z| < R$ the pair in the argument of f lies in $\mathcal{S}_j(U)|_{X_W^+}$: by (6.75) the complex part $z\alpha_1$ has norm less than $5R_Aa\delta_W = \mu_c a_{(j)}/2$; its covariant differences are less than $2R_Aa_2 = \mu_c a_{(j)}L^{-(k+1)}/(5\delta_W) \le \mu_c\,\alpha_{1,j}L^{-2j}/(10M)$, because $\delta_W \ge 2ML^{-m}$; by (6.63), (6.74), and the definition of μ_c its curvature is at most

$$c_G f_U + \frac{c_G\mu_c\lambda_{\mathrm{r}}}{10M}\alpha_{0,j}L^{-2j} + c_G a_{(j)}^2 \le \Big(\frac{1}{4}+\frac{1}{10}+\frac{1}{4}\Big)c_F\,\alpha_{0,j}L^{-2j}\ ;$$

and the current has norm less than $\mathsf{r}_{k+1} + 3R_J\mathsf{r}_{k+1} = \mathsf{r}_j$, because $|R(u_1)\mathbf{J}| \le 2\mathsf{r}_{k+1}$. Hence f is holomorphic on the disc of radius R. Along the interval the transport of $e^{iz\alpha_1}U$ is a product of at most $\mathsf{d}_W/2$ factors $e^{iz\alpha_1(b)}U(b)$, with $\pi(U(b))$ unitary and $\|\pi(e^{iX})\| \le e^{c'_\pi|X|}$ (Section 6.3), so that by (6.60) and (a)

$$\begin{aligned}\|f(z)\| &\le \|Y_W\|_{(j)}\exp(c'_\pi\tfrac{1}{2}\mu_c a_{(j)}\cdot\tfrac{1}{2}\mathsf{d}_W) = \|Y_W\|_{(j)}\exp(\tfrac{1}{2}c'_\pi\mu_c u_\star M\alpha_{1,j}\,\sigma(W))\\ &\le e^{\nu\sigma(W)/8192}\,\|Y_W\|_{(j)}\ ,\end{aligned}$$

the last for γ so small that $4096\,c'_\pi u_\star M\alpha_{1,n} \le \nu$ for all n. By Schwarz's lemma for $f - f(0)$, which is at most twice this bound on the disc, $\|f(1)-f(0)\| \le (2/R)e^{\nu\sigma(W)/8192}\|Y_W\|_{(j)}$. In the same way $\|\pi(\mathbf{U}(I_W))\| \le \exp(c'_\pi a\mathsf{d}_W/2) = \exp(\frac{1}{2}c'_\pi\alpha_{1,k+1}\delta_W) \le 2$, and $\|f(0)\| = \|Y_W(U,J)\|$ because $\pi(U(I_W))$ is unitary. By (6.76),

$$\|Y_W(\mathbf{U},\mathbf{J})\| \le 2(\|f(0)\| + \|f(1)-f(0)\|) \le 2\|Y_W(U,J)\| + \frac{4}{R}\,e^{\nu\sigma(W)/8192}\|Y_W\|_{(j)}\ ,$$

and $4/R \le 4/R_A + 4/R_J \le 40\varkappa_m L^{-m}\delta_W/\mu_c + 12\cdot\frac{20}{19}\varkappa_m L^{-3m}$ by (6.73), which is (6.72). *q.e.d.*

The sum over the generations. It remains to sum the bounds of the proposition over the groups at a traversal.

Proposition 6.27 (The Carried Groups in the Supremum Clause)**.** *Let the ensemble be dressed at the steps* $1, \dots, k$*, and suppose that on the term without large-field regions of the step* $k+1$ *the groups carried from these steps satisfy* (5.13) *and* (5.14) *at the step* $k+1$*, at every real background of the term, and vanish at its reference data. Let* γ *be sufficiently small and* $L > (1+\beta_0)^6$*. Then, for every traversal* $\mathfrak{n}'$ *of the step* $k+1$*,*

$$\sum_{W \ni \mathfrak{n}'} e^{\theta(W)+\lambda_c \nu \sigma(W)+\nu_e \operatorname{ext}(W)} \|Y_W\|_{(k+1)} \le 2C_\chi \tilde\rho_{k+1} \bar\delta^*_{\mathfrak{n}'} + 2\hat C\, C_{S'}\, \rho_{k+1} \,, \tag{6.77}$$

the sum over the groups of the deviation species made at the steps $j \le k$ *with* $\mathfrak{n}'$ *in the interval, where* $\bar\delta^*_{\mathfrak{n}'}$ *is the expression* (5.7) *of the step* $k+1$ *with each maximum replaced by its supremum over the real backgrounds of the term, and*

$$\hat C := \sum_{m \ge 1} (1+\beta_0)^{6m}\, \varkappa_m \Big[L^{-m} \Big(\frac{1280\, u_\star M}{e\, \nu \mu_c} + 14 L^{-m} \Big) + c_{\mathrm{pl}} \exp\Big(-\frac{\nu \mu_c L^m}{32 u_\star M} \Big) \Big] \,. \tag{6.78}$$

Here $\theta(W)$*,* $\sigma(W)$*, and* $\operatorname{ext}(W)$ *are the numbers fixed when* W *was made, as in* (5.22)*.*

The constant $\hat C$ is finite because $L > (1+\beta_0)^6$, and it depends on L, M, ν, β_0, $u_\star$, c_F, c_G, and λ_{r}, and not on k or on the coupling. The first term of (6.77) is at most $2C_\chi \tilde\rho_{k+1} \cdot 2c_\sigma B_3 \varepsilon_{k+1} W_1$, because the bound (6.1) holds for each maximum and at every real background, and this is less than ρ_{k+1} for γ small, since $\tilde\rho_{k+1}\varepsilon_{k+1} = \rho_{k+1}\varepsilon_{k+1}/\min(\alpha_{0,k+1}, \alpha_{1,k+1})$ and $q_0 > p_0$. With (5.22) at the step $k+1$, in which $\ell_W \ge 1$, $S_W \ge \|Y_W\|_{(k+1)}$, and $\lambda_c \le 1$, the proposition therefore gives (5.21) at the step $k+1$ with

$$C_c := 1 + 2(1+\hat C)\, C_{S'} \,. \tag{6.79}$$

The constant does not come from a recursion: the bound of the carried groups uses the production bounds of the earlier steps and no supremum clause.

Proof. The real points. Let W be a group of the sum, made at the step j, and (6.69) a point of the space of the step $k+1$, around U. We apply Proposition 6.26(ii) if $\delta_W \le \mu_c$ and (i) otherwise, and in both cases

$$\|Y_W(\mathbf{U}, \mathbf{J})\| \le 2\|Y_W(U, J)\| + \epsilon_W \,,$$

with ϵ_W the second term of (6.72), respectively the right side of (6.71); it does not depend on the point. If U is the trivial configuration, the first term vanishes by hypothesis. If $U = U_{k+1}$, let $\mathfrak{n}_j \subset \mathfrak{n}'$ be a traversal of the step j in the interval of W and $t := t^{(k+1)}_{\mathfrak{n}'}(W)$. As in the proof of Lemma 5.3, the comparison region of W at $\mathfrak{n}_j$ lies within the distance R_t of $\mathfrak{n}'$ in the units of the scale $k+1$, so that by (5.13) and (5.8) $e^{\theta(W)}\|Y_W(U,J)\| \le \beta_W \bar m_t$, where $\bar m_t$ is the supremum over the real backgrounds of the maximum of $|U(\partial p) - 1|$ over $p \in \mathcal{P}_t(\mathfrak{n}')$. Taking the supremum over the points we obtain

$$e^{\theta(W)}\|Y_W\|_{(k+1)} \le 2\beta_W\, \bar m_{t^{(k+1)}_{\mathfrak{n}'}(W)} + e^{\theta(W)} \epsilon_W \,. \tag{6.80}$$

The numbers $\bar m_t$ are nondecreasing in t, and the summation by parts of the proof of Lemma 5.3, with (5.14) at the step $k+1$ and $\lambda_c \le 1$, bounds the weighted sum of the first terms of (6.80) over the groups of the sum by $2C_\chi \tilde\rho_{k+1} \bar\delta^*_{\mathfrak{n}'}$.

The excess of one generation. Fix $j \le k$ and consider the groups of the sum which were made at the step j. Each of them has a traversal of the step j in its interval which is contained in $\mathfrak{n}'$, namely the one which contains a fine bond common to $\mathfrak{n}'$ and the interval, and $\mathfrak{n}'$ contains L^m traversals of the step j. By (5.22) at the step j, with $\ell_W \ge 1$ and $S_W \ge \|Y_W\|_{(j)}$,

$$\sum_W e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)} \|Y_W\|_{(j)} \le 2L^m C_{S'}\, \rho_j \;, \tag{6.81}$$

the sum over these groups. For those with $\delta_W \le \mu_c$ we have $\lambda_c\nu + \nu/8192 = \frac{15}{16}\nu$ and, by (6.70) and $te^{-\nu t/16} \le 16/(e\nu)$,

$$\begin{aligned} e^{\lambda_c\nu\sigma(W)}\epsilon_W &\le e^{\nu\sigma(W)} \varkappa_m L^{-m} \Big(\frac{80 u_\star M L^{-m}}{\mu_c} \sigma(W) e^{-\nu\sigma(W)/16} + 14 L^{-2m} \Big) \|Y_W\|_{(j)} \\ &\le e^{\nu\sigma(W)} \varkappa_m L^{-2m} \Big(\frac{1280\, u_\star M}{e\,\nu\mu_c} + 14 L^{-m} \Big) \|Y_W\|_{(j)} \;. \end{aligned}$$

For those with $\delta_W > \mu_c$ we have $\sigma(W) > \mu_c L^m/(2u_\star M)$, so that $e^{\lambda_c\nu\sigma(W)} \le e^{\nu\sigma(W)} e^{-\nu\sigma(W)/16} \le e^{\nu\sigma(W)} \exp(-\nu\mu_c L^m/(32 u_\star M))$ and

$$e^{\lambda_c\nu\sigma(W)}\epsilon_W \le e^{\nu\sigma(W)} c_{\mathrm{pl}}\, \varkappa_m L^{-m} \exp\Big(-\frac{\nu\mu_c L^m}{32 u_\star M} \Big) \|Y_W\|_{(j)} \;.$$

Multiplying by $e^{\theta(W)+\nu_e \operatorname{ext}(W)}$ and summing with (6.81), we find that the weighted sum of the ϵ_W over the groups made at the step j is at most the m-th term of (6.78) without the factor $(1+\beta_0)^{6m}$, times $2C_{S'}\rho_j$, and $\rho_j \le (1+\beta_0)^{6m}\rho_{k+1}$ by (6.55).

The sum. Summing the excesses over $j \le k$, that is, over $m \ge 1$, and adding the first terms of (6.80), we obtain (6.77). *q.e.d.*

6.11 The Regions of a Term and the Collar

On a term with large-field regions the step meets two things which the term without them does not have: fields which it does not integrate, and insertions which earlier steps have made near such fields. Both are confined to a neighbourhood of the large-field regions, and what makes them harmless is a matter of distances: "the fluctuation field is localized in Ω_{k+1}" [9, p. 276], the small-field region of the step, and Bałaban's construction keeps this region far from every earlier large-field region, in the units of the step. We collect these distances first. A labelled term of ϱ_{k+1} has the domains $\Omega_1 \supset \Lambda_1 \supset \Omega_2 \supset \cdots \supset \Omega_{k+1} \supset \Lambda_{k+1}$ of its steps, and $Z_j := \Lambda_j^c$ is the large-field region of the step j [9, p. 264].

Lemma 6.28 (The Regions of a Term)**.** *Consider a labelled term of ϱ_{k+1}, and let $1 \le j \le k$.*

(i) $\Lambda_{j+1} \subset \Omega_{j+1} \subset \Lambda_j$.

(ii) In the units of the scale j, every point whose ℓ^∞-distance from Λ_j^c is less than $8LMR_{j+1}$ lies in Ω_{j+1}^c.

(iii) Let the collar $\mathbf{Coll}_j$ *be the union of the cubes of π_j whose ℓ^∞-distance from Λ_j^c is at most $6M$ in the units of the scale j, the cubes of Λ_j^c included. Then, for every n with $j \le n \le k$, the collar $\mathbf{Coll}_j$ lies in Ω_{n+1}^c, and its ℓ^∞-distance from Ω_{n+1} is at least $7LMR_{n+1}$ in the units of the scale n.*

Proof. All three statements are read from the two formulae by which Bałaban defines the regions of a step, here the step $j \to j+1$. The first is [9, (3.5), p. 265]: with Z_j' the union of

the cubes of side LMR_{j+1} of the partition which meet Z_j, the set $B^{j+1}(P^1_{j+1})$ is contained in $(Z_j'^{\sim 5})^c$, and $\Omega_{j+1} = (Q_{j+1}'^{\sim 2} \cup (B^{j+1}(P^1_{j+1})^c)^{\sim 3})^c$, where the operation $\sim$ adds a layer of these cubes. Hence $\Omega^c_{j+1} \supset (Z_j'^{\sim 5})^{\sim 3} = Z_j'^{\sim 8}$. The set $Z_j'^{\sim 8}$ contains every point at a distance less than $8LMR_{j+1}$ from Z_j', since such a point lies in a cube of the partition which is joined to a cube of Z_j' by a chain of at most eight further cubes, consecutive ones having a point in common. With $Z_j' \supset Z_j = \Lambda_j^c$ this proves (ii), and the second inclusion of (i). The second formula is [9, (3.20), p. 269], $\Lambda_{j+1} = \Omega_{j+1}^{\sim -2} \cap (R_{j+1}'^{\sim})^c$, after which the components of Λ^c_{j+1} which are contained in a cube of the size $100LMR_{j+1}$ are replaced by parallelepipeds which contain them; both operations give a subset of Ω_{j+1}, which is the first inclusion of (i).

(iii) A point of $\mathbf{Coll}_j$ is at a distance of at most $6M + M$ from Λ_j^c in the units of the scale j, that is, of at most $7ML^{-(n-j)}$ in the units of the scale n, and $\Lambda_j^c \subset \Lambda_n^c$ by (i). By (ii) at the step n, a point of Ω_{n+1} is at a distance of at least $8LMR_{n+1}$ from Λ_n^c. The distance between the two sets is therefore at least $8LMR_{n+1} - 7M \geq 7LMR_{n+1}$, because $LR_{n+1} \geq 7$, and in particular they are disjoint. *q.e.d.*

In the sides of the cubes of π_n the distance of (iii) is $7LR_{n+1}$, a number which tends to infinity as the coupling tends to zero, since $R_{n+1} \geq (\log \hat{g}_{n+1}^{-2})^r$ (Definition 3.3). This is the distance which a polymer of the step $n \to n+1$ has to cross if it is to connect an insertion made in the collar of an earlier step with the field which that step integrates.

6.12 A New Large-Field Region: The Interior Groups

We begin with the step at which a large-field region first appears. Throughout this subsection the labelled term of ϱ_k is the term without large-field regions and the sets Ω_{k+1}, Λ_{k+1}, S_{k+1} of the step are arbitrary, which is the setting of Section 6.3 and of Proposition 6.8. We call the labelled term of ϱ_{k+1} which these sets define the *present term*, and we write ℓ_0 for the term of ϱ_{k+1} without large-field regions. Two things distinguish the present term from ℓ_0. The field A_k is integrated on Λ_{k+1} only, and its restriction σ_k to the bonds of $\Omega_{k+1} \cap \Lambda^c_{k+1}$, which we call the *outer field of the step*, is one of the variables which $\mathbf{T}_{k+1}$ holds pending [9, p. 270], so that the factors which the step makes may depend on it. And the test functions, the potentials of the fluctuation integral, and the regularity of the background are those of a small-field region only inside Λ_{k+1} and Ω_{k+1}. We show that both differences are confined to a neighbourhood of Λ^c_{k+1}: a group of the step whose domain stays away from that neighbourhood is, factor by factor, a group of ℓ_0, and the bounds of Sections 6.6–6.10 hold for it as they were proved. The groups at the neighbourhood, and the groups of the step k whose comparison regions leave Ω_{k+1}, become attached to the new region.

The neighbourhood and the spaces. The set Λ_{k+1} is a union of cubes of side LMR_{k+1} in the units of the scale k [9, (3.20), p. 269], hence a union of cubes of π_{k+1}. Let $\mathcal{N}_{k+1}$ be the union of the cubes of π_{k+1} which lie in Λ^c_{k+1} or have a point in common with a cube of Λ^c_{k+1}. A union of cubes of π_{k+1} none of which belongs to $\mathcal{N}_{k+1}$ lies in Λ_{k+1}, hence in Ω_{k+1} by Lemma 6.28(i), and its distance from Λ^c_{k+1} is at least LM in the units of the scale k. The background U_{k+1} of the present term is a background of the step k: the block field $V_k^* := M^k(U_{k+1})$ lies in the support of the term without large-field regions of ϱ_k, by Supplement, Proposition S.8, which is stated for every labelled term, and $U_k(V_k^*) = U_{k+1}$ (Section 6.1). On Ω_{k+1} it has the regularity of the scale $k+1$, $|U_{k+1}(\partial p) - 1| < 2B_3\varepsilon_{k+1}L^{-2(k+1)}$ for the fine plaquettes p of Ω_{k+1} [9, p. 268]. For a union $X \subset \Omega_{k+1}$ of cubes of π_{k+1} we call the *space of the step* $k+1$ *on* X the union of the sets $\mathcal{S}_{k+1}(U)|_X$ of Definition 5.2(iii) over the configurations U which are the trivial configuration, the background U_{k+1} of the present term at a block field

and outer variables of the support of its test functions, or a transform of one of these by a G-valued gauge transformation.

The groups of the present term. By Lemma 6.4 and Proposition 6.8 the expectation $\langle F \rangle$ has the form (6.24) on the present term. Its vacuum polymers are the localisation domains which meet Λ_{k+1} [9, p. 277]. Their activities, and those of the marked polymers, may depend on σ_k and on the sets of the step; we use them at the real fields σ_k of the support of the test functions and at $\sigma_k = 0$, and we use Bałaban's bound [10, (2.38), p. 20] for the vacuum polymers of the present term, in the form of Supplement, Lemma 17.24, on the strength of his statement that the expansion of the measure [9, (3.27)] is made "in exactly the same way" and that "the differences, like the restriction of the integration to Λ_{k+1}, or the larger class of localization domains, do not matter" [9, pp. 277, 279] (Section 14). The *reference data* of the present term are the trivial configuration with $\mathbf{J} = 0$ and $\sigma_k = 0$, and the cluster weights $\varphi_\circ^T$ and the numbers $\Psi_\circ(M)$ are formed at them as in Section 6.5. We write $\varphi_\circ^{T,0}$ and $\Psi_\circ^0$ for the same quantities of ℓ_0, and we give a cluster which does not occur on one of the two terms the weight 0 on that term. For a marked polymer M the numbers $\Psi_\circ(M)$ and $\Psi_\circ^0(M)$ differ by the clusters which feel the new region, and we expand the difference, as the soft interactions were expanded in Section 6.5. With $\delta\varphi(\mathcal{C}) \coloneqq \varphi_\circ^T(\mathcal{C}) - \varphi_\circ^{T,0}(\mathcal{C})$,

$$e^{-\Psi_\circ(M)} = e^{-\Psi_\circ^0(M)} \sum_{\mathcal{O}'} \prod_{\mathcal{C} \in \mathcal{O}'} z_{\mathcal{C}} \;, \qquad z_{\mathcal{C}} \coloneqq e^{-\delta\varphi(\mathcal{C})} - 1 \;, \tag{6.82}$$

the sum over the finite sets $\mathcal{O}'$ of clusters which are incompatible with M and have $\delta\varphi(\mathcal{C}) \neq 0$; it converges absolutely, because $\sum_{\mathcal{C} \not\sim M} (|\varphi_\circ^T(\mathcal{C})| + |\varphi_\circ^{T,0}(\mathcal{C})|)$ is finite by Supplement, Lemma 17.24(ii), on each of the two terms. We call the members of $\mathcal{O}'$ the *collar occurrences* of M. The *groups of the step $k+1$ on the present term* are those of Definition 6.10 with the collar occurrences added to the data of a term, and a collar occurrence is treated as a link is: it is carried by its marked polymer M, its polymers belong to the domain of M and to the body $B(M)$ of (6.30), and the conditions (ii) and (iv) of that definition and the conditions of the compatibility of two groups apply to it as to a link. The insertion of a group contains the tensors $e^{-\Psi_\circ^0(M)} \tilde{H}(M)$ of its marked polymers, in the place of $e^{-\Psi_\circ(M)} \tilde{H}(M)$, and the further factors $z_{\mathcal{C}}$ of their collar occurrences. With these changes the remarks after Definition 6.10 and Proposition 6.11, with its proof, hold on the present term, under the hypothesis of that proposition.

Lemma 6.29 (Interior Constituents)**.** *On the present term the following hold.*

(i) Let G be a group of Lemma 6.4 such that no cube of π_{k+1} which meets the set D_G of Section 6.4 belongs to $\mathcal{N}_{k+1}$. Then the tensor Y_G does not depend on σ_k, and it is the same function of the field A on the bonds of D_G and of the configuration as on ℓ_0.

(ii) Let M be a union of cubes of π_{k+1} none of which belongs to $\mathcal{N}_{k+1}$. If M is a vacuum polymer, its activity does not depend on σ_k or on the sets Ω_{k+1}, Λ_{k+1}, S_{k+1}, and it equals the activity of M on ℓ_0; and the same is true of the marked activity $\tilde{H}(M)$ of a marked polymer.

(iii) A cluster none of whose polymers has a cube in $\mathcal{N}_{k+1}$ has the same weight on the present term and on ℓ_0, at every configuration and at the reference data. Consequently every collar occurrence has a polymer with a cube in $\mathcal{N}_{k+1}$.

Proof. We go through the ingredients of the three objects and observe that each of them is determined by data in the set concerned.

(i) By Lemma 6.4 the tensor Y_G is built from the functions $\mathbf{H}_Y(s, B')$ of the components Y of G, from the insertions of the members which these components touch, and from transports. On the term without large-field regions of ϱ_k the operators of the step are those of one domain (Section 6.3), so that $\mathbf{H}_Y(s, B')$ is the same functional of the parameters, of B', and of the configuration on Y on both terms, and the insertions of the members are functions of the earlier steps. The variable B' on the bonds of a component is a function of the field A_k on the bonds in the interior of D_G (Section 6.4), given by the same formula [10, (1.19), p. 6] on both terms. By the hypothesis these bonds lie in Λ_{k+1}, at a distance of at least LM from its complement, so that the field on them is the integrated field A and not σ_k.

(ii) By the construction [10, (2.1)–(2.11), pp. 11–14] the activity of a union Z of cubes is a finite sum of integrals, over variables on the bonds of Z and over parameters, of products of four kinds of factors: the Mayer factors of potentials whose localisation domains lie in Z; the factors of the test functions of the bonds of Z; Gaussian densities and operators whose parameters [10, (2.8), (2.9)] vanish outside Z, so that the term is "localized in the interior of Z" [10, p. 14]; and, for a marked polymer, the factor $\mathcal{X}_M$. Let $Z = M$ be as in the statement. The localised action of the measure [9, (3.27)] consists of the terms of $\mathbf{P}^{(k)}$ and of the brackets of $\mathbf{E}_k$ and $\mathbf{R}_k$, of the terms of the boundary terms $\mathbf{B}_k$ and of $A(g_k^{-2}(\cdot) - g_k^{-2}, \cdot)$, and of the terms of $V^{(k)}(S_{k+1})$ [9, pp. 276–277]. Those of the first kind with a localisation domain $Y \subset \Lambda_{k+1}$ "satisfy exactly all the conditions of those lemmas" [9, p. 276], and each of them depends on the fluctuation field "restricted to the interior of Y" [10, p. 9], hence on A alone when $Y \subset M$; they are given by the formulae of [10, Sect. 1] at the operators of one domain, as on ℓ_0. The terms of the second kind are absent, because the term of ϱ_k has no large-field region: the boundary terms are sums over localisation domains which meet $Z_k^\sim$ [9, (2.41), p. 261], and $g_k^{-2}(\cdot) - g_k^{-2}$ is supported in Z_k [9, p. 277]. The terms of the third kind "have at least one localization in S_{k+1}" [9, p. 277], and $S_{k+1} \subset \Lambda_{k+1}^c$ by [9, (3.20)], so that none of them has its localisation domains in M. The test functions on Λ_{k+1} are "only the small field characteristic functions" of the single bonds [9, p. 269], as on ℓ_0. The operators with vanishing parameters outside M are sums of the walks of the expansions which stay in M, and they do not involve the field or the sets outside M; in particular the term $\langle A, C^* \Delta^{(k)} C A_k \rangle$ of the measure, which couples A to σ_k through the kernel between Λ_{k+1} and its complement, does not enter. Finally the factor $\mathcal{X}_M$ is the same function on both terms, by (i), since the sets D_G of its groups lie in M. This is the reasoning behind Bałaban's remark that the terms of his expansion with domains in Λ_{k+1} "do not depend on Λ_{k+1}, and coincide with the corresponding terms arising from the expressions defined on the whole lattice" [9, p. 278].

(iii) The weight (6.15) of a cluster is a number times the product of the activities of its polymers, to which (ii) applies. Hence $\delta\varphi(\mathcal{C}) = 0$ for such a cluster, and a collar occurrence, for which $\delta\varphi(\mathcal{C}) \neq 0$, is not of this kind. *q.e.d.*

Definition 6.30 (Interior Groups and Groups at the Collar)**.** A group $\mathcal{G}$ of the step $k+1$ on the present term is *interior* if no cube of its domain $X_\mathcal{G}$ belongs to $\mathcal{N}_{k+1}$, and it is *at the collar* otherwise.

The domain of a group contains the cubes which meet the bodies of its constituents, the polymers of its occurrences, or its intervals (Lemma 6.13). An interior group therefore has no collar occurrence, by Lemma 6.29(iii); its marked polymers, its links, and its vacuum occurrences have no cube in $\mathcal{N}_{k+1}$; and its intervals lie in Λ_{k+1}.

Proposition 6.31 (Interior Groups)**.** *Let $\mathcal{G}$ be an interior group of the present term.*

(i) The data of $\mathcal{G}$ define a group of the step $k+1$ on ℓ_0, with the same domain, intervals, size, and extent. The insertion $Y_\mathcal{G}$ of the present term does not depend on σ_k, and it

is the same function of the restriction of $(\mathbf{U}, \mathbf{J})$ *to* $X_{\mathcal{G}}$ *as the insertion of that group on* ℓ_0*; in particular the two have the same flat value.*

(ii) The insertion $Y_{\mathcal{G}}$ *is holomorphic on the space of the step* $k+1$ *on* $X_{\mathcal{G}}$ *and satisfies the covariance law* (5.19) *there. Let* $\hat{P}(\mathcal{G})$ *be the product* (6.38) *with each supremum taken over the union of the space of the step* $k+1$ *of* ℓ_0 *and of the space of the step* $k+1$ *on* $X_{\mathcal{G}}$*. Then* $\sup\|Y_{\mathcal{G}}\| \le \hat{P}(\mathcal{G})$ *on that union, and Lemma 6.16, Theorem 6.18, and Corollary 6.20 hold for the groups of* ℓ_0 *with* $\hat{P}$ *in the place of* P*.*

(iii) Consequently the bounds (6.48)*,* (5.22)*, and* (5.23) *of the step* $k+1$ *hold on the present term for the sums over its interior groups made at the step* $k+1$*, with the suprema over the spaces of the step* $k+1$ *on their domains and with the constants of* ℓ_0*.*

(iv) If $\mathcal{G}$ *is made at the step* $k+1$*, has one interval, embeds in the torus, and has its comparison regions in* Ω_{k+1}*, then the bound* (6.51) *holds for it at every block field and outer variables of the support of the present term, and the tail bound* (6.52) *holds for the sum over these groups.*

Proof. (i) A single group with its constituents, links, and occurrences is a set of pairwise compatible groups, and by the proof of Proposition 6.11 its data form a term whose only group it is, on the present term and on ℓ_0 alike: the relation of Definition 6.10 between two constituents, the arc sets, and the domain are defined by the constituents and their domains, in which the sets of the step do not occur. The factors of $Y_{\mathcal{G}}$ are the tensors $e^{-\Psi^0_\circ(M)}\tilde{H}(M)$ of its marked polymers, the factors u_λ of its links, the factors $v_{\mathcal{C}}$ of its vacuum occurrences, the numbers of its old occurrences, the insertions of its untouched members, and the transports along its intervals. The first three are the same on the two terms by Lemma 6.29, because the marked polymers and the clusters concerned have no cube in $\mathcal{N}_{k+1}$; the old occurrences and the members belong to the step k, whose term is the same; and the transports are those of the configuration along the intervals. None of the factors involves σ_k.

(ii) The proofs of Lemma 6.19, of Proposition 6.9, and of the bound (6.39) use of a point $(\mathbf{U}, \mathbf{J})$ two things: that it satisfies the pointwise conditions (5.17) around a real configuration U of Bałaban's spaces, on which his operators, potentials, and activities are analytic with uniform bounds; and that, for a fluctuation field of the support and parameters with $|s(\Delta)| \le e^{\kappa_1}$, the interpolated configuration lies in the space of the step k, where the insertions of the members are holomorphic and bounded by their suprema. Both hold at the points of the space of the step $k+1$ on $X_{\mathcal{G}}$: the first by the regularity of U_{k+1} on Ω_{k+1}, and the second by (6.6), because U_{k+1} is a background of the step k. The three proofs therefore apply to the function of (i) on that space, and so do the proofs of Lemma 6.16, of Proposition 6.17, of Theorem 6.18, and of Corollary 6.20, which bound the suprema of the factors by these results, by Supplement, Lemma 17.24(iv), and by the class at the step k.

(iii) By (i) the interior groups of the present term which are at a cube or at a traversal p are among the groups of ℓ_0 at p, with the same sizes and extents, and by (ii) each of them has $S_{\mathcal{G}} \le 3e^{\nu\sigma_{k+1}(\mathcal{G})}\hat{P}(\mathcal{G})$, as in the proof of Corollary 6.20. The three bounds are therefore instances of that corollary and of Lemma 6.37(i) for the groups of ℓ_0, with $\hat{P}$ in the place of P.

(iv) The proof of Proposition 6.21 uses the term at three places. It uses the holomorphy of $Y^{\mathrm{d}}_{\mathcal{G}}$ and the bound $S_{\mathcal{G}}$ on the sets $\mathcal{S}_{k+1}(1)|_{\mathcal{Q}^{\sim 2}}$ around the trivial configuration, which belong to the space of ℓ_0 and are covered by (ii). It applies Supplement, Lemma 17.22, to the cube $\mathcal{Q}$, which requires $\mathcal{Q}^{\sim 5} \subset \Omega_{k+1}$ and the background on that set only; here $\mathcal{Q}^{\sim 5}$ lies in the comparison region, hence in Ω_{k+1}. And in its trivial case it uses that the pair of U_{k+1} and its

current, restricted to $X_{\mathcal{G}}$, is a point of the space on which $S_{\mathcal{G}}$ bounds the insertion; this is the space of the step $k+1$ on $X_{\mathcal{G}}$, by the regularity of U_{k+1} on Ω_{k+1}, as in (b) of Section 6.10. The tail bound follows from (6.48) as in that proof, by (iii). *q.e.d.*

The groups attached to the new region. The groups of the step k which the step leaves alone are the old functions on the present term as on ℓ_0, and what is new for them is only the place of their comparison regions. Let W be either a group of the deviation species of the step k which is carried, made at a step $j \le k$, or an interior group made at the step $j = k+1$ which has one interval and embeds in the torus. We say that W is *attached to the new region* if one of its comparison regions meets Ω_{k+1}^c. If it is not, it is of the deviation species at the step $k+1$ on the present term: for an interior group this is Proposition 6.31(iv), and for a carried group Propositions 6.22 and 6.27 apply with their proofs, which use the background of the present term through Supplement, Proposition S.8, and on the dependence set of the group, a subset of Ω_{k+1}. The attached groups have no deviation bound at the scale $k+1$, since the background is not regular at that scale on their comparison regions. They are small in another way.

Proposition 6.32 (Attachment to the New Region)**.** *On the present term, let $\mathfrak{n}$ be a traversal of the step $k+1$, and let d be its ℓ^∞-distance from Ω_{k+1}^c in the units of the scale $k+1$. Then*

$$\begin{aligned} \sum_{W \ni \mathfrak{n}} e^{\theta(W)+\lambda_c \nu \sigma(W)+\frac{3}{4}\nu_e \operatorname{ext}(W)} \|Y_W\|_{(k)} &\le L C_c \, \rho_k \, e^{-\frac{1}{4}\nu_e d/M} \,, \\ \sum_{\mathcal{G} \ni \mathfrak{n}} e^{\theta(\mathcal{G})+\nu \sigma_{k+1}(\mathcal{G})+\frac{3}{4}\nu_e \operatorname{ext}(\mathcal{G})} S_{\mathcal{G}} \, \ell_{\mathcal{G}} &\le 2 C_{S'} \rho_{k+1} \, e^{-\frac{1}{4}\nu_e d/M} \,, \end{aligned} \tag{6.83}$$

the first sum over the carried groups of the deviation species of the step k and the second over the interior groups made at the step $k+1$ with one interval which embed, in both cases those which are attached to the new region and have $\mathfrak{n}$ in the interval.

Proof. Let W be one of these groups, made at the step $j \le k+1$, and let $\mathfrak{n}_j \subset \mathfrak{n}$ be a traversal of the step j in its interval. By the definition of the extent (Section 5.3) the comparison regions of W lie in the ball of radius $\operatorname{ext}(W)M$ about $\mathfrak{n}_j$, in the units of the scale j, and one of them meets Ω_{k+1}^c. The distance of $\mathfrak{n}_j$ from Ω_{k+1}^c in these units is at least d, because $\mathfrak{n}_j \subset \mathfrak{n}$ and a length in the units of the scale j is not smaller than in those of the scale $k+1$. Hence $\operatorname{ext}(W) \ge d/M$, and a quarter of the weight of the extent pays the factor $e^{-\frac{1}{4}\nu_e d/M}$. The second bound now follows from the production bound (5.22) for the interior groups (Proposition 6.31(iii)). For the first, a carried group with $\mathfrak{n}$ in its interval has one of the L traversals of the step k in $\mathfrak{n}$ in its interval, and we sum the clause (5.21) of the step k over these. *q.e.d.*

Since U_{k+1} is a background of the step k, the first sum of (6.83) bounds the same sum with $\|Y_W(U_{k+1}, J_{k+1})\|$, and the second bounds the sum with the deviation parts at the background. Because $\Omega_{k+1}^c \subset \Lambda_{k+1}^c = Z_{k+1}$, both are of the form of the second term of (5.11) at the step $k+1$, with the rate $\frac{1}{4}\nu_e \ge c_{\text{att}}$. What they lack is a bound at the scale of the later steps: a traversal of a later step n contains L^{n-k-1} traversals of the step $k+1$ near the region, each with its attached groups, and the sum over them is controlled only in total, by the number of the traversals near the region. The groups at the collar are in the same position, with the outer field as a further argument; we turn to them now.

6.13 A New Large-Field Region: The Groups at the Collar

For a group at the collar nothing can be quoted from ℓ_0. Its factors may depend on the outer field; the potentials at the boundary of Λ_{k+1} are of the first order in the fluctuation field, not of the second [9, p. 277]; and the measure of the step is not even under the reflection of the integrated field, since it contains the term $\langle A, C^* \Delta^{(k)} C A_k \rangle$, so that the gain of Supplement, Lemma 17.28, is lost. A marked polymer at the collar is therefore of the order of the threshold ϑ_k of the fluctuation field, and not of the order of ρ_k. This is still a quantity which tends to zero with the coupling, and it is all that the count of Section 6.6 requires. The setting and the notation are those of Section 6.12.

The collar space. At the collar the configurations are those of two scales: on Ω_{k+1} the background is regular at the scale $k+1$, and outside it at the scale k. For a connected union X of cubes of π_{k+1} we call the *collar space of the step* $k+1$ *on* X the set of the pairs $(e^{i\alpha} U, \mathbf{J})$ on X with the following three properties. The configuration U is the restriction of the trivial configuration, of a background U_{k+1} of the present term at a block field and outer variables of the support, or of a gauge transform of one of these. The conditions (5.17) hold at $n = k+1$ on the bonds and plaquettes of Ω_{k+1}, and at $n = k$ with their right sides multiplied by $1 - \frac{1}{2}\beta$ on the others, as in [9, (2.34), (2.36), (2.39), p. 261]. And the restriction of the pair to every connected union $Y \subset X$ of cubes belongs to Bałaban's space $\tilde{U}^c_{k+1}(Y, \tilde{\alpha}_0, \tilde{\alpha}_1)$ of [9, p. 261] for the sequence of the domains of the present term. The third property places the pair in the spaces [9, (3.43), p. 276] on which the potentials of the measure of the step are analytic, and the second gives, with (6.6), that the interpolated configurations of Section 6.3 lie in the space of the step k, as was noted there for the bonds of $\Omega_k \setminus \Omega_{k+1}$. On a set $X \subset \Omega_{k+1}$ the second property is that of the space of the step $k+1$ on X.

Proposition 6.33 (The Size of a Marked Activity at the Collar)**.** *Suppose that the hypotheses of Supplement, Lemma 17.27, hold, together with* (6.6)*,* $b_k \le \frac{1}{2}\min(\varepsilon_H, r_\alpha)$*,* $4\bar{\Lambda} M c'_\pi \alpha_{1,k+1} \le L$*, and* $4\bar{\Lambda} M c'_\pi \alpha_{1,k} \le 1$*, and let* γ *be so small that the remark which follows that lemma applies. Then, for every marked polymer* M *of the present term, at every real outer field of the support of the test functions and at* $\sigma_k = 0$*, and at every configuration of the collar space on* M*,*

$$\|\tilde{H}(M)\| \le \Big(\frac{b_k}{\min(\varepsilon_H, r_\alpha)}\Big)^{n(M)} e^{-(\kappa_1 - 2 - c_\kappa)\#(G_M)}\, e^{-(L+2)\kappa \#(M)/1296} \prod_{W \in \mathfrak{B}(M)} \|Y_W\|_{(k)}\,, \quad (6.84)$$

with the notation of Proposition 6.9.

Proof. This is the proof of the second and of the third case of Proposition 6.9. The counts (6.23) do not involve the term. The bound of $\|\tilde{H}(M)\|$ by $\|\mathcal{X}_M\|_{\sup}$ is Supplement, Lemma 17.27, in the form of the remark which follows it, with the supremum over the fluctuation fields of the support, the outer field included, and over the collar space. For the factor we use Lemmata 6.5 and 6.7, which are stated for the present term. On the support $|B'|_Y \le b_k$ for every component, on the bonds of $\Omega_{k+1} \cap \Lambda^c_{k+1}$ as on those of Λ_{k+1}, because the bound $|g_k A_k| \le r_y \vartheta_k$ of Section 6.3 holds on every bond of the fluctuation field. A configuration $e^{i\eta a'} U$ of the collar space has $|a'(b)| \le \alpha_{1,k+1}/L$ on the fine bonds of Ω_{k+1} and $|a'(b)| \le \alpha_{1,k}$ on the others, in the units of the step k, so that the factor $\exp(4\bar{\Lambda} M c'_\pi \|a'\|_\infty \#(G))$ of these lemmata is at most $e^{\#(G)}$ by the last two hypotheses; and the configurations over which Lemma 6.7 takes the suprema of the insertions of the members lie in the space of the step k, by the second property of the collar space. The even and odd parts of the factor are not separated. *q.e.d.*

The count. We now run the count of Section 6.6 on the present term. The vertices of a group are those of that section, with two changes for the polymer vertices. A polymer vertex has a further datum, a finite set $\mathcal{K}'_v$ of clusters which are incompatible with M, the collar occurrences; their supports are added to its cube set and to its reach, as those of the clusters of $\mathcal{K}_v$ are, and its activity (6.36) has the further factor $\prod_{\mathcal{C}\in\mathcal{K}'_v}\xi'(\mathcal{C})$ with $\xi'(\mathcal{C}) := 2\bar\varphi(\mathcal{C})e^{2\bar\varphi(\mathcal{C})}$, where $\bar\varphi$ now denotes the sum of the bounds of Supplement, Lemma 17.24(iv), for the present term and for ℓ_0, so that $|z_{\mathcal{C}}| \le \xi'(\mathcal{C})$ in (6.82). And the number $\mathfrak{a}_v$ of a polymer vertex without members is $c_a\rho_k$ only if M has no cube in $\mathcal{N}_{k+1}$; otherwise it is $\varpi_k = b_k/\min(\varepsilon_H, r_\alpha)$, by (6.84). Put

$$E'_\alpha := C'_\alpha\varpi_k\ , \qquad C'_\alpha := 2C_{\text{root}}L^4e^{-\kappa'}\ . \tag{6.85}$$

Proposition 6.34 (The Count at a New Region)**.** *Suppose that the hypotheses of Theorem 6.18 and of Proposition 6.33 hold, and that* $E_\alpha+E'_\alpha+E_\beta+E_o \le \frac12\nu$. *Then the hypothesis of Proposition 6.11 holds on the present term, at every real outer field of the support, and for every cube* Δ *of* π_{k+1}

$$\begin{aligned}\sum_{\mathcal{G}\colon \Delta\subset X_{\mathcal{G}}} & e^{\frac52\nu\sigma_{k+1}(\mathcal{G})+\nu_e\,\mathrm{ext}(\mathcal{G})}P(\mathcal{G})\,\ell_{\mathcal{G}} \\ &\le \frac{e^{2\nu}}{\nu}\Big[E_\alpha+E'_\alpha+E_\beta+\frac{2}{\nu}(E_\alpha+E'_\alpha+E_\beta+E_o)E_o+\frac{2}{\nu}(E_\beta+E_o)(E_\alpha+E'_\alpha)\Big]\ ,\end{aligned} \tag{6.86}$$

the sum over the groups of the step $k+1$ *of the present term, other than those which consist of a single member, where* $P(\mathcal{G})$ *is the product* (6.38) *with the factors of the collar occurrences, the suprema being taken over the outer fields of the support and over the collar space on* $X_{\mathcal{G}}$.

Proof. We go through the proofs of Section 6.6 and state what changes. Lemma 6.13 and Lemma 6.14 concern the geometry of the constituents; the support of a collar occurrence of M shares a cube or a wall with M, like the support of a cluster of a link, and the two lemmata hold with the sums over $\mathcal{K}(M)$ extended to the collar occurrences. In the proof of Lemma 6.16 the factors of the collar occurrences are bounded by the numbers $\xi'(\mathcal{C})$, and the marked activities by Proposition 6.9 for the polymers without a cube in $\mathcal{N}_{k+1}$, to which Lemma 6.29(ii) applies, and by (6.84) for the others; the bound $|\Psi^0_\circ(M)| \le c\,\#(M)$ is that of ℓ_0. In the proof of Proposition 6.17 the parts (i)–(iii) concern the groups of the step k, whose term has not changed. In (iv) and (v) the sum over the sets $\mathcal{K}'_v$ of the product of the numbers $\xi'(\mathcal{C})e^{N(\mathcal{C})/800} \le 2e\,\bar\varphi(\mathcal{C})e^{d(\mathcal{C})+N(\mathcal{C})}$ is at most $\exp(4e\,c\,\#(M))$, by Supplement, Lemma 17.24(iv), on the two terms, and this factor is absorbed as the factor $\exp(2e\,c\,\#(M))$ of the links is, because $(6e+1)c+\frac{1}{800} \le 1$. Part (v) is unchanged, and part (iv) holds as stated for the polymer vertices without members whose polymer has no cube in $\mathcal{N}_{k+1}$, while the same computation with ϖ_k in the place of $c_a\rho_k$ gives $\sum_{v\text{ at }\Delta} w(v)e^{2b(v)} \le E'_\alpha$ for those whose polymer has one. The proof of Theorem 6.18 then applies with $E_\alpha + E'_\alpha$ in the place of E_α for the roots of the third kind, and gives (6.86) and the convergence. *q.e.d.*

The new term E'_α is of the order of ϑ_k, and it is the largest of the five: by (6.46) the others are of the order of ρ_k or smaller. For γ_q small the right side of (6.86) is therefore at most $(4e^{2\nu}/\nu)E'_\alpha$.

Proposition 6.35 (The Groups at the Collar)**.** *Under the hypotheses of Proposition 6.34, with* $\mathsf{E}_k \le C_{\mathsf{E}}\rho_k$ *and* γ_q *sufficiently small, the following hold on the present term.*

(i) $\langle F\rangle$ *is the sum* (6.29) *over the finite sets of pairwise compatible groups of the step* $k+1$ *of the present term.*

(ii) The insertion of a group $\mathcal{G}$ at the collar is, at every real outer field of the support and at $\sigma_k = 0$, a holomorphic function of the restriction of $(\mathbf{U}, \mathbf{J})$ to $X_{\mathcal{G}}$ on the collar space on $X_{\mathcal{G}}$, and for every G-valued gauge transformation v each of its factors satisfies the law (5.19) *with $R(v)\sigma_k$ in the place of σ_k on the left side.*

(iii) For a group $\mathcal{G}$ at the collar let $S_{\mathcal{G}}$ be the supremum of $\|Y_{\mathcal{G}}\|$ over the outer fields and the configurations of (ii). Then, for every traversal $\mathfrak{n}$ of the step $k+1$,

$$\sum_{\mathcal{G} \ni \mathfrak{n}} e^{\theta(\mathcal{G}) + \nu\sigma_{k+1}(\mathcal{G}) + \frac{3}{4}\nu_e \operatorname{ext}(\mathcal{G})} S_{\mathcal{G}}\, \ell_{\mathcal{G}} \le C_{\text{coll}}\, \vartheta_k\, e^{-\frac{1}{4}\nu_e \operatorname{dist}(\mathfrak{n}, \mathcal{N}_{k+1})/M} ,$$
$$C_{\text{coll}} := \frac{8 e^{2\nu} C'_\alpha C_1 r_y}{\nu \min(\varepsilon_H, r_\alpha)} , \tag{6.87}$$

the sum over the groups at the collar with $\mathfrak{n}$ in an interval, for all numbers $0 \le \theta(\mathcal{G}) \le \nu\sigma_{k+1}(\mathcal{G})/16$, the distance being measured in the units of the scale $k+1$.

Proof. (i) is Proposition 6.11 on the present term (Section 6.12), whose hypothesis holds by Proposition 6.34.

(ii) The proof of Lemma 6.19 applies, with the collar space in the place of the space of the step. For the analyticity, the operators, the potentials, and the activities of the measure [9, (3.27)] are analytic on the spaces [9, (3.43)], which contain the collar space by its third property, and the insertions of the members are evaluated in the space of the step k, by its second property. For the covariance, the outer field is the restriction of the fluctuation field A_k to bonds of Ω_{k+1}, and it is transformed with it: under a gauge transformation v of the block field the variable B' becomes $R(v)B'$ [8, p. 265], on the bonds of Λ_{k+1} and on the others. The substitution of $R(v)B$ for the integration variable in the integrals which define the activities is then made for the field A on Λ_{k+1}, with $R(v)\sigma_k$ in the place of σ_k; the Gaussian form $\langle A, C^* \Delta^{(k)} C A_k \rangle$ and the potentials are invariant under the simultaneous transformations, the boundary terms by [9, (2.41)(iii), p. 261], and the rest of the proof is unchanged.

(iii) A group at the collar which has $\mathfrak{n}$ in an interval has a cube of $\mathcal{N}_{k+1}$ in its domain, and its domain lies in the ball of radius $\operatorname{ext}(\mathcal{G})M$ about $\mathfrak{n}$, so that $\operatorname{ext}(\mathcal{G}) \ge \operatorname{dist}(\mathfrak{n}, \mathcal{N}_{k+1})/M$, and a quarter of the weight of the extent pays the last factor of (6.87). Nothing is subtracted from the insertion of such a group, so that $S_{\mathcal{G}} \le P(\mathcal{G})$ by (6.39), and $\theta(\mathcal{G}) + \nu\sigma_{k+1}(\mathcal{G}) \le \frac{5}{2}\nu\sigma_{k+1}(\mathcal{G})$. The domain $X_{\mathcal{G}}$ contains one of the at most two cubes which contain an end of $\mathfrak{n}$. The left side of (6.87) is therefore at most $e^{-\frac{1}{4}\nu_e \operatorname{dist}(\mathfrak{n}, \mathcal{N}_{k+1})/M}$ times twice the right side of (6.86), which is at most $(4e^{2\nu}/\nu)E'_\alpha$ for γ_q small, as was noted after that proposition; and $E'_\alpha = C'_\alpha C_1 r_y \vartheta_k / \min(\varepsilon_H, r_\alpha)$, because $b_k = C_1 r_y \vartheta_k$. *q.e.d.*

A group at the collar is not split: its insertion is kept whole, with the number 0 in the place of a flat value, as that of a cross group is (Section 6.7), while an interior group is split as on ℓ_0, with the same flat value, whether or not it is attached to the new region. The reference gas of the present term is therefore the gas of its interior groups which lie on one loop, with the activities of ℓ_0, and the numbers $\theta(\mathcal{G})$ of clause (i) of Definition 5.2 are bounded for all the groups of the present term as in Section 6.7, by (6.48) for the interior groups. The bound (6.87) is the third term of (5.11) at the step $k+1$, with the mark 1 for every cube of $\mathcal{N}_{k+1}$: its right side is at most $C_{\text{coll}} \vartheta_k \sum_{\Delta \subset \mathcal{N}_{k+1}} e^{-\frac{1}{4}\nu_e \operatorname{dist}(\Delta, \mathfrak{n})/M}$, and $\vartheta_k \le g_k r_k \le (1+\beta_0)^2 g_{k+1} r_{k+1} \le (1+\beta_0)^2 \rho_{t,k+1}/(2C_t)$ by (3.17), (3.9), (3.10), and (5.6). Like the groups attached to the new region, the groups at the collar are bounded per traversal of

the step at which they are made. Their totals over the components of Λ_{k+1}^c, their dependence on a complex outer field, and their fate at the later steps are treated in Section 6.15.

6.14 The Step on a Term with Earlier Large-Field Regions

We now let the labelled term of ϱ_k have large-field regions of its own, so that Λ_k is not the whole torus. Three things are then new. The operators of the step are those of the sequence of the domains of the term [9, p. 276]. The term carries groups which earlier steps made at their large-field regions, and their insertions depend on the outer fields of those steps, which the gauge transformation of the present step rotates. And these groups are not few: a traversal of the step k which runs along the large-field region of a step $j < k$ contains L^{k-j} traversals of the step j, each with groups of the size of ϑ_{j-1} (Proposition 6.35), so that their sum over a traversal of the present step is not small, and the count of Section 6.6, in which the members at a cube are summed, cannot contain them. The resolution has two parts. A group of the step which contains none of them and stays away from the regions is, factor by factor, a group of ℓ_0, as in Section 6.12. And the others are never summed over the loop: in every term of the expansion they are fixed data, the step reaches them only across the distance of Lemma 6.28(iii), and their total grows at the step by a factor $1 + \delta_k$ with $\sum_k \delta_k < \infty$.

Regular and attached groups. Consider a labelled term of ϱ_n, and for $j \le n$ let $\mathcal{N}_j$ be the union of the cubes of π_j which lie in Λ_j^c or have a point in common with a cube of Λ_j^c, as in Section 6.12. We divide the groups of the term into regular and attached ones, by induction on the step at which they are made. A group made at a step $j \ge 2$ is *interior* if no cube of its domain belongs to $\mathcal{N}_j$ and every member of the step $j-1$ among its constituents is interior; at $j = 1$ the interior groups are those which are not attached to the large-field region of the first step in the sense of Definition 5.2(ii). For a group which the **R**-operation of a step j makes (Section 6.15) the condition on the domain is stronger: none of its cubes lies within six layers of cubes of the large-field region which the operation leaves, and its members, which are groups of the step j, are interior. Whether a group is interior is thus decided by the regions of the term within six cubes of the domains of the group and of its members of all generations. An interior group is *regular at the step* n if every member among its constituents is regular at the step $j-1$ and if it is a cross group, or does not embed in the torus, or embeds and has its comparison regions in Ω_m for every m with $j \le m \le n$. Every other group is *attached at the step* n, and it is then attached at all later steps; on the term without large-field regions every group is regular. The interior groups which lie on one loop are split, with the flat values which they have on ℓ_0 (Proposition 6.36(ii)), whether or not they are regular, and they form the reference gas of the term; the groups which are not interior are not split. An attached group is thus of one of three kinds: its domain has a cube of $\mathcal{N}_j$ at its production (Definition 6.30); or it is interior and one of its comparison regions has left the small-field region of a step m (Proposition 6.32); or one of its members is attached. In the first two cases we call it an *atom*, made at the step j or m at which the condition first holds, and we assign it to a component Z of the large-field region Λ_j^c, respectively Λ_m^c, of that step: the first, in a fixed order, among those which contain a cube of its domain or touch one, or which contain the component of Ω_m^c that the comparison region meets. A group with attached members belongs to the *species* of the first of them, in a fixed order, and the species of an atom is the pair (j, Z).

The outer fields. For $j < n$ let σ_j be the restriction of the fluctuation field A_j of the step $j \to j+1$ to the bonds of $\Omega_{j+1} \cap \Lambda_{j+1}^c$, the outer field of that step (Section 6.12; for $j = 0$ the field outside the region of integration of the first step, Supplement, Proposition S.7). These

fields are among the variables which $\mathbf{T}_n$ holds pending, and we write $\omega = (\sigma_j)_{j<n}$. A gauge transformation v acts on them by the adjoint representation, $(R(v)\sigma)(b) = v(b_-)\sigma(b)v(b_-)^{-1}$, as on the field A of Bałaban's boundary terms [9, (2.41)(iii), p. 261]. The insertion of an attached group is a function $Y_W(\mathbf{U}, \mathbf{J}, \omega)$ of the configuration, of the current, and of the outer fields on its dependence set, and its covariance law is (5.19) with $R(v)\omega$ on the left side (Proposition 6.35(ii)). On the support of the term $g_j|\sigma_j| \le r_y\vartheta_j$ (Section 6.3). A later step evaluates such an insertion at rotated outer fields, which are complex when the parameters of the localisation are, and the class therefore records its analyticity on polydiscs about the real outer fields, whose radii shrink from step to step as the analyticity radii of [9, (2.34)–(2.39), p. 261] do:

$$r_j^{(n)} \coloneqq g_j^{1/2}(\tfrac{1}{2} + 2^{-(n-j)}) \quad (j \ge 1) , \qquad r_0^{(n)} \coloneqq \tfrac{1}{2} + 2^{-n} . \tag{6.88}$$

The *attached space of the step n* is the set of the triples $(e^{i\alpha}U, \mathbf{J}, \omega)$ with the following properties, and of their transforms by G-valued gauge transformations. The configuration U is the background U_n of the term at a point of its support, with the outer fields ω_0 of that point, or the trivial configuration with $\omega_0 = 0$. The field α satisfies the first two conditions of (5.17) at n on every fine bond, on the earlier layers as on Ω_n, and the restriction of the pair $(e^{i\alpha}U, \mathbf{J})$ to every connected union Y of cubes belongs to Bałaban's space $\tilde{U}_n^c(Y, \tilde{\alpha}_0, \tilde{\alpha}_1)$ of [9, p. 261]. And $|\sigma_j(b) - \sigma_{j,0}(b)| < r_j^{(n)}$ for every $j < n$ and every bond. At its points the transport along a traversal of the step n has norm at most $e^{c'_\pi \alpha_{1,n}}$ on every layer, which is why the complex part is restricted to the radius of the top layer. The attached spaces decrease with n: the background U_{n+1} is a background of the step n (Supplement, Proposition S.8), $\alpha_{1,n+1}/L \le \frac{1}{2}\alpha_{1,n}$, and the radii (6.88) decrease. For an attached group W of the species (j, Z) let $\|Y_W\|_{[n]}$ be the supremum of the norm of its insertion over the attached space of the step n, let $\mathbf{Coll}_j$ be the collar of Lemma 6.28(iii), and let $\mathrm{ext}_{\mathrm{c}}^{(n)}(W)$ be the largest distance, in sides of the cubes of π_n, from a cube of π_n which meets the dependence set of W to $\mathbf{Coll}_j$. Its *weighted mass* at the step n is

$$\mu_n(W) \coloneqq e^{c_{\mathrm{sh}}}\, e^{\theta(W) + \nu\sigma(W) + \frac{3}{8}\nu_e \,\mathrm{ext}_{\mathrm{c}}^{(n)}(W)} \|Y_W\|_{[n]} , \qquad c_{\mathrm{sh}} \coloneqq \tfrac{1}{60} , \tag{6.89}$$

with the size $\sigma(W)$ of the step at which W was made; the constant c_{sh} is that of (6.44), the part of the share of a member in a new group which is not proportional to its size. For a component Z of Λ_j^c let $A_j(Z)$ be the sum of $\mu_j(D)$ over the atoms D of the species (j, Z).

The gauge transformation of the step. For real fluctuation fields of the support the background of the step k at the block field $e^{iB'}V^{(k)}$ is the transform of $e^{i\eta\mathbf{H}_k(B')}U_{k+1}$ by a gauge transformation w^{-1} [9, (3.6), p. 266] [8, p. 265], so that by the covariance of its factors the word of a family is the same word at the configuration $e^{i\eta\mathbf{H}_k(B')}U_{k+1}$ and at the outer fields $R(w)\omega$. For the regular groups, which do not depend on ω, the transformation drops out, as in Section 6.3. For the attached ones it does not, and three of its properties are needed (Supplement, Proposition S.9(vi)). It is local: its value at a site is a function $W_y(\mathbf{H}, U_{k+1})$ of the restrictions of $\mathbf{H} = \mathbf{H}_k(B')$ and of U_{k+1} to the block $B^m(y)$ of the layer $\Omega_m \setminus \Omega_{m+1}$ which contains the site, and $W_y = 1$ if $\mathbf{H}$ vanishes on the block. It is analytic in $\mathbf{H}$, so that $w(s) \coloneqq W(\mathbf{H}(s, B'), U_{k+1})$ is defined for the complex parameters of Section 6.3. And it is small on the earlier layers. The parameters and the scaled fluctuation fields at which the factors of the words are evaluated are those of Lemma 6.7, $|s(\Delta)| \le e^{\kappa_1}$ and $|t_Y||B'|_Y \le \min(\varepsilon_H, r_\alpha)$, for which $|\mathbf{H}(s, tB')| \le \hat{h}_k \coloneqq 4B_0 e^{16\kappa_1} r_\alpha$ by (6.4). With constants C'_w and c_G which depend on d, L, and G only, at these arguments and at the sites of the bonds of σ_j, $j < k$,

$$|\log w| \le C'_w\, 4^{-(k-j-1)}\hat{h}_k , \qquad |R(w)\sigma - \sigma| \le (e^{2c_G|\log w|} - 1)|\sigma| , \tag{6.90}$$

the second bond by bond; the factor $4^{-(k-j-1)}$ comes from the decay of $\mathbf{H}_k$ across the layers $\Omega_m \setminus \Omega_{m+1}$, $j + 1 < m \le k$, which separate the fluctuation field from the bonds of σ_j.

Consequently $R(w)$ maps the polydisc of the radius $r_j^{(k+1)}$ about a real outer field $\sigma_{j,0}$ of the support into the polydisc of the radius $r_j^{(k)}$, provided that

$$16\, c_G C'_w\, 2^{-(k-j-1)} \hat{h}_k \Big(\frac{r_y \vartheta_j}{g_j\, r_j^{(j+1)}} + 1 \Big) \le 1 \tag{6.91}$$

for all $0 \le j < k \le K'$. Indeed, the two radii differ by $2^{-(k+1-j)} r_j^{(j+1)}$, a point of the smaller polydisc has $|\sigma_j| \le |\sigma_{j,0}| + r_j^{(j+1)}$, and $e^x - 1 \le 2x$ for $x \le 1$. The condition holds for γ small. By (6.12), $\hat{h}_k \le \alpha_{1,k}$, a constant times $g_k (\log \hat{g}_k^{-2})^{q_1}$. For $j \ge 1$, $\vartheta_j \le g_j r_j$ by (3.17), $g_j^{-1/2} \le (1 + \gamma^2 \beta'(k-j))^{1/4} g_k^{-1/2}$ by (3.9), and the power of $\log g_j^{-2}$ in r_j is at most $(1 + \gamma^2 \beta'(k-j))^{\beta_0}$ times the same power of $\log g_k^{-2}$ by (3.10). The left side is therefore at most a constant times $g_k^{1/2} (\log g_k^{-2})^{q_1 + p_0}\, 2^{-(k-j)} (1 + \beta'(k-j))^{\beta_0 + 1/4}$, in which the last two factors are bounded uniformly and the others tend to zero with γ; for $j = 0$ the same holds with r_0 in the place of $r_j g_j^{-1/2}$.

Proposition 6.36 (The Step on a Term with Earlier Large-Field Regions)**.** *Let the ensemble be dressed at the steps $1, \dots, k$, suppose the hypotheses of Proposition 6.35 and of Supplement, Proposition S.9, together with* (6.91)*, and let γ_q be sufficiently small. On every labelled term of ϱ_{k+1} the following hold.*

(i) The expansion. *$\langle F \rangle$ is the sum* (6.29) *over the finite sets of pairwise compatible groups of the step $k+1$, which are defined as in Section 6.12 from the expansion of the words at the configurations $e^{i\eta \mathbf{H}(s,B')} U_{k+1}$ and the outer fields $R(w(s))\omega$; the sum converges absolutely.*

(ii) The interior groups. *The data of an interior group made at the step $k+1$ define a group of the step $k+1$ on ℓ_0, with the same domain, intervals, size, and extent, and its insertion is the same function, with the same flat value; if its members are regular, the statements (ii)–(iv) of Proposition 6.31 hold for it. Consequently the regular groups of the term at the step $k+1$ are among the groups of ℓ_0 with the same insertions, and they satisfy* (5.13), (5.14), (5.21), (5.22), *and* (5.23) *at the step $k+1$ with the constants of ℓ_0.*

(iii) The atoms. *The insertion of an atom made at the step $k+1$ is holomorphic on the attached space of the step $k+1$, restricted to its dependence set, and covariant, with $R(v)\omega$; the bounds* (6.83) *and* (6.87) *hold for the atoms made at the step $k+1$ with the suprema over that space, and with $2C_{\mathrm{coll}}$ in the place of C_{coll}.*

(iv) The carry. *At the points at which the step evaluates the insertion of an attached member D of the step k, the configuration and the outer fields belong to the attached space of the step k, so that $\|Y_D\|_{[k+1]} \le \|Y_D\|_{[k]}$ and $\mu_{k+1}(D) \le \mu_k(D)$ for a carried member. The sum, over the finite sets of pairwise compatible attached groups of the term at the step $k+1$, of the products of their weighted masses μ_{k+1} is at most*

$$\prod_D \Big(1 + \mu_j(D)\,(1+\epsilon_j) \prod_{m=j}^{k} (1+\delta_m) \Big) \le \exp\Big(2 \sum_{(j,Z)} A_j(Z) \Big)\,, \tag{6.92}$$

the product over the atoms D of the term, $j \le k+1$ being the step at which D is made, and the sum over the species, with the numbers ϵ_j and δ_m of Supplement, Proposition S.9, which satisfy $(1+\epsilon_j) \prod_{m \le K'} (1+\delta_m) \le 2$ for every j. The groups which

the **R***-operation of an earlier step* $n \le k$ *has left bound to components of the second class (Section 6.15) are not among these atoms; with them both sides of* (6.92) *have the further factor* $\prod b_n$*, one factor* $b_n := \exp(4\nu N_n + C_x)$*,* $N_n := (107R_n)^4$*, for each such step* n *and each component of the second class of the term at that step, with a constant* C_x *of the construction.*

Proof. (i) We go through Sections 6.3–6.5 and state what changes on a term with earlier large-field regions. The partition of the parameters consists of the cubes "in proper scales", of the side M of the scale j in $\Omega_j \setminus \Omega_{j+1}$, and the propagators are those of the random-walk expansions for the sequence of the domains; with these Bałaban repeats the localisation, and "the remaining considerations and bounds are the same" [9, p. 276], so that the properties (a)–(c) of the parameters hold, the bound (6.4) in the norms of each layer. The word of a family is evaluated at $(e^{i\eta \mathbf{H}(s,B')}U_{k+1}, R(w(s))\omega)$, with the argument of Section 6.3 in the place of $\mathbf{J}$; at $s = 1$ this is the word at the configuration of integration, by the identity which precedes (6.90), and at $s = 0$ it is the word at (U_{k+1}, ω), because $w = 1$ where $\mathbf{H}$ vanishes. If s vanishes outside a set σ, then $w(s) = 1$ on every block which does not meet a component of $Y(\sigma)$, by the locality of w. We therefore say that a component touches an attached member W if its interior meets the dependence set of W or a block which meets that set; a block of a layer $\Omega_m \setminus \Omega_{m+1}$, $m \le k$, has at most the side of a block of the scale k, which is less than the side of a cube of π_k, and we add these blocks to the dependence set of an attached group. With this definition Lemma 6.4 holds with its proof. The factors are evaluated in their domains of analyticity: for the regular members as in Section 6.3, and for the attached ones by (iv). Proposition 6.8 is stated for the measure [9, (3.27)], and its proof applies to every term, the expansion being that of [9, pp. 275–279]; the bounds of the activities are those of Supplement, Lemmata 17.24 and 17.27, on the strength of Bałaban's statement which was quoted in Section 6.12. The groups of the step and their compatibility are those of Definition 6.10 with the collar occurrences of Section 6.12, the reference gas of the term of ϱ_k being the gas of its interior groups on one loop, a part of the gas of ℓ_0 with the same activities. Proposition 6.11 holds with its proof once the convergence is known. For the terms without attached members the convergence is that of Proposition 6.34, whose proof uses the members through their sum E_k alone, which the regular members satisfy by (ii) at the step k. For the others it follows from (6.92): the sum over the terms with a given family of attached members is at most the product of their weighted masses, of the factors $1 + \delta_k$, and of the convergent sum over the sets of groups without attached members, and the sum of the weighted masses of the attached groups of the term of ϱ_k is finite by (6.92) at the step k.

(ii) The proof is that of Lemma 6.29 and of Proposition 6.31, with three additions. First, the members of an interior group are interior, hence groups of ℓ_0 with the same insertions, by (ii) at the step k, and the groups of its old occurrences are groups of the reference gas of ℓ_0; where Proposition 6.31 bounds the members through their sum E_k, the members are assumed regular. Secondly, the operators with vanishing parameters outside a union M of cubes none of which belongs to $\mathcal{N}_{k+1}$ are sums over the walks which stay in $M \subset \Lambda_{k+1} \subset \Omega_k$, and for these walks the operators of two sequences of domains coincide: "in the difference all terms for walks with localizations contained in Ω are cancelled" [5, Theorem 3.14 and its proof, pp. 426–427], with Ω the common domain; and the terms of the action with localisation domains in Λ_{k+1} "satisfy exactly all the conditions" of the lemmata of the term without large-field regions [9, p. 276] and "coincide with the corresponding terms arising from the expressions defined on the whole lattice" [9, p. 278]. Thirdly, the boundary terms $\mathbf{B}_k$ and the term $A(g_k^{-2}(\cdot) - g_k^{-2}, \cdot)$ are now present, but their localisation domains meet $Z_k^\sim$ [9, (2.41),

p. 261], respectively Z_k [9, p. 277], hence $\Lambda_k^c \subset \Lambda_{k+1}^c$, and none of them lies in M. The second statement follows by induction on the step: a regular group is interior at its production, and the clauses concerned are those of ℓ_0 for the same functions, with the suprema over the space of the step on their domains, on which the background of the term has the regularity of that step (Proposition 6.31(ii)), and with the deviations taken over comparison regions which lie in the small-field regions.

(iii) The atoms made at the step $k+1$ are the groups at the collar without attached members, the interior groups one of whose comparison regions meets Ω_{k+1}^c, and the carried regular groups of which the same is true. For the last two the bounds are Proposition 6.32, whose proof uses (5.22) and (5.21) for regular groups, which hold by (ii); these insertions do not depend on the outer fields. For the groups at the collar the proofs of Propositions 6.33, 6.34, and 6.35 apply, with the regular members in the place of the members, and it remains to pass from the real outer field σ_k to the polydisc of the radius $r_k^{(k+1)} = g_k^{1/2}$ about it. The outer field enters every factor through the variable B' and through entire quadratic forms; on the polydisc $|B'| \le 2b_k$, the potentials of the measure [9, (3.27)] whose localisation domains are not contained in Λ_{k+1} are analytic in σ_k, and their bounds are at most twice those at real outer fields (Supplement, Proposition S.9(vii)). The expansions use the potentials through their bounds and their localisation domains alone [10, (2.1)–(2.5), p. 12], so that the three propositions hold on the polydisc with $2b_k$ in the place of b_k, which doubles ϖ_k and C_{coll}.

(iv) Let $(e^{i\eta a'}U_{k+1}, \mathbf{J}, \omega)$ be a point of the attached space of the step $k+1$. The points of evaluation are $(e^{i\eta \mathbf{H}(s,tB')}e^{i\eta a'}U_{k+1}, R(w(s))\omega)$ with the corresponding argument in the place of $\mathbf{J}$, for $|s(\Delta)| \le e^{\kappa_1}$ and $|t_Y||B'|_Y \le \min(\varepsilon_H, r_\alpha)$. The configuration and the current belong to the attached space of the step k by the argument which gave (6.6): the complex part grows by at most $c_{\mathbf{J}} \cdot 4B_0 e^{16\kappa_1} r_\alpha \le \frac{1}{4}\min(\beta,1)\min(\alpha_{0,k}, \alpha_{1,k})$ in the norms of the step k, which is less than the difference of the radii of the top layers, and on the earlier layers the difference of the radii of Bałaban's spaces of the two steps is "still much greater than bounds on this function" [9, p. 277]. The outer fields belong to it by (6.90) and (6.91). The norm of the insertion of D at these points is therefore at most $\|Y_D\|_{[k]}$, and the extent $\mathrm{ext}_{\mathrm{c}}^{(n)}(D)$ does not increase with n. The bound (6.92) is Supplement, Proposition S.9(ii), which is proved there by the count of Section 6.6 with the attached members as fixed vertices; the atoms of the step $k+1$ enter it at $j = k+1$ with their weighted masses, which are bounded by (iii). *q.e.d.*

By (ii) the bounds of the class for the regular groups of every term are those of ℓ_0, and by (iv) the attached groups enter the bound of a term at the stopping step through the numbers $A_j(Z)$ alone: the sum over the families of a term is at most the product of the factors $1 + e^{\theta(W)}\|Y_W\|$ of its regular groups and of the left side of (6.92), because a family consists of a family of regular groups and of a family of attached ones, and $e^{\theta(W)}\|Y_W\| \le \mu_n(W)$ at the real points of the support.

6.15 Large Fields and the R-Operation

It remains to bound the numbers $A_j(Z)$ and to say what pays for them. The attached insertions carry no deviation, and nothing contracts them; their total over each component Z of the large-field region of the step j is paid once, by the small factor which Bałaban's large-field estimates attach to Z.

The payment. Bałaban attaches to every large-field region Z created at the step j the factor $\exp(-\frac{1}{2}\gamma_0 A_1^2 p_0^2(g_j)(d'_j(Z)+1) - 2p_0(g_j))$, with $d'_j(Z)$ the linear size of Z, "the fundamental large field factors, which control convergence of the expansion of the effective densities" [12,

p. 381], and he shows that its part $\exp(-\kappa_j(Z))$ "controls K renormalization steps, under the assumption that no large fields are created in these steps", in the sense of the inequality [12, (1.80), p. 384]. Two costs have to be paid from it. The first is the total of the weighted masses of the atoms of the species (j, Z),

$$A_j(Z) \le C\hat\rho_j N_j(Z) + C\rho_{t,j}\mathfrak{m}_j(Z) \le C_A(d'_j(Z)+1)\,, \tag{6.93}$$

where $N_j(Z) = \sum_{\mathfrak{n}_j} e^{-c_{\mathrm{att}} d_j^Z(\mathfrak{n}_j)/M}$ counts the traversals of the step j near Z, which by Lemma 5.5(i) are at most $C\Lambda(C)$ per cube of Z, and $\mathfrak{m}_j(Z)$ is the sum of the marks of the data assigned to Z, at most a multiple of the number of cubes of Z; the first term bounds the atoms of Proposition 6.32 and the second those of Proposition 6.35, with the weights of (6.89), since $\frac{3}{8}\nu_e \operatorname{ext}_{\mathfrak{c}}^{(j)} \le \frac{3}{4}\nu_e \operatorname{ext} + \frac{3}{8}\nu_e$ for an atom (Supplement, Proposition S.9(i)). By (6.92) this total enters the bound of a term through the factor $\exp(2A_j(Z))$, at every later step. The second cost recurs at every step n of the lifetime of Z: the reference gas of a term does not contain the groups at its large-field regions, and the difference $E_n^{\partial}(\ell)$ between the logarithms of the reference gases of ℓ and of ℓ_0 is a sum of local terms at the traversals near the region (Supplement, Proposition S.9(iii)), each of which costs at most $C_e\hat\tau_n^2$, $\hat\tau_n^2 \coloneqq \max(\tau_n^2, C\hat\rho_n)$, with a constant C_e of the construction. By Lemma 5.5(i) a cube of side MR_n meets at most $4\Lambda(C)MR_n$ traversals, and along Bałaban's chain for the regions $S^{n-j}(Z)$ which descend from Z [12, pp. 384–385], whose numbers of cubes are at most $C2^{-(n-j)}d'_j(Z)$ until the region fits into a cube of size $64MR_n$, after which at most R_j steps remain, each with a region of at most 64^d cubes,

$$D_j(Z) \coloneqq \sum_n 4C_e\Lambda(C)MR_n\hat\tau_n^2\,\#\mathsf{cubes}_n(S^{n-j}(Z)) \le C_d\Lambda(C)MLR_j^2\,\bar\tau^2\,(d'_j(Z)+1)\,, \tag{6.94}$$

with $\bar\tau^2 \coloneqq \sup_{n\le K'}\hat\tau_n^2 \le C\gamma_q^2(\log\gamma_q^{-2})^{6p_0}$, by $g_n \le \gamma_q$ for $n \le K'$ from (3.46), the sum running over the steps $n \le K'$ at which the loops are carried, and $R_n \le LR_j$ for $n \ge j$ by Lemma 3.4. Both costs are at most a polylogarithm of g_j^{-2} of fixed degree times $d'_j(Z)+1$, while $\kappa_j(Z)$ is of order $(\log g_j^{-2})^{2p_0}(d'_j(Z)+1)$, so that for p_0 large and γ_q small, depending on $c_{\mathcal{C}}$, the factor of Z pays both and leaves a reserve, since $j \le K'$ and $\log g_j^{-2} \ge \log\gamma_q^{-2}$; Bałaban's induction over the merging of regions [12, pp. 384–389] runs with the summand of [12, (1.80)] enlarged by these costs, since both are subadditive under the union of regions up to the slack $2d$ of [12, p. 386]. That the costs fit into Bałaban's induction, with a reserve, is Supplement, Proposition S.9(iv).

The **R***-operation.* From the step at which the depth condition (N1) first admits a component, the **R**-operation acts on the numerator. It integrates a term over the block field on a domain inside a small and old large-field component and replaces it by the normalised fraction of [11, (1.100), p. 201]; since the fraction integrates to one over the variables which it replaces, the operation has the normalisation property [11, (1.102), p. 201], $\int(\mathbf{R}'f)\,\mathrm{d}V_k = \int f\,\mathrm{d}V_k$, for every integrable f and not only for the density; the removal of the δ-functions on the components of the first class [12, p. 378] is an equivalence which needs the invariance of the integrand under gauge transformations, and the dressed loop is gauge invariant. On the components of the first class the term is expanded by Bałaban's localisation expansion [12, pp. 372–376, (1.66)–(1.67)], which he makes for "any term in the sum" and which applies to the words of the loop, entire functions of their arguments. The part of the resulting polymer gas which is linked to the loop is written exactly as a ratio of two polymer gases, and at flat reference data each group of polymers which interleaves with the loop contracts to a scalar by Schur's lemma, as in Section 6.7. The flat part of the ratio is therefore the cluster series of a one-dimensional gas along each loop; its perimeter and corner parts are added to the step factor κ_k (Definition 7.8), and its remainder $E_k^{\circ,R}$, bounded in Proposition 7.10, stays in $\mathcal{E}_k$.

Two kinds of factors of the term are not numbers of the term after the operation, and both are taken into the polymers. The local terms of the differences $E_n^{\partial}(\ell)$, $n \le k$, which read the labels of a component of the first class are summed by the operation of that component [12, p. 378]; each of them is written as $e^{eX} = 1 + (e^{eX} - 1)$, and the second summand becomes a factor of the polymer of the component, paid from the reserve which the component keeps. The terms of the clusters of the two gases depend on the background field, through the activities of Bałaban's polymers; each is split into its value at flat reference data, which stays in the gas, and a remainder, which is expanded in the same way and joins the group of polymers to which its cluster is attached, or forms a group of its own on the loop. The flat values are split off only from the clusters and from the groups of polymers which keep a distance of more than six layers of cubes from the components which the operation leaves in place; at that distance the activities are those of the term without large-field regions (Supplement, Proposition S.10(iv)), so that the gas of the flat values is a part of the gas of ℓ_0 with the same activities, as at the other steps. The groups so formed are the groups which the operation makes. Their sizes are counted in the cubes of π_k, as the class requires, with the unit constant $u_\star$ of Appendix A, and with these sizes they satisfy the clauses (i) and (vii) of Definition 5.2 and the bounds (5.22) and (5.21), with $C_{\mathbf{R}} e^{-p_0(g_k)/4}$ in the places of $2C_{S'}\rho_k$ and $C_c\rho_k$ and a constant $C_{\mathbf{R}}$ of the construction. Those which lie on one loop, meet no component which the operation leaves in place, have regular members only, and have their comparison regions in the small-field region are of the deviation species: each vanishes at pure-gauge data on its domain, and the argument of Proposition 6.21 gives (5.13) and the tail clause (5.14) for them, with a constant which is at most $C^{\mathbf{R}}(g_k) := \exp(-A_0(\log g_k^{-2})^{p_0}/4 + \log g_k^{-2} + C \log\log g_k^{-2})$ and which is added to C_{new} in (6.56). The other groups are attached, or of the cross or of the wrapping species. An attached group which contains an attached member belongs to the species of the first such member, and the further factors which the atoms receive at these steps have a product of at most $\frac{16}{15}$ (Supplement, Propositions S.9(ii) and S.10(ix)). The components of the first class leave the large-field region. Those which the operation leaves in place are the unchanged ones and the components of the second class, which lie in cubes of side $104MR_k$ and may lie inside the polymers. A group which comes within six layers of cubes of a component left in place, without containing one, is an atom of the step k, and the atoms of this kind satisfy (6.93) with a constant which tends to zero with γ_q. The polymers and the clusters which contain a component of the second class, and the parts of the letters of the loops which depend on its variables, are collected in one group for each class of such components which they tie together. The size of this group counts the cubes of each of its components once, at most $N_k = (107R_k)^4$ of them, and the sums over these groups are bounded by the product of a factor $b_k = \exp(4\nu N_k + C_x)$ for each component of the second class (Supplement, Proposition S.10(x)). This is the further factor of (6.92). Since $\log b_k \le \frac{1}{16} p_0(g_k)$ for γ_q small, it is paid by the factor $e^{-p_0(g_k)}$ which a component of the second class receives at the step k [12, pp. 385–386]. The details, including the depth N_k of Definition 3.9 and the commensurability of the loops with the period MR_k of the gas, are Supplement, Proposition S.10.

6.16 Several Loops, the Torus, and the Assembly

The groups of the deviation species are small because the background is close to flat. On the term without large-field regions two other species remain, and they are small for a reason of another kind, which is the same for both. A cross group joins two loops, and a group of the wrapping species has a comparison region which does not fit into the torus, so that in both cases the domain of the group is long; and the weight $e^{\nu\sigma}$ of the size of a group, of which the bounds of the class use only a part, pays for a factor which is exponentially small

in that length. Nothing is contracted for these groups, and nothing could be: the flat value of a cross group is not a scalar (Section 6.7), and for a group of the wrapping species the gauge of Supplement, Lemma 17.22, is not available, since that lemma requires a cube which is embedded in the torus. What makes their sums over the steps converge is that the two lengths in question, the distance d_k between two loops and the smallest period $N_{\min}^{(k)}$ of the torus, are larger by the factor L^{k-j} in the units of an earlier step j, whereas a traversal of the step k consists of only L^{k-j} traversals of that step.

Lemma 6.37 (The Cross and the Wrapping Species)**.** *Consider the terms without large-field regions, and let $L \geq 20(1+\beta_0)^2$, $\nu_\times \leq \nu_e$, and $c_{\text{att}} \leq \nu_e/8$.*

(i) Under the hypotheses of Corollary 6.20, those of its last statement included, the groups made at the step $k+1$ satisfy (5.23) *at the step $k+1$.*

(ii) Let the ensemble be dressed at the steps $1,\dots,k$, let γ be so small that Supplement, Proposition S.8, applies, and let γ_q be so small that $\min(\nu_\times, c_{\text{att}})L^2(\log\gamma_q^{-2})^r \geq 2$. Then, for every traversal $\mathfrak{n}$ of the step k,

$$\sum_{W\ni\mathfrak{n}} e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)}\|Y_W\|_{(k)} \leq \sum_{j=1}^{k} 2C_{S'}L^{k-j}\rho_j\, e^{-\nu_\times d_j/M} \leq 4C_{S'}\rho_k\, e^{-\nu_\times d_k/M}\,,$$
$$\sum_{W\ni\mathfrak{n}} e^{\theta(W)+\nu\sigma(W)+\nu_e \operatorname{ext}(W)}\|Y_W\|_{(k)} \leq \sum_{j=1}^{k} 2C_{S'}L^{k-j}\rho_j\, e^{-c_{\text{att}}N_{\min}^{(j)}/M} \leq 4C_{S'}\rho_k\, e^{-c_{\text{att}}N_{\min}^{(k)}/M}\,, \tag{6.95}$$

the first sum over the groups of the cross species and the second over those of the wrapping species of the term of ϱ_k which have $\mathfrak{n}$ in an interval.

(iii) Under the hypotheses of (ii), the number (6.41) *satisfies $\mathsf{E}_k \leq (C_c + 8C_{S'})\rho_k$, and for every $j \leq k$ the groups of the cross species of the term of ϱ_k which were made at the step j satisfy $\sum_W e^{\theta(W)+\nu\sigma(W)}\|Y_W\|_{(k)} \leq 2C_{S'}P_j(\mathcal{C})\rho_j e^{-\nu_\times d_j/M}$.*

Proof. For (i) we bound the size of a new group of either species from below by the length which defines the species, so that the further factor of Corollary 6.20 becomes the small factor of (5.23). For (ii) we reduce every group to the step at which it was made, where (5.23) holds, and sum over the steps; (iii) then collects the species.

(i) Let $\mathcal{G}$ be a cross group made at the step $k+1$. Its domain $X_\mathcal{G}$ is connected (Lemma 6.13(iv)), and it contains the cubes which meet its intervals, hence points of two loops, whose distance is at least d_{k+1} in the units of the scale $k+1$. Since a connected union of N cubes of side M has diameter at most NM, this gives $\ell_\mathcal{G} = \#(X_\mathcal{G}) \geq d_{k+1}/M$. Let now $\mathcal{G}$ be of the wrapping species, and let $\mathfrak{n}$ be a traversal of its interval with $N_{\min}^{(k+1)} \leq 2R_{t+1}+1$, where $t = t_\mathfrak{n}(\mathcal{G})$. The domain lies in the ball of radius $M\ell_\mathcal{G}+1$ about $\mathfrak{n}$, so that $R_t \leq \max(2C_{\text{box}}M, M\ell_\mathcal{G}+1+C_{\text{box}}M) \leq 2C_{\text{box}}M\ell_\mathcal{G}$ by the definition of t, as in the proof of Proposition 6.21. Hence $R_{t+1} = R_t + C_{\text{box}}M \leq 3C_{\text{box}}M\ell_\mathcal{G}$ and $N_{\min}^{(k+1)} \leq 6C_{\text{box}}M\ell_\mathcal{G}+1 \leq 7C_{\text{box}}M\ell_\mathcal{G}$. In both cases $\sigma_{k+1}(\mathcal{G}) \geq \ell_\mathcal{G}$. The further factor $e^{\frac{7}{16}\nu\sigma_{k+1}(\mathcal{G})}$ of Corollary 6.20 is therefore at least $e^{\frac{7}{16}\nu d_{k+1}/M} \geq e^{\nu_\times d_{k+1}/M}$ for a cross group, because $\nu_\times \leq \nu_e \leq \frac{7}{16}\nu$, and for a group of the wrapping species it is at least

$$\exp\Big(\frac{7\nu}{16}\cdot\frac{N_{\min}^{(k+1)}}{7C_{\text{box}}M}\Big) \geq e^{(\nu_e/8)N_{\min}^{(k+1)}/M} \geq e^{c_{\text{att}}N_{\min}^{(k+1)}/M}\,,$$

because $\nu_e \leq \nu/(2C_{\text{box}})$. By that corollary the sum (6.48) with the further factor is at most $2C_{S'}\rho_{k+1}$, for the numbers $\theta(\mathcal{G})$ of the class, and its terms for the groups of the two species

are the terms of (5.23) at the step $k+1$ multiplied by factors which are at least those just found.

(ii) Fix $j \le k$, and let W be a group of one of the two species which was made at the step j and has $\mathfrak{n}$ in an interval. We first compare its supremum at the step k with that at the step j. The background $U_k(V_k)$ of a block field in the support of the term of ϱ_k is the background of a block field in the support of the term of ϱ_j, by (b) of Section 6.10 with k in the place of $k+1$, and $\mathcal{S}_k(U) \subset \mathcal{S}_j(U)$ for every G-valued configuration U, by Lemma 6.23. The space of the step k is therefore contained in that of the step j, and since the insertion of W is the same function at both steps (Definition 5.2(iii)), $\|Y_W\|_{(k)} \le \|Y_W\|_{(j)} \le S_W$. Secondly, W has a traversal of the step j in an interval which is contained in $\mathfrak{n}$, namely the one which contains a fine bond common to $\mathfrak{n}$ and that interval, and $\mathfrak{n}$ consists of L^{k-j} traversals of the step j. Summing (5.23) at the step j over these traversals, with $\ell_W \ge 1$, we find that the groups of the cross species made at the step j contribute at most $2C_{S'}L^{k-j}\rho_j e^{-\nu_\times d_j/M}$ to the first sum of (6.95), and those of the wrapping species at most $2C_{S'}L^{k-j}\rho_j e^{-c_{\text{att}}N^{(j)}_{\min}/M}$ to the second. This is the first inequality in each line.

For the second inequality we put $m := k-j$. A length in the units of the scale j is L^m times the same length in those of the scale k, so that $d_j = L^m d_k$ and $N^{(j)}_{\min} = L^m N^{(k)}_{\min}$, and $\rho_j \le (1+\beta_0)^{6m}\rho_k \le 3^m\rho_k$ by (6.55) and $(1+\beta_0)^6 \le 3$ (Appendix A). With $x := \nu_\times d_k/M$ in the first line and $x := c_{\text{att}}N^{(k)}_{\min}/M$ in the second, each middle term is thus at most $2C_{S'}\rho_k \sum_{m\ge 0}(3L)^m e^{-xL^m}$. Here $x \ge 2$. Indeed, $d_k \ge a_k \ge L^2 M R_k$ by Lemma 3.14, the periods of the torus are $2L^{m_T+q}a_k \ge 2a_k$ in the units of the scale k (proof of that lemma), and $R_k \ge (\log \hat{g}_k^{-2})^r \ge (\log \gamma_q^{-2})^r$ by Definition 3.3 and (3.46), so that $x \ge \min(\nu_\times, c_{\text{att}})L^2(\log\gamma_q^{-2})^r \ge 2$ by the hypothesis on γ_q. For $m \ge 1$ we have $x(L^m-1) \ge 2(L^m-1) \ge L^m \ge mL \ge m(1+\log(3L))$, because $L^m \ge 2$, $L^{m-1} \ge m$, and $L \ge 1+\log(3L)$ for $L \ge 3$. Hence $(3L)^m e^{-xL^m} \le e^{-x}e^{-m}$, and

$$\sum_{m\ge 0}(3L)^m e^{-xL^m} \le e^{-x}\Big(1+\frac{1}{e-1}\Big) \le 2e^{-x}\,,$$

which is the second inequality in each line.

(iii) A group of the term is of the deviation, of the cross, or of the wrapping species, because a group which lies on one loop has one interval (Sections 5.4 and 6.7). By (5.21) the groups of the first species contribute at most $C_c\rho_k$ to E_k, and by (ii), in which the weight $e^{\nu\sigma}$ is at least the weight $e^{\lambda_c\nu\sigma}$ of (6.41), those of each of the other two at most $4C_{S'}\rho_k$. For the last statement, every group of the cross species made at the step j has a traversal of the step j in an interval, so that the sum of $e^{\theta(W)+\nu\sigma(W)}S_W$ over these groups is at most the sum of (5.23) at the step j over the $P_j(\mathcal{C})$ traversals of that step, and $\|Y_W\|_{(k)} \le S_W$ was shown in (ii). *q.e.d.*

Two remarks on the lemma. The size bound (6.48) holds for the cross groups in the norm (5.5), because the number $P(\mathcal{G})$ of (6.38) is a product of norms of tensors and a contraction of tensors does not increase such a product; the constituents with pieces on several loops are treated throughout Section 6.6. And nothing grows with the age of a group: a carried insertion is the function which was bounded when it was made, and whatever a later step ties to it forms a group made at that step, whose size is bounded anew.

The cross species on the other terms. On a term with large-field regions the regular cross groups are groups of ℓ_0 with the same insertions (Proposition 6.36(ii)), so that the lemma holds for them, and a cross group with an attached member belongs to the species of that

member. At the steps at which the **R**-operation acts, an insertion of the cross species may in addition be merged by the operation. Its mass is then multiplied by at most 2 at such a step (Supplement, Proposition S.10(v)), which is the origin of the factor 2^{k-j+1} of (5.15), and the cross groups which the **R**-operation makes add the second term of $\mathsf{A}_j^\times$ (Supplement, Proposition S.10(v)). The growth is paid at the stopping step by the decay of $\mathsf{A}_j^\times$ in the distance $d_j = L^{K'-j} d_{K'}$, which is exponential in $L^{K'-j}$ (Section 8.4).

The cross exponent. The cross exponent $E_{k+1}^\times$, the difference between the logarithm of the reference gas of the ensemble and the sum of those of its loops, is a sum over the traversals of terms which vanish unless a cluster reaches from one loop to another, and

$$|E_{k+1}^\times| \le C_{\mathrm{cr}} \mathfrak{a}_{k+1} \sum_{\mathfrak{n}} e^{-\nu_\times \delta_{\mathfrak{n}}/M} \le C_{\mathrm{cr}} \mathfrak{a}_{k+1} P_k(\mathcal{C}) e^{-\nu_\times a_{k+1}/M} \,, \tag{6.96}$$

the sum over the traversals of the step k, with the amplitude $\mathfrak{a}_{k+1}$ of (7.3) and the constants C_{cr}, $\nu_\times$ of Appendix A. The right side is summable over k, because $a_{k+1} = L^{j_q - q} L^{K'-k-1}$ grows geometrically as k decreases. The flat values are taken loop by loop, and a cross group keeps its insertion; no count depends on the number of loops, since every count is per traversal or per cube, and a cube meets one loop.

Assembly. We collect what has been proved on the term without large-field regions. By Proposition 6.3, Lemma 6.4, Proposition 6.8, and Proposition 6.11, whose hypothesis holds by Theorem 6.18, the dressed loop of the step $k+1$ is the sum (6.29) over the sets of pairwise compatible groups of the step $k+1$. The groups which consist of a single member are the carried groups, with the old insertions at the new background; the insertions of the others which lie on one loop are split (Section 6.7), the numbers $c_{\mathcal{G}}$ are the activities of the reference gas of the step $k+1$, and the resummation of Section 7 gives the form (5.10) at the step $k+1$, with the insertions $Y_{\mathcal{G}}^{\mathrm{d}}$ and the carried ones, the interactions Q_{k+1}, the normalisation $\prod_i \kappa^{(k)}(C_i, \chi_i) \kappa_{k+1}(C_i, \chi_i)$, and the exponent $\mathcal{E}_k$ plus the remainder E_{k+1}° of the flat part, the cross exponent, and the local terms of the step; the leading words are the terms $\mathfrak{Y} = \emptyset$ of (6.8) for the empty family. We take $\theta(W) := \nu\sigma(W)/16$ for every group, at the step at which it is made and at all later steps. The clauses of Definition 5.2 at the step $k+1$ then hold as follows.

Clause (i). For a group made at the step $k+1$ the sum of the absolute values of the weights of the clusters of the new reference gas which are incompatible with it is at most $a^\circ(\mathcal{G}) \le \theta(\mathcal{G})$ (Section 6.7). A group which is incompatible with a carried group W has a point of its domain in a cube which meets X_W^+ (Lemma 6.13(v)), so that it is at one of the at most $81 \cdot 625(1 + 81 u_\star \sigma(W)/L)$ cubes which meet X_W^+ or touch such a cube (Lemma 6.12), and by Lemma 7.3 and the theorem cited in its proof the weights of the clusters incompatible with W sum to at most that number times $C_{S'} \mathfrak{a}_{k+1}$, which is at most $\nu\sigma(W)/16$ for γ_q small, since $\sigma(W) \ge 1$. Clause (ii), first term: the bounds (5.13) and (5.14) hold for the carried groups by Proposition 6.22 and for the new ones by Proposition 6.21, with the constant C_χ of (6.57), and Lemma 5.3 gives the sum. Clause (ii), the wrapping and the cross species: the proof of Lemma 6.37(ii) and (iii) uses of the class at the last of its steps only the clause (5.23) and the analyticity (iii), which hold at the step $k+1$ by part (i) of that lemma and by what follows. By its part (ii) at the step $k+1$ the groups of the wrapping species satisfy the last term of (5.11) with their weights, and by its part (iii) the groups of the cross species made at a step $j \le k+1$ satisfy (5.15) at the step $k+1$ without the power of 2, because $2C_{S'}\rho_j \le C_{\mathrm{cr}} \mathfrak{a}_j$ by (7.3) and $C_{\mathrm{cr}} \ge C_{S'}$ (Appendix A). Clause (iii): the new insertions are holomorphic and covariant on the space of the step $k+1$ by Lemma 6.19, and the carried ones are the functions of the earlier steps, whose domains contain that space by Lemma 6.23. Clause (vii): the domain

of a new group is connected, its size is $\sigma_{k+1}(\mathcal{G}) = \#(X_{\mathcal{G}}) + n_{k+1}(\mathcal{G})/M$, and incompatible groups have domains with a common point, by Lemma 6.13(iv) and (v); the production bound (5.22) is Corollary 6.20, with the constant (6.49); its form (5.23) for the cross and the wrapping species is Lemma 6.37(i); and the supremum form (5.21) is Proposition 6.27 with the constant (6.79). The hypothesis $\mathsf{E}_k \le C_{\mathsf{E}}\rho_k$ of the corollary holds with $C_{\mathsf{E}} \coloneqq C_c + 8C_{S'}$, by Lemma 6.37(iii) at the step k. The constants are fixed in the order C_α, $C_{S'}$, C_c, C_{new}, C_χ, none of which depends on a later one, and then γ_q is taken small. The clauses (iv), (v), and (vi) are those of Section 7, which bounds the remainder and carries the torus consistency from the step k to the step $k+1$.

On the terms with large-field regions the same clauses hold for the regular groups, with the constants of ℓ_0, by Propositions 6.31 and 6.36(ii); the groups made at the collar of the step and those whose comparison regions leave its small-field region satisfy the second and the third term of (5.11) by Propositions 6.32, 6.35, and 6.36(iii); and the groups attached at earlier steps satisfy (6.92), with the totals of Section 6.15. At the steps at which the **R**-operation acts the corresponding statements are those of Section 6.15 and of Supplement, Proposition S.10. Together they prove Theorem 6.1.

7 The Flat Part and the Normalisation

We now determine the number which a step contributes to each loop, and this is the place at which the length of the loops has to be faced directly. At the trivial background the insertions of the step $k+1$ on one loop are scalars, by Schur's lemma, and the contribution of a loop C becomes the partition function of a hard-core gas of groups on C, whose logarithm is a sum over clusters. Every traversal carries clusters of total weight of order $\mathfrak{a}_{k+1}$, so that the logarithm is of order $P_k(C)\mathfrak{a}_{k+1}$, while $P_k(C)$ grows like L^{K-k}. This number can be neither bounded nor neglected, and it has to be extracted exactly, by a formula whose coefficients do not depend on the loop. Two properties of the gas make this possible: a cluster sees the loop only through the arcs of its groups, and the gas is invariant under the translations which preserve the partitions of the construction. Consequently a cluster which is small compared with the corner isolation cannot tell a loop from an infinite straight line away from the corners, nor from two perpendicular rays near a corner, and the density of the logarithm along the loop is the periodic density of the line or of the bent path, up to errors which decay exponentially in the corner isolation. The coefficients of the normalisation are the sums of these densities over one period, and the excess of the bent path over its two rays. Since the line and the bent path are infinite, these sums are defined on $\mathbb{Z}^4$, as limits over larger and larger tori, and the third property which we need is that the weights on a torus approach their limits exponentially fast in the period.

7.1 The Reference Gas

We fix a step $k+1 \le K'$ with $k \ge 1$, the step 1 being that of Section 5, and a labelled term ℓ of ϱ_{k+1}. The *trivial term* ℓ_0 is the term with $\Omega_j = \Lambda_j = \mathbb{T}$ for every $j \le k+1$. It has no large-field regions and no outer variables, and its reference data, the trivial field, lie in the support of all its test functions, so that the identities of Section 6 hold there. The traversals in this section are those of the step k, and lengths and distances are measured in units of the scale k unless the contrary is said.

The partitions and the period. The cubes of the partitions used by the construction have sides which are powers of L, and these partitions are compatible with each other [9, p. 245].

That R_j, M, and M_2 are powers of L is Bałaban's choice [9, p. 245], and that M_1 is one is a standing condition of [1, (3.1)]. Since every side s is odd, the cubes of a partition can be centred at sites, and we take every partition to consist of the cubes $sz + \{-\frac{s-1}{2}, \dots, \frac{s-1}{2}\}^4$, $z \in \mathbb{Z}^4$, of sites of the scale at which it is used, so that the origin is a site of every lattice and the centre of a cube of every partition; this is compatible with the blocks of the averages, which are centred at the sites of the coarser lattice [8, (0.3), p. 252]. For the families π_j it is Bałaban's own convention, whose cubes have their centres at the points of a lattice of spacing M [8, p. 257], and for the other partitions, which [9, p. 245] requires only to be compatible, it is our choice. Let Π_{k+1} be the largest side, in units of the scale k, of the cubes used at the steps $1, \dots, k+1$: the cubes of the families π_j, the cubes of side MR_j of which the domains of the step j are unions, the cubes of side $L^2 M_2 R_j$ of the test functions [9, (3.2), (3.3), (3.16)], and the cubes of side $R_1 M_1$ of the decoupling [10, p. 3]. Every translation by a vector of $\Pi_{k+1}\mathbb{Z}^4$ maps each of these partitions onto itself.

Lemma 7.1 (Period Points)**.** *Let $1 \le k < K'$. Then Π_{k+1} is a power of L with $\Pi_{k+1} \le L^2 M R_k \le a_k$, so that Π_{k+1} divides a_k. The corners of the loops of an ensemble at level q lie in $\Pi_{k+1}\mathbb{Z}^4$, the length of every segment is a multiple of Π_{k+1}, and so are the periods of the torus.*

Proof. At the step $j+1$ the cubes listed above have, in units of the scale j, the sides M for π_j, $LM \le MR_j$ for π_{j+1}, MR_j and $LMR_{j+1} \le LMR_j$ for the domains of the steps j and $j+1$, $L^2 M_2 R_{j+1} \le LMR_j$ for the test functions, and $R_1 M_1 \le M$ [10, p. 3]; here we used $R_{j+1} \le R_j$ from Lemma 3.4(ii), $LM_2 < M$ from [1, (3.1)], and $R_j \ge L$, which holds for γ small by Lemma 3.4(i). All of them are therefore at most $L^2 M R_j$ in units of the scale j, that is, at most $L^{j-k} L^2 M R_j$ in units of the scale k. For $j < k$ the second bound of (3.14) gives

$$L^{j-k} R_j \le (L+1)(k-j)^{\beta_0} L^{j-k} R_k \le (1 + L^{-1}) R_k \,, \tag{7.1}$$

because $m^{\beta_0} \le m \le L^{m-1}$ for every integer $m \ge 1$. A side which is a power of L and at most $(1+L^{-1})L^2 M R_k < L^3 M R_k$ is at most $L^2 M R_k$, since $L^2 M R_k$ is a power of L as well. Hence $\Pi_{k+1} \le L^2 M R_k$, and $L^2 M R_k \le a_k$ by (3.45). Both Π_{k+1} and $a_k = L^{j_q - q} L^{K'-k}$ are powers of L, so the first divides the second.

A loop at level q is a path of bonds of $\mathbb{T}_q$, so that its corners lie in $uL^{-q}\mathbb{Z}^4$, which is $a_k \mathbb{Z}^4$ in units of the scale k by the proof of Lemma 3.14, and its segments, each of which joins two corners along one axis, have lengths which are multiples of a_k. The period $T = 2L^{m_T} u$ is $2L^{m_T + q} a_k$ in units of the scale k. *q.e.d.*

The reference gas. By Section 6.7, the insertion $Y_{\mathcal{G}}$ of a group $\mathcal{G}$ of the step $k+1$ whose intervals lie on one loop takes at the reference data of ℓ the value $c_{\mathcal{G}}\,\mathrm{id}$, with a number $c_{\mathcal{G}} = c_{\mathcal{G}}(\ell)$ which is defined by the reference clause of Definition 5.2(iii); for a cross group we put $c_{\mathcal{G}} := 0$. We also put $c_{\mathcal{G}} := 0$ for an *attached* group, by which we mean a group which is attached to a large-field region of its step or which contains a part of an insertion attached at an earlier step. The insertion of an attached group is thus carried whole, as that of a cross group is. Nothing is lost by this, since no bound of Definition 5.2 for the attached insertions rests on the subtraction of a value at the reference data, and something is gained: the activities of the reference gas, and with them the interactions and the local terms of every later step, involve no attached insertion (Supplement, Lemma 18.2), so that the groups which exist on ℓ alone because they contain a part of an attached insertion do not enter the comparison of the reference gas of ℓ with that of the trivial term. The groups are those of Definition 6.10, in which the interactions of the step k and of the vacuum polymers have

been expanded and their clusters merged into the groups, the factor of each cluster becoming a factor of the insertion of its group; two groups interact only through the hard core of the compatibility defined there. The *reference gas* of ℓ is the hard-core gas of these groups with the activities $c_{\mathcal{G}}$, and its partition functions are

$$S^{\circ}(\ell) \coloneqq \sum_{\mathfrak{E}} \prod_{\mathcal{G}\in\mathfrak{E}} c_{\mathcal{G}} \ , \qquad S^{\circ,[D]}(\ell) \coloneqq \sum_{\mathfrak{E}\perp D} \prod_{\mathcal{G}\in\mathfrak{E}} c_{\mathcal{G}} \ , \tag{7.2}$$

the first sum over all families of pairwise compatible groups, the second over the families $\mathfrak{E}$ such that $D \cup \mathfrak{E}$ is a family and $D \cap \mathfrak{E} = \emptyset$, for a given family D. A *cluster* Γ is a finite multiset of groups whose incompatibility graph is connected, and its weight is $\Phi^{\circ}(\Gamma;\ell) \coloneqq \phi^{\mathrm{U}}(\Gamma)\prod_{\mathcal{G}\in\Gamma} c_{\mathcal{G}}$, with the Ursell function ϕ^{U} of that graph [15, Chapter 5]. The set J_Γ of the traversals in the arcs of the groups of Γ is nonempty, since every group has an arc, and the *network* $N(\Gamma)$ is the union of the domains, the arcs, and the merged clusters of its groups. On the trivial term we write $\Phi^{\circ}(\Gamma;T)$ for the weights on the torus T. The tori of the family $\mathfrak{T}$ are those whose periods are multiples of the period T of $\mathbb{T}$; on each of them the construction is defined by the same rules, with the partitions of $\mathbb{T}$ lifted, and every bound of the preceding sections holds on them with the same constants, since none of these constants depends on the torus.

Definition 7.2 (Amplitude)**.** For $j \geq 1$ the *amplitude* of the step j is

$$\mathfrak{a}_j \coloneqq \max(\tau_{j-1}^2,\, 2\rho_j) \ , \tag{7.3}$$

and the *weight* of a group of the step $k+1$ is $a^{\circ}(\mathcal{G}) \coloneqq (\nu/32)\sigma_{k+1}(\mathcal{G})$, extended additively to clusters.

The factor 2 is the value at $q_1 = q_0$ of the factor $1 + (\log g_k^{-2})^{(q_1-q_0)_+}$ which the exponents of the radii $\alpha_{0,k}$, $\alpha_{1,k}$ contribute to the sizes of the groups, and the construction takes $q_1 = q_0$ (Appendix A). Since $\tau(g)$ and $\rho(g)$ increase for small g and $g_j \leq (1+\beta_0)\bar{g}$ for $j \leq K$ by (3.9) and the window (4.1), the amplitudes are bounded by

$$\bar{\mathfrak{a}} \coloneqq \sup_{j\leq K} \mathfrak{a}_j \leq C\bar{g}^2(\log \bar{g}^{-2})^{6p_0} \ , \qquad \sup_{j\leq K'} \mathfrak{a}_j \leq C\gamma_q^2(\log \gamma_q^{-2})^{6p_0} \ , \tag{7.4}$$

the second because $g_j \leq \gamma_q$ for $j \leq K'$ by (3.46).

Lemma 7.3 (Root Bound and Tail)**.** *Let γ_q be sufficiently small, depending on $c_{\mathcal{C}}$. On every labelled term and every torus of $\mathfrak{T}$, for every traversal b and every $R \geq 0$,*

$$\sum_{\mathcal{G}\colon b\in J_{\mathcal{G}}} |c_{\mathcal{G}}| e^{2a^{\circ}(\mathcal{G})} \leq C_{S'}\mathfrak{a}_{k+1} \ , \qquad \sum_{\Gamma\colon b\in J_\Gamma} |\Phi^{\circ}(\Gamma)| e^{a^{\circ}(\Gamma)} \leq C_{S'}\mathfrak{a}_{k+1} \ , \tag{7.5}$$

where $J_{\mathcal{G}}$ is the set of the traversals in the arc of $\mathcal{G}$, and

$$\sum_{\Gamma\colon b\in J_\Gamma,\ \operatorname{diam} N(\Gamma)\geq R} |\Phi^{\circ}(\Gamma)| \leq C_{S'}\mathfrak{a}_{k+1} e^{-2c_E^T R} \ , \qquad c_E^T \coloneqq \frac{\nu}{64 A_{\mathrm{env}}^* ML} \ , \tag{7.6}$$

with $A_{\mathrm{env}}^* \coloneqq \max(2, A_{\mathrm{env}}+1)$ and A_{env} the constant of the envelopes of [1, Lemma E.8].

Proof. For the first bound we note that $|c_{\mathcal{G}}| \leq S_{\mathcal{G}}$ in (6.48), and that $e^{2a^{\circ}} = e^{(\nu/16)\sigma_{k+1}} \leq e^{\nu\sigma_{k+1}}$. Hence the first sum is at most $2C_{S'}\rho_{k+1} \leq C_{S'}\mathfrak{a}_{k+1}$ by Corollary 6.20 with $\theta = 0$, applied at the traversal of the step $k+1$ which contains b, and by (7.3); the same holds for the sums rooted at a traversal of the step $k+1$ or at a cube of π_{k+1}. On the terms with

large-field regions, and at the steps at which the **R**-operation acts, the bound (6.48) is that of Section 6.15.

For the second bound we verify the condition of Kotecký and Preiss with the weight a°. The domain of a group $\mathcal{G}'$ which is incompatible with $\mathcal{G}$ has a point in common with $X_\mathcal{G}$, by Lemma 6.13(v), so that $\mathcal{G}'$ is at a cube of π_{k+1} which belongs to $X_\mathcal{G}$ or touches it. There are at most $81\sigma_{k+1}(\mathcal{G}) \le 81M\sigma_{k+1}(\mathcal{G})$ such cubes, since σ_{k+1} counts the cubes of $X_\mathcal{G}$ and a cube has 81 cubes which touch it or are equal to it. By the first bound, rooted at these cubes,

$$\sum_{\mathcal{G}' \not\sim \mathcal{G}} |c_{\mathcal{G}'}| e^{2a^\circ(\mathcal{G}')} \le 81M\sigma_{k+1}(\mathcal{G}) C_{S'} \mathfrak{a}_{k+1} \le \frac{\nu}{32} \sigma_{k+1}(\mathcal{G}) = a^\circ(\mathcal{G}) \,, \tag{7.7}$$

where the second inequality is $2592\, MC_{S'}\mathfrak{a}_{k+1} \le \nu$, which holds for γ_q small, depending on $c_\mathcal{C}$, by the second bound of (7.4), because $k+1 \le K'$. This is the condition of [15, Theorem 5.4] with the functions $a = a^\circ$ and $d = a^\circ$. That theorem gives the convergence of the cluster expansion of the reference gas and of the gases with the activities $c_\mathcal{G}\mathbf{1}[\mathcal{G} \perp D]$, and it bounds the clusters which are incompatible with an external polymer by the sum of $|c_\mathcal{G}| e^{a^\circ(\mathcal{G}) + d(\mathcal{G})}$ over the groups incompatible with it. With the traversal b as the external polymer, incompatible with the groups whose arcs contain it, this is the second bound of (7.5).

For the tail we bound the diameter of the network by the weight. The groups of a cluster are connected under incompatibility, and two incompatible groups have networks at distance at most one cube side M, in units of the scale $k+1$. The network of one group is connected and has diameter at most $M\#_{k+1}(\mathcal{G}) + n_{k+1}(\mathcal{G}) = M\sigma_{k+1}(\mathcal{G})$ in units of the scale $k+1$. Hence, in units of the scale k, and since $\sigma_{k+1} \ge 1$ for a group with a domain,

$$\operatorname{diam} N(\Gamma) \le L \sum_{\mathcal{G} \in \Gamma} (M\sigma_{k+1}(\mathcal{G}) + M) \le 2ML \sum_{\mathcal{G} \in \Gamma} \sigma_{k+1}(\mathcal{G}) = \frac{64ML}{\nu}\, a^\circ(\Gamma) \,, \tag{7.8}$$

so that $a^\circ(\Gamma) \ge (\nu/(64ML)) \operatorname{diam} N(\Gamma) \ge 2c_E^T \operatorname{diam} N(\Gamma)$, because $A^*_{\text{env}} \ge 2$. The clusters in (7.6) therefore have $e^{a^\circ(\Gamma)} \ge e^{2c_E^T R}$, and the second bound of (7.5) gives (7.6). *q.e.d.*

The rate $2c_E^T$ is the one which the first step allows: there the network of a cluster has diameter at most $A^*_{\text{env}} LM \sum \sigma$ fine bonds, by the envelopes of [1, Lemma E.8], and the same argument gives (7.6) at the step 1 with the same constant.

Lemma 7.4 (Locality and Periodicity)**.** *On the trivial term the following hold.*

(i) Let Γ be a cluster of the gas of a loop C on a torus T all of whose groups have arcs of fewer than $P_k(C)/2$ traversals. Let C' be a loop on a torus $T' \in \mathfrak{T}$ which contains the arcs of the groups of Γ, with their orientation, each of them as an arc of fewer than $P_k(C')/2$ traversals of C', and let the dependence set of Γ embed isometrically in T'. Then Γ is a cluster of the gas of C' on T' with the same incompatibility graph, and its weight on T' is the weight on T evaluated on the configuration of T' on the image of the dependence set.

(ii) A translation by a vector of $\Pi_{k+1}\mathbb{Z}^4$ maps the reference gas of a loop C onto the reference gas of the translated loop, with the same activities and the same weights.

Proof. (i) The insertion of a group of the step $k+1$ is built from the activities of the polymers of its domains, the tensors Y_G of Lemma 6.4 and the old insertions in its arc, the transports

along its arc, and the clusters of the step k merged into it (Section 6.7 and the merging above). None of these depends on the loop outside the arc: the activities are functions of the fields on their polymers, the tensors Y_G and the transports are attached to the arc, and the old insertions and old cluster weights have the same property by induction, starting from the first step, where the groups are products along their intervals [1, Definition 3.24]. The length of the loop enters only through the rule which takes the smallest arc containing a set, and this rule is applied to sets S of traversals inside the arc A of a group of Γ, at this step or at an earlier one. If A has fewer than half of the traversals of a loop, its complement in that loop lies in one gap of S and is longer than half of the loop, so that this gap is the longest one; the smallest arc containing S is its complement, the part of A between the first and the last traversal of S. By the hypotheses on the lengths this description applies on C and on C', and it gives the same arc on both. The hard core depends only on the arcs, the domains, and the merged clusters, so that the incompatibility graph and the Ursell function are the same.

(ii) Every operation of Bałaban's step, the averages, the minimisers, the propagators and their random walk expansions, the test functions, the localisation, and the activities, is defined from the lattice, the partitions, and the fields alone. A translation by a vector of $\Pi_{k+1}\mathbb{Z}^4$ preserves the lattices and the partitions, and it maps the trivial field to itself, so it maps each object to the object of the translated data. For the sums of the regular part and for the **R**-part this covariance is printed [9, (2.29), (2.32), p. 260]; for the localised activities it follows from their definitions, provided the choices which the papers leave free are made by translation covariant rules, which we do (Section 14). *q.e.d.*

The comparison of two tori needs a condition which excludes that a small cluster wraps around the torus. Let $\rho_{\mathrm{dep}} < LMR_1/4$ be the dependence range of the objects of the first step in fine bonds [1, Sections E.9.2 and E.10.4], and put, for $1 \le j \le K'$,

$$\iota_{T,j} := \max\Big(0,\ \frac{N_{\min}^{(j)}}{4} - \rho_{\mathrm{dep}} L^{-j}\Big)\ , \tag{7.9}$$

in units of the scale j, so that $\iota_{T,j} = L\iota_{T,j+1}$ whenever both are positive. The dependence set of an object of the step $k+1$ lies within the distance

$$R_D^{(k)} \le 5M + LM + L^2 MR_k + \max_{j<k} L^{j-k} R_D^{(j)} \le 6L^2 MR_k \tag{7.10}$$

of its network: the walk pieces of the propagators depend on the field in the fifth enlargement of their domains by cubes of side M_1 [5, Theorem 3.10, p. 416], at most $5M$; the localisation of the step uses the cubes of π_{k+1} [10, p. 13], of side LM; the test functions use cubes of side $L^2 M_2 R_{k+1} \le L^2 MR_k$; and the old insertions have their own ranges. The last inequality follows by iteration from $5M + LM + L^2 MR_k \le 2L^2 MR_k$, from $R_{k-m} \le (L+1)m^{\beta_0} R_k$ by (3.14), and from $(L+1)\sum_{m\ge 1} mL^{-m} = (L+1)L/(L-1)^2 \le 2$ for $L \ge 5$, the range $\rho_{\mathrm{dep}} L^{-k}$ of the first step being smaller than the term $m = k-1$.

Lemma 7.5 (The Torus Condition)**.** *Let $L \ge 195$. For every loop ensemble at level q and every $1 \le k \le K'$,*

$$\iota_{T,k} \ge 16L \cdot 6L^2 MR_k \ge 16L\, R_D^{(k)}\ . \tag{7.11}$$

Proof. A loop at level q has a bond of $\mathbb{T}_q$, so that $uL^{-q} \le \mathrm{diam}(C)$, and $3\,\mathrm{diam}(C) < T = 2L^{m_T} u$ by Definition 1.1; hence $L^{m_T+q} > 3/2$ and $m_T + q \ge 1$. By Lemma 7.1 the smallest period in units of the scale K' is $N_{\min}^{(K')} = 2L^{m_T+q} a_{K'} \ge 2L \cdot L^3 MR_{K'-1} \ge 2L^4 MR_{K'}$, by (3.44) and Lemma 3.4(ii). The first-step range is $\rho_{\mathrm{dep}} L^{-K'} \le L^{1-K'} MR_1/4 \le MR_{K'}/2$,

by (3.14) with $m^{\beta_0} L^{-m} \le L^{-1}$ for $m = K' - 1 \ge 1$, and trivially for $K' = 1$. Hence $\iota_{T,K'} \ge (L^4 - 1) M R_{K'}/2$. For $k < K'$ we have $\iota_{T,k} = L^{K'-k} \iota_{T,K'}$ and $R_k \le (L+1)(K'-k)^{\beta_0} R_{K'} \le (1 + L^{-1}) L^{K'-k} R_{K'}$, so that in all cases

$$\iota_{T,k} \ge \frac{L^4 - 1}{2(1 + L^{-1})} M R_k \ge 96 L^3 M R_k \,, \tag{7.12}$$

where the last inequality is $L^4 - 1 \ge 192 L^3 + 192 L^2$, which holds for $L \ge 195$. The second inequality of (7.11) is (7.10). *q.e.d.*

A cluster Γ *embeds* in T if its dependence set, the $R_D^{(k)}$-neighbourhood of its network, has diameter less than $\iota_{T,k}/4$; the covering map of a torus $T' \in \mathfrak{T}$ over T is then injective on its lifts, and $\tilde{\Gamma}$ denotes a lift.

Proposition 7.6 (Torus Consistency)**.** *Let L satisfy* (C.1)*, let γ_q be sufficiently small, depending on $c_{\mathcal{C}}$, and assume that the ensemble is dressed at the step k. On the trivial term, for a loop C of the ensemble, at the reference data:*

(i) for tori $T \subset T'$ of $\mathfrak{T}$ and every traversal b,

$$\sum_{\Gamma \colon b \in J_\Gamma} |\Phi^\circ(\Gamma; T) - \Phi^\circ(\tilde{\Gamma}; T')| \le \bar{C}_Z \mathfrak{a}_{k+1} e^{-c^Z \iota_{T,k+1}} \,, \qquad c^Z := \frac{\nu}{256 A^*_{\text{env}} M} \,, \tag{7.13}$$

where $\Phi^\circ(\Gamma; T) := 0$ for a cluster of T' which is not the lift of a cluster embedding in T;

(ii) along the tori $T_m \in \mathfrak{T}$ of the periods $L^m T$, the weight of every cluster which embeds in some T_m converges to a limit $\Phi^{\circ,\infty}(\Gamma)$, which does not depend on the sequence of tori, and

$$\sum_{\Gamma \colon b \in J_\Gamma} |\Phi^\circ(\Gamma; T) - \Phi^{\circ,\infty}(\Gamma)| \le 3 \bar{C}_Z \mathfrak{a}_{k+1} e^{-c^Z \iota_{T,k+1}} \,; \tag{7.14}$$

(iii) the clause (vi) of Definition 5.2 holds at the step $k+1$.

Proof. We split the clusters at b into those which embed in T and those which do not.

The embedding clusters. Let Γ embed in T. The activities of its groups are composed of two kinds of factors. The factors made at the step $k+1$ are the same functions on T and on T' of the configuration on the dependence set: the walk pieces of the propagators with boundary conditions depend on the field in their domains only [5, p. 394; Theorem 3.10, p. 416]; a term of the localised expansion "depends on **U**, **J** restricted to $Y_0 \cup \square_0$" [10, p. 5], and the activity $H(Z)$ of the fluctuation integral "is localized in the interior of Z with respect to the external gauge fields" [10, p. 14]; the test functions are local in their cubes; and the couplings are the same numbers on every torus, because the coefficients β_{j+1} are defined through the limit of the vacuum polarisation "as $T^{(j+1)} \nearrow Z^d$" [8, p. 264] and the couplings of a tube are prescribed. The factors made at earlier steps, the insertions of the step k inside the fluctuation integral and the merged clusters of the step k, differ on the two tori by at most the amounts of the clause (vi) at the step k, at fields which coincide on the $\iota_{T,k}/4$-neighbourhood of the dependence set; the deformation of the background by the fluctuation field vanishes outside one lift of the domain of the integral [10, p. 4], so this hypothesis holds at every field of the integral. Replacing these factors by their differences one at a time, and summing over the clusters at b with one marked factor by the count which gives (7.5), we obtain for the embedding clusters the bound $2 C_{\text{rel}}^{(k+1)} \mathfrak{a}_{k+1} e^{-\iota_{T,k+1}/(32M)}$, with

$$C_{\text{rel}}^{(k+1)} \le (L(1+\beta_0)^7 + \tfrac{1}{8}) e^{-(L-1)\iota_{T,k+1}/(32M)} C_{\text{rel}}^{(k)} + C_* \,. \tag{7.15}$$

Here the factor L counts the traversals of the step k in one of the step $k+1$, $(1+\beta_0)^7$ compares the amplitudes of the two steps, the exponential is the gain $e^{-\iota_{T,k}/(32M)} = e^{-L\iota_{T,k+1}/(32M)}$ of the clause (vi) at the step k against the rate of the step $k+1$, and C_* bounds the clusters of Bałaban's gas of polymers inside the activities which do not embed, whose sets differ between the two tori. By (7.12), $(L-1)\iota_{T,k+1}/(32M) \geq 3L^2(L-1)R_{k+1}$, so the prefactor in (7.15) is at most $1/2$, and since $C^{(1)}_{\mathrm{rel}} = 0$ by Theorem 5.4, we get $C^{(k)}_{\mathrm{rel}} \leq 2C_*$ for every k. This is also the clause (vi) at the step $k+1$ for the cluster weights; for the insertions the same telescoping applies, with the deviation parts, which vanish at the reference data, bounded relative to the deviation by the Cauchy estimate of Section 6.9. This proves (iii).

The other clusters. If Γ does not embed in T, its dependence set has diameter at least $\iota_{T,k}/4$, so that its network has diameter at least $\iota_{T,k}/4 - 2R^{(k)}_D \geq \iota_{T,k}/8$ in units of the scale k, by Lemma 7.5. By (7.8) its weight is $a^\circ(\Gamma) \geq (\nu/(64ML))\iota_{T,k}/8 = \nu\iota_{T,k+1}/(512M) \geq c^Z\iota_{T,k+1}$, and the second bound of (7.5) shows that these clusters contribute at most $C_{S'}\mathfrak{a}_{k+1}e^{-c^Z\iota_{T,k+1}}$ on T, and the same on T', where the clusters at a lift of b which are not lifts of embedding clusters of T are exactly those whose dependence sets have diameter at least $\iota_{T,k}/4$.

Adding the two parts gives (i), with $\bar{C}_Z \coloneqq 2C_{S'} + 4C_*$, since $c^Z \leq 1/(32M)$. For (ii), any two tori of $\mathfrak{T}$ are contained in a third one, whose periods are common multiples of theirs, and $\iota_{T_m,k+1} = L^m\iota_{T,k+1}$. By (i) the weights of a cluster along (T_m) form a Cauchy sequence, whose limit does not depend on the sequence by the comparison through a common torus, and the telescoping sum gives the bound $\bar{C}_Z\mathfrak{a}_{k+1}\sum_{m\geq 0} e^{-c^Z L^m\iota_{T,k+1}} \leq 2\bar{C}_Z\mathfrak{a}_{k+1}e^{-c^Z\iota_{T,k+1}}$ for the clusters which embed in T, because $L^m - 1 \geq m(L-1)$ and $c^Z\iota_{T,k+1} \geq 1$; the latter follows from $\iota_{T,k+1} \geq 96L^3MR_{k+1}$, by (7.12), and from $R_{k+1} \geq (\log \hat{g}^{-2}_{k+1})^r \geq (\log \gamma_q^{-2})^r$, for γ_q small, since $k+1 \leq K'$. The clusters which do not embed in T contribute at most $C_{S'}\mathfrak{a}_{k+1}e^{-c^Z\iota_{T,k+1}}$ on T, and the same in the limit, by Fatou's lemma applied to the bound on each T_m, which holds with $\iota_{T,k+1}$ in place of $\iota_{T_m,k+1}$ because the condition is on the diameter of the dependence set. This gives (7.14). *q.e.d.*

The exact factorisation. With the reference gas we can now carry out the resummation of Section 6.7. On a labelled term ℓ we split every insertion as $Y_{\mathcal{G}} = c_{\mathcal{G}}T(\mathcal{G}) + Y^{\mathrm{d}}_{\mathcal{G}}$, where $T(\mathcal{G})$ is the product of the representations of the transports of the background field along the arc of $\mathcal{G}$. For a group on one loop, $Y^{\mathrm{d}}_{\mathcal{G}}$ vanishes at the reference data and is covariant with the endpoints of the arc. For a cross group $c_{\mathcal{G}} = 0$, so that $Y^{\mathrm{d}}_{\mathcal{G}} = Y_{\mathcal{G}}$ is the whole insertion, which does not vanish at the reference data in general and is bounded by (5.15). Since every word is multilinear in its insertions, and the transports of the scalar parts join the transports of the gaps,

$$w(\mathfrak{F}) = \sum_{D\subseteq\mathfrak{F}} \Big(\prod_{\mathcal{G}\in\mathfrak{F}\setminus D} c_{\mathcal{G}} \Big) w^{\mathrm{d}}(D) \,, \tag{7.16}$$

where $w^{\mathrm{d}}(D)$ is the word of the family D with the insertions $Y^{\mathrm{d}}_{\mathcal{G}}$, and $w^{\mathrm{d}}(\emptyset) = \prod_i f^{(i)}_{k+1}$. After the merging the families of groups interact only through the hard core, so that the sum over the families $\mathfrak{F}$ of (7.16) is a sum over the pairs $(D, \mathfrak{E})$ with $\mathfrak{F} = D \cup \mathfrak{E}$, and

$$\sum_{\mathfrak{F}} w(\mathfrak{F}) = S^\circ(\ell)\Big[\prod_i f^{(i)}_{k+1} + \sum_{D\neq\emptyset} Q_{k+1}(D)\, w^{\mathrm{d}}(D) \Big] \,, \tag{7.17}$$

with

$$Q_{k+1}(D) \coloneqq \frac{S^{\circ,[D]}(\ell)}{S^\circ(\ell)} = \exp\Big(- \sum_{\Gamma\not\supset D} \Phi^\circ(\Gamma;\ell) \Big) \,, \tag{7.18}$$

the last sum over the clusters which contain a group incompatible with a member of D. The rearrangements are justified by the absolute convergence of $\sum_{\mathfrak{F}} \prod_{\mathcal{G}\in\mathfrak{F}} (|c_{\mathcal{G}}| + \|Y^{\mathrm{d}}_{\mathcal{G}}\|)$, which follows from (6.48); the gas is a gas of groups together with their merged clusters, and the sum over the latter is infinite even on a finite torus. The cluster series of $S^{\circ}(\ell)$ and of $S^{\circ,[D]}(\ell)$ converge by Lemma 7.3, which gives $S^{\circ}(\ell) \neq 0$ and the second expression for $Q_{k+1}(D)$, and $\sum_{\Gamma \not\perp \mathcal{G}} |\Phi^{\circ}(\Gamma)| \le a^{\circ}(\mathcal{G}) \le \nu\sigma_{k+1}(\mathcal{G})/16$ by [15, Theorem 5.4] and (7.7), which is the clause (i) at the step $k+1$ with $\theta := a^{\circ}$. For an ensemble the groups of all loops and the cross groups form one gas; since cross groups have the activity 0, on the trivial term

$$\log S^{\circ}(\ell_0) = \sum_{i=1}^{n} F^{\circ}_{k+1}(C_i; T) + E^{\times}_{k+1} \,, \qquad F^{\circ}_{k+1}(C; T) := \sum_{\Gamma} \Phi^{\circ}(\Gamma; T) \,, \tag{7.19}$$

the second sum over the clusters of the gas of the single loop C, and this defines the cross exponent $E^{\times}_{k+1}$ of (6.96). On a general term the difference $E^{\partial}_{k+1}(\ell) := \log S^{\circ}(\ell) - \log S^{\circ}(\ell_0)$ consists of the clusters which see the large-field data of ℓ; its representation as a sum of local terms $e^{(k+1)}_X$ attached to these data, with the first bound of (5.20), is part of Supplement, Proposition S.9.

7.2 The Coefficients

For $v \in \mathcal{V}$ let $\ell_v := \{tv \,|\, t \in \mathbb{Z}\}$ be the line through the origin in the orientation v, with its traversals numbered by $b \in \mathbb{Z}$, the traversal b running from bv to $(b+1)v$. For a perpendicular ordered pair (v, w) let $B_{v,w}$ be the bent path (2.6) with $c = 0$, $v_c = v$, and $w_c = w$, with its traversals numbered by $b \in \mathbb{Z}$: for $b < 0$ the traversal from bv to $(b+1)v$, and for $b \ge 0$ the traversal from bw to $(b+1)w$. The *line gas* of ℓ_v consists of the clusters Γ on $\mathbb{Z}^4$ whose groups have finite arcs on ℓ_v, with the weights $\Phi^{\circ,\infty}_v(\Gamma)$ obtained as follows: we take a torus $T' \in \mathfrak{T}$ in which Γ embeds and on it a square C' at level q, in a coordinate plane containing v and with its corners in $\Pi_{k+1}\mathbb{Z}^4$, one side of which contains the arcs of Γ in the orientation v and is longer than twice the diameter of the network of Γ; then Γ is a cluster of the gas of C', and its weight does not depend on the choice of C' by Lemma 7.4(i), whose hypothesis on the lengths holds because an arc of Γ lies in the network and therefore has fewer traversals than a side of either square, a quarter of its perimeter. Further, $\Phi^{\circ,\infty}_v(\Gamma)$ is the limit of these weights along (T_m) given by Proposition 7.6(ii). The *bent-path gas* of $B_{v,w}$ is defined in the same way, with a square in the plane spanned by v and w which has a corner at the origin with the incoming orientation v and the outgoing orientation w, and whose two sides at this corner contain the arcs of Γ and are longer than twice the diameter of its network; an arc of Γ on these two sides, whose ends are within that diameter of each other, then has fewer traversals than one side. The squares used here need not be admissible. The proofs of Theorems 5.4 and 6.1 use the loops only through $\Lambda(C) = 2$, $P(C)/r(C) = 8$, and the torus conditions, and the torus conditions, which compare the loop with the periods, hold at the given K once the periods of T' are large compared with the square. The densities and the coefficients of the step $k+1$ are

$$a_v(b) := \sum_{\Gamma:\, b\in J_\Gamma} \frac{\Phi^{\circ,\infty}_v(\Gamma)}{|J_\Gamma|} \,, \qquad \kappa^{(k+1)}_v := \sum_{b=0}^{\Pi_{k+1}-1} a_v(b) \tag{7.20}$$

on the line, and, with $a_{v,w}(b)$ defined in the same way on $B_{v,w}$ and with $k_{v,w}(b) := a_v(b)$ for $b < 0$ and $k_{v,w}(b) := a_w(b)$ for $b \ge 0$,

$$\zeta^{(k+1)}_{v,w} := \sum_{b\in\mathbb{Z}} (a_{v,w}(b) - k_{v,w}(b)) \tag{7.21}$$

on the bent path.

Lemma 7.7 (The Coefficients)**.** *Let L and $\bar{g}$ be as in Proposition 7.6, and put $C_\zeta \coloneqq 2C_{S'}(1+2/(e^{2c_E^T}-1))$.*

(i) The density a_v is Π_{k+1}-periodic, $|a_v(b)| \le C_{S'}\mathfrak{a}_{k+1}$, and $|\kappa_v^{(k+1)}| \le \Pi_{k+1}C_{S'}\mathfrak{a}_{k+1}$.

(ii) $|a_{v,w}(b)-k_{v,w}(b)| \le 2C_{S'}\mathfrak{a}_{k+1}e^{-2c_E^T|b|}$ for every $b \in \mathbb{Z}$, so that the series (7.21) converges absolutely and $|\zeta_{v,w}^{(k+1)}| \le C_\zeta \mathfrak{a}_{k+1}$.

(iii) The numbers $\kappa_v^{(k+1)}$ and $\zeta_{v,w}^{(k+1)}$ do not depend on the loops, on the torus, or on the labelled term. They depend on k, K, and $\bar{g}$ through the couplings and the parameters of the tube, on the character, and on the constants of the construction.

Proof. (i) By Lemma 7.4(ii), applied on each torus T_m, the weights of a cluster and of its translate by $\Pi_{k+1}v$ agree, and so do their limits. The bound on $a_v(b)$ is (7.5) at $R=0$ with $|J_\Gamma| \ge 1$, valid on each T_m and passed to the limit by Fatou's lemma; summing over one period gives the bound on $\kappa_v^{(k+1)}$.

(ii) Let $b \ge 0$. A cluster of the bent-path gas all of whose arcs lie on the outgoing ray is, by Lemma 7.4(i), a cluster of the line gas of ℓ_w with the same weight, and conversely a cluster of the line gas at b all of whose arcs lie in $\{b' \ge 0\}$ is one of the bent-path gas. The terms which survive in $a_{v,w}(b) - a_w(b)$ are therefore the clusters of the bent path at b with an arc on the incoming ray, and the clusters of the line at b with an arc on $\{b' < 0\}$. The network of each of them contains the position bw of b and a site $-tv$ or $-tw$ with $t \ge 1$, at distance at least b, so that (7.6) with $R=b$, in each of the two gases, gives the bound. The case $b<0$ is the same with the two rays exchanged, the distance being at least $|b|$. Summing, with $y \coloneqq e^{-2c_E^T}$,

$$|\zeta_{v,w}^{(k+1)}| \le 2C_{S'}\mathfrak{a}_{k+1}\Big(1+2\sum_{b\ge 1} y^b\Big) = 2C_{S'}\mathfrak{a}_{k+1}\frac{1+y}{1-y} = C_\zeta \mathfrak{a}_{k+1}\,. \tag{7.22}$$

(iii) The coefficients are sums of weights of the line and bent-path gases on $\mathbb{Z}^4$, which by construction involve neither the loops of the ensemble, nor the torus $\mathbb{T}$, nor any term but the trivial one. The dependence on the step is through the objects of the steps $1,\dots,k+1$ on the trivial term. *q.e.d.*

The coefficients are kept separately for each orientation and each ordered pair of perpendicular orientations, since the covariance of the construction under the hypercubic group is not established; no relation between them is used. With them we define the factor of a step. At the step 1 the coefficients are those of the first step: $\Pi_1 \coloneqq L$, and $\kappa_v^{(1)}$ and $\zeta_{v,w}^{(1)}$ are the perimeter and corner coefficients of [1, (5.33), (6.4)] in the orientation v and at the corner (v,w) (Section 5.4). A step j is an **R**-*step* if $j > N_j$, so that the **R**-operation acts on the numerator at that step.

Definition 7.8 (Step Factor)**.** For a loop C with the character χ and $1 \le j \le K'$ let $N_v^{(j-1)}(C)$ be the number of traversals of C of the orientation v at the step $j-1$. The *step factor* is $\kappa_j(C,\chi) \coloneqq \exp\log\kappa_j(C,\chi)$ with

$$\log\kappa_j(C,\chi) \coloneqq \sum_{v\in\mathcal{V}} \frac{N_v^{(j-1)}(C)}{\Pi_j}\,\kappa_v^{(j)} + \sum_{c\in\mathcal{K}(C)} \zeta_{v_c,w_c}^{(j)} + \mathbf{1}_{\mathbf{R}}(j)\Big[\sum_{v\in\mathcal{V}} N_v^{(j-1)}(C)\,\mu_{j,v}^{\mathbf{R}} + \sum_{c\in\mathcal{K}(C)} \zeta_{j,v_c,w_c}^{\mathbf{R}}\Big]\,, \tag{7.23}$$

where $\mathbf{1}_{\mathbf{R}}(j)$ is 1 at an **R**-step and 0 otherwise, and $\mu_{j,v}^{\mathbf{R}}$, $\zeta_{j,v,w}^{\mathbf{R}}$ are the coefficients of Proposition 7.10. The coefficients depend on χ through the representation π in the insertions.

The factor is not zero, being an exponential, and $\log \kappa_j$ has exactly the structure announced in Section 1.3: a term proportional to the length of the segments of each orientation and a term for each corner, with coefficients which are numbers of the step.

7.3 The Remainder of a Step

Theorem 7.9 (The Flat Part of a Step)**.** *Let L and $\bar{g}$ be as in Proposition 7.6, let $1 \le k < K'$, and assume that the ensemble is dressed at the step k. For every loop C of the ensemble,*

$$F^{\circ}_{k+1}(C;T) = \sum_{v\in\mathcal{V}} \frac{N^{(k)}_v(C)}{\Pi_{k+1}}\, \kappa^{(k+1)}_v + \sum_{c\in\mathcal{K}(C)} \zeta^{(k+1)}_{v_c,w_c} + E^{\circ}_{k+1}(C,T)\,, \tag{7.24}$$

with a number $E^{\circ}_{k+1}(C,T)$ which satisfies, with $C^{\mathcal{K}}_E \coloneqq 2C_{S'} + 2C_\zeta$,

$$|E^{\circ}_{k+1}(C,T)| \le C^{\mathcal{K}}_E \mathfrak{a}_{k+1}(P_k(C) + |\mathcal{K}(C)|)e^{-2c^T_E r_k(C)} + 3\bar{C}_Z \mathfrak{a}_{k+1} P_k(C) e^{-c^Z \iota_{T,k+1}}\,. \tag{7.25}$$

Proof. We allocate every cluster equally to the traversals of J_Γ,

$$F^{\circ}_{k+1}(C;T) = \sum_{b\in C} a_C(b;T)\,, \qquad a_C(b;T) \coloneqq \sum_{\Gamma\colon b\in J_\Gamma} \frac{\Phi^{\circ}(\Gamma;T)}{|J_\Gamma|}\,, \tag{7.26}$$

the sum over the $P_k(C)$ traversals, which converges absolutely by (7.5). Every traversal is assigned to one corner (Section 2.4), and we compare the density at a traversal b assigned to the corner c with the density of the bent path $B_c = c + B_{v_c,w_c}$ at the corresponding traversal.

Small clusters. Let Γ be a cluster at b whose network has diameter less than $r_k(C)$. By Lemma 2.10 every site of C or of B_c outside $S_{\mathrm{in}}(c) \cup S_{\mathrm{out}}(c)$ is at distance at least $r_k(C)$ from the position of b, while the arcs of the groups of Γ lie in its network; so these arcs lie in $S_{\mathrm{in}}(c) \cup S_{\mathrm{out}}(c)$, within distance $r_k(C)$ of the position of b, hence in an arc of fewer than $4r_k(C) \le P_k(C)/2$ traversals, since the chord–arc constant of two consecutive segments is 2 by Lemma 5.5(iii). The cluster does not wrap around the torus, because $r_k(C) \le \mathrm{diam}_k(C) < N^{(k)}_{\min}/3$. By Lemma 7.4(i), whose hypothesis on the lengths holds on C by this count and on the squares of the bent-path gas by their definition, it is a cluster of the bent-path gas of B_c, and conversely a cluster of that gas at b whose network has diameter less than $r_k(C)$ is a cluster of the gas of C. By Lemma 7.1 the corner c lies in $\Pi_{k+1}\mathbb{Z}^4$, so by Lemma 7.4(ii) the weights of the gas of B_c are those of B_{v_c,w_c} at the translated traversal. Their weights on T and on $\mathbb{Z}^4$ differ by at most $3\bar{C}_Z \mathfrak{a}_{k+1} e^{-c^Z \iota_{T,k+1}}$ in the sum at b, by (7.14).

Large clusters. The clusters at b whose networks have diameter at least $r_k(C)$ contribute at most $C_{S'}\mathfrak{a}_{k+1} e^{-2c^T_E r_k(C)}$ in the gas of C and the same in the gas of B_c, by (7.6). This includes the clusters with a group whose arc has at least $P_k(C)/2$ traversals, whose networks have diameter at least $P_k(C)/(2\Lambda(C)) \ge r_k(C)$ by Lemma 2.9 and Lemma 2.5.

Together, writing b also for the corresponding traversal of B_{v_c,w_c},

$$|a_C(b;T) - a_{v_c,w_c}(b)| \le 2C_{S'}\mathfrak{a}_{k+1} e^{-2c^T_E r_k(C)} + 3\bar{C}_Z \mathfrak{a}_{k+1} e^{-c^Z \iota_{T,k+1}}\,. \tag{7.27}$$

The corner sums. Let ℓ and ℓ' be the lengths of $S_{\mathrm{in}}(c)$ and $S_{\mathrm{out}}(c)$. The traversals assigned to c are the traversals b of B_{v_c,w_c} with $-\lfloor \ell/2 \rfloor \le b \le \lceil \ell'/2 \rceil - 1$. By (7.21) the sum of $a_{v_c,w_c}(b)$ over them is the sum of $k_{v_c,w_c}(b)$ over them, plus $\zeta^{(k+1)}_{v_c,w_c}$, minus the sum of $a_{v_c,w_c}(b) - k_{v_c,w_c}(b)$

over the other b, which have $b \geq \lceil \ell'/2 \rceil \geq s_k(C)/2$ or $|b| \geq \lfloor \ell/2 \rfloor + 1 \geq (s_k(C)+1)/2$. By Lemma 7.7(ii), with $y = e^{-2c_E^T}$, this last sum is at most

$$2C_{S'}\mathfrak{a}_{k+1}\Big(\sum_{b \geq s_k(C)/2} y^b + \sum_{b \geq (s_k(C)+1)/2} y^b\Big) \leq \frac{4C_{S'}\mathfrak{a}_{k+1}}{1-y} e^{-c_E^T s_k(C)} \leq 2C_\zeta \mathfrak{a}_{k+1} e^{-2c_E^T r_k(C)} \,, \quad (7.28)$$

because $C_\zeta \geq 2C_{S'}/(1-y)$ and $s_k(C) \geq 2r_k(C)$.

The segment sums. A segment σ from the corner c to the corner c'', of orientation v and length ℓ_σ, receives the terms $k_{v_c,w_c}(b) = a_v(b)$, $0 \leq b < \lceil \ell_\sigma/2 \rceil$, from c, and the terms $k_{v_{c''},w_{c''}}(b) = a_v(b)$, $-\lfloor \ell_\sigma/2 \rfloor \leq b < 0$, from c'', read on the line $c + \ell_v$, since $c'' = c + \ell_\sigma v$ and both corners lie in $\Pi_{k+1}\mathbb{Z}^4$. Together these are the terms $a_v(b)$, $0 \leq b < \ell_\sigma$, of the line ℓ_v, by Lemma 7.4(ii), and since ℓ_σ is a multiple of Π_{k+1} (Lemma 7.1) and a_v is Π_{k+1}-periodic, their sum is exactly $(\ell_\sigma/\Pi_{k+1})\kappa_v^{(k+1)}$. Summing over the segments gives the first term of (7.24), since $\sum_{\sigma \text{ of orientation } v} \ell_\sigma = N_v^{(k)}(C)$.

Conclusion. Adding (7.27) over the $P_k(C)$ traversals and (7.28) over the $|\mathcal{K}(C)|$ corners gives (7.25), with $e^{-2c_E^T r_k(C)}$ for both corner terms. *q.e.d.*

At the steps at which the **R**-operation acts, its polymer gas linked to the loop has a flat part of the same structure, with the period MR_j of the cubes of the **R**-operation in place of Π_j, the decay $e^{-\kappa d_j}$ of its activities in place of (7.6), and a group's hull in place of its network.

Proposition 7.10 (The Flat Part of an **R**-Step)**.** *Let $j \leq K'$ be an* **R***-step. There are numbers $\mu_{j,v}^{\mathbf{R}}$, $v \in \mathcal{V}$, and $\zeta_{j,v,w}^{\mathbf{R}}$, (v,w) perpendicular, independent of the loops, the torus, and the labelled term, with $|\mu_{j,v}^{\mathbf{R}}|, |\zeta_{j,v,w}^{\mathbf{R}}| \leq C_E^R e^{-p_0(g_j)/4}$, such that for every loop C of a dressed ensemble the flat part of the exponent which the* **R***-operation of the step j contributes along C is $\sum_v N_v^{(j-1)}(C)\mu_{j,v}^{\mathbf{R}} + \sum_{c \in \mathcal{K}(C)} \zeta_{j,v_c,w_c}^{\mathbf{R}} + E_j^{\circ,\mathbf{R}}(C,T)$, with*

$$|E_j^{\circ,\mathbf{R}}(C,T)| \leq C_E^R MR_j e^{-p_0(g_j)/4}\Big[(P_j(C) + |\mathcal{K}(C)|)e^{-c_E^R r_j(C)/(MR_j)} + P_j(C)e^{-c_E^R \iota_{T,j}/(MR_j)}\Big] \,. \quad (7.29)$$

Proof. Supplement, Proposition S.10. The proof is that of Theorem 7.9 for the gas of the **R**-operation linked to the loop, whose flat activities are periodic along each segment with the period MR_j; every segment is a union of such periods starting at a corner, because MR_jL^j divides $MR_{K'}L^{K'}$, which divides the fine length $L^{j_q - q + K'}$ of a bond of $\mathbb{T}_q$ by (3.44). *q.e.d.*

The identity of the step. We can now complete the step of Section 6. On every labelled term, by (7.17), (7.19), Theorem 7.9, and Proposition 7.10,

$$\prod_i \kappa^{(k)}(C_i,\chi_i)\, e^{\mathcal{E}_k} S^\circ(\ell) = \prod_i \kappa^{(k+1)}(C_i,\chi_i)\, e^{\mathcal{E}_{k+1}} \,, \quad (7.30)$$

with $\mathcal{E}_{k+1} := \mathcal{E}_k + E_{k+1}^\partial(\ell) + \sum_i E_{k+1}^\circ(C_i,T) + E_{k+1}^\times + \mathbf{1}_{\mathbf{R}}(k+1)(\sum_i E_{k+1}^{\circ,\mathbf{R}}(C_i,T) + E_{k+1}^{\mathbf{R}\times})$, the last terms entering at an **R**-step together with the flat part of the **R**-gas, with the cross constant $E_{k+1}^{\mathbf{R}\times}$ of Supplement, Proposition S.10(iv)(c), and $E_{\text{const},k+1} := E_{\text{const},k} + \sum_i E_{k+1}^\circ(C_i,T) + E_{k+1}^\times + \mathbf{1}_{\mathbf{R}}(k+1)(\sum_i E_{k+1}^{\circ,\mathbf{R}}(C_i,T) + E_{k+1}^{\mathbf{R}\times})$, which are numbers on each labelled term. The identity is exact, and it is the normalisation clause of the step.

7.4 The Normalisation

The remainders of the steps are collected in the following bound, in which $P_{j-1}(\mathcal{C}) \coloneqq \sum_i P_{j-1}(C_i)$, $|\mathcal{K}(\mathcal{C})| \coloneqq \sum_i |\mathcal{K}(C_i)|$, and $r_{j-1}(\mathcal{C}) \coloneqq \min_i r_{j-1}(C_i)$: for $1 \le j \le K'$,

$$\begin{aligned}\epsilon_j(\mathcal{C},T) \coloneqq \mathfrak{a}_j \Big[& C_E^{\mathcal{K}}(P_{j-1}(\mathcal{C}) + |\mathcal{K}(\mathcal{C})|)e^{-2c_E^T r_{j-1}(\mathcal{C})} + 3\bar{C}_Z P_{j-1}(\mathcal{C}) e^{-c^Z \iota_{T,j}} + C_{\mathrm{cr}} P_{j-1}(\mathcal{C}) e^{-\nu_\times a_j/M} \Big] \\ & + \mathbf{1}_{\mathbf{R}}(j)\, C_E^R M R_j e^{-p_0(g_j)/4} \Big[(P_j(\mathcal{C}) + |\mathcal{K}(\mathcal{C})|) e^{-c_E^R r_j(\mathcal{C})/(MR_j)} + P_j(\mathcal{C}) e^{-c_E^R \iota_{T,j}/(MR_j)} \Big] \\ & + \mathbf{1}_{\mathbf{R}}(j)\, C_E^{\mathbf{R}\times} P_j(\mathcal{C}) e^{-p_0(g_j)/4} e^{-\kappa a_j/(6MR_j)} \,. \end{aligned} \tag{7.31}$$

At $j = 1$ the first line bounds the remainder of the first step, by (5.25), by the periodic comparison of [1, Appendix F], and by the cross terms of [1, Section 10]. By (7.25), (6.96), (7.29), and Supplement, Proposition S.10(iv)(c), the constant part of the scalar exponent satisfies the bound of Definition 5.2(iv) at every step.

Corollary 7.11 (Sum of the Remainders)**.** *Let L be as in Proposition 7.6, and let $\mathcal{C}$ be a loop ensemble at level q whose loops are admissible, with γ_q as in that proposition. For every $K > j_q$,*

$$\sum_{j=1}^{K'} \epsilon_j(\mathcal{C},T) \le B_E(\mathcal{C}) \coloneqq \sum_{i=1}^{n} \Big[\Big(\frac{P(C_i)}{r(C_i)} + P_q(C_i) \Big) (C'_E \bar{\mathfrak{a}} + C''_E S^{\mathbf{R}}(\bar{g})) \Big] \,, \tag{7.32}$$

where C'_E and C''_E depend on the constants of the construction, on Λ_*, and on $c_{\mathcal{C}}$, nondecreasingly in $c_{\mathcal{C}}$, and $S^{\mathbf{R}}(\bar{g}) \coloneqq \sup_K \sum_{j \le K} (MR_j)^2 e^{-p_0(g_j)/4}$ is finite and tends to 0 as $\bar{g} \to 0$. In particular $B_E(\mathcal{C})$ does not depend on K.

Proof. We sum the five parts of (7.31) over j, loop by loop, with $m \coloneqq K' - j + 1$, so that the lengths of the step $j-1$ are those of the stopping step multiplied by L^m.

The corner part. The exponent of this part contains the length $r_{j-1}(\mathcal{C}) = \min_i r_{j-1}(C_i)$ of the ensemble, which for a loop C of the ensemble may be smaller than its own $r_{j-1}(C)$, so that we sum with $r(\mathcal{C})$ throughout. The ratio $P_{j-1}(C)/r_{j-1}(\mathcal{C}) = P(C)/r(\mathcal{C})$ does not depend on j, and $x \coloneqq r_{K'}(\mathcal{C}) \ge a_{K'}/2 \ge 1$ by (3.45) for each loop, since $s_{K'}, \delta_{K'} \ge a_{K'}$ and $P_{K'} \ge 4a_{K'}$. Lemma 2.11 with $c = 2c_E^T$, together with $xe^{-2c_E^T x} \le 1/(2ec_E^T)$, $e^{-2c_E^T x} \le 1$, and $|\mathcal{K}(C)| \le P(C)/(2r(C)) \le P(C)/(2r(\mathcal{C}))$ from Lemma 2.5, gives

$$\sum_{m \ge 1} \Big(\frac{P(C)}{r(\mathcal{C})} x L^m + |\mathcal{K}(C)| \Big) e^{-2c_E^T x L^m} \le \frac{P(C)}{r(\mathcal{C})} \Big(\frac{1}{2ec_E^T} + \frac{1}{c_E^T} + \frac{1}{2} + \frac{1}{2c_E^T} \Big) \,. \tag{7.33}$$

Since every loop of an ensemble at level q has $r(C_i) \ge \frac{1}{2} u L^{-q}$, the same holds for $r(\mathcal{C})$, and $P(C)/r(\mathcal{C}) \le 2P_q(C)$, a term of $B_E(\mathcal{C})$.

The torus part. By the proof of Lemma 7.5, $\iota_{T,K'} \ge a_{K'}(L^{m_T+q} - L^{-3})/2 \ge a_{K'}$, and $\iota_{T,j} = L^{m-1} \iota_{T,K'}$, while $P_{j-1}(C) = P_q(C) a_{K'} L^m \le P_q(C) L\, L^{m-1} \iota_{T,K'}$. Lemma 2.11 with $x = \iota_{T,K'}$ and $c = c^Z$, and $c^Z x \ge 1$, give $\sum_m P_{j-1}(C) e^{-c^Z \iota_{T,j}} \le 3P_q(C) L x e^{-c^Z x} \le 3P_q(C) L/(ec^Z)$.

The cross part. With $a_j = a_{K'} L^{m-1}$ and $P_{j-1}(C) = P_q(C) L a_j$, Lemma 2.11 with $x = a_{K'}$ and $c = \nu_\times / M$, and $\nu_\times a_{K'}/M \ge 1$, which holds for γ_q small since $a_{K'} \ge L^3 M R_{K'}$ and $R_{K'} \ge (\log \hat{g}_{K'}^{-2})^r \ge (\log \gamma_q^{-2})^r$, give at most $3P_q(C) L a_{K'} e^{-\nu_\times a_{K'}/M} \le 3P_q(C) LM/(e\nu_\times)$.

The $\mathbf{R}$-part. Since $P_j(C) = (P(C)/r(\mathcal{C})) r_j(\mathcal{C})$ and $te^{-ct/(MR_j)} \le MR_j/(ec)$ for $t \ge 0$, the first bracket is at most $(P(C)/r(\mathcal{C}))MR_j(1/(ec_E^R) + 1/2)$; since $\iota_{T,j} \ge a_j$ and $P_j(C) = P_q(C) a_j$,

the second is at most $P_q(C)MR_j/(ec_E^R)$. The sum over j is therefore at most $C_E^R(P(C)/r(\mathcal{C})+P_q(C))(1/2+2/(ec_E^R))S^{\mathbf{R}}(\bar{g})$, and $P(C)/r(\mathcal{C}) \le 2P_q(C)$ as in the corner part. In the same way the cross part of the **R**-steps, with $P_j(\mathcal{C}) = P_q(\mathcal{C})a_j$ and $a_j e^{-\kappa a_j/(6MR_j)} \le 6MR_j/(e\kappa)$, sums to at most $6C_E^{\mathbf{R}\times}P_q(\mathcal{C})S^{\mathbf{R}}(\bar{g})/(e\kappa)$. The number $S^{\mathbf{R}}(\bar{g})$ is finite and tends to 0 with $\bar{g}$: by Lemma 3.2 and the definition of a tube, $g_j^{-2} \ge \hat{g}_j^{-2}/\rho \ge (2/3)(\bar{g}^{-2}+b'(K-j))$, with $n = K-j$, and $(MR_j)^2 \le 4L^2M^2(\log g_j^{-2})^{2r}$ by Lemma 3.4(i), so that $(MR_j)^2 e^{-p_0(g_j)/4} \le 4L^2M^2y^{2r}e^{-(A_0/4)y^{p_0}}$ at $y = \log g_j^{-2}$. The two factors move in opposite directions as y grows, and we bound their product: the function $y \mapsto y^{2r}e^{-(A_0/4)y^{p_0}}$ is decreasing for $y^{p_0} \ge 8r/(A_0p_0)$, which for $\bar{g}$ small contains every $y \ge \log((2/3)\bar{g}^{-2})$, so the product is at most its value at $y = \log((2/3)(\bar{g}^{-2}+b'n))$. The resulting series over $n \ge 0$ converges because $p_0 > 1$, uniformly in K, with terms which tend to 0 as $\bar{g} \to 0$.

The amplitudes are at most $\bar{\mathfrak{a}}$. Adding the five parts and using $P(C_i)/r(\mathcal{C}) \le 2P_q(C_i)$ gives (7.32), with the factor 3 which this costs contained in C'_E and C''_E. *q.e.d.*

Definition 7.12 (Normalisation)**.** The normalisation $\kappa_K^{(q)}(C,\chi)$ of a loop C at level q with the character χ is the product (1.5) of the step factors $\kappa_k(C,\chi)$ of Definition 7.8, $1 \le k \le K'$.

The normalisation removes from the expectation a perimeter term and a term for each corner, and nothing else. Since the lengths of the segments scale with the steps, $N_v^{(j-1)}(C) = L^{-(j-1)}N_v^{(0)}(C)$, the definitions give the following description, which shows which divergences are removed.

Proposition 7.13 (Structure of the Normalisation)**.** *For a loop C at level q with the character χ and $K > j_q$,*

$$\log \kappa_K^{(q)}(C,\chi) = \sum_{v\in\mathcal{V}} N_v^{(0)}(C)\, m_{v,K} + \sum_{c\in\mathcal{K}(C)} z_{v_c,w_c,K}\ , \tag{7.34}$$

with the numbers

$$m_{v,K} \coloneqq \sum_{j=1}^{K'} L^{-(j-1)}\Big(\frac{\kappa_v^{(j)}}{\Pi_j} + \mathbf{1}_{\mathbf{R}}(j)\mu_{j,v}^{\mathbf{R}}\Big)\ , \qquad z_{v,w,K} \coloneqq \sum_{j=1}^{K'}(\zeta_{v,w}^{(j)} + \mathbf{1}_{\mathbf{R}}(j)\zeta_{j,v,w}^{\mathbf{R}})\ , \tag{7.35}$$

which depend on K, $\bar{g}$, q, the character, and the constants of the construction, but not on C or on the torus, and satisfy $|m_{v,K}| \le \sum_{j\le K'} L^{-(j-1)}(C_{S'}\mathfrak{a}_j + C_E^R e^{-p_0(g_j)/4})$ and $|z_{v,w,K}| \le \sum_{j\le K'}(C_\zeta \mathfrak{a}_j + C_E^R e^{-p_0(g_j)/4})$.

Proof. Insert $N_v^{(j-1)}(C) = L^{-(j-1)}N_v^{(0)}(C)$ into (7.23) and sum over j. The bounds are Lemma 7.7, its counterpart at the step 1 [1, (5.31), (6.5)], and Proposition 7.10. The dependence on q is through $K' = K - j_q$ only. *q.e.d.*

The number $N_v^{(0)}(C)$ is the length of the segments of orientation v in fine units, of order L^K, so that the first sum in (7.34) is the self-energy of the loop, proportional to the physical length of each orientation divided by the lattice spacing, and the second sum, which grows at most like $\sum_j g_j^2(\log g_j^{-2})^{6p_0}$ in K, is the renormalisation of the corners.

8 The Identity at the Stopping Scale and the Uniform Bound

We now stop the induction and read off what it says about the renormalised expectation. Two features of the problem seem to stand in the way of a bound which is uniform in K.

For the normalisation $\prod_i \kappa_K^{(q)}(C_i, \chi_i)$ we have no bound, from above or from below, which is uniform in K. By Proposition 7.13 its logarithm is a sum of the counts $N_v^{(0)}(C)$, which are proportional to the lengths of the loops in the units of the fine lattice and grow like L^K, multiplied by the numbers $m_{v,K}$, and the bounds of that proposition on $m_{v,K}$ do not make these products small or even bounded. Whether the normalisation in fact tends to zero or to infinity is not decided by these estimates; for a trivial character, for which the observable is the constant one, it is constant. What matters is that an estimate of $\mathbb{E}_K[W_{\mathcal{C}}]$ which is not exact in the normalisation is of no use once it is divided by a number about which only this much is known. The terms of the density with large fields, on the other hand, are not small by counting: in four dimensions the factor which a large field contributes "does not give any positive power of ε" [9, p. 244], while the number of places at which a large field may occur grows like ε_K^{-4}. Both difficulties disappear at the stopping step K'. The normalisation is the same number in every term of the dressed loops, so that it can be divided out exactly; and the density at the step K' is a function of the block field on the lattice $\mathbb{T}_{j_q}$, whose spacing uL^{-j_q} does not depend on K, so that the terms with large fields can be compared with the term without them as whole integrals over a fixed space, at a price which is exponential in the number of sites of $\mathbb{T}_{j_q}$ and in nothing else. The outcome is an exact identity, Theorem 8.1, and from it the bound, Theorem 8.7.

Throughout this section $\mathcal{C} = ((C_1, \chi_1), \dots, (C_n, \chi_n))$ is an ensemble as in Theorem 1.5, and

$$K \geq K_0(\mathcal{C}) \coloneqq \max(j_q + 1,\, K_{\text{tor}}(\mathcal{C}))\ , \tag{8.1}$$

so that $K' = K - j_q \geq 1$ and, by Corollary 6.2, the ensemble is dressed at the step K'. We use the units of the step K', in which the lattice of the block field $V \coloneqq V_{K'}$ is $\mathbb{T}_{j_q}$ with the spacing 1 and

$$N_q \coloneqq \frac{T}{uL^{-j_q}} = 2L^{m_T + j_q} \tag{8.2}$$

sites per side, and the fine lattice $\mathbb{T}_K$ has the spacing $L^{-K'}$. Every loop of $\mathcal{C}$ is a path of bonds of $\mathbb{T}_q \subset \mathbb{T}_{j_q}$, with $P_{K'}(C_i) = P_q(C_i) L^3 M R_{K'-1}$ bonds of $\mathbb{T}_{j_q}$ by Lemma 3.14, and we write $P_{K'}(\mathcal{C}) \coloneqq \sum_i P_{K'}(C_i)$. We write $g_{K'}$ for the running coupling of the step K'; it depends on K, but by (3.9) with $n - m = j_q$ it satisfies $g_{K'} \leq (1 + \beta_0) g_K = (1 + \beta_0)\bar{g}$, and by Lemma 3.2(i) $\log g_{K'}^{-2} \leq \log \hat{g}_{K'}^{-2}$, where $\hat{g}_{K'} = \hat{g}_{K - j_q}$ does not depend on K. The parameters $\varepsilon_{K'}$, $R_{K'}$, $\alpha_{0,K'}$, $\alpha_{1,K'}$ of the step K' are functions of $\hat{g}_{K'}$ (Definition 3.11), and they do not depend on K either.

8.1 The Identity

A *labelled term* ℓ of $\varrho_{K'}$ is a term of the expansion (3.23) at $k = K'$ with given labels: the sequences of domains, with the refined labels of Section 5.2 at the first step, together with the labels of the **R**-operations, a renormalised component being removed from the sequences [11, (0.3), p. 176]. We write

$$\varrho_\ell(V) \coloneqq \chi_{K'}(\Omega_{K'})\, \mathbf{T}_{K'}(\ell)[\exp A_{K'}]\ , \qquad Z_\ell \coloneqq \int \varrho_\ell(V)\, \mathrm{d}V\ , \tag{8.3}$$

for the density of ℓ and its integral, $\mathrm{d}V$ being the normalised Haar measure on $G^{\mathcal{B}}$ with $\mathcal{B}$ the set of the positive bonds of $\mathbb{T}_{j_q}$. For a function F of V and of the outer variables of ℓ, and for $Z_\ell > 0$, we put

$$\langle F \rangle_\ell \coloneqq \frac{1}{Z_\ell} \int \chi_{K'}(\Omega_{K'})\, \mathbf{T}_{K'}(\ell)[F \exp A_{K'}]\, \mathrm{d}V\ . \tag{8.4}$$

The term ℓ_0 is the one whose large-field region $Z_{K'} = \Lambda^c_{K'}$ is empty; by the nesting (3.19) it has $\Omega_j = \Lambda_j = \mathbb{T}$ for every $j \le K'$, and it collects, by [11, (0.6), p. 176], all the histories whose large-field components have been renormalised. For a labelled term ℓ we write $Z(\ell)$ for the set of the connected components of its large-field region at the step K', so that $Z(\ell) = \emptyset$ if and only if $\ell = \ell_0$.

Theorem 8.1 (Identity at the Stopping Scale)**.** *Under the hypotheses of Theorem 1.5, let $K \ge K_0(\mathcal{C})$. The numbers $w_\ell := Z_\ell / \sum_{\ell'} Z_{\ell'}$ are nonnegative and $\sum_\ell w_\ell = 1$. With $\widehat{\mathcal{W}}_\ell$ the dressed loop of the step K' on ℓ divided by $\prod_i \kappa^{(q)}_K(C_i, \chi_i)$,*

$$W_K(\mathcal{C}) = \sum_{\ell \,|\, Z_\ell > 0} w_\ell \, \langle \widehat{\mathcal{W}}_\ell \rangle_\ell \,, \qquad \widehat{\mathcal{W}}_\ell = e^{\mathcal{E}_{K'}(\ell)} \Big[\prod_{i=1}^{n} f^{(i)}_{K'} + \sum_{\mathfrak{B} \neq \emptyset} Q_{K'}(\mathfrak{B}) \, w_{K'}(\mathfrak{B}) \Big] \,. \tag{8.5}$$

On the term ℓ_0 the exponent $\mathcal{E}_{K'}(\ell_0) = E_{\mathrm{const},K'}$ is a number, and ϱ_{ℓ_0} is a function of the field V on $\mathcal{B}$ alone. Consequently, with the measure μ_K on $G^{\mathcal{B}}$ of total mass w_{ℓ_0} defined by

$$\mathrm{d}\mu_K(V) := \frac{\varrho_{\ell_0}(V) \, \mathrm{d}V}{\sum_\ell Z_\ell} \,, \tag{8.6}$$

and with $W_{\mathcal{C}}(V)$ the observable (1.4) of the field V on $\mathbb{T}_{j_q}$,

$$W_K(\mathcal{C}) = e^{E_{\mathrm{const},K'}} \int W_{\mathcal{C}}(V) \, \mathrm{d}\mu_K(V) + r_K(\mathcal{C}) \,, \tag{8.7}$$

where

$$r_K(\mathcal{C}) := e^{E_{\mathrm{const},K'}} \int \Big[\prod_{i=1}^{n} f^{(i)}_{K'} - W_{\mathcal{C}}(V) + \sum_{\mathfrak{B} \neq \emptyset} Q_{K'}(\mathfrak{B}) \, w_{K'}(\mathfrak{B}) \Big] \mathrm{d}\mu_K(V) + \sum_{\ell \neq \ell_0 \,|\, Z_\ell > 0} w_\ell \, \langle \widehat{\mathcal{W}}_\ell \rangle_\ell \,. \tag{8.8}$$

The identity is exact, and it is a convex combination: $W_K(\mathcal{C})$ is an average, with the weights w_ℓ, of expectations of the normalised dressed loops. The measure μ_K lives on the space $G^{\mathcal{B}}$, which does not depend on K, and the function $W_{\mathcal{C}}(V)$ which it integrates in the leading term does not depend on K either; everything that moves with K is in the measure, in the number $E_{\mathrm{const},K'}$, and in the remainder.

Proof. We first show that every term is nonnegative. At real configurations the test functions of the construction take values in $[0, 1]$ (Lemma 3.8(i)), the operation $\mathbf{T}_{K'}(\ell)$ consists of integrals against nonnegative measures (Haar measures, δ-functions of the block averaging, Gaussian weights, and characteristic functions), and the effective action is real; Bałaban proves that the densities are positive, "hence the denominators are positive, and the operation $\mathbb{R}$ is well defined" [11, p. 176], and the terms which the **R**-operation produces are such a term multiplied by the exponential of a real sum [11, (0.6), p. 176]. Hence $Z_\ell \ge 0$, and (8.4) is the expectation with respect to a probability measure on the variables of ℓ. Since every operation of the construction preserves the integral of a density (Section 5.1), $\sum_\ell Z_\ell = \int \varrho_{K'} \, \mathrm{d}V = Z_K > 0$, and the w_ℓ are nonnegative numbers with sum one.

Next we use the numerator. By (5.2) at $k = K'$ and Definition 5.2,

$$Z_K(\mathcal{C}) = \int N_{K'}(V) \, \mathrm{d}V = \sum_\ell \int \chi_{K'}(\Omega_{K'}) \, \mathbf{T}_{K'}(\ell) [\mathcal{W}_{K'}(\ell) \exp A_{K'}] \, \mathrm{d}V = \sum_{\ell \,|\, Z_\ell > 0} Z_\ell \, \langle \mathcal{W}_{K'}(\ell) \rangle_\ell \,, \tag{8.9}$$

where a term with $Z_\ell = 0$ contributes nothing, since it is the integral of the function $\mathcal{W}_{K'}(\ell)$, which is bounded on the support of the term by Lemma 8.6 below, against a nonnegative measure of total mass zero. The normalisation in front of the dressed loop is the same number on every term, namely $\prod_i \kappa^{(K')}(C_i, \chi_i) = \prod_i \kappa_K^{(q)}(C_i, \chi_i)$ by (5.10) and (1.5), and it is not zero. Dividing (8.9) by Z_K and by this number, and using (5.1) and Definition 1.3, we obtain (8.5).

On the term ℓ_0 there are no large-field data at any step. The carrier of the loop's integral is therefore the whole torus at every step, the distances $d^{(j)}(x)$ in (5.20) are infinite, and every local term $e_X^{(j)}$ vanishes, so that $\mathcal{E}_{K'}(\ell_0) = E_{\text{const},K'}$; for $j \geq 2$ this is also the statement $E_j^{\partial}(\ell_0) = 0$ of the definition of E_j^{∂} in Section 7.1. The operation $\mathbf{T}_{K'}(\ell_0)$ holds no integrations, so that $\varrho_{\ell_0}(V) = \chi_{K'}(\mathbb{T}) \exp A_{K'}(U_{K'}(V))$, where the $\mathbf{R}$-part and the boundary terms left by renormalised components are functions of V [11, (0.6), p. 176], [12, p. 392]. The insertions on ℓ_0 are functions of V as well, since there are no outer variables. Separating the term ℓ_0 in (8.5) and adding and subtracting $e^{E_{\text{const},K'}} \int W_{\mathcal{C}} \, \mathrm{d}\mu_K$ gives (8.7) with the remainder (8.8). *q.e.d.*

8.2 The Coarse Loop

On the term ℓ_0 the leading words $f_{K'}^{(i)} = \chi_i(U_{K'}(V)(C_i))$ are the Wilson loops of the background field, a field on the fine lattice which depends on K, while the leading term of (8.7) contains the Wilson loops of V itself. The two differ by little, because V is the K'-fold block average of $U_{K'}(V)$ and $U_{K'}(V)$ is close to flat on the scale of a plaquette of $\mathbb{T}_{j_q}$. To quantify this we need a comparison between an iterated average and a transport, which we now prove.

The average of [8, (0.4), p. 253] maps a field U on a lattice to the field $\bar{U} = M(U)$ on the lattice whose spacing is L times larger. With $d = 4$, let $B(y)$ be the set of the L^d points of the finer lattice in the cube of side L centred at y [8, (0.3), p. 252], $\mathbf{G}(y, x)$ the family of the shortest paths from y to x which run through the coordinate directions in some order, one path for each order, $[x, x']$ the translate of c to x, and $U(c)$ the transport along the straight path of L bonds of c [8, p. 252]. For $x \in B(c_-)$, $\Gamma \in \mathbf{G}(c_-, x)$, and $\Gamma' \in \mathbf{G}(c_+, x')$ put $\ell(x, \Gamma, \Gamma') \coloneqq \Gamma \cup [x, x'] \cup (-\Gamma') \cup (-c)$, a closed path at c_-, and $v(x, \Gamma, \Gamma') \coloneqq L^{-d} |\mathbf{G}(c_-, x)|^{-1} |\mathbf{G}(c_+, x')|^{-1}$. Then

$$\bar{U}(c) = \exp\Big[i \sum_{x, \Gamma, \Gamma'} v(x, \Gamma, \Gamma') \, \frac{1}{i} \log U(\ell(x, \Gamma, \Gamma')) \Big] U(c) \, , \tag{8.10}$$

and the weights $v(x, \Gamma, \Gamma')$ are nonnegative with sum one. We write M^j for the j-fold iterate of M, the average of order j, as the averages of higher order are defined recursively in [4, (43), p. 24], and $T_c(U)$ for the transport of U along the straight path of the fine bonds of a bond c of a coarser lattice. Here and below $\|\cdot\|$ is the operator norm, $\mathfrak{g}$ is the Lie algebra of G realised as the Hermitian matrices X with $e^{iX} \in G$, $r_G \in (0, 1/10]$ is a radius such that the principal logarithm of every $W \in G$ with $\|W - 1\| \leq r_G$ lies in $i\mathfrak{g}$, which exists because the exponential map is a local diffeomorphism at 0, and for a unitary representation π of G and $Y \in \mathfrak{g}$ we write $\pi(e^{iY}) = e^{i\pi_*(Y)}$, so that $\pi_*(Y) = -i \, \mathrm{d}\pi(iY)$ and, with the number c_π of (1.7), $\|\pi(e^{iY}) - 1\| \leq \|\pi_*(Y)\| \leq c_\pi \|Y\|$.

Three elementary facts about unitary matrices will be used. For unitary W and g, $\|gWg^{-1} - 1\| = \|W - 1\| = \|W^{-1} - 1\|$, and for any matrices $a_1, \dots, a_m$,

$$\Big\| \prod_{l=1}^{m} (1 + a_l) - 1 \Big\| \leq \prod_{l=1}^{m} (1 + \|a_l\|) - 1 \, . \tag{8.11}$$

Secondly, if W is unitary and $\|W - 1\| \leq e^s - 1$ with $0 \leq s \leq 1/10$, then the principal logarithm $W = e^{iX}$ has

$$\|X\| \leq 1.06 \, s \ : \tag{8.12}$$

an eigenvalue $e^{i\phi}$ of W with $\phi \in (-\pi, \pi]$ has $2|\sin(\phi/2)| = |e^{i\phi} - 1| \le e^s - 1 \le 1.052\, s$, because $(e^s - 1)/s$ is increasing and equals $1.0517\ldots$ at $s = 1/10$, and $2\arcsin(y/2) \le y(1 - y^2/4)^{-1/2} \le 1.002\, y$ for $0 \le y \le 0.11$. Thirdly, $\|e^{iX} - 1\| \le \|X\|$ for Hermitian X, since $|e^{i\phi} - 1| \le |\phi|$; consequently, for $X, F \in \mathfrak{g}$ with $s := \|X\| + \|F\| \le \min(1/10, \log(1 + r_G))$, (8.11) gives $\|e^{iX}e^{iF} - 1\| \le e^s - 1 \le r_G$, so that $e^{iX}e^{iF} = e^{iE}$ with $E \in \mathfrak{g}$ and $\|E\| \le 1.06\, s$ by (8.12).

Lemma 8.2 (The Coarse Loop)**.** *Let $L \ge 21$ be odd and $k \ge 1$, let $a_4 := 7.2$, and let U be a G-valued field on a periodic lattice of spacing η, with periods which are multiples of $L^k\eta$, such that $\|U(\partial p) - 1\| \le t$ for every plaquette p and*

$$a_4 L^{2k} t \le \min(1/10,\ \log(1 + r_G))\ . \tag{8.13}$$

(i) For every $0 \le j \le k$ and every bond c of the lattice of spacing $L^j\eta$ there is $E_c \in \mathfrak{g}$ with $(M^j U)(c) = e^{iE_c} T_c(U)$ and $\|E_c\| \le a_4 L^{2j} t$.

(ii) For every closed path C of P bonds of the lattice of spacing $L^k\eta$ and every character $\chi = (\dim \pi)^{-1} \operatorname{Tr} \pi$ of an irreducible unitary representation π,

$$|\chi(U(C)) - \chi((M^k U)(C))| \le e^{c_\pi a_4 P L^{2k} t} - 1\ . \tag{8.14}$$

The constant $a_4 = 0.3(d-1)(d+4)$ at $d = 4$ does not grow with j: each level of averaging loses only the factor $(d+3)/L$ on the deviation of the previous level, and the deviation of the plaquettes of the new level, which is L^2 times larger, is what sets the scale.

Proof. (i) We argue by induction on j, the case $j = 0$ being $E_c = 0$. Let $0 \le j < k$, assume (i) at the level j, and write $W := M^j U$, a G-valued field on the lattice of spacing $L^j\eta$, which we use as the unit of length in this step. Let c be a bond of the lattice of spacing $L^{j+1}\eta$, from c_- to $c_+ = c_- + Le_\mu$, so that $(M^{j+1}U)(c) = e^{iX_c} W(c)$ by (8.10), with X_c the convex combination in the exponent and $W(c)$ the product of W over the L bonds b of c.

The transport. By the hypothesis of the induction, $W(c) = \prod_{b \subset c} e^{iE_b} T_b(U) = R_c\, T_c(U)$ with $R_c := \prod_{b \subset c} g_b e^{iE_b} g_b^{-1} \in G$, where g_b is the transport of U along the part of c before b. By (8.11) and the third fact, $\|R_c - 1\| \le e^{s_c} - 1$ with $s_c := a_4 L^{2j+1} t$, and $s_c \le a_4 L^{2k-1} t \le \min(1/10, \log(1 + r_G))/L$ by (8.13), since $2j + 1 \le 2k - 1$. Hence $R_c = e^{iF_c}$ with $F_c \in \mathfrak{g}$ and $\|F_c\| \le 1.06\, a_4 L^{2j+1} t$ by (8.12).

The loops of the average. Let $x = c_- + n$ with $n \in \mathbb{Z}^d$ and $|n_\nu| \le (L-1)/2$, which is the condition $x \in B(c_-)$ for odd L, let $\Gamma \in \mathbf{G}(c_-, x)$ and $\Gamma' \in \mathbf{G}(c_+, x')$, and let $\ell := \Gamma \cup [x, x'] \cup (-\Gamma') \cup (-c)$. The loop ℓ has at most $(d+2)L$ bonds: Γ and Γ' have $|n|_1 \le d(L-1)/2$ each, and $[x, x']$ and c have L each. As above, $W(\ell) = R_\ell\, U(\ell)$, where $U(\ell)$ is the transport of U around the fine loop obtained by replacing each bond of ℓ by its straight fine path, and $\|R_\ell - 1\| \le e^{(d+2)a_4 L^{2j+1} t} - 1$. To bound $U(\ell)$ we contract ℓ across plaquettes of the lattice of spacing $L^j\eta$. Let Γ_+ be the translate of Γ by Le_μ, which lies in $\mathbf{G}(c_+, x')$. The paths Γ' and Γ_+ have the same displacement n, and passing from the order of the directions in Γ' to that in Γ_+ by transpositions of adjacent straight segments, each pair of directions $\alpha \ne \beta$ is transposed at most once and the transposition sweeps $|n_\alpha||n_\beta|$ plaquettes. This replaces Γ' by Γ_+; we then move $[x, x']$ back along Γ one bond at a time, which sweeps L plaquettes for a bond of direction $\pm e_\nu$ with $\nu \ne \mu$, and none for a bond of direction $\pm e_\mu$, after which the

loop is trivial. The number of plaquettes swept is therefore at most

$$\begin{aligned} A_L &\coloneqq \sum_{\alpha<\beta} |n_\alpha||n_\beta| + L(|n|_1 - |n_\mu|) \\ &\le \frac{d(d-1)}{2}\Big(\frac{L-1}{2}\Big)^2 + \frac{(d-1)L^2}{2} \le \frac{(d-1)(d+4)}{8}L^2 = \frac{a_4}{2.4}L^2 \,. \end{aligned} \tag{8.15}$$

Each sweep multiplies the transport around the loop, at its base point, by a conjugate of the transport of U around the boundary of a plaquette of the lattice of spacing $L^j\eta$, and the latter is an ordered product of conjugates of the transports around the L^{2j} fine plaquettes inside it. By (8.11) and the first fact, $\|U(\ell)-1\| \le (1+t)^{A_L L^{2j}} - 1 \le e^{A_L L^{2j} t} - 1$, and therefore $\|W(\ell)-1\| \le e^{s_\ell} - 1$ with

$$s_\ell \coloneqq A_L L^{2j} t + (d+2) a_4 L^{2j+1} t \le \Big(\frac{1}{2.4} + \frac{d+2}{L}\Big) a_4 L^{2j+2} t \le 0.71\, a_4 L^{2(j+1)} t \le a_4 L^{2k} t \,, \tag{8.16}$$

where we used (8.15) and $(d+2)/L \le 6/21$. By (8.13), $s_\ell \le \min(1/10, \log(1+r_G))$, so that $\|W(\ell)-1\| \le r_G$; since $W(\ell) \in G$, its principal logarithm lies in $i\mathfrak{g}$, and $\|(1/i)\log W(\ell)\| \le 1.06\, s_\ell$ by (8.12). The convex combination X_c therefore lies in $\mathfrak{g}$ and $\|X_c\| \le 1.06\,(A_L L^{2j} + (d+2)a_4 L^{2j+1})t$.

The new level. We have $(M^{j+1}U)(c) = e^{iX_c} e^{iF_c} T_c(U)$, and

$$\begin{aligned} \|X_c\| + \|F_c\| &\le 1.06 (A_L L^{2j} + (d+3) a_4 L^{2j+1}) t \\ &\le 1.06 \Big(\frac{1}{2.4} + \frac{d+3}{L}\Big) a_4 L^{2(j+1)} t \le 0.795\, a_4 L^{2(j+1)} t \,, \end{aligned} \tag{8.17}$$

since $(d+3)/L \le 1/3$ for $L \ge 21$. The right side is at most $0.795 \min(1/10, \log(1+r_G))$ by (8.13), so that by the third fact $e^{iX_c} e^{iF_c} = e^{iE_c}$ with $E_c \in \mathfrak{g}$ and $\|E_c\| \le 1.06 \cdot 0.795\, a_4 L^{2(j+1)} t \le a_4 L^{2(j+1)} t$. This is (i) at the level $j+1$; in particular $M^{j+1}U$ is G-valued, as the induction requires.

(ii) Let $c_1, \dots, c_P$ be the bonds of C in their order, put $T_l \coloneqq T_{c_l}(U)$ and $H_l \coloneqq T_1 \cdots T_{l-1}$, and use (i) at $j = k$, with $E_{c^{-1}} = -T_c^{-1} E_c T_c$ for a bond traversed against its orientation, which has the same norm. Then

$$(M^k U)(C) = \prod_{l=1}^{P} e^{iE_{c_l}} T_l = \Big(\prod_{l=1}^{P} H_l e^{iE_{c_l}} H_l^{-1}\Big)\, U(C) \,, \tag{8.18}$$

and since $|(\dim\pi)^{-1} \operatorname{Tr}((\pi(R)-1)\pi(H))| \le \|\pi(R)-1\|$ for unitary $\pi(H)$, (8.11) and $\|\pi(e^{iE})-1\| \le c_\pi \|E\|$ give $|\chi(U(C)) - \chi((M^k U)(C))| \le \prod_l (1 + c_\pi a_4 L^{2k} t) - 1 \le e^{c_\pi a_4 P L^{2k} t} - 1$. *q.e.d.*

We apply the lemma at the stopping step. Bałaban states the regularity of the background in a norm $|\cdot|$ on matrices; we let $c_h \ge 1$ be a constant with $c_h^{-1}|Y| \le \|Y\| \le c_h |Y|$ for all $N \times N$ matrices Y, and B_3 is the constant of [7, Theorem 1, p. 279], which depends on d and L only.

Corollary 8.3 (The Background at the Stopping Step)**.** *Let $\bar g$ be so small that*

$$2 a_4 c_h (1 + 2\kappa_s) B_3\, \varepsilon_{K'} \le \min(1/10,\ \log(1 + r_G)) \,. \tag{8.19}$$

Then on the term ℓ_0, for every real V in the support of ϱ_{ℓ_0} and every i,

$$|f^{(i)}_{K'} - W_{\chi_i}(C_i; V)| \le e^{x(C_i)} - 1 \,, \qquad x(C) \coloneqq 2 c_\pi a_4 c_h (1 + 2\kappa_s) B_3\, P_{K'}(C)\, \varepsilon_{K'} \,, \tag{8.20}$$

and $|\prod_i f^{(i)}_{K'} - W_{\mathcal{C}}(V)| \le \sum_i (e^{x(C_i)} - 1)$. The numbers $x(C_i)$ do not depend on K.

Proof. On ℓ_0 the determining set is the whole lattice $\mathbb{T}_{j_q}$, so that the background field $U_{K'} = U_{K'}(V)$ is a minimiser of the action on the fine fields U with $M^{K'}(U) = V$ on every bond of $\mathbb{T}_{j_q}$ [9, (2.11), (2.12), p. 256]. Its plaquettes satisfy $|\partial U_{K'} - 1| < 2B_3\varepsilon_{K'}(L^{-1}\eta)^2$ on $\Omega_{K'} = \mathbb{T}$ [9, p. 268], where $\eta = L^{-(K'-1)}$ is the fine spacing in the units of the step $K'-1$, so that $(L^{-1}\eta)^2 = L^{-2K'}$ is the area of a fine plaquette in the units of the step K'. This bound is drawn from passed test functions; for the soft test functions of the construction it holds with $\varepsilon_{K'}$ replaced by $(1+2\kappa_s)\varepsilon_{K'}$ (Supplement, Proposition S.2). Hence the hypotheses of Lemma 8.2 hold with $k = K'$, $\eta = L^{-K'}$, and $t \coloneqq 2c_h(1+2\kappa_s)B_3\varepsilon_{K'}L^{-2K'}$, for which $a_4L^{2K'}t$ is the left side of (8.19), and the periods of $\mathbb{T}_K$ are multiples of the spacing of $\mathbb{T}_{j_q}$. On ℓ_0 the leading word is $f^{(i)}_{K'} = \chi_i(U_{K'}(C_i))$, and (8.14) with $P = P_{K'}(C_i)$ gives (8.20), because $c_\pi a_4 P_{K'}(C_i)L^{2K'}t = x(C_i)$. For the product we telescope: $|\prod_i a_i - \prod_i b_i| \le \sum_i |a_i - b_i|$ for complex numbers of modulus at most one, and $|\chi_i(g)| \le 1$ for $g \in G$. Finally $P_{K'}(C_i)$ and $\varepsilon_{K'}$ do not depend on K. *q.e.d.*

8.3 The Weight of the Large-Field Terms

We next show that the terms $\ell \neq \ell_0$ carry a small total weight, even after their integrals are multiplied by the factors which the dressed loop acquires near the large-field regions. Since $\mathbb{T}_{j_q}$ is fixed, this can be done by comparing whole integrals: the integral of ℓ_0 is bounded below by the Haar volume of a ball of configurations, and the integral of every other term above by the supremum of its small-field exponent, each at a price exponential in N_q^4. The large-field factors of Bałaban's construction then pay for the history of each component, with half of their gain left over.

We need the following quantities, for a component $X \in Z(\ell)$. The *history* of X consists of the components of the large-field regions Z_n, $n \le K'$, of the term which lie in X, together with the new large-field regions Z' created at the steps $j' \le K'$ which are merged into X; we write $|Y|_n$ for the number of points of the scale n in a set Y, $\#_n(Y) \coloneqq |Y|_n(MR_n)^{-4}$ for the number of cubes of side MR_n of the scale n in a union Y of such cubes, and $d'_n(Y)$ for its linear size in terms of these cubes [12, p. 380]. Bałaban's factor [12, (1.79), p. 383] contains for each new region Z' created at the step j' the gain $\exp(-\frac{1}{2}\gamma_0 A_1^2 p_0^2(g_{j'})(d'_{j'}(Z')+1))$, which holds for the construction of this paper with $\frac{1}{2}\gamma_0$ replaced by $\frac{1}{2}\gamma_0/\rho \ge \frac{1}{3}\gamma_0$ (Proposition 3.12); here γ_0, A_1, and $p_0(g)$ are those of [12]. The regions created at the first step carry instead the gain $\exp(-\frac{1}{2}c_4\vartheta_V^2 A_1^2(\log g_1^{-2})^{2p_0}(d'_1(Z')+1))$ of the first step at the retreated region, with two constants c_4, ϑ_V of that step (Supplement, Proposition S.7(iii)). We put

$$P(X) \coloneqq \sum_{n \le K'} L^4 |Z_n \cap X|_n \,, \qquad G(X) \coloneqq \sum_{Z'} \Upsilon_{j'}(d'_{j'}(Z')+1) \,, \tag{8.21}$$

with $\Upsilon_{j'} \coloneqq \frac{1}{6}\gamma_0 A_1^2 p_0^2(g_{j'})$ for $j' \ge 2$ and $\Upsilon_1 \coloneqq \frac{1}{4}c_4\vartheta_V^2 A_1^2(\log g_1^{-2})^{2p_0}$, the *history volume* of X and half of its gains, and we let $C_P \ge 2$ be the constant of Appendix A.

Proposition 8.4 (The Weight of the Large-Field Terms)**.** *Let $\bar{g}$ be sufficiently small, depending on the constants of the construction, and let $K > j_q$. Then $Z_{\ell_0} > 0$, and for every choice of numbers $\omega(X)$, one for each component X of each term $\ell \neq \ell_0$, which depend on X and its history only and satisfy $1 \le \omega(X) \le \exp(C_P P(X) + \frac{1}{2}G(X))$,*

$$\sum_{\ell \neq \ell_0} Z_\ell \prod_{X \in Z(\ell)} \omega(X) \le \delta_{\mathrm{LF}} Z_{\ell_0} \,, \qquad \delta_{\mathrm{LF}} \coloneqq e^{2C_K}(e^{s_{\mathrm{LF}}} - 1) \,, \tag{8.22}$$

with

$$C_K \coloneqq C_{\mathrm{fs}} N_q^4 (1 + \log g_{K'}^{-1}) \,, \qquad s_{\mathrm{LF}} \coloneqq 2N_q^4 (MR_{K'})^{-4} \exp(-c_{\mathrm{LF}}\, p_0(g_{K'})) \,, \tag{8.23}$$

$c_{\mathrm{LF}} \coloneqq (\rho(1+\beta_0))^{-1}$, *and* C_{fs} *the constant of* (8.29) *below. In particular* $1 - w_{\ell_0} \le \delta_{\mathrm{LF}}$, *and* $\delta_{\mathrm{LF}} \le \bar{\delta}_{\mathrm{LF}}$, *where* $\bar{\delta}_{\mathrm{LF}}$ *is the number obtained by replacing* $\log g_{K'}^{-1}$ *by* $\frac{1}{2}\log \hat{g}_{K'}^{-2}$ *and* $p_0(g_{K'})$ *by* $p_0((1+\beta_0)\bar{g})$, *which depends on* q, T/u, $\bar{g}$, *and the constants of the construction, and not on* K.

Proof. We follow the terms through four steps: a common normalisation, an upper bound for each Z_ℓ, a lower bound for Z_{ℓ_0}, and the sum over the terms. We write $g_a \coloneqq g_{K'}$, and assume $g_a \le 1$.

The normalisation. Every term is written in Bałaban's completed form: for its components renormalised at the step K' the missing terms of the effective action "with localization domains intersecting the domain Z" are added "to complete the effective action" and subtracted from the remaining expression [12, p. 377, (1.70)], and its surviving components are written with the normalised operations of [12, (1.79), p. 383], which holds for all large-field regions and not only for those which satisfy the conditions of the **R**-operation [12, p. 383–384]. A term then reads $\chi_{K'} \prod_{X\in Z(\ell)} \mathbf{T}'_{K'}(X) \exp A'_{K'}$, where the constants subtracted in the completion, "the sum of normalization terms and vacuum energy terms with localizations intersecting the large field domain Z" [12, p. 378], stand inside $\mathbf{T}'_{K'}(X)$. We divide every term by $e^{-E^{\ell_0}}$, with E^{ℓ_0} the constant of the effective action of ℓ_0, a single number for all terms, which changes neither the w_ℓ nor (8.22). The constant E^ℓ of a term is obtained by subtracting localised one-step vacuum energies from the initial constant [9, p. 262], so that $E^{\ell_0} - E^\ell$ is the sum of the localised normalisation and vacuum-energy terms whose localisations meet the history of a component of ℓ; a normalisation constant is $O(\log g_n^{-2})$ per point of the scale n and the vacuum energies are $O(1)$ per point [12, p. 380], [9, p. 264], and with a constant C_λ of d, G, and these bounds which absorbs the enlargement of the localisations by a layer of cubes and their exponential tails,

$$|E^{\ell_0} - E^\ell| \le \sum_{X\in Z(\ell)} \sum_{n\le K'} C_\lambda (1 + \log g_n^{-2}) L^4 |Z_n \cap X|_n \,. \tag{8.24}$$

The upper bound. The action is nonnegative, so that $-A(1/g_{K'}^2, U_{K'}) \le 0$. On the support of the term, the regular part, the **R**-part, and the boundary part are bounded by [9, (2.45), (2.46), (2.48), pp. 263–264] by $C_{\mathcal{E}} \sum_n |\Gamma_n|$, with $C_{\mathcal{E}} \coloneqq E_1(1 - L^{-\beta})^{-1} + 1 + 2B_\partial$ and B_∂ the constant printed B_1 in [9, (2.47)]. The couplings of the completed action to the exterior of a renormalised component are at most $\alpha e^{-(1+2\beta)\kappa d_{k,Z}(Y)}$, with α arbitrarily small for γ small [12, (1.68), p. 377], and summed over the localisations which meet a cube of side $MR_{K'}$ they are at most a constant $C_\partial \le 1$, so that they contribute at most $C_\partial N_q^4$. The last layer $\Gamma_{K'}$ of the determining set lies in $\mathbb{T}_{j_q}$ and has at most N_q^4 points, and every earlier layer $\Gamma_n \subset \Omega_{n+1}^c \subset Z_{n+1}$ lies in a component of $Z(\ell)$, with $|\Gamma_n \cap X|_n \le L^4 |Z_{n+1} \cap X|_{n+1}$, since the large-field regions grow and merge until they are renormalised, and a renormalised component leaves the sequence [11, (0.3), p. 176]. Together with (8.24), the exponent of the term, less the common constant, is at most $(C_{\mathcal{E}} + C_\partial) N_q^4 + \sum_{X\in Z(\ell)} P^c(X)$ with the *history cost*

$$P^c(X) \coloneqq \sum_{n\le K'} c_n L^4 |Z_n \cap X|_n \,, \qquad c_n \coloneqq C_{\mathcal{E}} + C_\lambda (1 + \log g_n^{-2}) \,. \tag{8.25}$$

Since $\mathbf{T}'_{K'}(X)$ is a positive operation, $\mathbf{T}'_{K'}(X) F \le (\mathbf{T}'_{K'}(X)1) \sup F$ for $F \ge 0$ [12, (1.73), p. 380], and integrating over V with the normalised Haar measure,

$$Z_\ell \le M_\ell \coloneqq e^{C_K} \prod_{X\in Z(\ell)} \left[\sup \mathbf{T}'_{K'}(X)1 \cdot e^{P^c(X)} \right] , \tag{8.26}$$

where we used $(C_{\mathcal{E}} + C_\partial)N_q^4 \le C_K$, which holds for the constant (8.29) below.

The lower bound. Let $\mathfrak{B} \coloneqq \{V \mid \|V(b)-1\| < r_{\mathfrak{B}} \text{ for every } b \in \mathcal{B}\}$ with $r_{\mathfrak{B}} \coloneqq g_a/(10c_hB_3)$. The averaged extension $M^{\cdot}(Q^{s*}_{K'}V)$ of [9, (2.16), p. 257], from which the minimisers are built, takes on each bond either the value 1 or a value $V(c)^{\pm 1}$: the field $Q^{s*}_{K'}V$ equals 1 on the bonds inside the blocks and $V(c)$ on every bond which joins the two blocks of c [9, (1.3), p. 246], and in (8.10) the paths Γ, Γ' stay inside blocks, while $[x, x']$ and the straight path of a bond each cross the face between two blocks at most once. Hence on $\mathfrak{B}$ every plaquette of the block data is a product of at most four factors within $r_{\mathfrak{B}}$ of 1, so that by (8.11) its deviation is at most $c_h((1 + r_{\mathfrak{B}})^4 - 1) \le 5c_h r_{\mathfrak{B}} = g_a/(2B_3)$ in the norm $|\cdot|$. Then [7, Theorem 1, p. 279], with its $\varepsilon_1 = g_a/(2B_3) \le a_1$ for g_a small, puts the local minimisers and the minimiser $U_{K'}(V)$ in the space of the configurations with $|U(\partial p)-1| < B_3\varepsilon_1\eta^2 = (g_a/2)\eta^2$ for every fine plaquette, $\eta = L^{-K'}$. So every test function of the term ℓ_0 passes, because $g_a/2 \le (\rho^{1/2}/2)\hat{g}_{K'} < \hat{g}_{K'}p_0(\hat{g}_{K'}) = \varepsilon_{K'}$ by Lemma 3.2(i), and $\chi_{K'}(\mathbb{T}) = 1$ on $\mathfrak{B}$. For unitary W, $1 - \operatorname{Re}\operatorname{tr} W = \frac{1}{2N}\operatorname{Tr}((W-1)^*(W-1)) \le \frac{1}{2}\|W-1\|^2$, and there are $6N_q^4L^{4K'}$ fine plaquettes, so that on $\mathfrak{B}$

$$\frac{1}{g_a^2}A(U_{K'}(V)) \le \frac{1}{g_a^2} \cdot 6N_q^4L^{4K'} \cdot \frac{c_h^2}{2}\Big(\frac{g_a}{2}\Big)^2 L^{-4K'} = \frac{3}{4}c_h^2N_q^4 \; ; \tag{8.27}$$

and $\mathbf{E}_{K'} + \mathbf{R}_{K'} \ge -C_{\mathcal{E}}N_q^4$ by [9, (2.45), (2.46)] with $\sum_n|\Gamma_n| = N_q^4$. The Haar measure of the ball of radius $r \le 1$ about 1 in G is at least $(c_G r)^{\dim G}$, with a constant c_G of G, by the exponential coordinates, and $\mathcal{B}$ has $4N_q^4$ bonds. Therefore

$$Z_{\ell_0} \ge \exp\Big(-\frac{3}{4}c_h^2N_q^4 - C_{\mathcal{E}}N_q^4\Big)(c_G r_{\mathfrak{B}})^{4N_q^4 \dim G} \ge e^{-C_K} \; , \tag{8.28}$$

with

$$C_{\mathrm{fs}} \coloneqq 1 + c_h^2 + C_\partial + C_{\mathcal{E}} + 4\dim G(1 + \log_+(10c_hB_3/c_G)) \; , \tag{8.29}$$

a constant of d, L, G, and the constants of Bałaban's theorems, independent of K; for the last inequality of (8.28) we used $\log(c_G r_{\mathfrak{B}}) \ge -\log g_a^{-1} - \log_+(10c_hB_3/c_G)$ and $g_a \le 1$.

The component factors. Let $X \in Z(\ell)$, and let $\kappa^{\mathrm{tr}}(X)$ be the exact exponent of the factor of X in [12, (1.79)], so that $\sup \mathbf{T}'_{K'}(X)1 \le \exp(-\kappa^{\mathrm{tr}}(X) - (2/\rho)p_0(g_{j(X)}))$, with $j(X)$ the step of the first large-field region in X [12, p. 384]. By Supplement, Proposition S.11, which is Bałaban's induction for [12, (1.80), p. 384] with half of the gains set aside and the cost of each step raised,

$$\kappa^{\mathrm{tr}}(X) = \hat{\kappa}(X) + G(X) + H(X) \; , \qquad H(X) \coloneqq \sum_{n \le K'} \lambda_n \sum_{Z''} M^4R_n^5 d'_n(Z'') \; , \tag{8.30}$$

with $\lambda_n \coloneqq 1 + 3 \cdot 2^3L^4C_\lambda(1 + \log g_n^{-2})$, the inner sum over the components Z'' of Z_n in X, where $\hat{\kappa}(X) \ge 0$, and $\hat{\kappa}(X) \ge \hat{c}_0 \#_{K'}(X)$ if X survives the stopping step, that is, if it fails one of the conditions of the **R**-operation, with $\hat{c}_0 \coloneqq M^4R_{K'+1}^5(3 \cdot 2^3)^{-1}(2L^2)^{-4}$.

The history volume and the history cost are paid by $H(X)$ and half of $G(X)$:

$$P^c(X) + C_P P(X) \le H(X) + \frac{1}{2}G(X) \; , \tag{8.31}$$

provided that, for every $n \le K'$,

$$3 \cdot 2^3L^4(C_{\mathcal{E}} + C_P) \le R_n \qquad \text{and} \qquad 2^4L^4M^4R_n^4(c_n + C_P) \le \frac{1}{2}\Upsilon_n \; . \tag{8.32}$$

Indeed, a component Z'' of Z_n in X with $d'_n(Z'') \neq 0$ has $\#_n(Z'') \leq 3 \cdot 2^3 d'_n(Z'')$ [12, p. 384], so that its points cost $(c_n + C_P)L^4|Z''|_n \leq 3 \cdot 2^3 L^4(c_n + C_P)M^4 R_n^4 d'_n(Z'') \leq \lambda_n M^4 R_n^5 d'_n(Z'')$, by the first condition of (8.32) and $R_n \geq 1$. A component with $d'_n(Z'') = 0$ has at most 2^4 cubes, because a tree of length zero is a point and at most 2^4 cubes of a partition contain a given point. It contains no image $S(\cdot)$ of an earlier region, since S adds ten layers of cubes [12, p. 384], so it consists of new regions created at the step n, whose gain enters $G(X)$ once; its points cost at most $2^4 L^4 M^4 R_n^4 (c_n + C_P) \leq \frac{1}{2}\Upsilon_n$, which is half of the contribution of one new region to $G(X)$, by the second condition. The conditions (8.32) hold for $\bar{g}$ small, since $R_n \geq (\log g_n^{-2})^r$ and $R_n \leq 2L(\log g_n^{-2})^r$ by Lemma 3.4, while Υ_n is a multiple of $(\log g_n^{-2})^{2p_0}$ and $2p_0 > 4r + 1$ (Appendix A).

By (8.30), (8.31), and $p_0(g_j) \geq (1 + \beta_0)^{-1} p_0(g_a)$ for $j \leq K'$, which follows from $g_j \leq (1 + \beta_0) g_a$ by (3.9) for $\bar{g}$ small,

$$\sup \mathbf{T}'_{K'}(X) 1 \cdot e^{P^c(X)} \leq \exp\Big(-\hat{\kappa}(X) - C_P P(X) - \frac{1}{2} G(X) - 2 c_{\mathrm{LF}}\, p_0(g_a) \Big) \,. \tag{8.33}$$

Let C_{ent} be a constant such that at most C_{ent}^m connected unions of m cubes contain a given cube. If X survives the stopping step, then $\hat{\kappa}(X) \geq \hat{c}_0 \#_{K'}(X) \geq (\log C_{\mathrm{ent}} + 1)\#_{K'}(X)$ for $\bar{g}$ small, since $\hat{c}_0$ grows with $R_{K'+1}$. If X is renormalised at the stopping step, then $\#_{K'}(X) \leq 100^4$ by the first condition of the **R**-operation, and $c_{\mathrm{LF}} p_0(g_a) \geq (\log C_{\mathrm{ent}} + 1) 100^4$ for $\bar{g}$ small. In both cases, for every $\omega(X) \leq \exp(C_P P(X) + \frac{1}{2} G(X))$,

$$\omega(X) \sup \mathbf{T}'_{K'}(X) 1 \cdot e^{P^c(X)} \leq \exp\Big(-c_{\mathrm{LF}}\, p_0(g_a) - (\log C_{\mathrm{ent}} + 1)\#_{K'}(X) \Big) \,. \tag{8.34}$$

The sum. The components of $Z(\ell)$ are disjoint connected unions of the $N_q^4 (M R_{K'})^{-4}$ cubes of side $M R_{K'}$ of $\mathbb{T}_{j_q}$, each of one of the two classes. The sum over the labels with given components and classes, the sequences of domains and the labels of the **R**-operations, is contained in Bałaban's bound, since "the summations over the admissible sequences can be replaced by the factors $\exp O(1)(M R_j)^{-d}|Z_j|$" in the first exponential of [12, (1.79)] [12, p. 383], and (8.34) is a bound on the resulting sum for each component. Dropping the disjointness of the components, and with the factor 2 for the two classes,

$$\begin{aligned} \sum_{\ell \neq \ell_0} M_\ell \prod_{X \in Z(\ell)} \omega(X) &\leq e^{C_K} \sum_{m \geq 1} \frac{1}{m!} \Big(\frac{2 N_q^4}{(M R_{K'})^4} e^{-c_{\mathrm{LF}} p_0(g_a)} \sum_{m' \geq 1} C_{\mathrm{ent}}^{m'} e^{-(\log C_{\mathrm{ent}} + 1) m'} \Big)^m \\ &\leq e^{C_K} (e^{s_{\mathrm{LF}}} - 1) \,, \end{aligned} \tag{8.35}$$

because $\sum_{m' \geq 1} e^{-m'} < 1$. With $Z_\ell \leq M_\ell$ from (8.26) and $e^{C_K} \leq e^{2C_K} Z_{\ell_0}$ from (8.28), this is (8.22). Taking $\omega = 1$ gives $\sum_{\ell \neq \ell_0} Z_\ell \leq \delta_{\mathrm{LF}} Z_{\ell_0} \leq \delta_{\mathrm{LF}} \sum_\ell Z_\ell$, that is, $1 - w_{\ell_0} \leq \delta_{\mathrm{LF}}$. Finally C_K increases with $\log g_a^{-1} \leq \frac{1}{2} \log \hat{g}_{K'}^{-2}$, and s_{LF} decreases with $p_0(g_a) \geq p_0((1 + \beta_0)\bar{g})$, while N_q and $R_{K'}$ do not depend on K; so $\delta_{\mathrm{LF}} \leq \bar{\delta}_{\mathrm{LF}}$. *q.e.d.*

At a fixed level the number $\bar{\delta}_{\mathrm{LF}}$ tends to zero with $\bar{g}$, although it is not needed for the bound. Indeed, $L^{j_q} = L^{q+3} M R_{K'-1}$ grows like $(\log \bar{g}^{-2})^r$, so that C_K is of the order of $(\log \bar{g}^{-2})^{4r+1}$, while $c_{\mathrm{LF}} p_0((1 + \beta_0)\bar{g})$ is of the order of $(\log \bar{g}^{-2})^{p_0}$ with $p_0 > 4r + 1$. This smallness is used in Section 12.

8.4 The Uniform Bound

It remains to bound the normalised dressed loops on each term. On ℓ_0 the bound is a number which does not depend on K, because the number of traversals at the stopping step

is fixed; on the other terms the dressed loop acquires, in addition, the costs of the insertions attached to the large-field regions and of the traversals near them, and these are paid as in Proposition 8.4. We put

$$\bar\rho \coloneqq C_\rho(1+\beta_0)^2\bar g^2\Big(1+A_1(\log((1+\beta_0)^{-2}\bar g^{-2}))^{p_0}\Big)^4 , \tag{8.36}$$

$S_w \coloneqq \sum_{t\ge1}(t+1)^3e^{-\eta_w t}$, and $S_{\mathrm{wr}}(b) \coloneqq \sum_{i\ge0} L^i e^{-bL^i}$ for $b>0$, so that $\rho_j \le \bar\rho$ for every $j\le K$ when $\bar g$ is small, because $g_j \le (1+\bar\beta_0)\bar g$ by (3.9) and $x\mapsto x(1+A_1(\log x^{-1})^{p_0})^4$ is increasing for small x. With the constants of (5.11), let

$$I_*(\mathcal C) \coloneqq P_{K'}(\mathcal C)\Big[\frac{2c_\sigma C_\chi(1+2\kappa_s)B_3S_w\,\bar\rho\,\varepsilon_{K'}}{\min(\alpha_{0,K'},\alpha_{1,K'})} + 2C_{S'}\bar\rho\,S_{\mathrm{wr}}\Big(\frac{c_{\mathrm{att}}N_q}{M}\Big)\Big] + A^\times(\mathcal C) , \tag{8.37}$$

where the last term is the total of the cross species,

$$A^\times(\mathcal C) \coloneqq P_{K'}(\mathcal C)\Big[10\,C_{\mathrm{cr}}\bar{\mathfrak a}\,e^{-\nu_\times d_\times/M} + 2C^{\mathbf R}_{\mathrm{cr}}S^{\mathbf R}(\bar g)\,e^{-\kappa d_\times/(24MR_{K'})}\Big] , \tag{8.38}$$

with $d_\times = d_{K'} \ge a_{K'}$ the least ℓ^∞-distance between two loops of $\mathcal C$ in units of the scale K', and $A^\times(\mathcal C)\coloneqq 0$ for a single loop. These numbers do not depend on K. The first two terms of (8.37) are sums over the loops, and $A^\times$ does not decrease when a loop is added to an ensemble.

Lemma 8.5 (The Total of the Cross Species)**.** *Let γ_q be sufficiently small, depending on $c_{\mathcal C}$. On every labelled term and at every step $k\le K'$, the groups of the cross species present at the step k satisfy $\sum_W e^{\theta(W)}\|Y_W\| \le A^\times(\mathcal C)$.*

Proof. By (5.15), summed over $j\le k$, the left side is at most $\sum_{j\le K'} 2^{K'-j+1}\mathsf A^\times_j$. In units of the scale j the lengths of the stopping step are multiplied by L^m, $m\coloneqq K'-j$, so that $P_j(\mathcal C)=L^mP_{K'}(\mathcal C)$ and $d_j=L^md_\times$. We use the following consequence of Lemma 2.11: for $c>0$ and $x>0$ with $cx\ge1$,

$$\sum_{m\ge0}(2L)^me^{-cxL^m} \le e^{-cx}\Big(1+\frac{4}{cx}\Big) \le 5e^{-cx} , \qquad (2L)^me^{-cxL^m} \le e^{-cx/2} . \tag{8.39}$$

Indeed $L^m-1\ge 2m$ and $cx\ge\log2$ give $2^m\le e^{cx(L^m-1)/2}$, hence $(2L)^me^{-cxL^m}\le e^{-cx/2}L^me^{-(c/2)xL^m}$. The first bound follows by the lemma with $c/2$ in place of c, after division by x, and the second from $te^{-(c/2)t}\le 2/(ec)$ at $t=xL^m$ and $2/(ecx)\le1$. With $x=d_\times$ and $c=\nu_\times/M$, where $cx\ge\nu_\times a_{K'}/M\ge1$ as in the proof of Corollary 7.11, the first terms of the $\mathsf A^\times_j$ contribute at most $2C_{\mathrm{cr}}\bar{\mathfrak a}P_{K'}(\mathcal C)\cdot 5e^{-\nu_\times d_\times/M}$. For the second terms put $y\coloneqq\kappa d_\times/(12MR_{K'})$, so that $y\ge\kappa L^3/12\ge 2\log(2L)$, because $d_\times\ge a_{K'}\ge L^3MR_{K'}$, $L\ge195$, and $\kappa\ge4^d$ as in Supplement, Proposition S.10. For $m\ge1$, (3.14) gives $R_j\le(L+1)m^{\beta_0}R_{K'}\le 2LmR_{K'}$, hence $\kappa d_j/(6MR_j)\ge yL^{m-1}/m\ge y(L/2)^{m-1}$, and for $m=0$ the exponent is $2y$. Since $(L/2)^{m-1}-\frac12\ge m$ for $m\ge2$ and $L\ge8$, and $y/2\ge\log(2L)$, we have $(2L)^me^{-y(L/2)^{m-1}}\le e^{-y/2}$ for every $m\ge1$. The second terms therefore contribute at most $2C^{\mathbf R}_{\mathrm{cr}}P_{K'}(\mathcal C)e^{-y/2}\sum_j\mathbf 1_{\mathbf R}(j)e^{-p_0(g_j)/4}$, and the last sum is at most $S^{\mathbf R}(\bar g)$ of Corollary 7.11, because $MR_j\ge1$. *q.e.d.*

For a component $X\in Z(\ell)$ we collect the costs of its history in

$$\omega_A(X) \coloneqq \exp\Big(\sum_{n\le K'} C_{A,n}\#_n(Z_n\cap X)\Big) , \qquad C_{A,n} \coloneqq 4C_A+8C_e\Lambda_*MR_n\hat\tau_n^2+\mathbf 1_{\mathbf R}(n)(4\nu N_n+C_x) , \tag{8.40}$$

with C_A the constant of (6.93), $\hat{\tau}_n$ as in (6.94), C_e the constant of the cost of a local term in Section 6.15, and N_n, C_x the numbers of Proposition 6.36(iv); $\mathbf{1}_{\mathbf{R}}(n)$ is 1 at the steps at which the **R**-operation acts and 0 at the others.

Lemma 8.6 (The Terms at the Stopping Step)**.** *Let $\bar{g}$ satisfy* (8.19)*. For every labelled term ℓ of $\varrho_{K'}$ and every real configuration of its variables in the support of its indicators,*

$$|\widehat{\mathcal{W}_\ell}| \le e^{B_E(\mathcal{C})+I_*(\mathcal{C})} \prod_{X\in Z(\ell)} \omega_A(X)\,, \tag{8.41}$$

with $B_E(\mathcal{C})$ of Corollary 7.11. On ℓ_0, moreover, $|\sum_{\mathfrak{B}\neq\emptyset} Q_{K'}(\mathfrak{B})w_{K'}(\mathfrak{B})| \le e^{I_*(\mathcal{C})}-1$. If, for every $n \le K'$,

$$C_{A,n} \le L^4(MR_n)^4\,, \tag{8.42}$$

which holds for γ_q small, depending on $c_{\mathcal{C}}$, then $1 \le \omega_A(X) \le \exp(C_P P(X))$.

Proof. We first bound the sum over the families. For a family $\mathfrak{B}$ the word $w_{K'}(\mathfrak{B})$ is the normalised trace of a product of unitary transports, which contain the background and, on large-field regions, outer variables of the term, and of the insertions of $\mathfrak{B}$, so that $|w_{K'}(\mathfrak{B})| \le \prod_{W\in\mathfrak{B}}\|Y_W\|$ by Lemma 5.1; with Definition 5.2(i), the sum over the finite families of groups of the term is bounded by a product over the groups. The groups of a term may be infinite in number, since their labels are not bounded; the product is then the limit of the products over finite sets of groups, which converge because $\sum_W e^{\theta(W)}\|Y_W\|$ is finite by the bounds below, and the sum over the families converges absolutely to a number bounded by it:

$$\Big|\prod_i f^{(i)}_{K'} + \sum_{\mathfrak{B}\neq\emptyset} Q_{K'}(\mathfrak{B})\, w_{K'}(\mathfrak{B})\Big| \le \prod_W (1+e^{\theta(W)}\|Y_W\|) \le \exp\Big(\sum_W e^{\theta(W)}\|Y_W\|\Big)\,, \tag{8.43}$$

where we used $|f^{(i)}_{K'}| \le 1$, and where the sum without the leading term is at most the right side minus one. Every group contains a traversal, so that the sum over the groups which are neither attached nor of the cross species is at most $\sum_{\mathfrak{n}} \nu^{(K')}_{\mathfrak{n}}$, the sum over the $P_{K'}(\mathcal{C})$ traversals of the step K', and the groups of the cross species contribute at most $A^\times(\mathcal{C})$ on every term, by Lemma 8.5.

The term ℓ_0. On ℓ_0 there are no large-field data at the step K', so that $d_{K'}(\mathfrak{n}) = \infty$ and $\mathfrak{m}_{K'}(\mathfrak{n}) = 0$ in (5.11), and there are no attached groups, since the attached parts of the renormalised components have become deviation insertions of the class (Section 6.15 and Supplement, Proposition S.10). By (6.1) at $k+1 = K'$, with $\varepsilon_{K'}$ enlarged by the factor $1+2\kappa_s$ of the soft test functions as in the proof of Corollary 8.3, $\delta^*_{\mathfrak{n}}(U_{K'}) \le 2c_\sigma(1+2\kappa_s)B_3\varepsilon_{K'}S_w$, and $\tilde{\rho}_{K'} \le \bar{\rho}/\min(\alpha_{0,K'},\alpha_{1,K'})$. The smallest period of the torus in units of the scale j is $N^{(j)}_{\min} = N_q L^{K'-j}$, so that with $i = K'-j$ the last term of (5.11) is at most $2C_{S'}\bar{\rho}\sum_{i\ge0} L^i e^{-c_{\mathrm{att}}N_qL^i/M} = 2C_{S'}\bar{\rho}\,S_{\mathrm{wr}}(c_{\mathrm{att}}N_q/M)$. Summing over the traversals and adding the cross species, $\sum_{\mathfrak{n}}\nu^{(K')}_{\mathfrak{n}} + A^\times(\mathcal{C}) \le I_*(\mathcal{C})$, and (8.43) gives both assertions on ℓ_0, since $|e^{\mathcal{E}_{K'}(\ell_0)}| = e^{\mathrm{Re}\,E_{\mathrm{const},K'}} \le e^{B_E(\mathcal{C})}$ by Definition 5.2(iv) and Corollary 7.11.

The other terms. On a term $\ell \neq \ell_0$ a family consists of a family of regular groups and of a family of attached ones (Section 6.14), and the sum over the families is at most the product of the two sums. The regular groups are groups of ℓ_0 with the same insertions, and their comparison regions lie in the small-field regions, where the background has the regularity used on ℓ_0 (Proposition 6.36(ii)); their sum is bounded as on ℓ_0. For the attached groups, $e^{\theta(W)}\|Y_W\|$ is at most the weighted mass (6.89) at the real points of the support, and by (6.92) at the step K' the sum over their families is at most $\exp(2\sum_{(n,Z)} A_n(Z))$, the sum over the

components Z of the regions Z_n, $n \le K'$, each of which lies in a component $X \in Z(\ell)$. Here $A_n(Z) \le C_A(d'_n(Z)+1) \le 2C_A \#_n(Z)$ by (6.93), where $d'_n(Z)+1 \le 2\#_n(Z)$ for a connected union of cubes, since a tree through the centres of adjacent cubes meets all of them and has edges of length at most twice the side. The further factor of (6.92) is a product of numbers b_n, one for each step $n \le K'$ at which the **R**-operation acts and each component of the second class of that step; such a component is a component of Z_n and contains a cube of side MR_n, so that its factor is at most $\exp((4\nu N_n + C_x)\#_n(Z))$. The exponent $\mathcal{E}_{K'}(\ell)$ contains in addition the local terms $e_X^{(n)}$ of the differences $E_n^{\partial}(\ell)$, $n \le K'$, which are attached to the regions Z_n (Supplement, Proposition S.9(iii)): the traversals within a cube of side MR_n of Z_n are at most $4\Lambda_* MR_n$ per cube by Lemma 5.5(i), since such a cube meets one loop by Lemma 3.14, and each carries local terms of total at most $2C_e\hat\tau_n^2$, which gives at most $8C_e\Lambda_* MR_n\hat\tau_n^2$ per cube. The constant part of $\mathcal{E}_{K'}(\ell)$ is bounded by $B_E(\mathcal{C})$ as on ℓ_0. These contributions add to the exponent in (8.43) and to $\operatorname{Re}\mathcal{E}_{K'}(\ell)$ at most $\sum_{X\in Z(\ell)}\sum_{n\le K'} C_{A,n}\#_n(Z_n \cap X)$, which gives (8.41).

Finally, $\omega_A(X) \ge 1$ trivially, and under (8.42) $\sum_n C_{A,n}\#_n(Z_n\cap X) \le \sum_n L^4 |Z_n \cap X|_n = P(X) \le C_P P(X)$. The condition (8.42) holds for γ_q small, depending on $c_{\mathcal{C}}$, because each of the three terms of $C_{A,n}$ is at most $\frac{1}{3}L^4(MR_n)^4$ for $n \le K'$. Indeed, $R_n \ge (\log \hat g_n^{-2})^r \ge (\log \gamma_q^{-2})^r$ by Definition 3.3 and (3.46), so that $12C_A \le L^4(MR_n)^4$ once $L^4M^4(\log\gamma_q^{-2})^{4r} \ge 12C_A$. Moreover $R_n \le 2L(\log g_n^{-2})^r$ by Lemma 3.4(i) and $\hat\tau_n^2 \le C g_n^2 (\log g_n^{-2})^{6p_0}$, so that $24C_e\Lambda_* MR_n\hat\tau_n^2 \le C'\gamma_q^2(\log\gamma_q^{-2})^{6p_0+r}$, since $x \mapsto x(\log x^{-1})^{6p_0+r}$ is increasing for small x, and the right side is at most $1 \le L^4(MR_n)^4$ for γ_q small. Finally $4\nu N_n = 4\cdot 107^4 \nu R_n^4 \le \frac{1}{6}L^4(MR_n)^4$, because $\nu \le 1$ and $L \ge 2^{52}$, and $C_x \le \frac{1}{6}L^4M^4(\log\gamma_q^{-2})^{4r}$ for γ_q small. *q.e.d.*

Theorem 8.7 (Uniform Bound)**.** *Under the hypotheses of Theorem 1.5, let $\bar g$ be so small that* (8.19) *and the hypothesis of Proposition 8.4 hold, and γ_q so small that* (8.42) *holds. For every $K \ge K_0(\mathcal{C})$,*

$$|W_K(\mathcal{C})| \le B(\mathcal{C}) := e^{B_E(\mathcal{C})+I_*(\mathcal{C})}(1+\bar\delta_{\mathrm{LF}})\,, \tag{8.44}$$

and the remainder (8.8) *satisfies*

$$|r_K(\mathcal{C})| \le e^{B_E(\mathcal{C})}\Big[\sum_{i=1}^{n}(e^{x(C_i)}-1) + e^{I_*(\mathcal{C})}-1\Big] + e^{B_E(\mathcal{C})+I_*(\mathcal{C})}\,\bar\delta_{\mathrm{LF}}\,. \tag{8.45}$$

The number $B(\mathcal{C})$ does not depend on K; it depends on $\mathcal{C}$ through $\sum_i (P(C_i)/r(C_i) + P_q(C_i))$, through $P_q(\mathcal{C})$, and through the characters, and on q, T/u, $\bar g$, and the constants of the construction.

Proof. By (8.5), the triangle inequality, and Lemma 8.6,

$$|W_K(\mathcal{C})| \le e^{B_E(\mathcal{C})+I_*(\mathcal{C})}\Big(w_{\ell_0} + \sum_{\ell\ne\ell_0} w_\ell \prod_{X\in Z(\ell)} \omega_A(X)\Big)\,, \tag{8.46}$$

where we used that $\omega_A(X)$ is a number of the label, so that $\langle \prod_X \omega_A(X)\rangle_\ell = \prod_X \omega_A(X)$. By Lemma 8.6 the numbers $\omega(X) := \omega_A(X)$ satisfy the hypothesis of Proposition 8.4, which gives $\sum_{\ell\ne\ell_0} w_\ell \prod_X \omega_A(X) \le \delta_{\mathrm{LF}} Z_{\ell_0}/\sum_\ell Z_\ell \le \bar\delta_{\mathrm{LF}}$; together with $w_{\ell_0} \le 1$ this is (8.44). For the remainder, the integrand of the first term of (8.8) is at most $\sum_i (e^{x(C_i)}-1) + e^{I_*(\mathcal{C})} - 1$ by Corollary 8.3 and Lemma 8.6, the measure μ_K has mass $w_{\ell_0} \le 1$, and $|e^{E_{\mathrm{const},K'}}| \le e^{B_E(\mathcal{C})}$; the second term is bounded as above. The dependence of $B(\mathcal{C})$ is read from Corollary 7.11, (8.37), and Proposition 8.4: $P_{K'}(\mathcal{C}) = P_q(\mathcal{C})L^3 MR_{K'-1}$, and N_q, $\varepsilon_{K'}$, $R_{K'}$, $\alpha_{i,K'}$ are functions of q, T/u, $\bar g$, and the constants of the construction. *q.e.d.*

The bound (8.44) is the first half of (1.10). Two features of it deserve a comment. First, no condition on the coupling in this section involves the size of the loops or their number: (8.19) and the conditions of Proposition 8.4 concern the construction alone, and (8.42) is a condition on γ_q which involves the shape constant Λ_* and the representations through $c_{\mathcal{C}}$. The size and the number of the loops enter only the value of $B(\mathcal{C})$, through $B_E(\mathcal{C})$ and $I_*(\mathcal{C})$, which are sums over the loops. Secondly, the bound does not say that the large-field terms are small: $\bar{\delta}_{\mathrm{LF}}$ is a fixed number at a fixed coupling. What makes the terms with large fields harmless is that their weights, multiplied by the costs of the dressed loop, are summable against the term without large fields on the fixed lattice $\mathbb{T}_{j_q}$.

9 The Difference of the Walk Factors

The limit $K \to \infty$ is obtained by comparing the construction for the target $(K, \bar{g})$ with the construction for $(K+1, \bar{g})$. The two share every lattice but the finest, and on a shared lattice they perform the same operations, with the same parameters, on data which differ by little; what differs is the fine lattice, on which every background field, every propagator, and every term of the effective action is computed. This section and the next prove the two estimates of the comparison which are not contained in Bałaban's analysis of a single run. The present one concerns the propagators. The effective actions are built from the inverses of the quadratic forms of the fluctuation integrals, the propagators of [5], through their expansions into generalised random walks, and the comparison needs a bound on the difference between the terms of one walk in the two constructions which is small in the ratio of the fine spacing to the spacing of the scale on which the walk lives. The difficulty is that the two terms do not act on the same space: one maps sources on the lattice of spacing ε to fields on that lattice, the other does the same on the lattice of spacing ε/L. We compare them through the one-step average and its right inverses. The comparison is then exact up to a bilinear defect, and this defect is small only between fields which are regular at the scale of their shell; it is the decay of the propagators across the shells which turns this into a bound of the order of a power of ε/a_j.

9.1 Two Runs on Common Lattices

Throughout this section and the next two, $\mathcal{C}$ is an ensemble as in Theorem 1.5, $K \geq K_0(\mathcal{C})$, and $K' = K - j_q$. *Run A* is the construction for the target $(K, \bar{g})$, started at the bare coupling $g_0(K, \bar{g})$ of Theorem 4.1 and stopped after K' steps; *run B* is the construction for $(K+1, \bar{g})$, started at $g_0(K+1, \bar{g})$ and stopped after $K'+1$ steps. Both are constructions for tubes with the same numbers b, b', and C_W, which Theorem 4.1 provides independently of K. We write $\varepsilon \coloneqq \varepsilon_K = uL^{-K}$ for the fine spacing of run A and

$$a_k \coloneqq \varepsilon L^k \,, \qquad \text{so that} \qquad \frac{\varepsilon}{a_k} = L^{-k} \,. \tag{9.1}$$

The step k of run A maps a density of the field on the lattice of spacing a_{k-1} to one on the lattice of spacing a_k, and so does the step $k+1$ of run B, because $(k+1)-(K+1) = k-K$. We call k the *shared scale* of these two steps, $0 \leq k \leq K'$; the step 1 of run B, which maps the lattice of spacing ε/L to that of spacing ε, is its *extra step*. The couplings at the shared scale k are g_k^A and g_{k+1}^B, and we put

$$\delta_k \coloneqq (g_{k+1}^B)^{-2} - (g_k^A)^{-2} \,. \tag{9.2}$$

Lemma 9.1 (Shared Parameters)**.** *For every integer $k \leq K$ the numbers u_k and ℓ_k of* (3.3) *of run A are those of run B at $k+1$. Consequently:*

(i) every parameter of Definition 3.3, the depth N_k of Definition 3.9, and every threshold, radius, cube size, and soft width of the construction at the shared scale k are the same in both runs;

(ii) the stopping index j_q, the stopping lattice $\mathbb{T}_{j_q}$, and the number N_q of (8.2) *are the same in both runs, and run B's stopping step is $K'+1$;*

(iii) for $0 \le k \le K'$, $|\delta_k| \le u_k - \ell_k \le \frac{1}{2}\min((g_k^A)^{-2}, (g_{k+1}^B)^{-2})$.

Proof. By (3.3), u_k for run A is $\bar{g}^{-2} + b'(K-k) + C_W$, and u_{k+1} for run B is $\bar{g}^{-2} + b'((K+1) - (k+1)) + C_W$, the same number; the same holds for ℓ. (i) The parameters of Definition 3.3 are functions of $\hat{g}_k = u_k^{-1/2}$, and $N_0(k)$ and N_k' in Definition 3.9 are defined through $\hat{g}_k$ and through the cube sizes $R_{k'}$ at the integers $k' \le k$, which correspond to each other in the same way; the remaining parameters are, by Definition 3.11(i), functions of $\hat{g}_k$ as well. (ii) By Definition 3.13, j_q is defined through the numbers $R^{[n]} = R_{K-n}$, which depend on n and not on K. (iii) The squared inverse couplings $(g_k^A)^{-2}$ and $(g_{k+1}^B)^{-2}$ are coordinates of histories of the two tubes (Theorem 4.1), so both lie in $[\ell_k, u_k]$, which gives the first inequality, and $u_k - \ell_k = (b'-b)(K-k) + 2C_W$. Since $b' \le \frac{3}{2}b$ and $C_W \le \beta_0 t/4 \le t/5$ by (3.2), this is at most $\frac{1}{2}(t + b(K-k) - C_W) = \frac{1}{2}\ell_k$, and both squared inverse couplings are at least ℓ_k. *q.e.d.*

The lemma is the reason for the tubes. With Bałaban's own parameters the cube size R_k of the two runs at a shared scale would differ wherever a power of L lies between $(\log(g_k^A)^{-2})^r$ and $(\log(g_{k+1}^B)^{-2})^r$, and then the two expansions would have different terms, which could not be compared one by one. With the parameters of the tubes the two runs perform the same decompositions on every shared lattice, and they differ only through the fine lattice and through the couplings, the latter by the numbers δ_k, whose size is determined in Section 11.

We need one more object, which carries a field of run A to the fine lattice of run B. For a field U on the lattice of spacing ε whose plaquettes are small, let $\iota(U)$ be the minimiser of the Wilson action on the lattice of spacing ε/L among the fields W with $M(W) = U$, the minimal configuration of [9, (2.11), (2.12), p. 256] for one averaging step. By [7, Theorem 1, p. 279] it exists, it is unique up to gauge transformations, it is regular, and it is an analytic and gauge covariant function of U; moreover $M\iota(U) = U$ and $\iota(1) = 1$. For a domain X we write ι_X for the minimiser localised to X in the sense of [9, (2.14)–(2.16), p. 257], which depends on U on a neighbourhood $X^\sim$ of X and satisfies $M\iota_X(U) = U$ there and $\iota_X(1) = 1$. Finally, for a field U^B on the fine lattice of run B we call

$$\tilde{U} \coloneqq MU^B \tag{9.3}$$

the *aligned background*: a field on the fine lattice of run A, which satisfies run A's constraints whenever U^B satisfies run B's, since the iterated averages of run B are those of run A composed with M.

9.2 Propagators and Their Walk Expansions

We recall the objects to be compared. The fluctuation integral of a step is Gaussian to leading order, with a quadratic form built from the covariant Laplacian at the background, the block averages, and the gauge fixing, and its inverse is expressed through three operators of [5]: G', the inverse of the scalar operator Δ_a'; $(Q'G'^2Q'^*)^{-1}$, an operator on the block data; and G, the inverse of the one-form operator Δ_a. Each of them is approximated by local inverses:

for a cube $\square$ of the scale j, Bałaban takes a sequence of cubes $\Omega_{j+1}(\square) \subset \Omega_j(\square) = \tilde{\square}^4 \subset \Omega_{j-1}(\square) \subset \cdots \subset \Omega_0(\square) \subset \tilde{\square}^5$ with the same centre and

$$\operatorname{dist}(\Omega_n(\square)^c, \Omega_{n+1}(\square)) = 2R_0M_0L^n\eta \,, \tag{9.4}$$

on which the operators are taken with Dirichlet conditions outside $\Omega_0(\square)$ and with the averaging of the scale n on the shell $\Omega_n(\square) \setminus \Omega_{n+1}(\square)$, and he calls $G'_\square$, $C_\square = (Q'G'^2_\square Q'^*)^{-1}$, and $G_\square$ the resulting operators [5, pp. 408–409]. Here η is the fine spacing and R_0M_0 the smallest size for which the results of [3] hold, with $M = KR_0M_0$ [5, p. 408]. With a partition of unity $\sum_\square h_\square^2 = 1$ he puts $G'_0 = \sum_\square h_\square G'_\square h_\square$, and similarly C_0 and G_0 [5, (3.87), p. 409], writes $\Delta'_a G'_0 = I - R'$ with the small remainder $R' = \sum_\square K(h_\square) G'_\square h_\square$ [5, (3.88)–(3.89), p. 409], and expands $G' = G'_0 \sum_n R'^n$ into a sum over walks of cubes [5, Theorem 3.7, (3.90), p. 409]. The same construction, with more kinds of factors, gives the expansion of G,

$$G = \sum_\omega R_0(X_0)R_{\alpha_1}(X_1)\cdots R_{\alpha_n}(X_n) \,, \tag{9.5}$$

over walks $\omega = ((0, X_0), (\alpha_1, X_1), \ldots, (\alpha_n, X_n))$ with $X_{i-1} \cap X_i \neq \emptyset$, whose term depends on the configuration on $\tilde{X}_0^5 \cup \cdots \cup \tilde{X}_n^5$ and satisfies

$$|(R_0(X_0)\cdots R_{\alpha_n}(X_n)J)(x)| \leq O(1)(L^j\eta)^2 O(M^{-1/2})^{|\omega|} M^{-\frac{1}{2}|\omega|} e^{-\frac{1}{2}\delta_0 d(\omega,y,y')} |J| \tag{9.6}$$

for $x \in \Delta(y)$, $y \in \Lambda_j$, and $\operatorname{supp} J \subset \Delta(y')$, together with the corresponding bounds in the norms of [5, (3.42)–(3.47)] [5, Theorem 3.10, (3.107), (3.108), p. 416]. Here d is the weighted distance on the set $\mathfrak{B}$ of the blocks of all scales [3, (2.46), p. 231], which [5, p. 397] cites as (2.36) (Section 14), and $d(\omega, y, y')$ is the infimum of $d(y, y_0) + d(y_0, y_1) + \cdots + d(y_{n-1}, y_n) + d(y_n, y')$ over the points $y_i \in X_i \cap \mathfrak{B}$ [5, (3.93), p. 410, and p. 413]. The operators H, H_0, and $\tilde{G}$ through which the fixed point of the fluctuation integral is written [10, (1.2)–(1.4), p. 2] have expansions of the same form, "a propagator is represented by the sum (3.107) [13]", and the term of a walk is bounded "as (3.108) [13]" in "all operator norms in formulations of theorems in [13]" [10, (1.6), (1.7), p. 3]. Bałaban states these bounds for whole terms of walks and not for their factors, of which he says only that each "satisfies a bound of the type (3.89), possibly with an additional power of $L^j\eta$" [5, p. 413]; the comparison below is accordingly a comparison of whole terms.

What the two runs share is read off from these definitions. The cubes of the scale j are cells of the lattice $ML^j\eta\mathbb{Z}^4$, and $ML^j\varepsilon = ML^{j+1}(\varepsilon/L)$, so that for $\square$ at the shared scale j the cubes $\square$, $\tilde{\square}^n$, the partitions of unity as functions on the torus, the localisation domains, and the walks are the same in both runs. The two local sequences are related by

$$\Omega^B_{n+1}(\square) = \Omega^A_n(\square) \,, \qquad 0 \leq n \leq j+1 \,, \tag{9.7}$$

because, by (9.4), $\operatorname{dist}(\Omega_n(\square)^c, \tilde{\square}^4) = 2R_0M_0L^n\eta(L^{j-n}-1)/(L-1)$ [5, p. 408], and the value for run B at $n+1$, with $\eta = \varepsilon/L$ and $j+1$ in place of j, is the value for run A at n. So the shells of the two runs coincide at every common spacing $L^n\varepsilon$, and run B has one more shell $\Omega^B_0 \setminus \Omega^B_1$, of width $2R_0M_0\varepsilon/L$ at the spacing ε/L, where run A's operators have their Dirichlet exterior. What the runs do not share is the space on which the operators act: on $\Omega_0(\square)$ every factor of run A acts on fields and sources on the lattice of spacing ε, every factor of run B on the lattice of spacing ε/L.

9.3 The Galerkin Identities

Fix $\square$ at the shared scale j and one of the local operators: $\mathcal{O}^X$ is $\Omega_0\Delta_a\Omega_0$ or $\Omega_0\Delta'_a\Omega_0$ of run $X \in \{A, B\}$, acting on the space $\mathfrak{F}^X$ of fields (or functions) on the bonds (or sites) of

the fine lattice of run X in $\Omega_0^X(\square)$, at run B's background U^B for $X = B$ and at the aligned background $\tilde{U}$ for $X = A$, and $G^X := (\mathcal{O}^X)^{-1}$ on $\mathfrak{F}^X$. The pairings $\langle\cdot,\cdot\rangle_X$ carry the weights η_X^4 of [5, p. 393] and are bilinear, so that $\mathcal{O}^X$ is symmetric on the complex domain of [5, (3.37)], as the analytic extension of a symmetric form. Let Q_1 be the linearised one-step average at U^B, which maps fields of run B to fields of run A. Two averages have to be kept apart here. The operators of [5] are built with the average of the fluctuation relative to the background [5, (3.13), p. 392; (3.113), p. 418], and Q_1 is always its linear part; the average M of the construction has the linear part Q_1^{pl}, the plain one of (11.2). The two differ by a gauge transformation of the sites of run A, $Q_1 = Q_1^{\mathrm{pl}} + \bar{D}\Lambda_1$ with $\Lambda_1 A$ the block means of the contour variables of A, and this term does not vanish at a flat background; but every gauge-invariant function of the averaged configuration is the same for the two averages, and the minimisers of the two constraints are gauge transforms of each other (Supplement, Proposition S.12(j)). Let χ_A be the characteristic function of $\Omega_0^A(\square)$, and let $\mathcal{L}$ be a right inverse of Q_1 with

$$\mathcal{L}\mathfrak{F}^A \subset \mathfrak{F}^B \ . \tag{9.8}$$

The *bilinear defect* of $\square$ is

$$\beta_\square(\mathcal{F},\mathcal{A}) := \langle \chi_A Q_1 \mathcal{F}, \mathcal{O}^A \chi_A Q_1 \mathcal{A}\rangle_A - \langle \mathcal{F}, \mathcal{O}^B \mathcal{A}\rangle_B \ , \qquad \mathcal{F},\mathcal{A} \in \mathfrak{F}^B \ , \tag{9.9}$$

and $\beta_\square(\cdot,\mathcal{A})^\sharp$ denotes the source of run B which represents the linear form $\mathcal{F} \mapsto \beta_\square(\mathcal{F},\mathcal{A})$ on $\mathfrak{F}^B$. The defect measures how far the quadratic form of run A, pulled back by the average, is from that of run B.

Lemma 9.2 (Galerkin Identities)**.** *Under* (9.8)*, for every source g of run B and every source f of run A,*

$$\chi_A Q_1 G^B g - G^A \chi_A \mathcal{L}^* g = G^A \chi_A \mathcal{L}^* \beta_\square(\cdot, G^B g)^\sharp \ , \tag{9.10}$$

$$\mathcal{L} G^A f - G^B Q_1^* \chi_A f = -G^B \beta_\square(\cdot, \mathcal{L} G^A f)^\sharp \ . \tag{9.11}$$

Proof. For (9.10) put $u_B := G^B g$ and $u_A := G^A \chi_A \mathcal{L}^* g$. For $X \in \mathfrak{F}^A$ we have $\mathcal{L}X \in \mathfrak{F}^B$ by (9.8) and $\chi_A Q_1 \mathcal{L} X = X$, so that

$$\begin{aligned} \langle X, \mathcal{O}^A u_A\rangle_A &= \langle \mathcal{L}X, g\rangle_B = \langle \mathcal{L}X, \mathcal{O}^B u_B\rangle_B \ , \\ \langle X, \mathcal{O}^A \chi_A Q_1 u_B\rangle_A &= \langle \chi_A Q_1 \mathcal{L}X, \mathcal{O}^A \chi_A Q_1 u_B\rangle_A \ . \end{aligned} \tag{9.12}$$

The difference of the two right sides is $\beta_\square(\mathcal{L}X, u_B) = \langle X, \chi_A \mathcal{L}^* \beta_\square(\cdot, u_B)^\sharp\rangle_A$. Both $\chi_A Q_1 u_B - u_A$ and $G^A \chi_A \mathcal{L}^* \beta_\square(\cdot, u_B)^\sharp$ lie in $\mathfrak{F}^A$, on which X is arbitrary and $\mathcal{O}^A$ is invertible, and (9.10) follows. For (9.11) put $u_A := G^A f$ and $u_B := G^B Q_1^* \chi_A f$. For $Y \in \mathfrak{F}^B$, since $\chi_A Q_1 \mathcal{L} u_A = u_A$,

$$\langle Y, \mathcal{O}^B u_B\rangle_B = \langle \chi_A Q_1 Y, f\rangle_A = \langle \chi_A Q_1 Y, \mathcal{O}^A \chi_A Q_1 \mathcal{L} u_A\rangle_A = \langle Y, \mathcal{O}^B \mathcal{L} u_A\rangle_B + \beta_\square(Y, \mathcal{L}u_A) \ , \tag{9.13}$$

so that $\langle Y, \mathcal{O}^B(\mathcal{L}u_A - u_B)\rangle_B = -\beta_\square(Y, \mathcal{L}u_A)$ for every $Y \in \mathfrak{F}^B$, and $\mathcal{L}u_A - u_B \in \mathfrak{F}^B$ by (9.8). *q.e.d.*

The identities are exact, and the whole difficulty is in the size of the defect. This is not small as a bilinear form: for a field $\mathcal{A}$ which alternates at the spacing ε/L inside one block, $Q_1\mathcal{A} = 0$ and $\beta_\square(\mathcal{A},\mathcal{A}) = -\langle \mathcal{A}, \mathcal{O}^B \mathcal{A}\rangle_B$, which is of the order $(\varepsilon/L)^{-2}$ times the square of the norm of $\mathcal{A}$. In the identities, however, the defect is paired with $G^B g$ or $\mathcal{L}G^A f$, which are regular at the scale of their shell by [5, (3.43)], and between regular fields it is small: on a shell of spacing $L^n\eta$ the lattices of the two runs are averages of each other, and the defect of

two regular fields is an oscillation of relative order L^{-n}. To make this precise we use, for a block y of the shell n, the norm

$$\|\mathcal{A}\|_{1,\beta;y} \coloneqq \sup((L^n\eta)^{-1}|\mathcal{A}| + |\nabla\mathcal{A}|) + (L^n\eta)^{\beta}\|\nabla\mathcal{A}\|_{\beta} \,, \tag{9.14}$$

the supremum and the Hölder quotient of [5, (3.40)] taken over the blocks within weighted distance 2 of y. We index the shells by run A's index n, so that run A's shell n is run B's shell $n+1$ by (9.7), and run B's extra shell has the index -1.

Proposition 9.3 (The Defect of the Local Operators)**.** *Let $\beta \in [\frac{3}{4}, 1)$, and let R_0M_0 satisfy the conditions of Appendix C. There is a constant $C_{\mathrm{df}}(\beta)$, independent of the shell, of j, k, K, the torus, the block, and the field, such that for each of the operators of $G'_{\square}$, $C_{\square}$, and $G_{\square}$, at aligned backgrounds on the complex domain of [5, (3.37)], and for each $\mathcal{A} \in \mathfrak{F}^B$, there are fields s_0, s_1, s_2 of run B with*

$$\beta_{\square}(\mathcal{F}, \mathcal{A}) = \langle D^*\mathcal{F}, s_2\rangle_B + \langle \nabla\mathcal{F}, s_1\rangle_B + \langle \mathcal{F}, s_0\rangle_B \qquad \text{for every } \mathcal{F} \in \mathfrak{F}^B \,, \tag{9.15}$$

and, on every block y of the shell n,

$$\sup_{\Delta(y)}(|s_1| + |s_2| + L^n\eta|s_0|) \le C_{\mathrm{df}}(\beta)\varpi_n\|\mathcal{A}\|_{1,\beta;y} \,, \qquad \varpi_n \coloneqq \begin{cases} L^{-\beta n}, & n \ge 1, \\ 1, & n \in \{0, -1\}, \end{cases} \tag{9.16}$$

together with a Hölder bound of s_1 and s_2 at the scale of their shell; the same holds with the two arguments of $\beta_{\square}$ exchanged.

Proof. We show the mechanism on the averaging terms, which can be computed exactly, and refer to Supplement, Proposition S.12, for the other terms. The quadratic form of [5, (3.16)] contains Q^*aQ with the weight $(L^n\eta)^{d-2}$ on the bonds of the shell n. Run B's shell $n+1$ is run A's shell n, with the weight $(L^{n+1}\varepsilon/L)^{d-2} = (L^n\varepsilon)^{d-2}$, and run B's averaging of the scale $n+1$ is run A's averaging of the scale n composed with the one-step average, $Q^B_{n+1}(U^B) = Q^A_n(\tilde{U})Q_1$ (Supplement, Proposition S.12(a)), while $\chi_A = 1$ on run A's shells. Hence the averaging part of $\beta_{\square}$ is exactly $-a\sum_b(\varepsilon/L)^{d-2}\langle\mathcal{F}(b), \mathcal{A}(b)\rangle$, the sum over the bonds of run B's extra shell, which has the form (9.15) with s_0 supported there and $\varpi_{-1} = 1$. For the scalar operator the coefficients of the averaging on the shells are a_n, with $1/a_n = a^{-1}\sum_{i=0}^{n-1} L^{-2i}$ for $n \ge 1$ [5, p. 394]; run A's shell n carries a_n and run B's the same shell a_{n+1}, and $|a_{n+1} - a_n|/a_n \le L^{-2n}/(1 - L^{-2}) \le 2L^{-\beta n}$, which is again of the form (9.16). The covariant Laplacian, the curvature term, and the gauge-fixing term are not compositions of averages; on run A's shells $n \ge 1$ their defect between regular fields is an oscillation of relative order L^{-n}, of which the Hölder exponent retains $L^{-\beta n}$, and on the two shells of the finest spacing nothing relates the two runs and $\varpi = 1$. The derivation of these bounds, for $G'_{\square}$, for $C_{\square}$, which needs an identity of its own because the data spaces of the two runs differ by run B's extra layer of data, and for $G_{\square}$, whose gauge-fixing part is nonlocal through $P = G'Q'^*(Q'G'^2Q'^*)^{-1}Q'G'$, is Supplement, Proposition S.12(b)–(d). *q.e.d.*

The following lemma shows why the shells of the finest spacing, on which the defect is of order one, cost only a factor $(\varepsilon/a_j)^{\beta}$: a propagator of the scale j reaches them only after crossing j shells, each of width $2R_0M_0$ in its own units.

Lemma 9.4 (Reduction over the Shells)**.** *Let $\mathcal{L}$ satisfy* (9.8)*, have the range of one block of run A, and satisfy $\sum_{z\in\Delta(y)}\eta_B^4(|\nabla\mathcal{L}v(z)| + (L^n\eta)^{-1}|\mathcal{L}v(z)|) \le C_{\mathcal{L}}\sum_{y'''}\sum_{w\in\Delta(y''')}\eta_A^4(|\nabla v(w)| + (L^n\eta)^{-1}|v(w)|)$, the sum over the 3^4 blocks y''' next to y. Assume $\frac{1}{6}\delta_0R_0M_0 > 8\log c_0(\frac{1}{6}) + 1$ and $4\delta_0R_0M_0 \ge 3(1+\beta)\log L$, with c_0 the constant of [3, (2.58)]. Then for blocks y, y' of*

$\square$ in the layers j and $j+1$ of the local sequence, $x \in \Delta(y)$, and a source g of run B with $\operatorname{supp} g \subset \Delta(y')$,

$$|(\chi_A Q_1 G^B_\square - G^A_\square \chi_A \mathcal{L}^*) g(x)| \le C_\square(\beta) \Big(\frac{\varepsilon}{a_j}\Big)^\beta (L^j\eta)^2 e^{-\frac{1}{3}\delta_0 d(y,y')} |g| \,, \tag{9.17}$$

with $C_\square(\beta) \coloneqq 6c_0^4(\frac{1}{6}) c_\beta C_3$, $C_3 \coloneqq 9 \cdot 3^4 e^{4\delta_0}(1+L) C_{\mathcal{L}} C_{\mathrm{df}}(\beta) B_0 B_0(\beta)$, and $c_\beta \coloneqq L^{1+\beta}(3 + L^\beta + L^{1+\beta}/(L-1))$, where B_0 and $B_0(\beta)$ are the constants of [5, (3.42), (3.43)].

Proof. By (9.10) and Proposition 9.3 with $\mathcal{A} = u_B \coloneqq G^B_\square g$, the left side is $|(G^A \chi_A \mathcal{L}^*(Ds_2 + \nabla^* s_1 + s_0))(x)|$, and we split the sources over the blocks y'' of the shells. We bound the contribution of a block y'' of the shell n in two legs.

The leg from y' to y''. The bounds [5, (3.42), (3.43)] for $G^B_\square$, which hold for the local sequences by [5, Corollary 3.6, p. 408], give $\|u_B\|_{1,\beta;y''} \le 3e^{2\delta_0} B_0(\beta)(L^n\eta) e^{-\delta_0 d(y'',y')} |g|$, where $e^{2\delta_0}$ accounts for the blocks within distance 2 and $B_0 \le B_0(\beta)$. By (9.16),

$$\sup_{\Delta(y'')} (|s_1| + |s_2| + L^n\eta |s_0|) \le 3e^{2\delta_0} C_{\mathrm{df}}(\beta) \varpi_n B_0(\beta)(L^n\eta) e^{-\delta_0 d(y'',y')} |g| \,. \tag{9.18}$$

The leg from y'' to y. For a source σ on $\Delta(y'')$, the symmetry of G^A gives $(G^A \chi_A \mathcal{L}^* \nabla^* \sigma)(x) = \langle \nabla \mathcal{L}[G^A(\cdot, x)], \sigma \rangle_B$, which is at most $\sup|\sigma|$ times the block sum of $\nabla \mathcal{L} G^A(\cdot, x)$. By the hypothesis on $\mathcal{L}$ and [5, (3.42)] for G^A, this is at most

$$\sup|\sigma| \cdot 3^4 e^{2\delta_0} C_{\mathcal{L}} B_0 [(L^j\eta) + (L^j\eta)^2 (L^n\eta)^{-1}] e^{-\delta_0 d(y,y'')} \,, \tag{9.19}$$

and the same holds with D in place of ∇^*, while for s_0 the factor is $3^4 e^{2\delta_0} C_{\mathcal{L}} B_0 (L^j\eta)^2 e^{-\delta_0 d(y,y'')}$, that is $(L^j\eta)^2 (L^n\eta)^{-1}$ times $L^n\eta|s_0|$. Since $n \le j+1$, $L^j\eta \le L(L^j\eta)^2(L^n\eta)^{-1}$.

The product and the sum. The factor $(L^n\eta)^{-1}$ of the second leg cancels the factor $L^n\eta$ of the first, so that the shell n contributes at most $C_3 \varpi_n (L^j\eta)^2 \sum_{y''} e^{-\delta_0 [d(y,y'') + d(y'',y')]} |g|$. One third of the decay gives $e^{-\frac{1}{3}\delta_0 d(y,y')}$ by the triangle inequality [3, (2.54)]; one third gives the count $\sum_{y''} e^{-\frac{1}{3}\delta_0 d(y,y'')} \le 6c_0^4(\frac{1}{6})$ over one shell, by [3, (2.58)] with $\alpha = \frac{1}{3}$ under the first condition; and the last third is at most $e^{-\frac{4}{3}\delta_0 R_0 M_0 (j-n-1)_+} \le L^{-(1+\beta)(j-n-1)_+}$, because the weighted distance from a block of $\square$ to the shell n is at least $2R_0M_0(j-n-1)$ [3, (2.60)], both legs cross the shells in between, and the second condition holds. Summing over the shells with (9.16): the shells j and $j+1$ give at most $2L^{-\beta j}$; the shells $1 \le n \le j-1$ give $\sum_n L^{-\beta n} L^{-(1+\beta)(j-n-1)} \le L^{1+\beta} L^{-\beta j}/(L-1)$; the shell 0 gives $L^{-(1+\beta)(j-1)} \le L^\beta L^{-\beta j}$; and run B's extra shell, at distance at least $2R_0M_0 j$, gives at most $L^{-\beta j}$. Hence $\sum_n \varpi_n L^{-(1+\beta)(j-n-1)_+} \le c_\beta L^{-\beta j}$, with room for the intersected sequences of the cubes next to a coarser layer [5, p. 408], and $L^{-j} = \varepsilon/a_j$ by (9.1). *q.e.d.*

9.4 The Walk Terms

We can now state the comparison of the walk terms. Let W^X_ω be the term of the walk ω in run X's expansion of one of the operators H, H_0, or $\tilde{G}$ of [10, (1.2)–(1.4), p. 2], at run X's background, on a set of the **R**-operation whose layers lie at shared scales; for these operators the walks of run A and of run B are the same, because the cubes are. Let Y be a localisation domain of the walk, and let $\mathcal{L}_\iota$ be the operator H_1 of [5, (3.129), p. 421] of the one-step problem on the neighbourhood $Y^\sim$ at the background U^B, which assigns to a field of run A the minimiser of the linearised action among the fields of run B with this average. It is a right inverse of Q_1, and when U^B is the localised lift $\iota_Y(\tilde{U})$ of its own average it is the derivative of

ι_Y at $\tilde{U}$ up to a linear gauge transformation, the part of this derivative in the Landau gauge (Supplement, Proposition S.12(e), (j)). The operator $\mathcal{J}'$ is the identity on data, for H and H_0, and $\mathcal{L}_\iota^*$ on sources, for $\tilde{G}$; $m(\omega)$ is the finest scale of a layer of the set which meets $\bigcup_i \tilde{X}_i^5$.

Theorem 9.5 (Walk-Factor Difference)**.** *Let the constants satisfy the conditions of Appendix C, let $\bar{g}$ be sufficiently small, and let $K' \geq 2m_{\mathrm{rc}}+2$, where m_{rc} is the least integer with $L^{m_{\mathrm{rc}}} \geq (8\log L+\delta_0)/\delta_0'$ and δ_0' is the rate of Supplement, Proposition S.12. For every walk ω of the expansions of H, H_0, and $\tilde{G}$ at the backgrounds of the two runs on a set whose layers lie at the shared scales $j \geq m_{\mathrm{rc}}$, among them all the scales $j \geq K'/2-1$,*

$$|(W_\omega^B - \mathcal{L}_\iota W_\omega^A \mathcal{J}')\lambda(x)| \leq \delta_w(\omega)\cdot O(1)(L^j\eta)^p O(M^{-1/2})^{|\omega|} M^{-\frac{1}{2}|\omega|} e^{-\frac{1}{6}\delta_0 d(\omega,y,y')}|\lambda|\,, \tag{9.20}$$

in the supremum norm, where $(L^j\eta)^p$ is the power of (9.6) *for the operator in question, and*

$$\delta_w(\omega) \leq C_w\Big(\frac{\varepsilon}{a_{m(\omega)}}\Big)^{\gamma_w}\,, \qquad \gamma_w := \min\Big(\beta,\, \frac{1-\varepsilon_0}{3}\Big)\,, \tag{9.21}$$

with a constant C_w of the constants of the construction and of ε_0, ε_1, and β, which does not depend on K, k, ε, or the torus. If run A's term is evaluated at the aligned background $\tilde{U}$ in place of its own, the same holds with the exponent β in place of γ_w. The averaged difference $Q_1 W_\omega^B - W_\omega^A \mathcal{J}'$ satisfies the same bound, with the same map $\mathcal{J}'$, the identity on the data for H and H_0 and $\mathcal{L}_\iota^*$ on the sources for $\tilde{G}$. The first differences of the two differences satisfy these bounds with the power of the scale lowered by one and with $\delta_w(\omega)^{\beta/(1+\beta)}$ in place of $\delta_w(\omega)$.

Proof. The proof has three parts: the factors at the aligned background, the change from the aligned background to run A's own, and the nonlinear functions through which H, H_0, and $\tilde{G}$ are built from the propagators.

The factors. A term of a walk is a product of factors, each of which is a local operator $G'_\square$, $C_\square$, $G_\square$, multiplied by partition functions, commutators with them, and averaging operators [5, pp. 409–416]. For two products $a_1\cdots a_n$ and $b_1\cdots b_n$,

$$a_1\cdots a_n - b_1\cdots b_n = \sum_{i=1}^n a_1\cdots a_{i-1}(a_i-b_i)b_{i+1}\cdots b_n\,, \tag{9.22}$$

and we apply this first to the averaged difference $Q_1 W_\omega^B - W_\omega^A\mathcal{J}'$. Between two consecutive factors of a walk term stands a fine field, a fine source, or data, and the two runs are compared with Q_1 on the fine fields, with the adjoint $\mathcal{L}^*$ of the cut-off lift $\mathcal{L} := \chi_B\mathcal{L}_\iota$ on the fine sources, and with the identity on the data. The difference is then the sum, over the factors c of the term, of the product of run A's factors to the left of c, the *intertwining defect* of c, and run B's factors to the right. For a local propagator the intertwining defect is $Q_1G^B - G^A\mathcal{L}^*$, the left side of (9.10). For a factor $\mathcal{K}$ which maps fine fields to fine sources, a commutator with a partition function, it is $\mathcal{L}^*\mathcal{K}^B - \mathcal{K}^A Q_1$, and $\langle X, (\mathcal{K}^A Q_1 - \mathcal{L}^*\mathcal{K}^B)\mathcal{A}\rangle_A$ is the bilinear defect of $\mathcal{K}$ at the pair $(\mathcal{L}X, \mathcal{A})$, because $\chi_A Q_1 \mathcal{L}X = X$. For the averages it vanishes, by Supplement, Proposition S.12(a). The only property of the lift which this uses is that it is an exact right inverse on $\mathfrak{F}^A$ (Supplement, Proposition S.12(e)); the product $\mathcal{L}Q_1$ is not the identity on the fields of run B and does not occur. Each defect of a local propagator is bounded by Lemma 9.4, which costs the factor $C_\square(\beta)(\varepsilon/a_j)^\beta$ and lowers the rate $\frac{1}{2}\delta_0$ to $\frac{1}{6}\delta_0$; the explicit factors, multiplications by the partition functions and commutators with them, are functions on the torus which both runs take to be the same, and their defect between the two lattices is of relative order ε/a_j or $(\varepsilon/a_j)^\beta$, by the oscillation of these functions over

the reach of the kernel of Q_1 (Supplement, Proposition S.12(f)). The sum (9.22) has $|\omega|+1$ terms, which the factor $O(M^{-1/2})^{|\omega|}$ absorbs after a change of the constant. Since a cube of the walk lies at a scale $j \geq m(\omega)$, $(\varepsilon/a_j)^\beta \leq (\varepsilon/a_{m(\omega)})^\beta$, and the shells at the scales $n < j$ inside $\tilde{\square}^5$, which are not layers of the set, enter only through the constant c_β of Lemma 9.4. This gives the bound of the averaged difference at the aligned background, with the exponent β.

The fine form (9.20) follows from the averaged one, and it is here that the product $\mathcal{L}_\iota Q_1$ enters, applied to a field of run B which is regular. We write

$$(W_\omega^B - \mathcal{L}_\iota W_\omega^A \mathcal{J}')\lambda = (1 - \mathcal{L}_\iota Q_1) W_\omega^B \lambda + \mathcal{L}_\iota (Q_1 W_\omega^B - W_\omega^A \mathcal{J}')\lambda \,. \tag{9.23}$$

The second term is the lift of the averaged difference, and the kernel of $\mathcal{L}_\iota$ is bounded and decays exponentially at the scale ε [5, (3.133), p. 422]. The first term is the difference between a field of run B and the lift of its average. For an arbitrary field this difference is as large as the field, but for the lift $\mathcal{L}_\iota = G_1 Q^* (Q G_1 Q^*)^{-1}$ it has an exact expression,

$$(1 - \mathcal{L}_\iota Q_1) u = \tilde{C}_1 (D^* D + DRD^* + \Delta' - \Delta'_\pi - \Delta^{(2)}_\pi) u \,, \qquad \tilde{C}_1 \coloneqq G_1 - \mathcal{L}_\iota Q G_1 \,, \tag{9.24}$$

in which $\tilde{C}_1$ is the propagator of the fluctuations of one step, of the size ε^2 and of the range ε, and the operator in the bracket is of the second order with small terms of lower order. Since $u = W_\omega^B \lambda$ satisfies the bound of its walk also in the norm of the first differences, with one power of a_j less [5, (3.108)], the first term of (9.23) is of the relative order ε/a_j (Supplement, Proposition S.12(i)). The weights of both estimates decay at the scale ε, while the bounds of the walk vary at the scales $a_j \geq \varepsilon L^{m_{\mathrm{rc}}}$ of the layers, which ensures that the former are summed against the latter without a loss in the rate (Supplement, Proposition S.12(h)); the condition $K' \geq 2m_{\mathrm{rc}} + 2$ places the scales $j \geq K'/2 - 1$ of the large-field terms in this range. This gives (9.20) at the aligned background, with the exponent β.

The first differences are bounded by interpolation. The averaged difference is the difference of two fields of run A, each of which satisfies the bound of its walk in the Hölder norm of the first differences, and a field f at the scale a satisfies $\sup|\nabla f| \leq 4(\sup|f|)^{\beta/(1+\beta)}([\nabla f]_\beta)^{1/(1+\beta)} + 4a^{-1}\sup|f|$; this turns the factor $\delta_w(\omega)$ of the supremum into $\delta_w(\omega)^{\beta/(1+\beta)}$ for the first differences. For the fine difference the first differences of the two terms of (9.23) are bounded separately: those of the first carry the factor $(\varepsilon/a_j)^\beta$, because the two leading terms of (9.24) vanish on covariantly constant fields up to the curvature, and those of the second are bounded by the first differences of the averaged difference, because $\mathcal{L}_\iota$ maps regular fields of run A to regular fields of run B (Supplement, Proposition S.12(e), (h), (i)).

The background. Run A's own background is $U^A = e^{i\varepsilon A}\tilde{U}$ with $a_{j(x)}|A(x)| + a_{j(x)}^2|\nabla A(x)| \leq \bar{d}(x) \coloneqq C_U(\varepsilon/a_{j(x)})^{(1-\varepsilon_0)/3}\varepsilon_1$, where $j(x)$ is the layer of x and C_U a constant of M (Proposition 10.6). Put $\bar{d}(\omega) \coloneqq \sup \bar{d}(x)$ over $\bigcup_i \tilde{X}_i^5$ and $U(t) \coloneqq e^{it\varepsilon A}\tilde{U}$. For $|t| \leq r \coloneqq \alpha'_*/(2\bar{d}(\omega))$, with $\alpha'_* \coloneqq \min(\alpha_1, \alpha_J/\hat{x}_J)$ the radius of the class of pairs of a configuration and a current of Supplement, Proposition S.12(g), the configuration $U(t)$ lies in the domain of [5, (3.37)] layer by layer, where the local propagators are analytic with the bounds of [5, Theorems 3.1–3.3] [5, Theorem 3.4, p. 400], and so is the term of a walk, since the proof of (9.6) uses only these bounds [5, p. 410]. Under $4\bar{d}(\omega) \leq \alpha'_*$ we have $r \geq 2$, and Cauchy's estimate gives

$$|W_\omega^A(U^A) - W_\omega^A(\tilde{U})| \leq \sup_{|t|\leq 1} |\partial_t W_\omega^A(U(t))| \leq \frac{\sup_{|t|=r}|W_\omega^A(U(t))|}{r-1} \leq \frac{4\bar{d}(\omega)}{\alpha'_*} \tag{9.25}$$

times the bound (9.6), in each of its norms. With $\bar{d}(\omega) \leq C_U\varepsilon_1(\varepsilon/a_{m(\omega)})^{(1-\varepsilon_0)/3}$ this is of the form (9.21).

The nonlinear functions. The operators H, H_0, and $\tilde{G}$ are built from the propagators through the fixed point $\mathbf{A}_0$ of [10, (1.4), p. 2], the operator D of [10, (1.3)], and the solution of a Neumann series; their walk expansions are composite walks with vertices at which three chains meet [5, p. 427]. The comparison of these functions between the two runs, at complex values of the decoupling parameters $s(\Delta)$ on the circles $|s(\Delta)| = e^{\kappa_1}$ on which they enter Cauchy integrals [10, (1.23), p. 7], is Supplement, Proposition S.12(h); it costs the further constants collected in C_w and no further power of ε/a_j. *q.e.d.*

The proof shows where the exponent comes from. The local operators are compared at the rate $(\varepsilon/a_j)^\beta$ for every $\beta < 1$, and the loss is in the change of the background, whose rate $(1-\varepsilon_0)/3$ is that of the distance between the minimisers of the two runs in the norm of the analyticity domains. This distance is the subject of the next section.

10 The Large-Field Terms of Two Cutoffs

We now turn to the terms of the two densities which carry large fields. Here the obvious approach, a comparison of the two runs term by term, meets two obstacles. The first is that most of these terms cannot be compared at all with an error that is small relative to the term: a large-field region which was created at a fine scale has been integrated with the test functions, cutoffs, and Gaussian weights of the fine lattice of its run, and the two fine lattices are different. The second is that the test functions are evaluated on fields of the two runs which differ, so that a term of one run may be supported where the corresponding term of the other vanishes. The resolution of the first obstacle is that we do not need to compare the terms with an old large-field region at all: such a term carries the gain of every scale through which the region has lived, and the weight of Proposition 8.4 shows that the terms of this kind have a total weight which tends to zero with K, in each run separately. The terms that remain have their large fields at scales coarser than $K'/2$, where the fine lattice is invisible up to a power of $\varepsilon/a_{K'/2} = L^{-K'/2}$. The second obstacle is removed by the soft test functions, whose change is proportional to that of their arguments.

10.1 The Matched Decomposition

Both densities at the stopping step have the form (3.23), a sum over the admissible sequences of domains, with one pair (Ω_j, Λ_j) for each step, and with the components which the **R**-operation has renormalised removed from the sequences [11, (0.3), p. 176]. The sequences of run B have one entry more, that of the extra step.

Lemma 10.1 (Matched Decomposition)**.** *Call a labelled term of either run, at its stopping step,* early *if its pair of domains differs from* $(\mathbb{T}, \mathbb{T})$ *at a step whose shared scale is at most* 1*: for run B the extra step or the step* 2*, for run A the first step. Then every labelled term is of exactly one of two kinds:*

(M) it is not early; deleting the pair of the extra step from a term of run B of this kind gives a term of run A of this kind, and this is a bijection $\tau^+ \mapsto \tau$ between the terms of kind (M) of the two runs;

(U) it is early; then it has a component of its large-field region at its stopping step whose history contains a large-field region created at a shared scale at most 1*.*

Proof. A component which is renormalised at some step leaves the sequences, so a term whose pair at some step is not $(\mathbb{T},\mathbb{T})$ has a large-field region created at that step which belongs to a component alive at the stopping step; for an early term this is the kind (U). The shared scale 1 has to be excluded together with the extra step because the two runs do not perform the same operation there: it is the first step of run A, which is carried out at the retreated region, with labels refined by that region (Supplement, Proposition S.7), whereas for run B it is the second step, an ordinary one. A term of run A whose pair at the first step is $(\mathbb{T},\mathbb{T})$ has the empty retreat, because the retreat removes cubes near the boundary of the region of integration and the torus has none, so that its label at this step is the pair alone; and a term of run B whose pairs at the extra step and at the step 2 are $(\mathbb{T},\mathbb{T})$ has no further label at these steps either. For the later steps, the admissibility conditions on the sequences [9, pp. 255–256] and the conditions of the **R**-operation [11, p. 177, (i), (ii)] are conditions on the domains, stated with the cube sizes, the distances MR_j, and the depth N_j of the scale; by Lemma 9.1 these are the same at a shared scale, so that the conditions of run B at the steps $3,\dots,K'+1$ are those of run A at the steps $2,\dots,K'$. This gives the bijection. *q.e.d.*

For a matched pair (τ^+,τ) the two terms are the same operations applied to ingredients which depend on the run. The pair is the *all-small-field pair* if $\tau=\ell_0$. Otherwise let j_0 be the least scale at which one of its terms has a nonempty large-field region; since every nonempty large-field region of a term belongs, after the **R**-operations, to the history of a component which is alive at the stopping step, j_0 is the scale at which the oldest of these components was born, and $\Lambda_i=\Omega_i=\mathbb{T}$ for $i<j_0$ in both terms. With

$$m \coloneqq \lfloor K'/2 \rfloor \,, \tag{10.1}$$

the pair is *old* if $K'-j_0\ge m$ and *young* otherwise. A young pair has all its large fields at the shared scales $i\ge j_0-1\ge K'-m\ge K'/2$; this is the case in which a comparison is possible.

The terms which are not compared are disposed of by the following lemma, which uses only the weight of Section 8.3. We write

$$\mathfrak{b}(\mathcal{C}) \coloneqq e^{B_E(\mathcal{C})+I_*(\mathcal{C})} \,, \tag{10.2}$$

so that $\mathfrak{b}(\mathcal{C})\le B(\mathcal{C})$, and we use the notation of Theorem 8.1 for either run, with Z the integral of the density and $N\coloneqq\sum_\ell Z_\ell\langle\widehat{\mathcal{W}}_\ell\rangle_\ell$, so that $W=N/Z$.

Lemma 10.2 (Terms with an Old Component)**.** *Let $\bar g$ be as in Theorem 8.7, let $p\ge 0$, and let $\mathcal{S}$ be a set of labelled terms of one run, none of them ℓ_0, each of which has a component X of its large-field region at the stopping step with $P(X)\ge p$. Then*

$$\sum_{\ell\in\mathcal{S}} Z_\ell \le e^{-C_P p}\delta_{\mathrm{LF}} Z_{\ell_0} \,, \qquad \sum_{\ell\in\mathcal{S}} Z_\ell |\langle\widehat{\mathcal{W}}_\ell\rangle_\ell| \le \mathfrak{b}(\mathcal{C}) e^{-(C_P-1)p}\delta_{\mathrm{LF}} Z_{\ell_0} \,, \tag{10.3}$$

and removing the terms of $\mathcal{S}$ from N and Z changes $W=N/Z$ by at most $2B(\mathcal{C})e^{-(C_P-1)p}\bar\delta_{\mathrm{LF}}$.

Proof. For each $\ell\in\mathcal{S}$ choose a component X_ℓ with $P(X_\ell)\ge p$. For the first bound let $\omega(X_\ell)\coloneqq e^{C_P P(X_\ell)}$ and $\omega(X)\coloneqq 1$ for the other components; these numbers satisfy the hypothesis of Proposition 8.4, so that

$$\sum_{\ell\in\mathcal{S}} Z_\ell \le e^{-C_P p}\sum_{\ell\in\mathcal{S}} Z_\ell\,\omega(X_\ell) \le e^{-C_P p}\sum_{\ell\ne\ell_0} Z_\ell \prod_{X\in Z(\ell)} \omega(X) \le e^{-C_P p}\delta_{\mathrm{LF}} Z_{\ell_0} \,. \tag{10.4}$$

For the second we use Lemma 8.6, $|\langle\widehat{\mathcal{W}}_\ell\rangle_\ell|\le \mathfrak{b}(\mathcal{C})\prod_X \omega_A(X)$, and the numbers $\omega(X_\ell)\coloneqq \omega_A(X_\ell)e^{(C_P-1)P(X_\ell)}$ and $\omega(X)\coloneqq\omega_A(X)$ otherwise. Since $1\le\omega_A(X)\le e^{P(X)}$ under (8.42), as

the proof of Lemma 8.6 shows, they satisfy $1 \le \omega(X) \le e^{C_P P(X)}$, and the same computation gives the second bound. Finally, if $N_{\mathcal{S}}$ and $Z_{\mathcal{S}}$ are the contributions of $\mathcal{S}$, then $Z - Z_{\mathcal{S}} \ge Z_{\ell_0}$ and

$$\Big|\frac{N - N_{\mathcal{S}}}{Z - Z_{\mathcal{S}}} - \frac{N}{Z}\Big| = \Big|\frac{N_{\mathcal{S}} - W Z_{\mathcal{S}}}{Z - Z_{\mathcal{S}}}\Big| \le \frac{|N_{\mathcal{S}}| + |W| Z_{\mathcal{S}}}{Z_{\ell_0}} \le (\mathfrak{b}(\mathcal{C}) + B(\mathcal{C})) e^{-(C_P - 1)p} \delta_{\mathrm{LF}} \,, \qquad (10.5)$$

by (10.3) and Theorem 8.7, and $\delta_{\mathrm{LF}} \le \bar{\delta}_{\mathrm{LF}}$. *q.e.d.*

Corollary 10.3 (The Unmatched Terms and the Old Pairs)**.** *Under the hypotheses of Lemma 10.2, removing the terms of kind (U) from both runs changes $W_{K+1}(\mathcal{C}) - W_K(\mathcal{C})$ by at most $4B(\mathcal{C}) e^{-(C_P-1)K'} \bar{\delta}_{\mathrm{LF}}$, and removing the old pairs from both runs changes $W_{K+1}(\mathcal{C}) - W_K(\mathcal{C})$ by at most $4B(\mathcal{C}) e^{-(C_P-1)(m+1)} \bar{\delta}_{\mathrm{LF}}$.*

Proof. The large-field regions grow and merge until they are renormalised [11, (0.3), p. 176], so a component alive at the stopping step whose history begins at the scale j meets the large-field region at every scale from j to the stopping step, and its history volume (8.21), in which every scale with a nonempty intersection contributes at least L^4, is at least the number of these scales. For a term of kind (U) this number is at least K', the number of the steps from the shared scale 1 to the stopping step, and for each term of an old pair it is at least $m + 1$. Lemma 10.2 gives the assertions. *q.e.d.*

10.2 The Shells of the Test Functions

On a young pair the two terms are compared ingredient by ingredient, and the first ingredients are the test functions. Let Π_X be the product of the test functions and characteristic functions of run X's term and ϱ_X the rest of its integrand. Since all factors of Π_X take values in $[0, 1]$,

$$\Pi_A \varrho_A - \Pi_B \varrho_B = \min(\Pi_A, \Pi_B)(\varrho_A - \varrho_B) + (\Pi_A - \Pi_B)_+ \varrho_A - (\Pi_B - \Pi_A)_+ \varrho_B \,, \qquad (10.6)$$

which puts each of the last two pieces on the support of one run's own term, where that run's bounds apply.

Lemma 10.4 (Shells of the Soft Test Functions)**.** *Let a soft test function $\bar{\chi}(f_\square/t)$ of a cube $\square$ have the same threshold t in both runs, and let $|f^A_\square - f^B_\square| \le \delta_\square t$ wherever $\min(f^A_\square, f^B_\square) < (1 + 2\kappa_s)t$. Then $|\bar{\chi}(f^A_\square/t) - \bar{\chi}(f^B_\square/t)| \le \delta_\square/\kappa_s$, and the same holds for $1 - \bar{\chi}$. Consequently, if Π_A and Π_B consist of such soft test functions and of characteristic functions which are the same in both runs,*

$$(\Pi_A - \Pi_B)_+ \varrho_A \le \Big(\sum_\square \frac{\delta_\square}{\kappa_s}\Big) \Pi^\sharp_A \varrho_A \,, \qquad (10.7)$$

where $\Pi^\sharp_A$ is the product of the sharp test functions of run A at the thresholds $(1 + \kappa_s)t$ for the passed and t for the failed test functions; for a test function of the cascade of [11, (1.22)–(1.24), p. 181] the factor $\delta_\square/\kappa_s$ is replaced by $(N + 1)\delta_\square/\kappa_s$ and the thresholds by those of Supplement, Proposition S.2.

Proof. Where $\min(f^A_\square, f^B_\square) \ge (1+2\kappa_s)t$ both values of $\bar{\chi}$ vanish. Elsewhere $|f^A_\square/t - f^B_\square/t| \le \delta_\square$, and $\bar{\chi}$ is κ_s^{-1}-Lipschitz by Lemma 3.8(iii). Replacing the factors of Π_A by those of Π_B one at a time, with all factors in $[0, 1]$, gives $(\Pi_A - \Pi_B)_+ \le \mathbf{1}\{\Pi_A > 0\} \sum_\square \delta_\square/\kappa_s$, and on $\{\Pi_A > 0\}$ every passed test function of run A has $f^A_\square < (1 + \kappa_s)t$ and every failed one $f^A_\square > t$, by Lemma 3.8(ii). The cascade test functions are soft with the width $\kappa_s/(N + 1)$ (Supplement, Proposition S.2), which gives the factor $N + 1$. *q.e.d.*

The right side of (10.7) is a term of run A with sharp test functions at enlarged thresholds, and Bałaban's bounds hold for it by Proposition 3.12: a shell costs, relative to Bałaban's bound of the term, the factor $\delta_\square/\kappa_s$. The characteristic functions which restrict the fluctuation fields are sharp, and their thresholds differ between the runs through the factor g_j of Definition 3.11(ii). For them we use the density of the Gaussian weight.

Lemma 10.5 (Shells of the Coupling). *Let $|\delta_j| \le \frac{1}{2}(g_j^A)^{-2}$, and let the fluctuation field $A_j(b)$ of a bond be restricted in the two runs by $|A_j(b)| < t$ and $|A_j(b)| < t'$ with $t = \vartheta_j/g_j$ and $t' = \vartheta_j/g'_j$, g_j and g'_j the two couplings at the shared scale j. Then $|t' - t| \le 2tg_j^2|\delta_j|$, and the mass of the shell between the two radii under the normalised Gaussian weight of the step, with the other variables fixed, is at most*

$$C_{\mathrm{sh}}\, p_0(\hat g_j)^{\dim G} g_j^2 |\delta_j| \,, \tag{10.8}$$

with a constant C_{sh} of L, G, and A_1/A_0. The same holds for the field B_j of [11, (1.55), p. 188] with δ'_j in place of ϑ_j.

Proof. With $x := g_j^{-2}$ and $x' := g_j'^{-2}$ we have $t'/t = g_j/g'_j = (x'/x)^{1/2}$ and $|x'-x| = |\delta_j| \le \frac{1}{2}x$, so that $x'/x \in [\frac{1}{2}, \frac{3}{2}]$, and $|u^{1/2} - 1| \le |u - 1|$ for $u > 0$ gives $|t'/t - 1| \le |\delta_j|/x = g_j^2|\delta_j|$; exchanging the runs gives the bound with either coupling and the factor 2. The Gaussian weight of the fluctuation field has the precision $Q = C^*\Delta^{(j)}C - C^*\Delta^{(j)}CC^{(j)}C^*\Delta^{(j)}C \le C^*\Delta^{(j)}C$ of [9, (2.21), p. 258], because $C^{(j)} \ge 0$. The conditional variance of $A_j(b)$, the other variables fixed, is $(Q_{bb})^{-1} \ge \|C^*\Delta^{(j)}C\|^{-1}$, so that the conditional density of $A_j(b) \in \mathfrak{g}$ is at most $(\|C^*\Delta^{(j)}C\|/2\pi)^{\dim G/2}$, a constant of the lattice. The shell $\{\min(t,t') \le |A_j(b)| < \max(t,t')\}$ has Lebesgue measure at most $C|t'-t|\max(t,t')^{\dim G-1}$, and $t \le (A_1/A_0)p_0(\hat g_j)\cdot \hat g_j/g_j \le (A_1/A_0)p_0(\hat g_j)$ by Lemma 3.2(i), which gives (10.8). The same argument applies to B_j under the Gaussian form of [11, p. 188], and to the complement of a characteristic function, whose shell is the same set. *q.e.d.*

The positivity of the quadratic form is the same fact by which Bałaban bounds the integrals over the fluctuation fields in the large-field operations [12, p. 383], and so a shell term is at most (10.8) times his bound of the term. Two features of these shell bounds matter for what follows. The plaquette shells cost $\delta_\square/\kappa_s$, which will be a power of ε/a_j; the coupling shells cost a multiple of $g_j^2|\delta_j|$, which is controlled in Section 11 together with the other differences of the couplings.

10.3 The Configurations of the Two Runs

The functions $f_\square$ of the plaquette and reference test functions, and all integrands of the large-field operations, are computed from minimisers of the action on the fine lattice of the run. At equal data the minimisers of the two runs differ by a lattice artefact, which is the content of the following proposition. For a field W on the fine lattice of run B we write MW for its average on the fine lattice of run A, and for a point x of a determining set $j(x)$ is the layer of x.

Proposition 10.6 (Minimiser Distance). *Let $0 < \varepsilon_0 < 1/125$, let the constants satisfy the conditions of Appendix C, and let $\bar g$ be sufficiently small. There are constants $C_G(\varepsilon_0)$ and C_U, independent of K, k, ε, and the torus, with the following properties.*

(i) Let U^A and U^B be the minimisers of the two runs for a determining set at the shared scales with the same data V, which satisfy the hypothesis (7) of [7, Theorem 1, p. 279]

with the parameter ε_1. Then, in a gauge, $U^A = e^{i\varepsilon A'} MU^B$ with

$$a_{j(x)}|A'(x)| + a_{j(x)}^2|\nabla A'(x)| \le 2\bar\rho(x)\,, \qquad \bar\rho(x) := C_G(\varepsilon_0)\Big(\frac{\varepsilon}{a_{j(x)}}\Big)^{1-\varepsilon_0}\varepsilon_1\,. \tag{10.9}$$

(ii) For each of the configurations U_k'', U_k^0, $U_{k,\Lambda}$, U_k of the **R***-operation of [12, §1] on a young history born at the scale n, at the data of the two runs, $c^A = e^{i\varepsilon A_c} Mc^B$ with*

$$a_{j(x)}|A_c(x)| + a_{j(x)}^2|\nabla A_c(x)| \le \bar d(x) := C_U\Big(\frac{\varepsilon}{a_{\min(j(x),n)}}\Big)^{(1-\varepsilon_0)/3}\varepsilon_1\,. \tag{10.10}$$

Here $\varepsilon_1 \le \bar\varepsilon_1$, the number of (10.11) *below, which depends on $\bar g$ and the constants of the construction and not on K.*

The proof is Supplement, Proposition S.13. In (i), which is the case of a single run of [7, Theorem 1] on two lattices, MU^B satisfies run A's constraint, and the difference of the two actions at MU^B is the blocking artefact $b(W) := A_\varepsilon(MW) - A_{\varepsilon/L}(W)$, whose first variation is a tangential current of order ε/a_j; the gap of the action at the minimum and the bounds of [5] for the Green operator of the constrained problem turn it into (10.9), with the loss ε_0 in the Hölder norm. In (ii) the data themselves differ, on the region Λ where the **R**-operation integrates, through the minimiser U_0 of [12, (1.3), p. 357]; the minimiser of the nested problem of [11, Proposition 1, p. 194] is compared by the growth of its action away from the minimum. If Δ is the distance between the minimising data of the two runs, the quadratic growth $\frac12\kappa_\Lambda\Delta^2$ is at most the change of the difference of the two runs' actions between the two minimisers, a sum over blocks of the smaller of a bound of order $(\varepsilon/a_n)^{1-\varepsilon_0}$ and a bound linear in Δ; with $\min(u,v) \le (uv)^{1/2}$ this gives $\Delta^{3/2} \le C(\varepsilon/a_n)^{(1-\varepsilon_0)/2}$, whence the exponent $(1-\varepsilon_0)/3$.

The data parameter of the regularity of the history is

$$\begin{aligned}\varepsilon_1 &:= \max\Big(C\max_{j\le K'}(\varepsilon_j + \vartheta_j),\, C_M R_{K'}\varepsilon_{K'}\Big)\\ &\le \bar\varepsilon_1 := \max(C(1+A_1/A_0)\bar g p_0(\bar g),\, C_M L A_0 \bar g(\log \bar g^{-2})^{p_0+r})\,,\end{aligned} \tag{10.11}$$

where C is the constant with which [11, p. 178] bounds the data of a large-field region by its thresholds and C_M a constant of M [11, (1.83), p. 197]; the bound holds because $\varepsilon_j + \vartheta_j = (1 + A_1/A_0)\hat g_j p_0(\hat g_j)$ with $\hat g_j \le \bar g$, because $R_{K'}\varepsilon_{K'} < LA_0\hat g_{K'}(\log \hat g_{K'}^{-2})^{p_0+r}$ by Lemma 3.4(i), and because $g \mapsto gp_0(g)$ and $g \mapsto g(\log g^{-2})^{p_0+r}$ increase for small g. The point of (i) is the exponent: the tested quantities of a cube at the scale j differ between the runs by a power of ε/a_j, which is small precisely because the young histories live at the scales $j \ge K'/2 - 1$. In terms of the thresholds:

Corollary 10.7 (The Tested Quantities)**.** *Under the hypotheses of Proposition 10.6, let $f_\square^X$ be the quantity which a plaquette or reference test function of a cube $\square$ at the scale j of a young history evaluates in run X, at the threshold t. Wherever $\min(f_\square^A, f_\square^B) < (1+2\kappa_s)t$, at equal data,*

$$|f_\square^A - f_\square^B| \le \delta_j^s t\,, \qquad \delta_j^s := \delta_j^f + 7g_j^2|\delta_j|\,, \qquad \delta_j^f := C_f\Big(\frac{\varepsilon}{a_j}\Big)^{1-2\varepsilon_0}\,, \tag{10.12}$$

with a constant C_f which is independent of K, k, ε, and the torus, and at most polynomial in j_q.

Proof. For the test functions of the local minimisers of the cubes [11, (1.3), (1.4), (1.7), (1.8), p. 178], [9, (2.17), p. 257], the plaquettes of U^A and of MU^B differ by the curl of $\varepsilon A'$ with transports, at most $a_j^2|\nabla A'| + C(a_j|A'|)^2 + C(a_j|A'|)B_3\varepsilon_1 \leq 3\bar{\rho}$ in the units of the scale for $\bar{g}$ small, by (10.9) and [7, (8)]; the plaquettes of MU^B and of U^B differ by the blocking artefact, of the same order. Relative to the threshold, which is at least $(1-\beta)\varepsilon_j \geq \delta'_j$, this is at most a constant times $(\varepsilon/a_j)^{1-\varepsilon_0}\varepsilon_1/\delta'_j$. On a young history the scales satisfy $K' - j + 1 \leq 8(1+j_q)(j+1)$, and ε_1/δ'_j is at most a polynomial in j and $\log \hat{g}_j^{-2}$ with coefficients polynomial in j_q, by (10.11), Lemma 3.2, and $p_1 < p_0$; this polynomial is absorbed by $(a_j/\varepsilon)^{\varepsilon_0} = L^{\varepsilon_0 j}$, which gives δ_j^f. The test functions of the multiscale minimisers, those of the cascade [11, (1.22)–(1.24), p. 181], of the reference configurations [11, (1.27), p. 182], and of [11, (1.75), p. 193], are bounded in the same way, with the layer of the tested plaquette in place of j (Supplement, Proposition S.13(c)). The coupling part $7g_j^2|\delta_j|$ enters only where a tested quantity contains the scaled field $g_j B_j$, as in [11, (1.27), (1.59)], by the computation of Lemma 10.5. *q.e.d.*

10.4 The Large-Field Comparison

We can now state the comparison of the large-field terms of a young pair. Let $\tilde{\theta} \in (0,1)$ be the rate of (11.22) in Section 11; in this section we use only $\tilde{\theta} \geq L^{-(1-\varepsilon_0)/4}$ and $\tilde{\theta} \geq L^{-\theta_V/8}$, with θ_V of (11.8). The ingredients of a term are the factors of the integrands of its one-step operations and the terms of its effective action. For a young pair we distinguish three kinds of them at the scales $i \geq j_0 - 1$:

(α) the ingredients of the small-field integration: the frozen terms of the effective action, the Gaussian weights and their normalisations, the function [11, (1.6), p. 178] of the block variables, and the localised constants;

(β) the test functions of the plaquettes and of the reference configurations, and the characteristic functions of the fluctuation fields;

(γ) the large-field integrands: the action and the quadratic form of [12, (1.70), p. 377], the minimiser U_0 and the operators $H_{1,k}$ and $\Delta_1(\zeta_0)$ of the denominators of the renormalisation quotients [12, (1.1)–(1.2), p. 356], and the factors $V(Y)$ of [12, (1.91)] and the boundary terms of the steps at which a large-field region existed, as functions of the configurations and the propagators of the run, the terms of the effective action from which they are built being held fixed.

The dependence of the factors $V(Y)$ and of the boundary terms on the terms of the effective action from which they are built is of the kind (α). The ingredients (α) are compared in Section 11, by the recursion which also compares the all-small-field pair. The ingredients (β) and (γ) are compared by the following theorem, which says that their run dependence costs, per point of the history of a component, an amount which decays geometrically in the scale at which the component was born. For a component X of a young pair let $n(X)$ be its birth scale and $|Z_i \cap X|_i$ the number of points of the scale i of its history at the scale i, as in (8.21). Write M_τ^A for the majorant (8.26) of run A's term and $M_{\tau^+}^B$ for that of run B's.

Theorem 10.8 (Large-Field Comparison)**.** *Let the constants satisfy the conditions of Appendix C, and let $\bar{g}$ be sufficiently small. There are a constant C_M of L and G, a constant C_h which is independent of K, ε, and the torus and at most polynomial in j_q, and an index k_{LF} with the following property. For $K' \geq k_{\mathrm{LF}}$ and every young matched pair (τ^+, τ), let $\varrho^{A,B}[\tau]$*

be run A's term with its ingredients of the kinds (β) and (γ) replaced by run B's, except for the factors (10.6) *of the shells. Then*

$$\int |\varrho^{A,B}[\tau]-\varrho^{A}[\tau]|\,\mathrm{d}V \le \Big(\exp\Big(\sum_X \sum_i C_M s^{\mathrm{LF}}_{n(X)} L^4 |Z_i\cap X|_i\Big)-1+\sigma_\delta\Big) M^A_\tau\ , \qquad s^{\mathrm{LF}}_n \coloneqq C_h\tilde\theta^n\ , \tag{10.13}$$

the sum over the components X of the pair and the scales i of their histories, where $\sigma_\delta \coloneqq \sum_X P(X)\sup_{i\ge K'-m}|\delta_i|$ collects the coupling parts of the shells; the shell pieces of (10.6) *satisfy the same bound, with $M^B_{\tau^+}$ for the piece on run B's support. The same bounds hold for the numerators and the denominators of the renormalisation quotients of [12, (1.1), (1.2), p. 356] at the complex arguments of the inductive class, relative to Bałaban's majorants of their moduli, and linearly in the norms of the classes of the effective actions of the two runs.*

Proof. We treat the two kinds of ingredients in turn and then sum.

The ingredients (β). By the redundancy of Bałaban's test functions [9, (3.6)–(3.9), p. 265–266; p. 268–269], a test function of the scale j which is not identically one on the support of the other characteristic functions of the term lies in the large-field region of the term or in its collar, so on a young pair it lies on a cube of a history at a scale $j \ge j_0-1$, or on a cube of the stopping scale. By Lemma 10.4 and Corollary 10.7 each of them contributes the factor δ^s_j/κ_s, and $(N+1)\delta^s_j/\kappa_s$ for a cascade test function; by Lemma 10.5 each characteristic function of a fluctuation field contributes at most $C_{\mathrm{sh}}p_0(\hat g_j)^{\dim G}g_j^2|\delta_j|$. A cube of the scale j carries the test functions of the step j and of the **R**-operations of the steps $k' \ge j$ with $k'-N_{k'} \le j$, at most 14 of each step and two of the cascade, and there are at most $s_T(j) \coloneqq 2+N_j$ such steps, because N is nonincreasing (Lemma 3.10). Since $N_j \le \log_L C_{\mathrm{dep}} + \nu_L \log_L \log g_j^{-2}$ by (3.39), the first parts of all the test functions of a cube are at most a constant times $(j+1)s_T(j)L^{-(1-2\varepsilon_0)j}$, and with $L^{-(1-2\varepsilon_0)j} \le \tilde\theta^j L^{-(3-7\varepsilon_0)j/4}$, which holds because $\tilde\theta \ge L^{-(1-\varepsilon_0)/4}$, and $j \ge n(X)-1$, this is at most $s^{\mathrm{LF}}_{n(X)}$ once C_h contains $\tilde\theta^{-1}\sup_j(j+1)s_T(j)L^{-(3-7\varepsilon_0)j/4}$, which is finite since $s_T(j)$ grows like $\log\log j$. The coupling parts of all the test functions and characteristic functions of a cube sum to at most $(7/\kappa_s)16s_T(j)^2g_j^2|\delta_j| + 8s_T(j)C_{\mathrm{sh}}p_0(\hat g_j)^{\dim G}g_j^2|\delta_j| \le |\delta_j|$ for $\bar g$ small, and the cubes of the histories are counted by $P(X)$.

The ingredients (γ). By Proposition 10.6, the configurations of the **R**-operation of the two runs differ, in the units of their layers, by $(\varepsilon/a_n)^{(1-\varepsilon_0)/3}$ times a constant, where $a_n \ge a_{n(X)-1}$ is the spacing of the finest layer of the history. Each large-field integrand is a function of these configurations whose two-run difference is bounded per point of the history by a constant times $g_j^{-2}(\varepsilon/a_n)^{(1-\varepsilon_0)/3}\varepsilon_1^2$, for the actions and quadratic forms, by a constant times $(\varepsilon/a_n)^{(1-\varepsilon_0)/3}N_k$, for the forms of the denominators, and, for the factors $V(Y)$ relative to their bound $|V(Y)| \le \alpha e^{-(1+2\beta)\kappa d(Y)}$ of [12, (1.68), p. 377], by a sum over the generations j of the terms which they re-sum of powers θ_V of the configuration distance relative to the radii of analyticity, with the factor $L^{-(\beta'/2)(m-j)}$ of the age, where Theorem 9.5 compares the propagators through which they are built; the boundary terms are compared through the recursion of their generation, under the third condition of (C.2). These bounds are Supplement, Proposition S.14. The factors g_j^{-2}, $\varepsilon_1^2/\varepsilon_j^2$, N_k, and the polynomial dependence on the scale are absorbed into the geometric factor, as in the first part, since $(1-\varepsilon_0)/3 > (1-\varepsilon_0)/4$, and for the factors $V(Y)$ since $\tilde\theta \ge L^{-\theta_V/8}$.

The sum. Every ingredient of the two kinds enters per point of the history of a component at a scale $i \ge n(X)-1$, and its contribution relative to Bałaban's bound is at most $C_M s^{\mathrm{LF}}_{n(X)}$ per point plus the coupling parts, with C_M the number of ingredients per point, a constant of L and G. Bałaban's bound of a term is proved by estimating the integrals over the

group variables, with their normalised weights, by one, and those over the fluctuation fields by the positivity of their quadratic forms [12, p. 383]. A product of factors which each change by a relative amount $x_\iota \geq 0$ changes by at most $\prod_\iota (1 + x_\iota) - 1 \leq \exp(\sum_\iota x_\iota) - 1$ relative to the product of the bounds, which gives (10.13). For the renormalisation quotients the same argument applies to the moduli of the numerators and denominators at complex arguments, where the shells are those of the real integration variables, and the dependence on the effective actions of the two runs enters linearly through the norms of their classes (Supplement, Proposition S.14(e)). *q.e.d.*

The theorem is the per-point form in which the large fields enter the recursion of Section 11: the factor $s^{\mathrm{LF}}_{n(X)}$ is attached to the birth scale of the component and not to the scale of the point, so that its sum over all points of a history, counted by $P(X)$, is controlled by the weight of Proposition 8.4. The stopping-scale test functions of the all-small-field pair are the one place where a shell does not lie on a history, and they are treated directly.

Proposition 10.9 (The Stopping-Scale Shells)**.** *Under the hypotheses of Theorem 10.8, replacing the stopping-scale test function $\chi_{K'}(\mathbb{T})$ of run A's all-small-field term by run B's, in the numerator and in the normalisation, and the same for run B, changes $W_{K+1}(\mathcal{C}) - W_K(\mathcal{C})$ by at most $b^{(S)}_K B(\mathcal{C})$, with*

$$b^{(S)}_K := 4\kappa_s^{-1} e^{2C_K} N_q^4 \delta^f \,, \qquad \delta^f := C_f L^{-(1-2\varepsilon_0)K'} \,, \tag{10.14}$$

which is summable in K.

Proof. The term ℓ_0 carries the test function (3.22) of the stopping step, a product over at most N_q^4 cubes, built on each run's fine lattice at the common threshold $\varepsilon_{K'}$, and no other test function which is not identically one (Supplement, Proposition S.2). By Corollary 10.7 at $j = K'$, where $\varepsilon/a_{K'} = L^{-K'}$ and $\delta_{K'}$ does not enter, and by Lemma 10.4, the shell of each run is at most $N_q^4 \delta^f/\kappa_s$ times that run's term with sharp test functions at the enlarged thresholds, whose integral is at most e^{C_K} by the proof of Proposition 8.4. Relative to $Z_{\ell_0} \geq e^{-C_K}$ this is at most $N_q^4 \delta^f \kappa_s^{-1} e^{2C_K}$, and the change of the numerator is at most $\mathfrak{b}(\mathcal{C}) \leq B(\mathcal{C})$ times this by Lemma 8.6. Arguing as in (10.5) for the numerator and the normalisation of each run gives the factor $4B(\mathcal{C})$. The sum over K converges because C_K and N_q do not depend on K. *q.e.d.*

11 The Limit along the Full Sequence

We can now compare $W_{K+1}(\mathcal{C})$ with $W_K(\mathcal{C})$. By Theorem 8.1 each of them is the average of the normalised dressed loops against the density of its run on the common space $G^{\mathcal{B}}$, so that the comparison is a comparison of two densities and of two families of functions on one space. The difference between the densities is, at first sight, of order one: the exponents of the two runs contain the effective actions of all the steps, and the terms of the first steps, produced on the fine lattices, differ by quantities which do not tend to zero. What saves the comparison is the decay of Bałaban's effective actions in the age of their terms, by which a term produced m steps before the stopping step contributes at most $E_1 L^{-\beta m}$ per point of the stopping lattice [9, (2.43), p. 263]. The old terms can therefore be bounded without comparing them, and only the recent ones, produced at scales at which the fine lattice is far away, have to be compared. The comparison of the recent terms is a recursion: the difference of the terms produced by one step is bounded by the differences of the terms which enter it, through a contraction which is Bałaban's, and by sources which are the lattice artefacts of

Sections 9 and 10. Its solution decays geometrically in the scale, up to a term from the fine scales which is a negative power of K.

Theorem 11.1 (Comparison of Two Cutoffs)**.** *Under the hypotheses of Theorem 1.5 there are an index $K_1 \geq K_0(\mathcal{C})$ and nonnegative numbers a_K, given for $K \geq K_1$ by* (11.32) *and for $K_0(\mathcal{C}) \leq K < K_1$ by $a_K := 2B(\mathcal{C})$, such that $\sum_K a_K < \infty$ and $|W_{K+1}(\mathcal{C}) - W_K(\mathcal{C})| \leq a_K$ for every $K \geq K_0(\mathcal{C})$. The index K_1 depends on $\mathcal{C}$ through the quantities on which $B(\mathcal{C})$ depends, and on q, T/u, $\bar{g}$, and the constants of the construction.*

For the finitely many $K < K_1$ there is nothing to prove, since $|W_K(\mathcal{C})| \leq B(\mathcal{C})$ for every $K \geq K_0(\mathcal{C})$ by Theorem 8.7. The rest of this section proves the bound for $K \geq K_1$, first for the all-small-field pair and then for the other terms. Throughout, the two runs are those of Section 9.1.

11.1 The Exponents of the All-Small-Field Pair

On the term ℓ_0 of either run the large-field regions are empty, the coupling (3.26) is constant on the last domain, and the exponent of the density at V is, by (3.25)–(3.27),

$$-\frac{1}{g_{\text{stop}}^2} A_{\text{fine}}(U(V)) + \sum_j \text{irr}_j(U(V)) + \sum_j [\mathbf{R}^{(j)}(U(V)) - \mathbf{R}^{(j)}(1)] - E \,, \tag{11.1}$$

with $\text{irr}_j(U) := \mathbf{E}^{(j)}(U) - \mathbf{E}^{(j)}(1) - \beta_j A(\phi_j, U)$ the quantity bounded by (3.31), where for run A $g_{\text{stop}} = g_{K'}^A$, $A_{\text{fine}} = A_\varepsilon$, and $U = U^A(V)$ is the minimiser on the lattice of spacing ε with $M^{K'}U = V$, and for run B $g_{\text{stop}} = g_{K'+1}^B$, $A_{\text{fine}} = A_{\varepsilon/L}$, and $U = U^B(V)$ with $M^{K'+1}U = V$. To compare the terms of the same shared scale we carry run A's arguments to run B's fine lattice. For a localisation domain X and a term $\mathbf{E}^{B,(j+1)}(X, \cdot, z)$ of run B let

$$\mathcal{G}_j(X, (\mathbf{U}, \mathbf{J}), z) := \mathbf{E}^{B,(j+1)}(X, (\iota_X \mathbf{U}, Q_1^{\text{pl}}(\iota_X \mathbf{U})^* \mathbf{J}), z) \,, \qquad \Delta_j := \mathcal{G}_j - \mathbf{E}^{A,(j)} \,, \tag{11.2}$$

where $Q_1^{\text{pl}}(W)$ is the plain linearised one-step average at W and $\mathbf{J}$ is the current variable of the spaces of [8, (1.10)–(1.16), p. 262]; both are evaluated at the current $\mathbf{J} = J(U)$ of the configuration, so that Δ_j is a functional of fields on the lattice of spacing ε. We write $\text{irr}(\Delta_j)(U) := \Delta_j(U) - \Delta_j(1) - \Pi^\Delta A(h_z, U) - N_j(U)$, with Π^Δ the isotropic coefficient of Δ_j in the sense of [9, (3.57)–(3.61), pp. 281–282], h_z the tent functions of [9, (3.65), p. 283], and N_j the part of the second moment of $\mathcal{G}_j$ off its isotropic coefficient: since $\mathcal{G}_j$ evaluates every domain at its own lift, the Euclidean covariance by which Bałaban shows that this part vanishes [9, (3.60)–(3.61), p. 282] is not available for it, and N_j is kept as a term of its own, a sum of the 36 plane-pair actions $\operatorname{tr} F_{\kappa\mu} F_{\lambda\nu}$ of [9, (3.57)]. Its coefficients are small for a reason which is specific to the lift: for affine data the minimisers of both runs are affine fields and the global lift carries the affine field of run A to that of run B, so that the second moment of $\mathcal{G}_j$ would be that of run B's own family if every domain were evaluated at the global lift; what remains is the difference between the localised and the global lift on a domain, which decays exponentially in $M_1 L^j$, and each coefficient of N_j, as well as $\Pi^{\mathcal{G}} - \beta_{j+1}^B$, is at most $C_\Pi E_0 e^{-\delta_c M_1 L^j}$ (Supplement, Proposition S.15(d)). Similarly $\text{irr}_j^{\mathcal{G}}(U) := \mathcal{G}_j(U) - \mathcal{G}_j(1) - \Pi^{\mathcal{G}} A(h_z, U) - N_j(U)$, and $\mathcal{G}_j(1) = \mathbf{E}^{B,(j+1)}(1)$ because $\iota_X 1 = 1$ and $J(1) = 0$.

Lemma 11.2 (Exponent Difference)**.** *At every V in the support of both terms ℓ_0, the exponent*

of run B minus that of run A is

$$D(V) = -\delta_{K'} A_{\varepsilon/L}(U^B) - (g^A_{K'})^{-2}[A_{\varepsilon/L}(U^B) - A_\varepsilon(U^A)] + \mathrm{irr}^B_1(U^B) + \sum_{j=1}^{K'} \mathrm{irr}(\Delta_j)(U^A) + \sum_{j=1}^{K'} [\mathrm{irr}^B_{j+1}(U^B) - \mathrm{irr}^{\mathcal{G}}_j(U^A)] + D^{\mathbf{R}}(V) - (E^B - E^A)\,, \quad (11.3)$$

where $D^{\mathbf{R}}$ consists of the same three groups for the **R***-terms and $E^B - E^A$ is a number.*

Proof. The coupling of run B at its stopping step is $(g^B_{K'+1})^{-2} = (g^A_{K'})^{-2} + \delta_{K'}$ by (9.2), which gives the first two terms. The regular part of run B consists of its terms of the extra step, irr^B_1, and of those of the steps $j+1$, $1 \le j \le K'$. For the latter, $\mathrm{irr}^B_{j+1}(U^B) - \mathrm{irr}^A_j(U^A) = [\mathrm{irr}^B_{j+1}(U^B) - \mathrm{irr}^{\mathcal{G}}_j(U^A)] + [\mathrm{irr}^{\mathcal{G}}_j(U^A) - \mathrm{irr}^A_j(U^A)]$, and the second bracket is $\mathrm{irr}(\Delta_j)(U^A)$, because the subtractions at $U = 1$, of the isotropic part, and of N_j are linear and $\Pi^\Delta = \Pi^{\mathcal{G}} - \beta^A_j$, the coefficient of run A being absorbed into the coupling by (3.26). The same holds for the **R**-terms. *q.e.d.*

The groups of (11.3) are of four kinds: the coupling term, which is controlled by the recursion through $\delta_{K'}$; the difference of the two constrained minima of the action, a lattice artefact; the terms of the extra step and the old terms, which are bounded directly; and the differences of the recent terms of the same shared scale, which are the subject of the recursion. Since MU^B satisfies run A's constraint, the second group is

$$A_{\varepsilon/L}(U^B) - A_\varepsilon(U^A) = [A_\varepsilon(MU^B) - A_\varepsilon(U^A)] - b(U^B)\,, \qquad b(W) \coloneqq A_\varepsilon(MW) - A_{\varepsilon/L}(W)\,, \quad (11.4)$$

the gap between two configurations of run A's problem and the blocking artefact.

Proposition 11.3 (Classical Depth)**.** *Let $0 < \varepsilon_0 < 1/2$. There are $k_0(\varepsilon_0)$ and C_b such that for $K' \ge k_0(\varepsilon_0)$ and every V in the support of both terms ℓ_0,*

$$|A_{\varepsilon/L}(U^B(V)) - A_\varepsilon(U^A(V))| \le 2C_b L^{-K'} \varepsilon_1^2 N_q^4\,. \quad (11.5)$$

Proof. The blocking artefact is a sum over the coarse plaquettes of local Taylor remainders, each bounded by the regularity of its cube [7, Theorem 1, (9), (10), p. 279], which gives $|b(U^B)| \le C_b L^{-K'} \varepsilon_1^2 N_q^4$, and the gap is at most $C_{\mathrm{gap}}(\varepsilon_0) L^{-(2-2\varepsilon_0)K'} \varepsilon_1^2 N_q^4$, which is at most the same bound for $K' \ge k_0(\varepsilon_0)$; both are Supplement, Proposition S.13(a), (b). *q.e.d.*

11.2 The Direct Bounds

The terms of the extra step and the terms of large age are bounded without comparing them.

Lemma 11.4 (The Extra Step and the Old Tail)**.** *Let $\bar g$ be as in Theorem 1.5, and let $1 \le m \le K'$. On the support of both terms ℓ_0,*

$$|\mathrm{irr}^B_1(U^B)| + |\mathbf{R}^{B,(1)}(U^B) - \mathbf{R}^{B,(1)}(1)| \le (E_1 L^{-\beta K'} + R_1(bK - C_W)^{-\kappa_0/2}) N_q^4\,, \quad (11.6)$$

and the differences of the regular and **R***-parts of the shared scales $j \le K' - m$ satisfy*

$$\sum_{j \le K'-m} |\mathrm{irr}^B_{j+1}(U^B) - \mathrm{irr}^A_j(U^A)| \le \frac{2E_1 N_q^4 L^{-\beta m}}{1 - L^{-\beta}}\,, \qquad \sum_{j \le K'-m} |\Delta \mathbf{R}_j| \le C_{\mathbf{R}} N_q^4 (1+m)^{1-\kappa_0/2}\,, \quad (11.7)$$

with $\Delta\mathbf{R}_j$ the difference of the $\mathbf{R}$-parts of the shared scale j and $C_{\mathbf{R}}$ a constant of R_1, b, and κ_0.

Proof. On ℓ_0 the determining set is the stopping lattice itself, $\Gamma_{K'} = \mathbb{T}_{j_q}$ with N_q^4 points, so that in (3.31) at the generation j only the term $n = K'$ (respectively $K'+1$ for run B) occurs, and it gives $E_1 L^{-\beta(K'-j)} N_q^4$ for the regular part and $R_1 g_j^{\kappa_0} N_q^4$ for the $\mathbf{R}$-part. At the extra step $j = 1$ of run B this is $E_1 L^{-\beta K'} N_q^4$, and $(g_1^B)^{-2} \geq \bar{g}^{-2} + bK - C_W$ by (4.1) for the target $(K+1, \bar{g})$, which gives (11.6). For (11.7) the triangle inequality and the same bound for each run give $\sum_{j \leq K'-m} 2E_1 L^{-\beta(K'-j)} N_q^4$, and the geometric series gives the first bound. For the second, $g_j^{-2} \geq g_{K'}^{-2} + b(K'-j) - C_W$ by (4.1) gives $g_j^{\kappa_0} \leq C(1+K'-j)^{-\kappa_0/2}$ for $\bar{g}$ small, and $\sum_{i \geq m} (1+i)^{-\kappa_0/2} \leq C(1+m)^{1-\kappa_0/2}$ since $\kappa_0 > 2$. *q.e.d.*

We use (11.7) at the age $m_K := \lfloor K^{1/2} \rfloor$, where it is summable in K because $\kappa_0 \geq 13$ (Appendix A), and it uses no information on the differences of the terms. The terms of the shared scales $j > K' - m_K$ are compared, and this is where Bałaban's proof of (3.31) has to be run for a difference.

11.3 The Frozen Differences

The terms $\mathbf{E}^{A,(j)}$ and $\mathcal{G}_j$ are two families of terms of the generation j on the same lattices, with the localisation, analyticity, gauge invariance, and bound of Section 3.2, for $\mathcal{G}_j$ on a common polydisc of reduced radii $\alpha'_{i,m} := \alpha_{i,m}/(4B_3)$. For their difference we use the *class norm* d_j, the least number with $|\Delta_j(X, (\mathbf{U}, J(\mathbf{U})), z)| \leq d_j E_0 e^{-\kappa d_j(X)}$ on the evaluation sets of this polydisc, the sets of its pairs at which the constructions below read the terms, namely the minimisers of real and of complex data with their currents and the configurations which the steps build from them (Supplement, Proposition S.15), and $D_k := \sum_{j \leq k} L^{-\beta'(k-j)} d_j$ for its carried sum. Here

$$\beta' := \tfrac{1}{2} \min(\beta_H, 1 - \beta_E) \geq \tfrac{19}{40} , \qquad \theta_V := \frac{\beta'}{2(4+\beta')} , \tag{11.8}$$

with $\beta_H \geq 19/20$ the Hölder index in the regularity bounds of [8, (3.31), p. 276], which Bałaban states for every index less than one and which hold up to the index one for the minimisers [7, Theorem 1, (9), p. 279], and $\beta_E \leq 1/20$ the positive number of [9, (3.67), p. 283], which may be taken arbitrarily small at the cost of a larger constant. We write $d_j^{\mathbf{R}}$ for the class norm of the difference of the $\mathbf{R}$-terms of the shared scale j and $S_k := \sum_{j \leq k} d_j^{\mathbf{R}}$.

Proposition 11.5 (Theorem 2 for a Difference)**.** *Let the constants satisfy the conditions of Appendix C and $\bar{g}$ be small. There are constants E'_1, E_V, and C_α, independent of j, k, K, and the torus, such that for $z \in \Lambda_j^0 \cap (\Omega_n \setminus \Omega_{n+1})$ and every regular background U of the class of the step,*

$$|\mathrm{irr}(\Delta_j)(U)| \leq d_j \, C_\alpha E_0 (L^{j-n})^{4+\beta'} , \tag{11.9}$$

and on the all-small-field pair, at $n = K'$,

$$|\mathrm{irr}^B_{j+1}(U^B) - \mathrm{irr}^{\mathcal{G}}_j(U^A)| \leq E_V \Big[\Big(\frac{\hat{\sigma}_{K',j}}{r_{0,K'}} \Big)^{\theta_V} + (C_\Pi \varepsilon_1^2 e^{-\delta_c M_1 L^j})^{\theta_V} \Big] (L^{j-K'})^{4+\beta'/2} , \tag{11.10}$$

where $\hat{\sigma}_{K',j}$ is the distance between U^B and the lift $\iota(U^A)$ in the norm of the evaluation sets, bounded by Proposition 10.6 and the tail of the localised lift (Supplement, Proposition S.15(d)), $r_{0,K'}$ the radius of these sets, and $C_\Pi E_0 e^{-\delta_c M_1 L^j}$ the bound of $\Pi^{\mathcal{G}} - \beta^B_{j+1}$ and of the coefficients of N_j, with δ_c the rate of Supplement, Proposition S.12. The same holds for the $\mathbf{R}$-terms with $d_j^{\mathbf{R}}$ and without the factor of the age.

Proof. For (11.9) we run Bałaban's proof of [9, Theorem 2, (2.43)], which occupies [9, pp. 279–283] and rests on [8, Sects. 3–4], with the difference Δ_j in place of a single family; the proof is linear in the family, and it uses of the family only its localisation, its analyticity on the polydiscs of [8, (3.13)–(3.14)] with the radii of the step, and its bound, all of which Δ_j has with $d_j E_0$ in place of E_0, except the covariance which removes the plane-resolved moment, which is kept as N_j. The Hölder regularity of the background, of index β_H, gives the power $4+\beta'$ in place of 4. This is Supplement, Proposition S.15(a)–(b). For (11.10) the two values are the irrelevant parts of one term of run B at two configurations, U^B and the lift $\iota(U^A)$, each bounded by the age bound for one term; interpolating the configuration along a line of length $\hat\sigma_{K',j}$ in the norm of the evaluation sets and using Cauchy's estimate on a disc of radius $r_{0,K'}$, then interpolating between the bound of the age and the bound of the distance with the weight θ_V, gives (11.10); this is Supplement, Proposition S.15(c). *q.e.d.*

11.4 The Coefficients of the Two Runs

The recursion of the next subsection contains the differences δ_k of the couplings of the two runs, which are sums of differences of the coefficients of the flow. The coefficient of a step is the second moment of its regular term at the trivial configuration [9, (3.57)–(3.64), pp. 281–283], and one is tempted to bound the difference of two coefficients by the class norm of the difference of the two terms. This fails. The class norm d_j bounds Δ_j on polydiscs whose radii $\alpha'_{i,j}$ are proportional to $\hat g_j$, up to powers of its logarithm, and Cauchy's estimate for a second derivative on such a polydisc costs the inverse square of the radius,

$$|\Pi_j^\Delta| \le C \hat g_j^{-2}\, d_j E_0 \;. \tag{11.11}$$

Nothing better follows from d_j: the function $z \mapsto z^2$ has the supremum r^2 on the disc of radius r and the second coefficient one, and in the same way a multiple of the action $A(h_z,\cdot)$ with a coefficient of the order $\hat g_j^{-2} d_j E_0$ has a class norm of the order d_j. With (11.11) the recursion would not close, because the differences of the couplings would feed back into every step with a kernel which is not summable over the later steps. For a single run the coefficient is bounded all the same (Supplement, Proposition S.4), and the reason is a structure of the regular terms which we now use for the difference of two runs.

For a run and a complex number s let $\mathbf{E}^{(j)}(s)$ be the family of regular terms which the steps of the run produce when the **R**-terms which enter the regular part [9, p. 271] are multiplied by s, the couplings, the cutoffs, and all other terms being those of the run (Supplement, Lemma 3.4). At $s=1$ these are the regular terms of the run. At $s=0$ they are the terms of the small-field recursion of [8, 10] along the couplings of the run, and by Bałaban's Theorem 3 [8, p. 264] they are analytic and bounded on spaces whose radii α_0, α_1 do not depend on the coupling. For $|s| \le \varrho_{j-1}$, where

$$\varrho_k := 1 + \hat g_k^{-2} \;, \tag{11.12}$$

they satisfy the bounds of the regular terms on the spaces of the tube, and they differ from the terms at $s=0$ by at most $C_\rho g_{j-1}^{\kappa_0-4} E_0 e^{-\kappa d_j(X)}$, because the **R**-terms are of the order g^{κ_0} (Supplement, Lemma 3.5). Let $\mathcal{G}_j(s)$ be the family of run B at the lift, as in (11.2), and $\Delta_j(s) := \mathcal{G}_j(s) - \mathbf{E}^{A,(j)}(s)$, so that $\Delta_j(1) = \Delta_j$. Besides d_j we use two class norms:

- $d_j^\sharp$, the least number with $|\Delta_j(X,\cdot,z;s)| \le d_j^\sharp E_0 e^{-\kappa d_j(X)}$ on the evaluation sets of the polydisc of the tube, for every $|s| \le \varrho_{j-1}$;
- $d_j^\bullet$, the least number with $|\Delta_j(X,\cdot,z;0)| \le d_j^\bullet E_0 e^{-\kappa d_j(X)}$ on the evaluation sets of the polydisc of the fixed radii $\alpha_i^\bullet := \alpha_i/(4B_3)$.

Since $\varrho_{j-1} \geq 1$, $d_j \leq d_j^\sharp$. We write $D_k^\sharp$ and $D_k^\bullet$ for the carried sums, defined as D_k, and $u_j \coloneqq \max(d_j^\bullet, d_j^\sharp)$.

Lemma 11.6 (The Coefficients of the Two Runs)**.** *There is a constant C_β, which does not depend on j, K, the couplings, or the torus, such that at every shared scale j*

$$|\beta_{j+1}^B - \beta_j^A| \leq C_\beta u_j + C_\Pi E_0 e^{-\delta_c M_1 L^j} \ . \tag{11.13}$$

Proof. By the definition of Δ_j, $\Pi_j^\Delta = \Pi_j^{\mathcal{G}} - \beta_j^A$, so that $\beta_{j+1}^B - \beta_j^A = \Pi_j^\Delta - (\Pi_j^{\mathcal{G}} - \beta_{j+1}^B)$, and the bracket, the isotropic part of the moment defect of the lifted family, is at most $C_\Pi E_0 e^{-\delta_c M_1 L^j}$ (Supplement, Proposition S.15(d)). The coefficient is linear in the family, and we split it as

$$\Pi_j^\Delta = \Pi(\Delta_j(0)) + \Pi(\Delta_j(1) - \Delta_j(0)) \ , \tag{11.14}$$

where $\Pi(F)$ denotes the isotropic coefficient of a family F. The family $\Delta_j(0)$ is bounded by $d_j^\bullet E_0 e^{-\kappa d_j(X)}$ on a polydisc about the trivial configuration whose radius is a fixed multiple of $\min(\alpha_0^\bullet, \alpha_1^\bullet)$. Cauchy's estimate for its second moment costs therefore a constant which does not depend on the coupling, and $|\Pi(\Delta_j(0))| \leq C_\beta^\bullet d_j^\bullet$. For the second term we use the analyticity in s. At every point of the evaluation sets of the tube the function $s \mapsto \Delta_j(X; s)$ is bounded by $d_j^\sharp E_0 e^{-\kappa d_j(X)}$ on the disc of radius ϱ_{j-1}, and for $s \in [0,1]$ the disc of the radius $\hat{g}_{j-1}^{-2}$ about s lies in it, so that Cauchy's estimate for the derivative in s gives

$$|\Delta_j(X;1) - \Delta_j(X;0)| \leq \hat{g}_{j-1}^2 \, d_j^\sharp E_0 e^{-\kappa d_j(X)} \ . \tag{11.15}$$

The second moment of this family on the polydisc of the tube costs a constant times $\hat{g}_j^{-2}$, as in (11.11), and $\hat{g}_{j-1}^2 \hat{g}_j^{-2} \leq 1$ because $\hat{g}_j$ increases with j; hence $|\Pi(\Delta_j(1) - \Delta_j(0))| \leq C_\beta^\sharp d_j^\sharp$. This is (11.13) with $C_\beta \coloneqq 2\max(C_\beta^\bullet, C_\beta^\sharp)$, and the constants are those of Supplement, Proposition S.15(e). *q.e.d.*

The lemma replaces the radius of the tube by two radii which the construction provides: that of Bałaban's small-field spaces, for the part of the regular terms which does not involve the **R**-terms, and that of the disc of s, for the part which does. The second is available because the **R**-terms are small: a disc of the radius $\hat{g}^{-2}$ costs two of the κ_0 powers of the coupling which they carry. The price is that the two norms $d^\bullet$ and $d^\sharp$ have to be carried through the recursion in the place of d, and that the **R**-terms enter the recursion of $d^\sharp$ with the factor ϱ_k.

11.5 The Generated Difference of One Step

At a shared scale k the two runs apply the same step to inputs on the same lattice, with the same block data and the same fluctuation variable, and the step's output is the logarithm of a fluctuation integral. The two integrals differ by the interaction, which collects the fluctuation parts of the frozen terms and of the fine actions, by the Gaussian form, which is built from the propagators of the run, and by the cutoff of the fluctuation field. Let $\mathbf{P}^A$ and $\mathbf{P}^B$ be the two interactions, $\Delta_A^{(k)}$ and $\Delta_A^{(k)} + \delta\Delta^{(k)}$ the two quadratic forms of the fluctuation field, and for $t \in [0,1]$ put $\mathbf{P}_t \coloneqq \mathbf{P}^A + t(\mathbf{P}^B - \mathbf{P}^A)$, $\Delta_t \coloneqq \Delta_A^{(k)} + t\delta\Delta^{(k)}$, and

$$Z_t \coloneqq \int \chi^{A,(k)} \exp\Big[-\frac{1}{2}\langle A, C^* \Delta_t C A\rangle + \mathbf{P}_t\Big] \, \mathrm{d}A \ , \tag{11.16}$$

with run A's cutoff in both, so that run B's integral is Z_1 plus the shell term of Lemma 10.5.

Lemma 11.7 (Interpolation Identity)**.** *Let k be a shared scale at which the number r_k of* (11.18) *satisfies $C_{\mathrm{cov}} r_k \le \alpha_4^G$, with α_4^G the room of the Gaussian integral of Supplement, Proposition S.16(a). On the class of the scale $k+1$ and for every $t \in [0,1]$ the integral Z_t does not vanish, and with $\langle\cdot\rangle_t$ the expectation of the normalised integrand of Z_t and $\log Z_t$ the branch which is continuous in t,*

$$\log Z_1 - \log Z_0 = \int_0^1 \Big[\langle \mathbf{P}^B - \mathbf{P}^A \rangle_t - \frac{1}{2} \langle\langle A, C^* \delta\Delta^{(k)} C A \rangle\rangle_t \Big] \, \mathrm{d}t \, . \tag{11.17}$$

For every t the normalised integrand is of the kind which Bałaban's cluster expansion of the step treats: its interaction $\mathbf{P}_t$ satisfies the inductive bounds, and its quadratic form is run A's plus the form $t\,\delta\Delta^{(k)}$, which is a small quadratic term of the exponent.

Proof. We begin with the last statement, on which the others rest. The interaction $\mathbf{P}_t$ is a convex combination of two interactions which satisfy the inductive bounds [8, (1.18), p. 263], [9, (2.31), (2.42)], and these bounds define convex sets. For the quadratic form we do not argue through its positivity, which says nothing at the complex backgrounds of the class. We write $\Delta_t = \Delta^A + t\,\delta\Delta^{(k)}$ and treat the second term as a part of the exponent: by Supplement, Proposition S.17, its kernel is at most r_k times the bound of the kernel of the form itself, with exponential decay, so that $|\frac{1}{2} t \langle A, C^* \delta\Delta^{(k)} C A\rangle| \le \frac{1}{2} C_{\mathrm{cov}} r_k \sum_b |A(b)|^2$, and a quadratic term with a coefficient within the room α_4^G is absorbed into the Gaussian integral of run A by the steps [10, (2.14)–(2.25), pp. 15–17], as in Supplement, Proposition S.16(a). The integrand of Z_t is thus one to which Bałaban's expansion applies, with run A's covariance, at real and at complex backgrounds of the class.

In this expansion the integral which normalises the expectation, taken with the Gaussian measure of run A, "is represented as $\exp \tilde{\mathbf{E}}^{(k+1)}(\Lambda_{k+1})$", with $\tilde{\mathbf{E}}^{(k+1)}$ the convergent sum of analytic terms of the exponentiated cluster expansion [9, p. 277]; since the Gaussian normalisation of run A is the inverse square root of the determinant of an invertible form, $Z_t \neq 0$. The integrand is differentiable in t, with the derivative equal to the integrand times $\mathbf{P}^B - \mathbf{P}^A - \frac{1}{2}\langle A, C^* \delta\Delta^{(k)} C A\rangle$, which on the support of the cutoff is the sum of a bounded function and a quadratic form in A and is dominated, uniformly in t, by an integrable function, because on the support of the cutoff $|A|$ is bounded. So Z_t is continuously differentiable, $\partial_t \log Z_t = \partial_t Z_t / Z_t$ is the expectation of the derivative of the exponent, and integrating over t gives (11.17), the device by which Bałaban separates the terms of a step [9, (3.26)–(3.27), p. 271]. *q.e.d.*

At the steps at which $C_{\mathrm{cov}} r_k > \alpha_4^G$ the lemma is not used. By (11.18) below these are the steps $k \le k_r$ with a number k_r which does not depend on K, and there each of the two outputs satisfies the bounds of its class, so that the class norm of their difference is at most 2; this is the bound (11.19) below with a depth source $e_{k+1} \ge 2$, and the constant C_e of the depth sources is chosen at least $2\tilde{\theta}^{-k_r-1}$.

The right side of (11.17) is a cluster expansion with one marked activity, the piece of $\mathbf{P}^B - \mathbf{P}^A$ or of the quadratic insertion on a domain, and the localised terms of the difference of the step's outputs are the localised terms of this marked expansion. The marked activity consists of the fluctuation parts of the frozen differences of the generations $j \le k$, of size $D_k + S_k$ in the class norm; of the coupling, through the factors g_k in the backgrounds and the coefficients g_k^{n-2} of the terms of order $n \ge 3$, whose two values differ relatively by at most $g_k^2 |\delta_k|$; of the depth sources e_{k+1}, which are the differences of the two runs in the objects of the fine lattices; and of the shell of the cutoff. The depth sources need a word. One of them is the difference $\delta\Delta^{(k)}$ of the quadratic forms. The others are the differences of the parts of the

third and higher orders in the fluctuation field of the two actions, of the current terms, and of the configurations at which the frozen terms are read. The classical depth of Proposition 11.3 is not among them: it bounds the difference of the two actions at the two minimisers, which enters an exponent with the factor g_k^{-2}, and this factor is bounded at the stopping scale only. At the earlier scales the two actions are compared through their expansions in the fluctuation field, in which the terms of the orders zero and one are absent and the term of the second order is the quadratic form; the remaining terms are at most $g_k p_0(\hat{g}_k)^3$ per point, uniformly in the regularity of the background, and their two-run differences are smaller by a power of ε/a_k (Supplement, Proposition S.16(b)). The quadratic forms differ by at most

$$r_k \le C_{\delta\Delta}[L^{-\gamma_H(k-1)} + L^{-(1-\varepsilon_0)k}] , \qquad \gamma_H := \frac{\beta\gamma_w}{1+\beta} , \tag{11.18}$$

times the bound of the form itself, by Theorem 9.5 and Supplement, Proposition S.17, and the shell of the cutoff contributes at most $C|\delta_k| g_k^2 e^{-cp_0(\hat{g}_k)^2}$ per point, by Lemma 10.5 and the Gaussian tail beyond the radius ϑ_k/g_k. All of these sources are bounded at backgrounds whose regularity is measured by constants and not by the coupling, and they satisfy $e_{k+1} \le C_e \tilde{\theta}^{k+1}$ with a constant which does not depend on K. The marked-cluster bound of Supplement, Proposition S.16, then bounds the generated difference per point of the scale $k+1$ by $\varepsilon_k^*(D_k + S_k) + C_g \varepsilon_k^* g_k^2 |\delta_k| + e_{k+1}$. The argument reads the regular terms of the two runs only through their localisation, their analyticity, and their bounds, and it applies in the same way to the two families of Section 11.4. For the families $\Delta(s)$ on the spaces of the tube, uniformly in $|s| \le \varrho_k$, the couplings, the cutoffs, and the propagators are those of the two runs, so that δ_k and e_{k+1} do not depend on s, and the only change is that the **R**-terms which enter the regular part are multiplied by s:

$$d_{k+1}^\sharp \le \varepsilon_k^*(D_k^\sharp + \varrho_k S_k) + C_g \varepsilon_k^* g_k^2 |\delta_k| + e_{k+1} , \qquad \varepsilon_k^* := C_X C_{\mathrm{pt}} C_{\mathrm{KP}} L^4 C_\alpha^{\mathbf{A}} [C_T \mu_k + 4a_k] . \tag{11.19}$$

For the families at $s = 0$ on the spaces of the fixed radii no **R**-term enters, the expansion of the step is that of [10, Sect. 2] on these spaces, and the sources are the same differences of propagators, minimisers, and Gaussian forms, which Sections 9 and 10 bound at backgrounds whose regularity is measured by constants and not by the coupling:

$$d_{k+1}^\bullet \le \varepsilon_k^* D_k^\bullet + C_g \varepsilon_k^* g_k^2 |\delta_k| + e_{k+1} . \tag{11.20}$$

In both inequalities ε_k^* and e_{k+1} denote the larger of the coefficients and of the sources of the two settings (Supplement, Proposition S.16(b)). Here μ_k is the square of the ratio of the cutoff's supremum to the radius of analyticity of the frozen terms, a negative power of $\log g_k^{-2}$, a_k the bound of the interaction's activities, a positive power of g_k times a power of $\log g_k^{-2}$, and the other constants are those of Supplement, Proposition S.16, independent of k and K. The coefficient of the flow is the marginal part of the step's output, and the difference of the coefficients of the two runs is bounded by Lemma 11.6. From $\delta_K = 0$, which holds because both runs are pinned at $g_K^A = g_{K+1}^B = \bar{g}$, and from $\delta_j = \sum_{i>j} (\beta_{i+1}^B - \beta_i^A)$, the coupling differences are therefore $|\delta_j| \le C_\beta \sum_{i>j} u_i + 2C_\Pi E_0 e^{-\delta_c M_1 L^{j+1}}$, and the second term, which decays faster than any geometric sequence, is counted among the depth sources e_i of (11.19); the steps between the stopping step and the unit scale, which both runs perform for the coupling prescription, obey the same recursion. The **R**-terms have no decay in their age beyond the volume factor [9, p. 283], and their recursion, Supplement, Proposition S.16(c), together with the renormalisation quotients of the **R**-operation, in which Theorem 10.8 enters through $h_n := C_h \tilde{\theta}^n$, gives

$$S_k \le C_S \Big[\sum_{j \le k} c_j (\mathcal{D}'_{[j]} + D_{j-1}) + \sum_{j \le k} c_j^B \mathcal{D}_j^{*\prime} + \sum_{j \le k} c_j^V \mathcal{D}_j^{V\prime} + C_\nu w_K \Big] , \tag{11.21}$$

with c_j, c_j^B, c_j^V constants times $(100MR_j)^4 g_j^{\kappa_0}$, whose sum over j is bounded uniformly in K for $\kappa_0 > 2$, $\mathcal{D}'_{[j]}$, $\mathcal{D}_j^{*\prime}$, $\mathcal{D}_j^{V\prime}$ the maxima over the history of the integrands' differences per point, built from the d_i, the $|\delta_i|$, the e_i, and the h_i (Supplement, Proposition S.16(d)), and w_K of (11.23) below.

We now solve the recursion. Put

$$\theta \coloneqq L^{-\beta'} + \varepsilon^* , \qquad \tilde\theta \coloneqq \max\Big(\frac{1+\theta}{2}, L^{-\gamma_H}, L^{-(1-2\varepsilon_0)/3}, L^{-(1-\varepsilon_0)/4}, L^{-\theta_V/8}\Big) , \tag{11.22}$$

where $\varepsilon^* \coloneqq \sup_{g \le (1+\beta_0)\gamma} \varepsilon^*(g)$ is the supremum of the coefficient of (11.19) as a function of the coupling, which bounds ε_k^* at every shared scale because $g_k \le (1+\beta_0)\bar g$ by (3.9); and put

$$\ell_K \coloneqq (\log(bK+\bar g^{-2}))^{4r+2p_0} , \qquad \mu_K \coloneqq K^{3-\kappa_0/2}\ell_K , \qquad w_K \coloneqq (K+1)^2(\tilde\theta^{K'/8}+2^{-K'/16})+\mu_K . \tag{11.23}$$

Proposition 11.8 (Solution of the Recursion)**.** *Let γ satisfy the conditions (T1) of* (C.2)*. Then $\tilde\theta < 1$, and for every shared scale i*

$$d_i \le u_i \le 2C_s(i\tilde\theta^i + w_K) , \qquad S_i \le C_S'' w_K , \qquad |\delta_i| \le 2C_sC_\beta[2(1-\tilde\theta)^{-2}(i+1)\tilde\theta^{i+1} + (K+1-i)w_K] , \tag{11.24}$$

with constants C_s and C_S'' which do not depend on K. In particular, at the stopping step,

$$D_{K'} + S_{K'} + C\varepsilon_{K'}^2|\delta_{K'}| + e_{K'} \le C_a[(K+1)\tilde\theta^{K'} + (j_q+2)w_K] , \tag{11.25}$$

with a constant C_a which does not depend on K and depends on j_q at most as $L^{C\varepsilon_0 j_q}$ times a polynomial.

Proof. In (11.21) the differences of the regular terms are those of the two runs, which are bounded by $d_i \le u_i$. Substituting (11.21) and the bound on $|\delta_i|$ into (11.19) and (11.20) gives $u \le \mathcal{T}u + s$ for $u_i = \max(d_i^\bullet, d_i^\sharp)$, with an operator $\mathcal{T}$ on sequences which is positive, monotone, and positively homogeneous, and a source s built from the e_i, the h_i, and $C_\nu w_K$, with $s_1 \coloneqq 2$ for the first step, at which both runs' outputs satisfy the bound of their class. Put $m_i \coloneqq i\tilde\theta^i + w_K$ and $\|u\|_m \coloneqq \sup_i u_i/m_i$. Since $\tilde\theta > L^{-\beta'}$, $\sum_{j\le i} L^{-\beta'(i-j)}m_j \le (1 - L^{-\beta'}/\tilde\theta)^{-1}m_i$, and $m_i \le \tilde\theta^{-1}m_{i+1}$. The factor ϱ_i of (11.19) is paid by the **R**-terms: for $j \le i$, $\varrho_i \le 1 + \hat g_j^{-2} \le \frac{5}{2}g_j^{-2}$ by Lemma 9.1(iii), so that $\varrho_i c_j$ is at most a constant times $(100MR_j)^4 g_j^{\kappa_0-2}$, whose sum over j is still bounded uniformly in K, while the direct bounds of the fine scales in the source are multiplied by $\varrho_i \le (1 + \bar g^{-2} + b' + C_W)(K+1)$, which is the reason for the powers of K in (11.23). The sums over the fine scales are paid by the factors $\varrho_i c_j$, which for $j \le K/2$ are at most $C(bK/2)^{1-\kappa_0/2}\ell_K$, and the sums over the coarse scales by $\tilde\theta^j$, which together give at most a constant times $\mu_K \le w_K$; and the coupling differences enter through $g_i^2\sum_{l>i} m_l \le (2(1-\tilde\theta)^{-2} + b^{-1})m_{i+1}$, which uses $g_i^2(K-i) \le b^{-1}$ by (4.1). Hence $\|\mathcal{T}\|_m \le \varepsilon^* C_{\mathcal{T}} \le \frac14$ by the first condition of (C.2), and since the system is finite, $\|u\|_m < \infty$ and $\|u\|_m \le \|\mathcal{T}\|_m\|u\|_m + \|s\|_m \le \frac14\|u\|_m + C_s$. This gives the first bound of (11.24), and the others follow from (11.21) and the bound on $|\delta_i|$. The first condition implies $\varepsilon^* \le (1 - L^{-\beta'})/4$, because $C_{\mathcal{T}} \ge (1 - L^{-\beta'})^{-1}$, so that $(1+\theta)/2 \le (3 + L^{-\beta'})/4 < 1$, and the other entries of (11.22) are less than one. At $i = K'$, $K'\tilde\theta^{K'} \le (K+1)\tilde\theta^{K'}$ and $(K+1-K')w_K = (j_q+1)w_K$, which gives (11.25); the details, the constants, and the depth sources $e_i \le C_e\tilde\theta^i$ are Supplement, Proposition S.16(e). *q.e.d.*

The bound (11.25) is summable in K, because $(K+1)\tilde\theta^{K'}$ and $(j_q+2)(K+1)^2(\tilde\theta^{K'/8} + 2^{-K'/16})$ decay exponentially at fixed j_q, and $\mu_K = K^{3-\kappa_0/2}\ell_K$ with $\kappa_0 \ge 13$. The second

condition of (C.2) is the one which allows the recent **R**-terms, which are carried through $N'_k \le N'(\hat{g}_k)$ steps without contraction [9, pp. 270–271], to be summed by a window lemma; the memory N' grows as a double logarithm of the inverse coupling, while $\hat{\varepsilon}^*$ decays as a negative power of its logarithm. The same recursion holds for the deviation insertions of the dressed loops, which are produced by the same fluctuation integral with the loop's words as further marked activities, with their class norms d_j^{ins} in place of d_j and an additional term $\varepsilon_k^Q D_k^{\mathrm{ins}}$ from the old vertices of the groups, which enter as numbers, with $\varepsilon^Q(g)$ of order g^2 times a power of $\log g^{-2}$ (Supplement, Proposition S.16(f)). The coefficient ε^Q depends on the representations, but the recursion of the insertions runs over the steps $k \le K'$ only, at which $g_k \le \gamma_q$ by (3.46); under the condition

$$\varepsilon^Q(g) C_{\mathcal{T}} \le \frac{1}{4} \tag{11.26}$$

for every $g \le \gamma_q$, which holds for γ_q small, depending on $c_{\mathcal{C}}$, the operator of this recursion has the norm at most $(\varepsilon^* + \sup_{g \le \gamma_q} \varepsilon^Q(g)) C_{\mathcal{T}} \le \frac{1}{2}$, by the first condition of (C.2), and its solution is $d_i^{\mathrm{ins}} \le 2 C_{\mathrm{ins}}^* m_i$.

11.6 The All-Small-Field Pair

On the common support of the two terms ℓ_0 the groups of (11.3), less the number $E^B - E^A$, are bounded at every V by

$$X_K \coloneqq X_K^{\mathrm{dir}} + N_q^4 C_a [(K+1)\tilde{\theta}^{K'} + (j_q+2) w_K] + X_K^{\mathrm{var}} + X_K^{\mathbf{R}} + X_K^{\mathrm{cl}} \,, \qquad x_K \coloneqq e^{X_K} - 1 \,, \tag{11.27}$$

where X_K^{dir} is the sum of the right sides of (11.6) and (11.7) at $m = m_K$, the second term is (11.25) summed over the N_q^4 points of the stopping lattice, X_K^{var} is the sum over j of (11.10), $X_K^{\mathbf{R}}$ the bound of the corresponding group of the **R**-terms of the ages less than m_K, and $X_K^{\mathrm{cl}} \coloneqq 2 C_b (g_{K'}^A)^{-2} L^{-K'} \varepsilon_1^2 N_q^4$ by (11.5), together with its counterpart on the young pairs (Supplement, Proposition S.13(d)). Each of these is summable in K: the variation group by $\hat{\sigma}_{K',j} \le C(\varepsilon/a_{K'})^{(1-\varepsilon_0)/3} \varepsilon_1$ and the factor of the age, and the coupling factor $(g_{K'}^A)^{-2} \le \bar{g}^{-2} + b' j_q + C_W$ of the classical group is bounded. The loop's terms are compared on the same support in four parts.

The leading words. By Proposition 10.6(i), $U^A = e^{i\varepsilon A'} M U^B$ in a gauge, with $\varepsilon |A'(c)| \le 4 C_G(\varepsilon_0)(\varepsilon/a_{K'})^{2-\varepsilon_0} \varepsilon_1$ on every bond c of the fine lattice of run A, and by (8.10) $MU^B(c) = e^{iY_c} W(c)$, with $W(c)$ the transport of U^B along the L fine bonds of c and $|Y_c| \le C_Y(\varepsilon/a_{K'})^2 \varepsilon_1$ by Lemma 8.2. So the holonomies of C_i in the two backgrounds are $\prod_c W(c)$ and, up to a gauge transformation which leaves the trace unchanged, $\prod_c e^{i\varepsilon A'(c)} e^{iY_c} W(c)$. Telescoping inside the normalised trace, with $\|\pi(e^{iX}) - 1\| \le c_\pi \|X\| \le c_\pi c_h |X|$ for the unitary representation π (Section 8.2), and summing over the $P_{K'}(C_i) L^{K'}$ fine bonds of C_i,

$$\Big| \prod_i f^{(i),B} - \prod_i f^{(i),A} \Big| \le \lambda_K \coloneqq \sum_i c_{\pi_i} c_h (4 C_G(\varepsilon_0) + C_Y) \varepsilon_1 P_{K'}(C_i) L^{-(1-\varepsilon_0)K'} \,, \tag{11.28}$$

because every factor has modulus at most one and $L^{K'}(\varepsilon/a_{K'})^{2-\varepsilon_0} = L^{-(1-\varepsilon_0)K'}$. The words with insertions are covariant at the slots of their insertions, and the same telescoping with the insertions held fixed changes the normalised dressed loop by at most $\lambda_K \prod_b (1 + \|Y_b\|) \le \lambda_K \mathfrak{b}(\mathcal{C})$.

The deviation insertions. By the recursion for the insertions their difference changes the normalised dressed loop by at most $\mathfrak{b}(\mathcal{C}) X_K$.

The wrap insertions and the interaction weights. The insertions which wrap around the torus have no deviation bound and are not contracted by the recursion, but their mass decays as $e^{-cN_{\min}^{(j)}/M}$ in the smallest period of the torus at their scale, $N_{\min}^{(j)} \geq c(T/u)L^{K'-j+j_q}$. Those of age at least m_K^w, the least m with $a_w L^m \geq 3\log(K+2)$, are not compared and have total mass at most $\tau_K^w := 2A_w \max(L, 3L\log(K+2)/a_w)(K+2)^{-3}$ in each run; the younger ones are compared at their birth, by the interpolation of the constituents of their step described below for the loop constant, and then carried, with a difference which grows by at most $2L$ per step. This gives a change of the normalised dressed loop of at most $x_K^w \mathfrak{b}(\mathcal{C})$, with x_K^w summable in K; the interaction weights $Q_{K'}(\mathfrak{B})$ of the families, cluster activities of the reference gas of the stopping step, are compared in the same way, with a change at most $x_K^Q \mathfrak{b}(\mathcal{C})$. Both are Supplement, Proposition S.16(g).

The loop constant. The constant of the dressed ensemble is $E_{\text{const}} = \sum_{j \leq K'} E_j$, where $E_j := E_{\text{const},j} - E_{\text{const},j-1}$ collects, by (7.30), the flat remainders of the loops and the cross term of the step j, and at an **R**-step also the remainders $E_j^{\circ,\mathbf{R}}(C_i, T)$ and the cross constant $E_j^{\mathbf{R}\times}$. We index these terms by the age $m := K' + 1 - j$ in run A and $m := K' + 2 - j$ in run B, so that the two terms of the same shared scale have the same m. Each E_j is at most $\epsilon_j(\mathcal{C}, T)$ of (7.31), and each of the five parts of $\epsilon_j(\mathcal{C}, T)$ is at most $A_E L^{m/2} e^{-a_E L^{m/2}}$, with constants A_E and a_E which do not depend on K. Indeed, in the first three parts the lengths are those of the stopping step multiplied by L^m, as in the proof of Corollary 7.11, and $te^{-ct} \leq (2/c)e^{-ct/2}$ for $t \geq 0$. In the two parts of the **R**-steps the lengths $r_j(C_i)$, $\iota_{T,j}$, and a_j are at least $a_j/2$, by (3.45) and the proof of Lemma 7.5, and $a_j = a_{K'} L^{m-1} \geq L^{m+2} M R_{K'}$; since $R_{K'} \geq (\log \hat{g}_{K'}^{-2})^r$, $R_j < L(\log \hat{g}_j^{-2})^r$ by Lemma 3.4(i), and $\log \hat{g}_j^{-2} = \log(\hat{g}_{K'}^{-2} + b'(m-1)) \leq (1+\log(1+b'm)) \log \hat{g}_{K'}^{-2}$, we get $a_j/(MR_j) \geq L^{m+1}(1+\log(1+b^*m))^{-r} \geq c_L L^{m/2}$, with $c_L > 0$ a constant of L, r, and b^*. Since moreover $MR_j e^{-p_0(g_j)/4}$ is bounded uniformly in j, and $P_j(C_i) = P_q(C_i)a_j$, the bound $te^{-ct} \leq (2/c)e^{-ct/2}$ with $t = a_j/(MR_j)$ gives the claim for these parts as well. The terms of age $m \geq m_K^E$, the least m with $a_E L^{m/2} \geq 2\log(K+2)$, are not compared; their sum is at most $\tau_K^E := 2A_E \max(L^{1/2}, 2L^{1/2}\log(K+2)/a_E)(K+2)^{-2}$ in each run, and for $K \geq K_1$ the extra step of run B is among them. For the younger terms, which are produced at **R**-steps of both runs, we interpolate: every constituent x of the step, an activity of the effective action, a marked activity of the loop, a factor $V(Y)$, or the operation of a young component, is replaced by $x^\varsigma := x^A + \varsigma(x^B - x^A)$, $\varsigma \in \mathbb{C}$, after the old components, those born at a scale at most $K' - \lfloor K'/2 \rfloor$, have been removed by multiplying their operations by a parameter $\mathfrak{z}$ with $|\mathfrak{z}| \leq e^{C_P K'/4}$. This is possible because the bound (8.33) holds at every step and for every component separately: an old component has lived through at least $K'/4$ steps at a young step, so that its operation has a total mass at most $e^{-C_P K'/4}$ times the bound under which the expansion of the step converges, and the multiplication by $\mathfrak{z}$ keeps each activity within that bound. The relative differences of the constituents are at most the number ρ_K^E defined in the proof of Supplement, Proposition S.16(g)–(h), which is at most a constant times $(K+1)w_K + \ell_K L^{-\gamma_H K'/4}$ plus a multiple of x_K^w, so that it tends to zero and is summable in K after multiplication by any power of $\log(K+2)$; for $|\varsigma| \leq 1/(2\rho_K^E)$ the constituents are at most twice their bounds, the cluster expansion of the step converges, and the per-step term is analytic in ς with the bound $A^{E,2} L^{m/2} e^{-a^{E,2} L^{m/2}}$ at the enlarged amplitudes and the lowered decay; for the cross constant this holds for each of the cluster expansions, of the gas linked to $\mathcal{C}$ and of the gases linked to the single loops C_i, whose constants it subtracts. Cauchy's estimate on the disc of radius $1/(2\rho_K^E) \geq 2$ gives

$$|E_j(1) - E_j(0)| \leq 4\rho_K^E A^{E,2} L^{m/2} e^{-a^{E,2} L^{m/2}} , \tag{11.29}$$

and the old components cost at most $2e^{-C_P K'/4} A_E L^{m/2} e^{-a_E L^{m/2}}$ in each run, by Cauchy's estimate in $\mathfrak{z}$. Summing over m with $\sum_{m \geq 1} \Lambda^m e^{-a\Lambda^m} \leq e^{-a}(1 + 6/a)$ for $\Lambda \geq 2$, we obtain

$|E^B_{\text{const}} - E^A_{\text{const}}| \le \Delta E_K$, with $\Delta E_K \le \min(2\tau^E_K + \Delta^E_K, 2B_E(\mathcal{C}))$ summable in K. The details are Supplement, Proposition S.16(h).

Lemma 11.9 (The All-Small-Field Pair)**.** *For $K \ge K_1$, with run A's constant on both runs, the terms ℓ_0 change $W_{K+1}(\mathcal{C}) - W_K(\mathcal{C})$ by at most $(3x_K + \lambda_K + x^w_K + x^Q_K)\, B(\mathcal{C})$ on their common support, and by at most $b^{(S)}_K B(\mathcal{C})$ through the shells of the stopping-scale test functions.*

Proof. On the common support, run B's density is $e^{D(V)+E^B-E^A}$ times run A's, and $|D(V) + E^B - E^A| \le X_K$, so that $\int |\varrho^B_{\ell_0} - \varrho^A_{\ell_0}|\, \mathrm{d}V \le x_K Z^A_{\ell_0}$, where the number $E^B - E^A$ cancels between the numerator and the normalisation. The four parts above give $\int |\widehat{\mathcal{W}}^B \varrho^B_{\ell_0} - \widehat{\mathcal{W}}^A \varrho^A_{\ell_0}|\, \mathrm{d}V \le \mathfrak{b}(\mathcal{C})(x_K + X_K + \lambda_K + x^w_K + x^Q_K) Z^A_{\ell_0}$. As in (10.5), $N'/Z' - N/Z = (\Delta N - W'\Delta Z)/Z$ with $|W'| \le B(\mathcal{C})$ and $Z \ge Z^A_{\ell_0}$, and $X_K \le x_K$. The shells are Proposition 10.9. *q.e.d.*

11.7 The Pairs with Large Fields

For a young matched pair Theorem 10.8 compares the ingredients (β) and (γ); the ingredients (α) are compared by (11.24) per point of a history at a scale $i \ge K' - m$: the frozen exponents and the depth terms by $C_a[(i+1)\tilde{\theta}^i + w_K]$; the coupling term of the action by $C\varepsilon_i^2 |\delta_i| \le |\delta_i|$; the function [11, (1.6), p. 178], $z^{-1} \exp[-g_i^{-2}(1 - \operatorname{Re}\operatorname{tr} V_i)]$, relatively by at most $e^{4|\delta_i|} - 1 \le 10|\delta_i|$, because $0 \le 1 - \operatorname{Re}\operatorname{tr} V \le 2$ and the normalisation z has the common cutoff ε_i; the Gaussian form by $Cr_i |A_i(b)|^2$ per variable, by (11.18), which on the cutoff is at most $Cr_i p_0(\hat{g}_i)^2$ and where a large-field function leaves A_i unbounded is absorbed by the positivity of the form; and the localised normalisation constants, of order $\log g_i^{-2}$ per variable [12, p. 380], whose dependence on the coupling is the term $-\frac{1}{2} \dim G \log g_i^{-2}$ per variable of the normalisation of a Gaussian integral over a variable scaled by g_i, by $\frac{1}{2} \dim G |\log(1 + g_i^2 \delta_i)| \le |\delta_i|$. Collecting, with C_M the number of ingredients per point,

$$\begin{aligned} s'_K \coloneqq {} & 2C_a[(K'+1)\tilde{\theta}^{K'-m} + w_K] + 10 \sup_{i \ge K'-m} |\delta_i| \\ & + 4C(1 + C_{\text{cov}}) C_{\delta\Delta} \ell_K [L^{-\gamma_H (K'-m-1)} + L^{-(1-\varepsilon_0)(K'-m)}] \,, \end{aligned} \tag{11.30}$$

$s_K \coloneqq s'_K + L^4 C_h \tilde{\theta}^{K'/2}$, and $\omega^s(X) \coloneqq (1 + C_M P(X)) e^{C_M s_K P(X)}$, which is at most $e^{C_P P(X)}$ when $C_P \ge 2C_M$ and $s_K \le 1$.

Proposition 11.10 (The Young Pairs)**.** *Let k_* be the least integer with $k_* \ge k_{\text{LF}}$ such that $s'_K \le \frac{1}{2}$ and $L^4 C_h \tilde{\theta}^{K'/2} \le \frac{1}{2}$ for $K' \ge k_*$. For $K' \ge k_*$ the young matched pairs change $W_{K+1}(\mathcal{C}) - W_K(\mathcal{C})$ by at most $\bar{\delta}_{\text{LF}}(5s_K + 3x_K + s^W_K) B(\mathcal{C})$, where s^W_K, summable in K, bounds the change of the dressed loops near the large-field regions (Supplement, Proposition S.14(f)). The threshold k_* exists, and s_K is summable in K.*

Proof. By (10.6), (10.13), and the bounds on the ingredients (α), each of whose contributions enters per point of a history and is counted by $P(X)$, with the small-field exponent on the last domain bounded by X_K as for the all-small-field pair,

$$\begin{aligned} \int |\varrho^{B'}[\tau^+] - \varrho^A[\tau]|\, \mathrm{d}V &\le (s_K + x_K) \prod_X \omega^s(X)\, M^A_\tau \,, \\ \int (\Pi_B - \Pi_A)_+ \varrho_B\, \mathrm{d}V &\le s_K \prod_X \omega^s(X)\, M^B_{\tau^+} \,, \end{aligned} \tag{11.31}$$

where $\varrho^{B'} \coloneqq \min(\Pi_A, \Pi_B)\varrho_B$ and we used $(e^a - 1) + (e^b - 1) \le e^{a+b} - 1$, $e^{\sum \phi_X} - 1 \le (\sum \phi_X) e^{\sum \phi_X}$, and $\sum_X C_M P(X) \le \prod_X (1 + C_M P(X))$. Since $Z_\ell \le M_\ell$ and the weights $\omega^s(X)$

are admissible in Proposition 8.4, by the proof of that proposition the sums over the young pairs are at most $\delta_{\mathrm{LF}}(s_K + x_K)Z^A_{\ell_0}$ and $\delta_{\mathrm{LF}} s_K Z^B_{\ell_0}$. Against the normalised dressed loops the first bound holds with $\mathfrak{b}(\mathcal{C})(2s_K+2x_K+s^W_K)$ in place of s_K+x_K, the deviation insertions at the points of the histories being compared by the recursion and the dressings near the large-field regions by Supplement, Proposition S.14(f). Removing run B's own-support pieces changes W_{K+1} by at most $2B(\mathcal{C})s_K\delta_{\mathrm{LF}}$, as in (10.5), and the rest changes $W_{K+1} - W_K$ by at most $B(\mathcal{C})\delta_{\mathrm{LF}}(3s_K + 3x_K + s^W_K)$ through the numerators and the normalisations. Finally $s'_K \to 0$: $(K'+1)\tilde{\theta}^{K'-m} \to 0$ since $K' - m \geq K'/2$, the covariance terms decay geometrically against the polylogarithm ℓ_K, and for $i \geq K'/2$ the bound (11.24) gives $|\delta_i| \leq C[K'\tilde{\theta}^{K'/2} + Kw_K]$, where Kw_K is summable because $K\mu_K = K^{4-\kappa_0/2}\ell_K$ and $\kappa_0 > 10$. *q.e.d.*

11.8 Proof of the Comparison

Let $K_1 \geq K_0(\mathcal{C})$ be the least index such that every $K \geq K_1$ satisfies $K' \geq \max(k_0(\varepsilon_0), k_*, 2m_{\mathrm{rc}} + 2)$, with m_{rc} the integer of Theorem 9.5, $\max(m^E_K, m^w_K) \leq \lfloor K'/4 \rfloor$, $\rho^E_K \leq \frac{1}{4}$, and the further conditions of Supplement, Proposition S.16, each of which holds for K large because m^E_K and m^w_K grow as $\log\log K$ while ρ^E_K, x^*_K, and s_K tend to zero. For $K \geq K_1$ put

$$a_K := \Big[3x_K + \lambda_K + x^w_K + x^Q_K + b^{(S)}_K + \bar{\delta}_{\mathrm{LF}}(a^{(U)}_K + b^{(M)}_K)\Big]B(\mathcal{C}) + (e^{\Delta E_K} - 1)B(\mathcal{C})\,, \quad (11.32)$$

with $a^{(U)}_K := 4e^{-(C_P-1)K'}$ and $b^{(M)}_K := 4e^{-(C_P-1)(\lfloor K'/2\rfloor+1)} + 5s_K + 3x_K + s^W_K$.

Proof of Theorem 11.1. For $K < K_1$ the bound $|W_{K+1} - W_K| \leq 2B(\mathcal{C})$ is Theorem 8.7. Let $K \geq K_1$. We first replace run B's constant E^B_{const} by run A's: with $W^{B,A} := e^{E^A_{\mathrm{const}} - E^B_{\mathrm{const}}} W_{K+1}(\mathcal{C})$,

$$W_{K+1} - W_K = (W^{B,A} - W_K) + (e^{E^B_{\mathrm{const}} - E^A_{\mathrm{const}}} - 1)W^{B,A}\,, \quad (11.33)$$

and $|W^{B,A}| \leq B(\mathcal{C})$ because the bound of Lemma 8.6 holds for run B's dressed loop with run A's constant, so that the second term is at most $(e^{\Delta E_K} - 1)B(\mathcal{C})$. For the first, the terms of kind (U) of the two runs change the difference by at most $\bar{\delta}_{\mathrm{LF}} a^{(U)}_K B(\mathcal{C})$ and the old pairs change the difference by at most $4e^{-(C_P-1)(m+1)}\bar{\delta}_{\mathrm{LF}}B(\mathcal{C})$, by Corollary 10.3; the young pairs change it by at most $\bar{\delta}_{\mathrm{LF}}(5s_K + 3x_K + s^W_K)B(\mathcal{C})$, by Proposition 11.10; and the all-small-field pair by at most $(3x_K + \lambda_K + x^w_K + x^Q_K + b^{(S)}_K)B(\mathcal{C})$, by Lemma 11.9. This gives $|W_{K+1} - W_K| \leq a_K$. Each term of (11.32) is summable in K: $x_K \leq X_K e^{X_K}$ with X_K summable and bounded; λ_K, $a^{(U)}_K$, and the first term of $b^{(M)}_K$ decay geometrically; $b^{(S)}_K$ by Proposition 10.9; s_K and s^W_K by Proposition 11.10; x^w_K, x^Q_K, and ΔE_K by the bounds above; and $e^{\Delta E_K} - 1 \leq \Delta E_K e^{2B_E(\mathcal{C})}$. Since $B(\mathcal{C})$ and $\bar{\delta}_{\mathrm{LF}}$ do not depend on K, $\sum_K a_K < \infty$. *q.e.d.*

Proof of Theorem 1.5. The thresholds. We sort the requirements on the coupling of Sections 3–12 into the two kinds of Section 1.3. Those of the first kind are finitely many upper bounds on $\bar{g}$ in terms of the constants of the construction and of Λ_*, among them $\bar{g} \leq \gamma'$ of Theorem 4.1, and we let $\gamma_0 > 0$ be the smallest of these bounds. Each requirement of the second kind, for an ensemble with $c_{\mathcal{C}} = c$, bounds a nondecreasing function of γ_q from above by a positive quantity which does not increase with c, because the constants which it contains are nondecreasing functions of $c_{\mathcal{C}}$ (Definition 5.2(v)); it therefore holds whenever γ_q lies below a positive number which does not increase with c. We let $\gamma_\sharp(c) > 0$ be the smallest of these numbers, a nonincreasing function of c which depends only on the constants of the

construction and on Λ_*, and we put

$$q_\sharp(c,\bar g) \coloneqq \max\Big(0,\ \Big\lceil \frac{\gamma_\sharp(c)^{-2} - \bar g^{-2} + C_W}{b} \Big\rceil\Big)\ , \tag{11.34}$$

with the constants b and C_W of Theorem 4.1. This function is nondecreasing in c, because $\gamma_\sharp$ does not increase, and nondecreasing in $\bar g$; it vanishes as soon as $\bar g^{-2} \geq \gamma_\sharp(c)^{-2} + C_W$, and it depends only on the constants of the construction and on Λ_*. If $q \geq q_\sharp(c,\bar g)$, then $bq \geq \gamma_\sharp(c)^{-2} - \bar g^{-2} + C_W$, and the first inequality of (3.47) gives

$$\gamma_q^{-2} \geq \bar g^{-2} + bq - C_W \geq \gamma_\sharp(c)^{-2}\ , \tag{11.35}$$

that is, $\gamma_q \leq \gamma_\sharp(c)$.

The bound and the limit. Let $\bar g \in (0,\gamma_0]$, and let $\mathcal{C}$ be a loop ensemble with admissible loops at a level $q \geq q_\sharp(c_{\mathcal{C}},\bar g)$. Then $\bar g$ satisfies the requirements of the first kind, and γ_q satisfies those of the second kind for $\mathcal{C}$ by (11.35), so that $\bar g$ and q are as in Theorem 1.5 for $\mathcal{C}$ in the sense of Section 1.3. The bound of (1.10) is therefore Theorem 8.7 with $K_0 \coloneqq K_0(\mathcal{C})$ and $B \coloneqq B(\mathcal{C})$, and the summability of the differences is Theorem 11.1. A sequence with summable differences is a Cauchy sequence, which gives the limit (1.11). The dependence of K_0 and B is read from (8.1), Theorem 5.4, and Theorem 8.7. *q.e.d.*

11.9 Several Loops

The proof has been written for an ensemble of n loops throughout, and it remains to say where the number of loops enters. Every comparison above is made per traversal of a bond of the stopping step, per insertion, per cube, or per point of a history, and a cube of side less than the separation a_k of Lemma 3.14 meets one loop. The number of loops therefore enters only through sums over the loops: the leading words through λ_K, a sum over the loops, because every factor of $\prod_i f^{(i)}$ has modulus at most one; the transports and the insertions through $\mathfrak{b}(\mathcal{C})$, the exponential of a sum over the loops; the wraps through the total perimeter $P_{K'}(\mathcal{C})$ in their constants A_w; and the loop constant through the sum over the loops of the single-loop constants and the constants of the cross groups, whose per-step terms are bounded as those of one loop with the total perimeter in place of the perimeter (Corollary 7.11). The insertions of the cross species are not contracted, so that the recursion of the deviation insertions does not apply to them, and they are compared as the wraps are. Those born at the step $K'-m$ have total mass at most $2^{m+1}\mathsf{A}^\times_{K'-m}$ in each run, by (5.15), which decays doubly exponentially in m; those of age at least $m_K^\times$, an index which grows like $\log\log K$ and is not the age $m_K = \lfloor K^{1/2}\rfloor$ of the old terms of the effective action, are not compared, and their total $\tau_K^\times$ in each run is summable in K. The younger ones are produced by the fluctuation integrals and the **R**-operations of the last $m_K^\times$ steps, are compared at their birth by the interpolation of the constituents of their step, and are then carried. Their differences cannot be bounded separately from those of the wraps: a difference of the cross species can pass into a wrap, and the wraps inside the groups of the reference gas change the cluster occurrences which every new group merges, the deviation groups among them. The differences of the young insertions of the two species, of the activities of the reference gas which contain wraps, and of the part of the deviation insertions which comes from these occurrences therefore satisfy one linear recursion over the last $m_K^\times$ steps, with a factor per step which does not depend on K and with sources which are summable in K; its solution is at most a power of $\log K$ times these sources. Together with the weights this changes the normalised dressed loop by at most $x_K^\times \mathfrak{b}(\mathcal{C})$ with $x_K^\times$ summable in K, and for several loops $x_K^\times$ takes the place of $x_K^w + x_K^Q$ (Supplement, Proposition S.16(i)). The counts of the positions of a cross group are those of

one loop with the total perimeter, because two loops are at distance at least a_k. No condition on the coupling or on L involves n; the index K_1 involves $\mathcal{C}$ through $B(\mathcal{C})$, the total perimeter, and the thresholds of the single loops.

12 Symmetries, Reflection Positivity, and Nonvanishing

The symmetries and the reflection positivity are properties of the lattice measures at every cutoff, and one expects them to pass to the limits without effort, since a limit of equal numbers is an equality and a limit of positive semidefinite matrices is positive semidefinite. The difficulty lies in the normalisation. The renormalised expectations are the expectations divided by the products $\prod_i \kappa_K^{(q)}(C_i, \chi_i)$, so that a symmetry h of the lattice measures survives the division only if $\kappa_K^{(q)}(hC, \chi) = \kappa_K^{(q)}(C, \chi)$ for every loop; and the positivity of a matrix $(\mathbb{E}_K[F_a\, \Theta F_b])_{a,b}$ survives it only if the number by which the entry (a, b) is divided factorises as a number of a times the complex conjugate of the same number of b, which for the ensembles $\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b$ means that the normalisation of a reflected loop with the conjugate character must be the complex conjugate of the normalisation of the loop. The coefficients of the normalisation are defined by cluster expansions in which these symmetries are not visible, so we obtain them from the exact values of the flat parts, whose symmetries follow from those of the construction, and from the extraction of the coefficients along dilations of the loop. The construction is symmetric under the reflections in coordinate hyperplanes, but it is not known to be symmetric under the exchange of two axes, and this is where our covariance statements stop. The nonvanishing is then a consequence of the identity at the stopping scale: at small coupling the measure μ_K has almost full mass, the background is almost flat on its support, and all the corrections tend to zero with $\bar{g}$.

12.1 Symmetries and Reflection Positivity on the Lattice

A symmetry h of level q (Section 1.3) acts on the bonds of $\mathbb{T}_K$, $K \geq q$, by $h(x, y) := (hx, hy)$, on paths bond by bond, and on gauge fields by $(hU)(b) := U(hb)$, so that $(hU)(\gamma) = U(h\gamma)$ for every path γ. For a function F of the gauge field we put $(F \circ h)(U) := F(hU)$; in particular $W_\chi(C) \circ h = W_\chi(hC)$ for a loop C at level q, and $W_{\mathcal{C}} \circ h = W_{h\mathcal{C}}$ for an ensemble.

Lemma 12.1 (Invariance of the Lattice Measures)**.** *Let $K \geq q$ and let h be a symmetry of level q. For every bounded measurable function F of the gauge field on $\mathbb{T}_K$, $\mathbb{E}_K[F \circ h] = \mathbb{E}_K[F]$ and $\mathbb{E}_K[\bar{F}] = \overline{\mathbb{E}_K[F]}$.*

Proof. We write $s_U(p) := 1 - \operatorname{Re}\operatorname{tr} U(\partial p)$. The holonomy $(hU)(\partial p) = U(h\partial p)$ is that of U around the plaquette hp, from one of its corners and in one of its two orientations, and $\operatorname{Re}\operatorname{tr}$ is invariant under conjugation and, on unitary matrices, under inversion; hence $s_{hU}(p) = s_U(hp)$. Since h permutes the plaquettes, the action of (1.2) is invariant under $U \mapsto hU$. The map h permutes the unoriented bonds, and the value of hU on a positive bond b is the value of U on hb if hb is positive and the inverse of its value on $(hb)^{-1}$ otherwise; since the Haar measure is invariant under inversion, the bijection $U \mapsto hU$ preserves the product Haar measure. The change of variables $U \mapsto hU$ in the integral defining $\mathbb{E}_K[F \circ h]$ therefore gives the first assertion, and the second holds because μ_{K,g_0} is a positive measure. *q.e.d.*

For the reflection positivity we use the reflections of level q, for which the positivity of the Wilson action was observed by Osterwalder and Seiler [17]; we give the short proof, since it is the basis of everything in this section. Let $\mu \in \{0, \dots, 3\}$ and $a \in uL^{-q}\mathbb{Z}$. Since a

and $T/2 = L^{m_T} u$ are multiples of ε_K for $K \geq q$, the reflection $\theta_{\mu,a}$ is an involution of the sites of $\mathbb{T}_K$, which fixes the two hyperplanes $P_{\mu,a} \coloneqq \{x_\mu \equiv a\} \cup \{x_\mu \equiv a + T/2\}$ pointwise and exchanges the open half-tori $H_{\mu,a} \coloneqq \{a < x_\mu < a + T/2\}$ and $\theta_{\mu,a} H_{\mu,a}$. For a bounded measurable function F of the gauge field we put $(\Theta_{\mu,a} F)(U) \coloneqq \overline{F(\theta_{\mu,a} U)}$; the map $\Theta_{\mu,a}$ is antilinear and multiplicative.

Let $\bar{\Lambda}_+$ be the set of the sites with $x_\mu - a \in [0, T/2]$ modulo T, and $\bar{\Lambda}_- \coloneqq \theta_{\mu,a} \bar{\Lambda}_+$, so that $\bar{\Lambda}_+ \cap \bar{\Lambda}_- = P_{\mu,a}$. Let E_0 be the set of the bonds with both ends in $P_{\mu,a}$, E_+ the set of the other bonds with both ends in $\bar{\Lambda}_+$, and $E_- \coloneqq \theta_{\mu,a} E_+$. Every bond belongs to exactly one of these sets: a bond orthogonal to e_μ has both ends on one hyperplane $\{x_\mu = \text{const}\}$, and a bond along e_μ joins two neighbouring hyperplanes, which lie both in $\bar{\Lambda}_+$ or both in $\bar{\Lambda}_-$, because the two hyperplanes of $P_{\mu,a}$ are hyperplanes of $\mathbb{T}_K$ at distance $T/2 \geq 2\varepsilon_K$. The bonds of E_0 are orthogonal to e_μ, so $\theta_{\mu,a}$ fixes each of them with its orientation. We write $\mathcal{A}_+$ for the set of the bounded measurable functions of the restriction of U to $E_+ \cup E_0$; for $F \in \mathcal{A}_+$ the function $\Theta_{\mu,a} F$ depends on the restriction to $E_- \cup E_0$.

Lemma 12.2 (Reflection Positivity on the Lattice)**.** *Let $K \geq q$, $\mu \in \{0, \dots, 3\}$, and $a \in uL^{-q}\mathbb{Z}$. For every bounded measurable F, $\mathbb{E}_K[\Theta_{\mu,a} F] = \overline{\mathbb{E}_K[F]}$, and for every $F \in \mathcal{A}_+$, $\mathbb{E}_K[F\, \Theta_{\mu,a} F] \geq 0$.*

Proof. We write θ for $\theta_{\mu,a}$, which is a symmetry of level q, and $s_U(p) \coloneqq 1 - \operatorname{Re} \operatorname{tr} U(\partial p)$ as in the proof of Lemma 12.1. That lemma, applied to $\bar{F}$, gives $\mathbb{E}_K[\Theta F] = \mathbb{E}_K[\bar{F} \circ \theta] = \mathbb{E}_K[\bar{F}] = \overline{\mathbb{E}_K[F]}$, which is the first assertion.

For the second, a plaquette lies in $\bar{\Lambda}_+$ or in $\bar{\Lambda}_-$, by the argument given for the bonds, and in both exactly when it lies in $P_{\mu,a}$. Put

$$B(U) \coloneqq \sum_{p \subset \bar{\Lambda}_+,\ p \not\subset P_{\mu,a}} s_U(p) + \frac{1}{2} \sum_{p \subset P_{\mu,a}} s_U(p)\,, \tag{12.1}$$

a real function in $\mathcal{A}_+$. The reflection maps the plaquettes of $\bar{\Lambda}_+$ which do not lie in $P_{\mu,a}$ bijectively onto those of $\bar{\Lambda}_-$, and each plaquette of $P_{\mu,a}$ onto itself, so that, since $s_{\theta U}(p) = s_U(\theta p)$, $B(\theta U)$ is the sum (12.1) with $\bar{\Lambda}_-$ in place of $\bar{\Lambda}_+$, and the action is $B + B \circ \theta$. With $f \coloneqq F e^{-g_0^{-2} B} \in \mathcal{A}_+$ and since B is real, the integrand of $\mathbb{E}_K[F\, \Theta F]$ against the Haar measure is therefore

$$F(U) \overline{F(\theta U)}\, e^{-g_0^{-2}(B(U) + B(\theta U))} = f(U)\, \overline{f(\theta U)}\,. \tag{12.2}$$

Write $U = (U_+, U_0, U_-)$ for the restrictions to E_+, E_0, and E_-. The restriction of θU to $E_+ \cup E_0$ is $(\varphi(U_-), U_0)$, where $\varphi(U_-)(b) \coloneqq U(\theta b)$ for $b \in E_+$ defines a map $\varphi \colon G^{E_-} \to G^{E_+}$ which maps the product Haar measure to the product Haar measure, as in the proof of Lemma 12.1. By Fubini's theorem, the integral of (12.2) is

$$\int \Big(\int f(U_+, U_0)\, \mathrm{d}U_+ \Big) \overline{\Big(\int f(\varphi(U_-), U_0)\, \mathrm{d}U_- \Big)}\, \mathrm{d}U_0 = \int \Big| \int f(U_+, U_0)\, \mathrm{d}U_+ \Big|^2 \mathrm{d}U_0 \geq 0\,, \tag{12.3}$$

and $\mathbb{E}_K[F\, \Theta F]$ is this number divided by $Z_{K,g_0} > 0$. *q.e.d.*

The lemma makes $\langle F, G \rangle \coloneqq \mathbb{E}_K[F\, \Theta_{\mu,a} G]$ a positive semidefinite sesquilinear form on $\mathcal{A}_+$, linear in F and antilinear in G; such a form is Hermitian, by polarisation. If $F \in \mathcal{A}_+$ has the mean $c \coloneqq \mathbb{E}_K[F]$, then the constant c lies in $\mathcal{A}_+$, $\mathbb{E}_K[\Theta_{\mu,a} F] = \bar{c}$ by the lemma, and

$$0 \leq \langle F - c, F - c \rangle = \mathbb{E}_K[F\, \Theta_{\mu,a} F] - |\mathbb{E}_K[F]|^2\,. \tag{12.4}$$

The Wilson loops behave as follows. Let C be a loop whose sites lie in $H_{\mu,a}$, with the character χ of the representation π. Its bonds have both ends in $H_{\mu,a}$, hence lie in E_+, and $W_\chi(C) \in \mathcal{A}_+$. Since π is unitary, $\overline{\chi(g)} = \bar{\chi}(g)$ for the character $\bar{\chi}$ of the conjugate representation, which is irreducible and unitary, and therefore

$$(\Theta_{\mu,a} W_\chi(C))(U) = \overline{\chi(U(\theta_{\mu,a} C))} = W_{\bar{\chi}}(\theta_{\mu,a} C; U) \,, \tag{12.5}$$

where $\theta_{\mu,a} C$ is the image of C with the image orientation, a loop at level q with the same chord–arc constant, ratio P/r, and number P_q as C (Section 1.3). For an ensemble $\mathcal{D}$ all of whose loops lie in $H_{\mu,a}$, the ensemble $\theta_{\mu,a}\bar{\mathcal{D}}$ consists of the images with the conjugate characters, and by multiplicativity $\Theta_{\mu,a} W_{\mathcal{D}} = W_{\theta_{\mu,a}\bar{\mathcal{D}}}$.

12.2 The Covariance of the Normalisation

The division by the normalisations keeps the symmetries of the lattice measures only where the normalisations share them, and the following proposition says where they do.

Proposition 12.3 (Covariance of the Normalisation)**.** *Let L and Λ_* be as in Theorem 1.5, let C be an admissible loop at level q with the character χ of a representation π, and let $\bar{g}$ and q be as in Theorem 1.5 for $((C,\chi))$. For every $K > j_q$ and every $g \in \mathfrak{G}_q$,*

$$\kappa_K^{(q)}(gC,\chi) = \kappa_K^{(q)}(C,\chi) \,, \qquad \kappa_K^{(q)}(C,\bar{\chi}) = \overline{\kappa_K^{(q)}(C,\chi)} \,, \qquad \kappa_K^{(q)}(C^{-1},\chi) = \kappa_K^{(q)}(C,\bar{\chi}) \,. \tag{12.6}$$

In particular $\kappa_K^{(q)}(\theta_{\mu,a} C,\bar{\chi}) = \overline{\kappa_K^{(q)}(C,\chi)}$ for every μ and every a with $2a \in uL^{-q}\mathbb{Z}$.

For the translations in $\mathfrak{G}_q$ the first identity can be read off from Proposition 7.13, since a translate of C has the same numbers $N_v^{(0)}$ and the same pairs (v_c, w_c) at its corners as C. The structure of the normalisation does not help with the reflections θ_μ, nor with the other two identities: the loops $\theta_\mu C$ and C^{-1} have different numbers of segments of a given orientation and different pairs at their corners, and the coefficients of χ and of $\bar{\chi}$ are different numbers, while nothing relates the coefficients of different orientations, or of conjugate characters, to each other. The coefficients of Definition 7.8 are sums of cluster weights, and (12.6) would follow from symmetries of the individual weights; but the expansions which define the weights involve many choices whose behaviour under the reflections we do not wish to follow. We use instead the exact values of the flat parts, which do not depend on any choice. For a torus $T' \in \mathfrak{T}$, a loop C at level q on T', and $0 \le j \le K'$, let $\widehat{\mathcal{W}}_j(C,\chi;T')$ be the value at the reference data of the dressed loop of the step j of the single-loop ensemble $((C,\chi))$ on the trivial term, with $\widehat{\mathcal{W}}_0 := 1$, whenever this ensemble is dressed at the step j.

Lemma 12.4 (Flat Values under the Symmetries)**.** *Let $T' \in \mathfrak{T}$, let C be a loop at level q on T' with the character χ, let $1 \le j \le K'$, and let $\mu \in \{0,\dots,3\}$. If the ensemble $((C,\chi))$ is dressed at the steps $1,\dots,j$, then*

$$\widehat{\mathcal{W}}_j(C,\chi;T') = \kappa^{(j)}(C,\chi)\, e^{E_{\mathrm{const},j}(C,\chi;T')} \,, \tag{12.7}$$

where $E_{\mathrm{const},j}(C,\chi;T')$ is the constant of the scalar exponent of the step j on the trivial term, and

$$\begin{gathered} \widehat{\mathcal{W}}_j(\theta_\mu C,\chi;T') = \widehat{\mathcal{W}}_j(C,\chi;T') \,, \qquad \widehat{\mathcal{W}}_j(C,\bar{\chi};T') = \overline{\widehat{\mathcal{W}}_j(C,\chi;T')} \,, \\ \widehat{\mathcal{W}}_j(C^{-1},\chi;T') = \widehat{\mathcal{W}}_j(C,\bar{\chi};T') \,, \end{gathered} \tag{12.8}$$

each identity holding whenever the single-loop ensembles which occur in it are dressed at the steps $1,\dots,j$.

Proof. At the reference data the leading word is $\chi(1) = 1$, and the insertions of the nonempty families vanish: after the exact factorisation (7.17), and after its counterpart at the first step in the proof of Theorem 5.4, the insertions of the families are the parts $Y^{\mathrm{d}}_{\mathcal{G}}$, which vanish at the reference data, because for a single loop every insertion takes a scalar value there, by Schur's lemma. On the trivial term the scalar exponent is the number $E_{\mathrm{const},j}$, as in the proof of Theorem 8.1. This gives (12.7).

For (12.8) we express the flat values through the construction itself. The reference data of the trivial term are those at the block field $V_j = 1$, which lies in the support of all its test functions, and on the trivial term the operation $\mathbf{T}_j$ holds no integrations. Let $N_j[F]$ be the density obtained from $F\varrho_0$ by the operations of the first j steps on T', and $N_j[F](1;\ell_0)$ the value of its term ℓ_0 at $V_j = 1$. Since the operations which produce the terms of the numerator are those which produce the terms of the density, applied to a function which carries one more factor, Definition 5.2 gives

$$\widehat{\mathcal{W}}_j(C,\chi;T') = \frac{N_j[W_\chi(C)](1;\ell_0)}{N_j[1](1;\ell_0)} \,, \tag{12.9}$$

where $N_j[1](1;\ell_0) = \exp A_j(1) > 0$. The right side depends on the loop and its character only through the function $W_\chi(C)$. The map $F \mapsto N_j[F](1;\ell_0)$ has two properties, which are proved in Supplement, Proposition S.18. It is *real*: $N_j[\bar{F}](1;\ell_0) = \overline{N_j[F](1;\ell_0)}$, because at real fields every operation of the construction is a sum, a multiplication by a real function, or an integral against a real kernel (Haar and Gaussian measures, δ-functions of the block averaging, test functions, and exponentials of real actions [9, (3.27), p. 271], [12, p. 380]). And it is *invariant under each* θ_μ: $N_j[F\circ\theta_\mu](1;\ell_0) = N_j[F](1;\ell_0)$, because θ_μ maps every lattice and every partition of the construction to itself, the partitions being centred at the origin (Section 7.1), maps the trivial term and the trivial field to themselves, and commutes with every operation: with the averages, which Bałaban defines so as to be "symmetric with respect to lattice Euclidean transformations" [8, p. 252], with the gauge fixing, introduced "to preserve the Euclidean invariance" [9, p. 247], with the fluctuation integral [8, (2.17), (2.18), p. 269], whose fluctuation fields are written first in an axial gauge [6, (1.19), p. 79] with contours which start at the centre of a block and change the coordinates in a fixed order [2, (1.7), p. 18], an order which the reflection of one coordinate about the centre does not change, with the soft test functions, which are functions of plaquette deviations in norms invariant under conjugation and inversion, and with the **R**-operation, whose gauge-fixing graphs are averaged over "2^d reflections in subsets of coordinates" [11, (1.100), p. 201], [12, p. 379].

The three identities now follow from (12.9). Since $W_\chi(C)\circ\theta_\mu = W_\chi(\theta_\mu C)$, the invariance gives $N_j[W_\chi(\theta_\mu C)](1;\ell_0) = N_j[W_\chi(C)](1;\ell_0)$. Since π is unitary, $\overline{\chi(V)} = \chi(V^{-1}) = \bar{\chi}(V)$ for $V \in G$, so that $W_{\bar\chi}(C) = \overline{W_\chi(C)}$, and the reality gives $N_j[W_{\bar\chi}(C)](1;\ell_0) = \overline{N_j[W_\chi(C)](1;\ell_0)}$. Finally $U(C^{-1}) = U(C)^{-1}$, so that $W_\chi(C^{-1}) = \chi(U(C)^{-1}) = \bar\chi(U(C)) = W_{\bar\chi}(C)$ is the same function as $W_{\bar\chi}(C)$. Dividing by the positive number $N_j[1](1;\ell_0)$ gives (12.8). *q.e.d.*

Proof of Proposition 12.3. We first reduce the first identity to the reflections θ_μ. Every $g \in \mathfrak{G}_q$ is a composition $g = g_1\cdots g_r$ of translations by vectors of $uL^{-q}\mathbb{Z}^4$ and reflections θ_μ. Each loop $g_{i+1}\cdots g_r C$ is an admissible loop at level q with the character χ (Section 1.3), so that $\bar g$ and q are as in Theorem 1.5 for it, the conditions of that theorem involving the loop only through c_π and Λ_*; and the normalisation of a loop is unchanged by a translation in $\mathfrak{G}_q$, as noted after the proposition. It therefore suffices to prove the first identity for $g = \theta_\mu$, and the last assertion follows from the first two identities, since $\theta_{\mu,a} \in \mathfrak{G}_q$ when $2a \in uL^{-q}\mathbb{Z}$.

The three remaining identities have a common form. Let $(C',\chi',\chi^\circ,\epsilon)$ be one of the three

quadruples

$$(\theta_\mu C, \chi, \chi, \mathrm{id}) \,, \qquad (C, \bar\chi, \chi, \mathrm{conj}) \,, \qquad (C^{-1}, \chi, \bar\chi, \mathrm{id}) \,, \tag{12.10}$$

where id is the identity of $\mathbb{C}$ and conj the complex conjugation, so that the identity to be proved reads $\kappa_K^{(q)}(C', \chi') = \epsilon(\kappa_K^{(q)}(C, \chi^\circ))$. The pairs (C, χ°) and (C', χ') satisfy the hypotheses of the proposition, because $c_{\bar\pi} = c_\pi$ and $\Lambda(C') = \Lambda(C)$. It suffices to prove $\kappa_j(C', \chi') = \epsilon(\kappa_j(C, \chi^\circ))$ for every step $1 \le j \le K'$, and since κ_j is the exponential of (7.23) and ϵ commutes with the exponential, it suffices to prove that

$$D_j(C) \coloneqq \log \kappa_j(C', \chi') - \epsilon(\log \kappa_j(C, \chi^\circ)) \in 2\pi i \mathbb{Z} \,. \tag{12.11}$$

Fix j. Let c_0 be a corner of C, and for an integer $m \ge 1$ let C_m be the image of C under the dilation $x \mapsto c_0 + m(x - c_0)$, read on a torus $T'_m \in \mathfrak{T}$ whose period we choose below. The path C_m visits no site twice, its corners lie in $c_0 + uL^{-q}\mathbb{Z}^4$, its segments have the orientations of those of C and m times their lengths, and its corners have the same pairs (v_c, w_c) as those of C. Hence C_m is a loop at level q on T'_m as soon as $3\,\mathrm{diam}(C_m) < T'_m$, with $\Lambda(C_m) = \Lambda(C)$, $P(C_m)/r(C_m) = P(C)/r(C)$, $P_{j-1}(C_m) = mP_{j-1}(C)$, and $r_{j-1}(C_m) = mr_{j-1}(C)$. We choose the period of T'_m so large that, in addition, $K \ge K_{\mathrm{tor}}$ for C_m on T'_m, which is possible because the torus conditions of the first step read $(P_0(C_m)/L)e^{-c\iota_{T'_m}} \le 1$ in the proof of Theorem 5.4, and that $mP_{j-1}(C)e^{-c_2\iota_{T'_m,j}} \le 1/m$, with the constant $c_2 > 0$ of (12.15) below. Let C'_m be obtained from C_m as C' is obtained from C, that is, C'_m is $\theta_\mu C_m$, C_m, or C_m^{-1}; it is the dilation of C' by the factor m about $\theta_\mu c_0$, c_0, or c_0, since θ_μ is linear. Then the ensembles $((C_m, \chi^\circ))$ and $((C'_m, \chi'))$ are dressed at every step up to K', by Theorems 5.4 and 6.1, applied on T'_m; their hypotheses hold because $\Lambda(C'_m) = \Lambda(C_m) = \Lambda(C)$, because C'_m has the same numbers P_i and r_i as C_m at every step i, so that the torus conditions hold for both, and because $c_{\bar\pi} = c_\pi$, so that $\bar g$ and q are as in Theorem 1.5 for both ensembles.

By (7.23), $\log \kappa_j(C_m, \chi^\circ)$ is an affine function of m, and so is $\log \kappa_j(C'_m, \chi')$, because C'_m is a dilation of C' by the factor m. Hence

$$D_j(C_m) = mA + B \tag{12.12}$$

with numbers A and B which do not depend on m, and $D_j(C) = A + B$. On the other hand, by (12.7) at the steps j and $j-1$, for a single loop on the trivial term,

$$\frac{\widehat{\mathcal{W}}_j(C_m, \chi^\circ; T'_m)}{\widehat{\mathcal{W}}_{j-1}(C_m, \chi^\circ; T'_m)} = \exp(\log \kappa_j(C_m, \chi^\circ) + E_j(C_m, \chi^\circ; T'_m)) \,, \tag{12.13}$$

where $E_j \coloneqq E_{\mathrm{const},j} - E_{\mathrm{const},j-1}$ is the remainder of the step, which for a single loop is $E_j^\circ + \mathbf{1}_{\mathbf{R}}(j) E_j^{\circ,\mathbf{R}}$ by (7.30) at $j \ge 2$, and the remainder bounded in (5.25) and by the periodic comparison of [1, Appendix F] at $j = 1$. The same identity holds for (C'_m, χ'), and by (12.8) at the steps j and $j-1$ the left side for (C'_m, χ') is the image under ϵ of the left side for (C_m, χ°). Since ϵ commutes with the exponential, taking exponentials apart gives

$$D_j(C_m) \in 2\pi i \mathbb{Z} + \epsilon(E_j(C_m, \chi^\circ; T'_m)) - E_j(C'_m, \chi'; T'_m) \,. \tag{12.14}$$

The remainders tend to zero as $m \to \infty$. Indeed, by (7.25) and (7.29) at $j \ge 2$, and by (5.25) and the periodic comparison of [1, Appendix F] at $j = 1$, and since $P_j \le P_{j-1}$ and $r_{j-1} \ge r_j$, each of them is at most a constant times

$$(mP_{j-1}(C) + |\mathcal{K}(C)|)e^{-c_1 m r_j(C)} + mP_{j-1}(C)\, e^{-c_2 \iota_{T'_m,j}} \,, \tag{12.15}$$

with positive constants c_1 and c_2 which depend on j but not on m (at $j \ge 2$ the smaller of $2c_E^T$ and $c_E^R/(MR_j)$, and of c^Z and $c_E^R/(MR_j)$), and with the constants of χ and $\bar\chi$; the first

term tends to zero because $r_j(C) > 0$, and the second is at most $1/m$ by the choice of T'_m. Let δ_m be the modulus of the difference of the two remainders in (12.14), so that $\delta_m \to 0$. By (12.12), $A = D_j(C_{m+1}) - D_j(C_m)$ lies within the distance $\delta_m + \delta_{m+1}$ of $2\pi i\mathbb{Z}$ for every m, and since $2\pi i\mathbb{Z}$ is closed, $A \in 2\pi i\mathbb{Z}$. Then $B = D_j(C_m) - mA$ lies within the distance δ_m of $2\pi i\mathbb{Z}$ for every m, and $B \in 2\pi i\mathbb{Z}$. Hence $D_j(C) = A + B \in 2\pi i\mathbb{Z}$, which is (12.11). *q.e.d.*

The argument determines the logarithms of the normalisations only modulo $2\pi i$, which is all that (12.6) asks for; it does not show that the coefficients themselves are real, nor that they are invariant under the reflections, since the dilations of one loop see only the combinations of the coefficients which occur in that loop.

12.3 Proof of the Symmetries

Proof of Corollary 1.6. Each of the ensembles $\mathcal{C}'$ of the statement, namely $\mathcal{C}$ with its loops in another order, $g\mathcal{C}$, $\bar{\mathcal{C}}$, $\mathcal{C}$ with one loop reversed and its character conjugated, and $h\mathcal{C}$, consists of admissible loops at level q which are pairwise vertex-disjoint, the maps g and h being bijections, and has $c_{\mathcal{C}'} = c_{\mathcal{C}}$, because $c_{\bar\pi} = c_\pi$. So $q \geq q_\sharp(c_{\mathcal{C}'}, \bar g)$, the limit $W(\mathcal{C}')$ exists by Theorem 1.5, and $\bar g$ and q are as in that theorem for every single loop of $\mathcal{C}$ with its character, since $c_{\pi_i} \leq c_{\mathcal{C}}$ and $q_\sharp$ is nondecreasing. It suffices to prove the identities for W_K at a fixed $K > j_q$.

(i) Both the observable (1.4) and the product of the normalisations in (1.6) are products over the loops, which do not depend on the order of the factors.

(ii) By Lemma 12.1, $\mathbb{E}_K[W_{g\mathcal{C}}] = \mathbb{E}_K[W_{\mathcal{C}} \circ g] = \mathbb{E}_K[W_{\mathcal{C}}]$, and by Proposition 12.3, $\kappa_K^{(q)}(gC_i, \chi_i) = \kappa_K^{(q)}(C_i, \chi_i)$ for every i. Dividing gives $W_K(g\mathcal{C}) = W_K(\mathcal{C})$.

(iii) Since $W_{\bar\chi_i}(C_i) = \overline{W_{\chi_i}(C_i)}$, as in the proof of Lemma 12.4, Lemma 12.1 gives $\mathbb{E}_K[W_{\bar{\mathcal{C}}}] = \overline{\mathbb{E}_K[W_{\mathcal{C}}]}$, and Proposition 12.3 gives $\kappa_K^{(q)}(C_i, \bar\chi_i) = \overline{\kappa_K^{(q)}(C_i, \chi_i)}$; dividing gives $W_K(\bar{\mathcal{C}}) = \overline{W_K(\mathcal{C})}$. If C_i is replaced by C_i^{-1} and χ_i by $\bar\chi_i$, the observable does not change, since $W_{\bar\chi_i}(C_i^{-1}) = W_{\chi_i}(C_i)$, and neither does the normalisation, since $\kappa_K^{(q)}(C_i^{-1}, \bar\chi_i) = \kappa_K^{(q)}(C_i, \chi_i)$ by the third identity of (12.6) applied to $(C_i, \bar\chi_i)$.

(iv) By Lemma 12.1, $\mathbb{E}_K[W_{h\mathcal{C}}] = \mathbb{E}_K[W_{\mathcal{C}}]$, and by Proposition 12.3, $\kappa_K^{(q)}(hC_i, \chi_i) = \kappa_K^{(q)}(g_iC_i, \chi_i) = \kappa_K^{(q)}(C_i, \chi_i)$ for every i. *q.e.d.*

The proof of (iv) shows what is missing for a general symmetry h which exchanges two axes. Writing $h(x) = Ox + t$, the image hC has $N_v^{(0)}(hC) = N_{O^{-1}v}^{(0)}(C)$ and the pairs (Ov_c, Ow_c) at its corners, so that Lemma 12.1 and Proposition 7.13 give

$$\frac{W_K(h\mathcal{C})}{W_K(\mathcal{C})} = \prod_{i=1}^{n} \exp\Big(\sum_{v \in \mathcal{V}} N_v^{(0)}(C_i)(m_{v,K} - m_{Ov,K}) + \sum_{c \in \mathcal{K}(C_i)} (z_{v_c,w_c,K} - z_{Ov_c,Ow_c,K}) \Big) \quad (12.16)$$

whenever $W_K(\mathcal{C}) \neq 0$, with the coefficients of the characters χ_i. The covariance of the limits under h therefore follows from the equality of the coefficients of the orientations v and Ov and of the corner pairs (v, w) and (Ov, Ow), and the method of Proposition 12.3 would give it from the invariance $N_j[F \circ h](1; \ell_0) = N_j[F](1; \ell_0)$ of the flat values. We do not know whether the construction has this invariance. The averages, the first gauge fixing, the designated bonds [8, p. 267, ll. 19–20], and the **R**-operation, whose graphs are averaged over the permutations of the coordinates as well [11, p. 201, ll. 34–38], are symmetric under the exchange of two axes; but the fluctuation fields of the steps $k \geq 2$ are written in the axial gauge of [6, (1.19), p. 79],

whose contours change the coordinates in one fixed order [2, (1.7), p. 18], and the Landau gauge into which they are then transformed is defined relative to this preliminary choice, since the restricted gauge transformations do not form a group and the orbit "depends on an element U chosen to generate the set" [6, pp. 80–81]. The test functions of the fluctuation fields [9, (3.16), p. 268], and with them the term ℓ_0, are therefore not known to be invariant under the exchange of two axes. Bałaban states the covariance of his effective actions under every Euclidean transformation which preserves the lattices [8, p. 269, ll. 9–26], [9, (3.39), p. 274], but the operations from which we would derive the invariance of the numerator include this axial gauge, and we have not found an argument which removes it. An axial gauge in the averaged variables, as for the first gauge fixing, would give the symmetry, at the price of proving Bałaban's existence, uniqueness, and bounds for the Landau gauge again from it.

The positivity has no counterpart for the reflections which exchange two axes, for a reason which has nothing to do with the normalisations. Such a reflection is $x \mapsto Ox + t$ with O the reflection in the hyperplane orthogonal to $e_\mu - e_\nu$ or to $e_\mu + e_\nu$. Its fixed points, if there are any, satisfy one equation $x_\mu \mp x_\nu \equiv c$ modulo T, so that they form a single hyperplane of $\mathbb{T}$, whose complement is connected: it is the image of the connected set $(0,T) \times (\mathbb{R}/T\mathbb{Z})^3$ under the continuous map which sends (s,y) to the point whose coordinates other than x_μ are those of y and for which $x_\mu \mp x_\nu = c + s$. There are therefore no two half-tori which the reflection exchanges and whose union with the fixed points is the torus, and the proof of Lemma 12.2 has nothing to apply to.

12.4 Proof of the Reflection Positivity

Proof of Corollary 1.7. Let $\theta = \theta_{\mu,s}$ with $s \in uL^{-q}\mathbb{Z}$. The two half-tori bounded by its fixed points are $H_{\mu,s}$ and $H_{\mu,s+T/2}$, and $\theta_{\mu,s+T/2} = \theta_{\mu,s}$, because $2s + T \equiv 2s$ modulo T; since $T/2 = L^{m_T} u \in uL^{-q}\mathbb{Z}$, we may therefore assume $H_+ = H_{\mu,s}$, and we write $\Theta \coloneqq \Theta_{\mu,s}$ and $H_- \coloneqq \theta H_+$. For $0 \le a, b \le m$ the family $\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b$ is a loop ensemble at level q with admissible loops: the loops of $\mathcal{D}_a$ lie in H_+ and those of $\theta\bar{\mathcal{D}}_b$ in H_-, which is disjoint from H_+, so that they are pairwise vertex-disjoint, and the images are admissible loops at level q by Section 1.3. Its number $c_{\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b} = \max(c_{\mathcal{D}_a}, c_{\mathcal{D}_b})$ is at most c, because $c_{\theta\bar{\mathcal{D}}_b} = c_{\mathcal{D}_b}$, so that $q \ge q_\sharp(c, \bar{g}) \ge q_\sharp(c_{\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b}, \bar{g})$. By Theorem 1.5 the limits $W(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b)$ exist, with $W_K(\emptyset) = 1$ for every K.

Fix $K > j_q$, write $F_a \coloneqq W_{\mathcal{D}_a}$, with $F_0 \coloneqq 1$, and κ_a for the product of the normalisations $\kappa_K^{(q)}$ of the loops of $\mathcal{D}_a$, with $\kappa_0 \coloneqq 1$. By the last assertion of Proposition 12.3, which applies to each loop of $\mathcal{D}_b$ because its number $c_\pi \le c$ gives $q \ge q_\sharp(c_\pi, \bar{g})$, the product of the normalisations of the loops of $\theta\bar{\mathcal{D}}_b$ is $\bar{\kappa}_b$. Since $F_a \in \mathcal{A}_+$ and $\Theta F_b = W_{\theta\bar{\mathcal{D}}_b}$ by Section 12.1, the functions $G_a \coloneqq F_a/\kappa_a \in \mathcal{A}_+$ satisfy, by Definition 1.3,

$$W_K(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b) = \frac{\mathbb{E}_K[F_a\, \Theta F_b]}{\kappa_a \bar{\kappa}_b} = \mathbb{E}_K[G_a\, \Theta G_b] = \langle G_a, G_b \rangle \;, \tag{12.17}$$

because Θ is antilinear. For $\xi \in \mathbb{C}^m$ we obtain $\sum_{a,b} \xi_a \bar{\xi}_b W_K(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b) = \langle \sum_a \xi_a G_a, \sum_b \xi_b G_b \rangle \ge 0$ by Lemma 12.2, and the matrix is Hermitian because the form is. Letting $K \to \infty$ entry by entry, the limit matrix is Hermitian and positive semidefinite, which is (i).

For (ii), $W_K(\theta\bar{\mathcal{D}}_a) = \mathbb{E}_K[\Theta F_a]/\bar{\kappa}_a = \overline{\mathbb{E}_K[F_a]/\kappa_a} = \overline{W_K(\mathcal{D}_a)}$ by Lemma 12.2, and the limit gives $W(\theta\bar{\mathcal{D}}_a) = \overline{W(\mathcal{D}_a)}$. The mean of G_a is $w_a \coloneqq W_K(\mathcal{D}_a)$, and the mean of ΘG_b is $\bar{w}_b$, so that

$$\langle G_a - w_a, G_b - w_b \rangle = W_K(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b) - w_a \bar{w}_b \;, \tag{12.18}$$

and the matrix of the right sides is Hermitian and positive semidefinite by the same argument; its limit is the matrix of (ii). *q.e.d.*

12.5 Nonvanishing and the Growth Bound

The connected functions of Corollary 1.7(ii), and the logarithms of Theorem 1.8, are useful only if the limits do not vanish, and we now show that they do not, on the ensembles whose loops are not too long for the coupling.

Proposition 12.5 (Nonvanishing)**.** *Let L and Λ_* be as in Theorem 1.5, and let $q \geq 0$. There are $\bar g_1 \in (0, \gamma_0]$ and a function $\ell\colon [0,\infty) \times (0, \bar g_1] \to [0,\infty]$, nonincreasing in its first argument and with $\ell(c, \bar g) \to \infty$ as $\bar g \to 0$ for every c, which depend only on the constants of the construction, Λ_*, q, and T/u, such that the following holds for every $\bar g \in (0, \bar g_1]$. If $\mathcal{C}$ is a loop ensemble at level q whose loops are admissible and whose lengths satisfy*

$$\sum_{i=1}^{n} \Big(P_q(C_i)^2 + \frac{P(C_i)}{r(C_i)} \Big) \leq \ell(c_{\mathcal{C}}, \bar g) \,, \tag{12.19}$$

then $q \geq q_\sharp(c_{\mathcal{C}}, \bar g)$, the limit $W(\mathcal{C})$ exists, and, for every $K \geq K_0(\mathcal{C})$,

$$|W_K(\mathcal{C})| \geq \frac{1}{2} \,, \qquad \text{so that} \qquad |W(\mathcal{C})| \geq \frac{1}{2} \,. \tag{12.20}$$

The condition (12.19) bounds the lengths of the loops by a number which depends on their representations through $c_{\mathcal{C}}$ and grows without limit as the coupling decreases. Unlike the conditions of Theorem 1.5, it does restrict the representations at a given coupling: since a loop has at least four bonds, it fails whenever $\ell(c_{\mathcal{C}}, \bar g) < 16$. It involves the number of loops only through the sum, and for the ensembles which are generated from a fixed one by reflections in coordinate hyperplanes it is needed only for the ensemble from which they are generated (Proposition 12.8).

For the lower bound we need that the Wilson loop of a nearly flat field is close to one, with a bound in terms of the length of the loop and the size of a box containing it.

Lemma 12.6 (Holonomy of a Nearly Flat Field)**.** *Let U be a G-valued field on a periodic lattice of spacing η, let B be a box of side $D\eta$ in it, with $D \geq 1$ an integer smaller than every period in lattice units, and let C be a closed path of P bonds in B. If $\|U(\partial p) - 1\| \leq t$ for every plaquette p of B and $s := 3DPt \leq \min(1/10, \log(1 + r_G))$, then for every irreducible unitary representation π of G with the character χ,*

$$\|\pi(U(C)) - 1\| \leq 1.06\, c_\pi s \qquad \text{and} \qquad |\chi(U(C)) - 1| \leq 1.06\, c_\pi s \,. \tag{12.21}$$

Proof. Let o be the corner of B with the smallest coordinates, and for a site y of B let γ_y be the path from o to y which changes first the coordinate 0, then 1, 2, and 3. For a bond $b = (y, y')$ of B put $\ell_b := \gamma_y\, b\, \gamma_{y'}^{-1}$, a closed path at o, and $U^\gamma(b) := U(\ell_b)$. Then U^γ is the gauge transform of U by $y \mapsto U(\gamma_y)^{-1}$, so that $U^\gamma(C)$ is conjugate to $U(C)$ and $\chi(U^\gamma(C)) = \chi(U(C))$.

We first bound $U^\gamma(b)$. It suffices to take b in a positive direction e_ν, since $U^\gamma(b^{-1}) = U^\gamma(b)^{-1}$. Let p be the path formed by the legs of γ_y in the directions $\nu' > \nu$, a path of $n \leq 3D$ bonds from a site z to y. Then $\gamma_y = \gamma_z\, p$ and $\gamma_{y'} = \gamma_z\, e\, (p + e_\nu)$, with e the bond from z to $z + e_\nu$, so that $\ell_b = \gamma_z\, p\, b\, (p + e_\nu)^{-1} e^{-1} \gamma_z^{-1}$, and $U(\ell_b)$ is conjugate to the holonomy H of the boundary of the strip swept by p in the direction e_ν. With $p = (p_1, \dots, p_n)$ and q_l

the plaquette spanned by p_l and e_ν, telescoping along the strip writes $H = \lambda_n \cdots \lambda_1$, where λ_l is the holonomy of q_l, from one of its corners and in one of its orientations, conjugated by the transport along $p_1 \cdots p_{l-1}$. Each λ_l satisfies $\|\lambda_l - 1\| = \|U(\partial q_l) - 1\| \le t$, since the operator norm of $W-1$ is invariant under unitary conjugation and inversion, and (8.11) gives

$$\|U^\gamma(b) - 1\| = \|H - 1\| \le (1+t)^{3D} - 1 \,. \tag{12.22}$$

The path C has P bonds, and (8.11) applied to $U^\gamma(C) = \prod_{b\in C} U^\gamma(b)$ gives $\|U^\gamma(C)-1\| \le (1+t)^{3DP}-1 \le e^s - 1$. Since $s \le \min(1/10, \log(1+r_G))$, the principal logarithm $U^\gamma(C) = e^{iX}$ has $X \in \mathfrak{g}$ and $\|X\| \le 1.06\, s$ by (8.12), so that $\|\pi(U^\gamma(C))-1\| \le c_\pi \|X\|$ by (1.7). Since $\pi(U^\gamma(C))$ is conjugate to $\pi(U(C))$ by a unitary matrix, $\|\pi(U(C)) - 1\| = \|\pi(U^\gamma(C)) - 1\| \le 1.06\, c_\pi s$. Finally $|\chi(W) - 1| = |(\dim \pi)^{-1} \operatorname{Tr}(\pi(W) - 1)| \le \|\pi(W) - 1\|$. *q.e.d.*

At the stopping step we apply the lemma to the background field. For a loop C at level q with the character of π put

$$y(C) := 3.18\, c_\pi c_h (1 + 2\kappa_s) B_3\, P_{K'}(C)^2\, \varepsilon_{K'} \,, \tag{12.23}$$

which does not depend on K, since $P_{K'}(C)$ and $\varepsilon_{K'}$ do not (Section 8).

Proof of Proposition 12.5. We first bound the leading words on ℓ_0. On the support of ϱ_{ℓ_0} every fine plaquette of the background satisfies $\|U_{K'}(\partial p) - 1\| \le t := 2c_h(1+2\kappa_s) B_3 \varepsilon_{K'} L^{-2K'}$, as in the proof of Corollary 8.3. In fine units the loop C_i has $P := P_{K'}(C_i) L^{K'}$ bonds, and it lies in a box of side $D \le P/2$, since a closed path of P bonds has ℓ^∞-diameter at most $P/2$; the box embeds in the torus because $3\,\mathrm{diam}(C_i) < T$. Hence $3DPt \le 3c_h(1 + 2\kappa_s) B_3 P_{K'}(C_i)^2 \varepsilon_{K'}$, and Lemma 12.6 gives $|f^{(i)}_{K'} - 1| \le 1.06\, c_{\pi_i} \cdot 3DPt \le y(C_i)$ whenever $3c_h(1 + 2\kappa_s) B_3 P_{K'}(C_i)^2 \varepsilon_{K'} \le \min(1/10, \log(1 + r_G))$. Since $|f^{(i)}_{K'}| \le 1$, telescoping gives $|\prod_i f^{(i)}_{K'} - 1| \le \sum_i y(C_i)$.

Next we bound the term ℓ_0 of the identity (8.5) from below. On ℓ_0 the normalised dressed loop is $e^{E_{\mathrm{const},K'}} [\prod_i f^{(i)}_{K'} + \sum_{\mathfrak{B}\neq\emptyset} Q_{K'}(\mathfrak{B}) w_{K'}(\mathfrak{B})]$, where $|E_{\mathrm{const},K'}| \le B_E(\mathcal{C})$ by Definition 5.2(iv) and Corollary 7.11, and the sum over the families has modulus at most $e^{I_*(\mathcal{C})} - 1$ by Lemma 8.6. Hence

$$|\langle \widehat{\mathcal{W}}_{\ell_0} \rangle_{\ell_0}| \ge e^{-B_E(\mathcal{C})} \Big(2 - e^{I_*(\mathcal{C})} - \sum_i y(C_i) \Big) \,. \tag{12.24}$$

The other terms are bounded as in the proof of Theorem 8.7: $\sum_{\ell\neq\ell_0} w_\ell |\langle \widehat{\mathcal{W}}_\ell \rangle_\ell| \le e^{B_E(\mathcal{C}) + I_*(\mathcal{C})} \bar\delta_{\mathrm{LF}}$, and $w_{\ell_0} \ge 1 - \bar\delta_{\mathrm{LF}}$ by Proposition 8.4. By (8.5) and the triangle inequality, for every $K \ge K_0(\mathcal{C})$,

$$|W_K(\mathcal{C})| \ge (1 - \bar\delta_{\mathrm{LF}}) e^{-B_E(\mathcal{C})} \Big(2 - e^{I_*(\mathcal{C})} - \sum_i y(C_i) \Big) - e^{B_E(\mathcal{C}) + I_*(\mathcal{C})} \bar\delta_{\mathrm{LF}} \,. \tag{12.25}$$

It remains to choose $\bar g_1$ and ℓ so that the four quantities $B_E(\mathcal{C})$, $I_*(\mathcal{C})$, $\sum_i y(C_i)$, and $\bar\delta_{\mathrm{LF}}$ are small. Each of them is at most a sum over the loops of a number of the loop, multiplied by a constant which is a nondecreasing function of $c_{\mathcal{C}}$ and by a function of $\bar g$ which tends to zero, at fixed q and T/u, except for $\bar\delta_{\mathrm{LF}}$, which does not depend on the ensemble. For $B_E(\mathcal{C})$ this is (7.32), with the constants C'_E and C''_E, since $\bar{\mathfrak{a}} \le C \bar g^2 (\log \bar g^{-2})^{6p_0}$ by (7.4) and $S^{\mathbf{R}}(\bar g) \to 0$; for $\bar\delta_{\mathrm{LF}}$ it is the remark after Proposition 8.4. For the other two we note that $\hat g^{-2}_{K'} = \bar g^{-2} + b' j_q + C_W$ by (3.3), so that $\hat g_{K'} \le \bar g$, and $\log \hat g^{-2}_{K'} \le 2 \log \bar g^{-2}$ for $\bar g$ small, because

j_q grows only like $\log\log \bar g^{-2}$ by (3.44). The numbers $P_{K'}(C) = P_q(C) L^3 M R_{K'-1}$ are at most a constant times $P_q(C)(\log \hat g_{K'}^{-2})^r$ by Lemma 3.4; by Definition 3.11, $\varepsilon_{K'} = \hat g_{K'} p_0(\hat g_{K'})$ is $\hat g_{K'}$ times a power of $\log \hat g_{K'}^{-2}$, and $\varepsilon_{K'}/\min(\alpha_{0,K'}, \alpha_{1,K'})$ is a power of $\log \hat g_{K'}^{-2}$; the number $\bar\rho$ of (8.36) is $\bar g^2$ times a power of $\log \bar g^{-2}$; and $S_{\mathrm{wr}}(b) \le e^{-b}(1 + 2/b)$ by Lemma 2.11 with $x = 1$. The cross total (8.38) is at most $\sum_i P_{K'}(C_i)$ times $10\, C_{\mathrm{cr}} \bar{\mathfrak{a}} + 2 C_{\mathrm{cr}}^{\mathbf{R}} S^{\mathbf{R}}(\bar g)$, which tends to zero with $\bar g$ as in the bound on $B_E(\mathcal{C})$. Hence, apart from the factors $P_q(C_i)$, c_{π_i}, and the constants C_χ and C of (5.11), the numbers (12.23) and (8.37) are at most a positive power of $\bar g$ times a power of $\log \bar g^{-2}$. Collecting, there are functions h_E, h_I, and h_y on $(0, \gamma_0]$ which tend to zero as $\bar g \to 0$, and nondecreasing functions F_E and F_I on $[0, \infty)$, all of which depend only on the constants of the construction, Λ_*, q, and T/u, such that

$$\begin{aligned} B_E(\mathcal{C}) &\le F_E(c_{\mathcal{C}}) h_E(\bar g) \sum_i \Big(\frac{P(C_i)}{r(C_i)} + P_q(C_i) \Big) , \\ I_*(\mathcal{C}) &\le F_I(c_{\mathcal{C}}) h_I(\bar g) \sum_i P_q(C_i) , \qquad y(C_i) \le c_{\pi_i} h_y(\bar g) P_q(C_i)^2 , \end{aligned} \tag{12.26}$$

and such that $3 c_h (1 + 2\kappa_s) B_3 P_{K'}(C_i)^2 \varepsilon_{K'} \le h_y(\bar g) P_q(C_i)^2$. We let $\bar g_1 \in (0, \gamma_0]$ be so small that $\bar\delta_{\mathrm{LF}} \le 1/20$ for $\bar g \le \bar g_1$, put $m_G := \min(1/10, \log(1 + r_G))$, and define $\ell(c, \bar g) := 0$ if $q < q_\sharp(c, \bar g)$ and otherwise

$$\ell(c, \bar g) := \min\Big(\frac{1}{10 F_E(c) h_E(\bar g)}, \ \frac{1}{20 F_I(c) h_I(\bar g)}, \ \frac{1}{20\, c\, h_y(\bar g)}, \ \frac{m_G}{h_y(\bar g)} \Big) , \tag{12.27}$$

with the convention $1/0 := \infty$. Each entry of the minimum does not increase with c, and neither does the indicator of $q \ge q_\sharp(c, \bar g)$, since $q_\sharp$ is nondecreasing in c; so ℓ does not increase with c. For fixed c we have $q_\sharp(c, \bar g) = 0 \le q$ for $\bar g$ small, by Theorem 1.5, and every entry of the minimum tends to infinity as $\bar g \to 0$; so $\ell(c, \bar g) \to \infty$.

Let now $\bar g \le \bar g_1$ and let $\mathcal{C}$ satisfy (12.19). A closed path which visits no site twice has at least four bonds, so that the left side of (12.19) is at least 16; hence $\ell(c_{\mathcal{C}}, \bar g) > 0$, which by definition means $q \ge q_\sharp(c_{\mathcal{C}}, \bar g)$, and Theorem 1.5 gives the limit $W(\mathcal{C})$. Since $P_q(C_i) \le P_q(C_i)^2$, the bounds (12.26) and (12.27) give $B_E(\mathcal{C}) \le 1/10$, $I_*(\mathcal{C}) \le 1/20$, $\sum_i y(C_i) \le c_{\mathcal{C}} h_y(\bar g) \sum_i P_q(C_i)^2 \le 1/20$, and the condition of Lemma 12.6 for every loop, and $\bar\delta_{\mathrm{LF}} \le 1/20$ by the choice of $\bar g_1$. Then (12.25) gives

$$|W_K(\mathcal{C})| \ge 0.95\, e^{-0.1}(2 - e^{0.05} - 0.05) - 0.05\, e^{0.15} \ge 0.71 \ge \frac{1}{2} , \tag{12.28}$$

and in the limit $|W(\mathcal{C})| \ge 1/2$. For a loop C at level q in H_+ for which $\mathcal{D} := ((C, \chi))$ satisfies (12.19), we have $q \ge q_\sharp(c_\pi, \bar g)$, as just shown, and Corollary 1.7 with $m = 1$, $\mathcal{D}_1 = \mathcal{D}$, and $c = c_\pi$ gives $W(\mathcal{D} \cup \theta\bar{\mathcal{D}}) - |W(\mathcal{D})|^2 \ge 0$, so that $W(\mathcal{D} \cup \theta\bar{\mathcal{D}}) \ge |W(\mathcal{D})|^2 \ge 1/4$. *q.e.d.*

The same quantities give an upper bound which factorises over the loops: each loop contributes a factor which is exponential in its length in units of $\mathbb{T}_q$ and in its ratio P/r, with a rate which tends to zero with the coupling, and the number of loops enters only through the number of factors.

Proposition 12.7 (Growth Bound)**.** *Let L, Λ_*, and $q \ge 0$ be as in Proposition 12.5, and let $\bar g_1$, F_E, F_I, h_E, and h_I be as in its proof. Put $a(c, \bar g) := F_E(c) h_E(\bar g) + F_I(c) h_I(\bar g)$, a function which is nondecreasing in c and tends to zero as $\bar g \to 0$ for every c. For every $\bar g \in (0, \bar g_1]$ and every loop ensemble $\mathcal{C}$ at level $q \ge q_\sharp(c_{\mathcal{C}}, \bar g)$ whose loops are admissible, and for every $K \ge K_0(\mathcal{C})$,*

$$|W_K(\mathcal{C})| \le \frac{21}{20} \prod_{i=1}^{n} \exp\Big(a(c_{\mathcal{C}}, \bar g) \Big(\frac{P(C_i)}{r(C_i)} + P_q(C_i) \Big) \Big) , \tag{12.29}$$

and the limit $W(\mathcal{C})$ satisfies the same bound.

Proof. By Theorem 8.7, $|W_K(\mathcal{C})| \le e^{B_E(\mathcal{C})+I_*(\mathcal{C})}(1+\bar{\delta}_{\mathrm{LF}})$, and $\bar{\delta}_{\mathrm{LF}} \le 1/20$ for $\bar{g} \le \bar{g}_1$ by the choice of $\bar{g}_1$. The bounds (12.26) on $B_E(\mathcal{C})$ and $I_*(\mathcal{C})$ were derived in the proof of Proposition 12.5 from the definitions (7.32) and (8.37) alone, without the condition (12.19), and $P_q(C_i) \le P(C_i)/r(C_i) + P_q(C_i)$ gives $B_E(\mathcal{C}) + I_*(\mathcal{C}) \le a(c_{\mathcal{C}}, \bar{g}) \sum_i (P(C_i)/r(C_i) + P_q(C_i))$. The limit exists by Theorem 1.5, and the bound passes to it. *q.e.d.*

Since $P_q(C) u L^{-q}$ is the length of C, the bound (12.29) is a perimeter bound with the rate $a(c_{\mathcal{C}}, \bar{g}) L^q/u$ per unit length, together with a factor for the shape of each loop; with Proposition 12.5 it brackets the limits between $\frac{1}{2}$ and this bound on the ensembles of short loops. Continuity of W in the positions of the loops is not asserted, since the limits are defined on the loops of the lattices $\mathbb{T}_q$ only.

The condition (12.19) involves the number of loops, because the bounds on $B_E(\mathcal{C})$, $I_*(\mathcal{C})$, and $\sum_i y(C_i)$ are bounds on sums over the loops. For ensembles which are generated from a fixed one by reflections this dependence disappears. We say that an ensemble $\mathcal{D}$ at level q is *generated by reflections from* an ensemble $\mathcal{B}$ if $\mathcal{D} = \mathcal{B}$, or if $\mathcal{D} = \mathcal{D}' \cup \theta_{\mu,a}\bar{\mathcal{D}}'$ for an ensemble $\mathcal{D}'$ which is generated by reflections from $\mathcal{B}$ and all of whose loops lie in $H_{\mu,a}$, for some μ and some $a \in uL^{-q}\mathbb{Z}$. The *blocks* of $\mathcal{B}$ are $\{\mathcal{B}\}$, and those of $\mathcal{D}' \cup \theta_{\mu,a}\bar{\mathcal{D}}'$ are the blocks of $\mathcal{D}'$ and their images under $\theta_{\mu,a}$ with the conjugate characters. Every block is the image of $\mathcal{B}$ under a composition of reflections, with all its characters conjugated or none, so that it has the same chord–arc constants, ratios P/r, and numbers P_q as $\mathcal{B}$, and either the characters of $\mathcal{B}$ or their conjugates. In particular $c_{\mathcal{B}_i} = c_{\mathcal{B}} = c_{\mathcal{D}}$ for every block $\mathcal{B}_i$ of $\mathcal{D}$, because $c_{\bar{\pi}} = c_\pi$, and every block satisfies (12.19) if $\mathcal{B}$ does.

Proposition 12.8 (Products over Reflections)**.** *Let L and Λ_* be as in Theorem 1.5, let $\mathcal{B}$ be a loop ensemble at level q whose loops are admissible, with $\bar{g}$ and q as in Theorem 1.5 for $\mathcal{B}$, and let $\mathcal{D}$ be generated by reflections from $\mathcal{B}$, with the blocks $\mathcal{B}_1, \dots, \mathcal{B}_N$. Then $|W_K(\mathcal{D})| \ge \prod_{i=1}^N |W_K(\mathcal{B}_i)|$ for every $K > j_q$. If $\bar{g} \le \bar{g}_1$ and $\mathcal{B}$ satisfies* (12.19)*, then $|W(\mathcal{D})| \ge 2^{-N}$; the condition does not involve $\mathcal{D}$, and in particular not its number of loops.*

Proof. We argue by induction along the generation. For $\mathcal{D} = \mathcal{B}$ there is nothing to prove. Let $\mathcal{D} = \mathcal{D}' \cup \theta_{\mu,a}\bar{\mathcal{D}}'$, with the loops of $\mathcal{D}'$ in $H_{\mu,a}$, and write $\Theta := \Theta_{\mu,a}$ and $F := W_{\mathcal{D}'} \in \mathcal{A}_+$. Then $W_{\mathcal{D}} = F\,\Theta F$ by Section 12.1, and (12.4) with Lemma 12.2 gives

$$|\mathbb{E}_K[W_{\mathcal{D}}]| = \mathbb{E}_K[F\,\Theta F] \ge |\mathbb{E}_K[F]|^2 = |\mathbb{E}_K[W_{\mathcal{D}'}]|\,|\mathbb{E}_K[W_{\theta_{\mu,a}\bar{\mathcal{D}}'}]|\,. \tag{12.30}$$

The normalisation of $\mathcal{D}$ is the product of those of $\mathcal{D}'$ and $\theta_{\mu,a}\bar{\mathcal{D}}'$, and dividing by its modulus gives $|W_K(\mathcal{D})| \ge |W_K(\mathcal{D}')||W_K(\theta_{\mu,a}\bar{\mathcal{D}}')|$. By the induction hypothesis $|W_K(\mathcal{D}')|$ is at least the product over the blocks of $\mathcal{D}'$; and since $\theta_{\mu,a}\bar{\mathcal{D}}'$ is generated by reflections from $\theta_{\mu,a}\bar{\mathcal{B}}$, with the reflections conjugated by $\theta_{\mu,a}$, and has the images of the blocks of $\mathcal{D}'$ as its blocks, the same holds for it. This proves the first assertion. If $\bar{g} \le \bar{g}_1$ and $\mathcal{B}$ satisfies (12.19), then so does every block, so that $|W_K(\mathcal{B}_i)| \ge 1/2$ for $K \ge K_0(\mathcal{B}_i)$ by Proposition 12.5, and the limit gives $|W(\mathcal{D})| \ge 2^{-N}$; the limit exists by Theorem 1.5, because the loops of $\mathcal{D}$ are admissible and $c_{\mathcal{D}} = c_{\mathcal{B}}$, so that $\bar{g}$ and q are as in Theorem 1.5 for $\mathcal{D}$. *q.e.d.*

The lower bound decays with the number of blocks, but it is positive at a coupling which is fixed once $\mathcal{B}$ is. The number of loops which can be generated in this way is limited only by the torus: starting from one loop of diameter h and doubling in each direction at a hyperplane at distance h from the current configuration, the extent in the direction μ after k_μ doublings

is at most $2^{k_\mu} \cdot 3h - 2h$, so that of the order of $4 \log_2(T/h)$ doublings fit, and the number of loops is of the order of $(T/h)^4$.

13 Non-Gaussianity

We now prove Theorem 1.8. The proof computes the number $\mathfrak{Y}$ of (1.13) in two ways. For a Gaussian charge model it vanishes identically, by an algebraic cancellation which uses nothing but two facts: $\log|W|$ is a quadratic form in the current of the loop, linear in the covariance, and the logarithm of a normalisation is a linear function of the lengths and the corners (Section 13.2). For the limits of Yang–Mills theory we compute $\mathfrak{Y}$ to the order $\bar{g}^2$ at the stopping scale, where the identity of Theorem 8.1 expresses W_K through a measure on the fixed lattice $\mathbb{T}_{j_q}$ which, at small coupling, is concentrated near the flat fields.

Three things have to be shown, and each meets a difficulty of its own. The normalisations have to drop out of $\mathfrak{Y}$, which they do because $\mathfrak{Y}$ was built to annihilate every function of the form of Proposition 7.13 (Section 13.3). The pair terms have to be of the fourth order. Under the measure of the stopping scale the Wilson loop of each of two separated rectangles depends, up to corrections which decay with their distance, on the field near that rectangle only, and its deviation from one has an L^2-norm of the order $\bar{g}^2$ rather than $\bar{g}$, because its part of first order is a trace of an element of the Lie algebra and vanishes; the covariance of two such deviations is then of the order $\bar{g}^4$ (Section 13.5). Finally the third difference of the rectangles has to be bounded below by a positive multiple of $\bar{g}^2$, and this is the heart of the matter. After the insertions are shown to cancel in the third difference, by the translation invariance which made the coefficients of the normalisation independent of the loop (Section 13.6), the third difference is a combination of the second moments of the fluxes of the background field through the rectangles I_a. An integration by parts against the density of the stopping scale identifies these second moments, to the leading order, with a quadratic form evaluated at a flat field near the given one (Section 13.8). At a flat field this form is the covariance of the fluxes of the block field of a free field, which differs from the covariance of the free field itself by a term that decays on the scale of the blocks (Section 13.7), and the third difference of the free covariances is computed explicitly (Section 13.4).

Throughout the section q, the plane spanned by e_μ and e_ν, t, s, o, and π are fixed as in Section 1.3, with $\Lambda_* \geq 1 + \max(s/t, 4t/s)$ and L as in Theorem 1.5, and the eleven diagrams are those defined there. Every requirement in this section that $\bar{g}$ be small is an upper bound on $\bar{g}$ in terms of q, t/u, s/u, T/u, π, and the constants of the construction, and $\bar{g}_0$ in Theorem 1.8 is the least of these bounds.

13.1 The Witness

For an ensemble $\mathcal{D}$ we write $|\mathcal{D}|$ for the number of its loops, $N_v(\mathcal{D}) \coloneqq \sum_{C \in \mathcal{D}} N_v(C)$ for the total length of its segments of orientation $v \in \mathcal{V}$, and, for a perpendicular pair (v, w), $K_{v,w}(\mathcal{D})$ for the number of the corners c of its loops with $(v_c, w_c) = (v, w)$. The number (1.13) is

$$\mathfrak{Y}(W) = \sum_{\mathcal{D}} n_{\mathcal{D}} \log|W(\mathcal{D})| = \mathfrak{y}_S(W) - \mathfrak{y}_2(W) + \mathfrak{y}_3(W) \,, \tag{13.1}$$

the sum over the six diagrams R_1, R_2, R_3, R_4, P_2, P_3 with the weights $n_{\mathcal{D}} = 1, -3, 3, -1, -1, 1$, where $\mathfrak{y}_S(W) \coloneqq \sum_{n=1}^{4} n_{R_n} \log|W(R_n)|$ is the *shape term* and $\mathfrak{y}_k(W) \coloneqq \log|W(P_k)/W(R_1)^2|$ are the pair terms. To each diagram we assign the set $\mathrm{ind}(\mathcal{D})$ of the indices a of the rectangles I_a which tile it, or which are its loops:

$\mathrm{ind}(I_a) \coloneqq \{a\}$, $\mathrm{ind}(R_n) \coloneqq \{0,\dots,n-1\}$, $\mathrm{ind}(R_n^{(1)}) \coloneqq \{1,\dots,n\}$, and $\mathrm{ind}(P_k) \coloneqq \{0,k\}$. The Gaussian models are tested by a second set of weights on all eleven diagrams,

$$n_{\mathcal{D}}^{\mathrm{sym}} \coloneqq \begin{cases} 0,1,1,-1 & \text{for } \mathcal{D} = I_0, I_1, I_2, I_3 \ , \\ -1,-2,2,1 & \text{for } \mathcal{D} = R_2, R_2^{(1)}, R_3, R_3^{(1)} \ , \\ -1,-1,1 & \text{for } \mathcal{D} = R_4, P_2, P_3 \ , \end{cases} \tag{13.2}$$

and we put $\mathfrak{Y}^{\mathrm{sym}}(W) \coloneqq \sum_{\mathcal{D}} n_{\mathcal{D}}^{\mathrm{sym}} \log|W(\mathcal{D})|$, the sum over the eleven diagrams.

Lemma 13.1 (Counts and Index Sums)**.** *(i) A rectangle of width at and height s among the nine has $N_{\pm e_\mu} = at$, $N_{\pm e_\nu} = s$, and $N_v = 0$ for the other v, and it has exactly one corner of each of the types (e_μ, e_ν), $(e_\nu, -e_\mu)$, $(-e_\mu, -e_\nu)$, $(-e_\nu, e_\mu)$ and no other corner.*

(ii) Let f be a function on the eleven diagrams such that $f(\mathcal{D})$ depends only on $\mathcal{D}$ through $|\mathcal{D}|$, the numbers $N_v(\mathcal{D})$, and the numbers $K_{v,w}(\mathcal{D})$, and linearly. Then $\sum_{\mathcal{D}} n_{\mathcal{D}} f(\mathcal{D}) = 0$, $\sum_n n_{R_n} f(R_n) = 0$, $f(P_k) = 2f(R_1)$, and $\sum_{\mathcal{D}} n_{\mathcal{D}}^{\mathrm{sym}} f(\mathcal{D}) = 0$.

(iii) If $f(I_a) = f(I_0)$ for every a and $f(R_n^{(1)}) = f(R_n)$ for $n \in \{2,3\}$, then $\sum_{\mathcal{D}} n_{\mathcal{D}}^{\mathrm{sym}} f(\mathcal{D}) = \sum_{\mathcal{D}} n_{\mathcal{D}} f(\mathcal{D})$, the left sum over the eleven diagrams and the right one over the six.

(iv) For $0 \le a \le b \le 3$, the sum $\mathsf{s}(a,b)$ of $n_{\mathcal{D}}^{\mathrm{sym}}$ over the diagrams with $a, b \in \mathrm{ind}(\mathcal{D})$ vanishes.

Proof. (i) The rectangle is traversed from its corner of least coordinates in the directions e_μ, e_ν, $-e_\mu$, $-e_\nu$, in this order, which gives the four corners and the lengths.

(iii) The weights n^{sym} sum to $0+1+1-1 = 1$ over $I_0,\dots,I_3$, to $-1-2 = -3$ over R_2, $R_2^{(1)}$, and to $2+1 = 3$ over R_3, $R_3^{(1)}$, which are the weights n of R_1, R_2, and R_3, and they agree with n on R_4, P_2, P_3.

(ii) It suffices to take for f each of the functions $|\mathcal{D}|$, $N_v(\mathcal{D})$, $K_{v,w}(\mathcal{D})$. On the six diagrams $R_1, R_2, R_3, R_4, P_2, P_3$, by (i), $|\mathcal{D}| = 1,1,1,1,2,2$, $N_{\pm e_\mu}(\mathcal{D}) = t, 2t, 3t, 4t, 2t, 2t$, $N_{\pm e_\nu}(\mathcal{D}) = s, s, s, s, 2s, 2s$, the other N_v vanish, and $K_{v,w}(\mathcal{D}) = |\mathcal{D}|$ for the four types of (i) and 0 for the others. Hence

$$\sum_{\mathcal{D}} n_{\mathcal{D}} |\mathcal{D}| = 1-3+3-1-2+2 = 0 \ , \qquad \sum_{\mathcal{D}} n_{\mathcal{D}} N_{\pm e_\mu}(\mathcal{D}) = (1-6+9-4-2+2)t = 0 \ , \tag{13.3}$$

and $\sum_{\mathcal{D}} n_{\mathcal{D}} N_{\pm e_\nu}(\mathcal{D}) = (1-3+3-1-2+2)s = 0$. On the rectangles alone the same sums are $1-3+3-1 = 0$, $(1-6+9-4)t = 0$, and 0. The loops of P_k have the lengths and the corners of R_1, which gives $f(P_k) = 2f(R_1)$. Finally f takes the same value on a rectangle and on its translates, so that (iii) and the first assertion give the last one.

(iv) The diagrams which contain a given pair of indices, with their weights n^{sym}, are: for $(0,0)$, I_0, R_2, R_3, R_4, P_2, P_3, with $0-1+2-1-1+1$; for $(1,1)$, I_1, R_2, $R_2^{(1)}$, R_3, $R_3^{(1)}$, R_4, with $1-1-2+2+1-1$; for $(2,2)$, I_2, $R_2^{(1)}$, R_3, $R_3^{(1)}$, R_4, P_2, with $1-2+2+1-1-1$; for $(3,3)$, I_3, $R_3^{(1)}$, R_4, P_3, with $-1+1-1+1$; for $(0,1)$, R_2, R_3, R_4, with $-1+2-1$; for $(0,2)$, R_3, R_4, P_2, with $2-1-1$; for $(0,3)$, R_4, P_3, with $-1+1$; for $(1,2)$, $R_2^{(1)}$, R_3, $R_3^{(1)}$, R_4, with $-2+2+1-1$; and for $(1,3)$ and $(2,3)$, $R_3^{(1)}$, R_4, with $1-1$. Each sum is zero. *q.e.d.*

13.2 Gaussian Charge Models

The Gaussian charge models of Section 1.3 fail this criterion for an algebraic reason, which the next proposition records.

Proposition 13.2 (Gaussian Charge Models)**.** *For every Gaussian charge model, its function W does not vanish on the eleven diagrams and $\mathfrak{Y}^{\mathrm{sym}}(W) = 0$. If W_m are the functions of a sequence of Gaussian charge models, each with its own charge, Gaussian vector, and normalisation, and $W_m(\mathcal{D}) \to W(\mathcal{D}) \neq 0$ for each of the eleven diagrams, then $\mathfrak{Y}^{\mathrm{sym}}(W) = 0$.*

Proof. For a real Gaussian variable Z we have $|\mathbb{E}[e^{iZ}]| = e^{-\operatorname{Var} Z/2}$. Hence $W(\mathcal{D}) \neq 0$ and

$$\log|W(\mathcal{D})| = -\frac{\mathsf{g}^2}{2} \sum_{a,b \in \mathrm{ind}(\mathcal{D})} \operatorname{Cov}(\mathsf{X}_a, \mathsf{X}_b) - \sum_{C \in \mathcal{D}} \log|\kappa(C)| \; . \tag{13.4}$$

We multiply by $n_{\mathcal{D}}^{\mathrm{sym}}$ and sum over the eleven diagrams. In the first term the coefficient of $\operatorname{Cov}(\mathsf{X}_a, \mathsf{X}_b)$, $a \leq b$, is $-\frac{1}{2}\mathsf{g}^2$ or $-\mathsf{g}^2$ times $\mathsf{s}(a,b)$, which vanishes by Lemma 13.1(iv). The second term is a function of $\mathcal{D}$ which is linear in the numbers $N_v(\mathcal{D})$ and $K_{v,w}(\mathcal{D})$, by the definition of a Gaussian charge model, so that its weighted sum vanishes by Lemma 13.1(ii). Hence $\mathfrak{Y}^{\mathrm{sym}}(W) = 0$. For the limit, $\mathfrak{Y}^{\mathrm{sym}}(W) = \lim_m \mathfrak{Y}^{\mathrm{sym}}(W_m) = 0$, since $\log|\cdot|$ is continuous on $\mathbb{C} \setminus \{0\}$. *q.e.d.*

The cancellation can be read off in the models whose covariance is invariant under the translation by te_μ, $\operatorname{Cov}(\mathsf{X}_a, \mathsf{X}_b) = c^{\mathrm{G}}_{|a-b|}$. There $\operatorname{Var} \mathsf{X}(R_n) = n c_0^{\mathrm{G}} + 2\sum_{1 \leq k < n} (n-k) c_k^{\mathrm{G}}$ and $\operatorname{Var} \mathsf{X}(P_k) = 2c_0^{\mathrm{G}} + 2c_k^{\mathrm{G}}$, so that, the normalisations cancelling by Lemma 13.1(ii),

$$\mathfrak{y}_S(W) = \mathsf{g}^2(c_3^{\mathrm{G}} - c_2^{\mathrm{G}}) \; , \qquad \mathfrak{y}_k(W) = -\mathsf{g}^2 c_k^{\mathrm{G}} \; , \qquad \mathfrak{Y}(W) = \mathfrak{Y}^{\mathrm{sym}}(W) = 0 \; : \tag{13.5}$$

the covariances of the separated rectangles which make up the shape term also make up the pair terms.

Remark 13.3 (Gaussian Fields with Values in the Lie Algebra). The models of Proposition 13.2 are abelian, and the number $\mathfrak{Y}$ does not distinguish Yang–Mills theory from Gaussian fields with values in $\mathfrak{g}$ and path-ordered holonomies. Let (T_α) be a basis of $\mathfrak{g}$ which is orthonormal for an Ad-invariant inner product, with $\sum_\alpha \pi_*(T_\alpha)^2 = C \cdot 1$, let $B = \sum_\alpha B^\alpha T_\alpha$ on the bonds of a lattice containing the rectangles, with B^α real, centred, and jointly Gaussian, $\operatorname{Cov}(B^\alpha(e), B^\beta(f)) = \delta^{\alpha\beta} G(e,f)$, and G invariant under the translation by te_μ, and let $W(\mathcal{D}) \coloneqq \mathbb{E}[\prod_{C \in \mathcal{D}} \chi(\prod_{e \in C} e^{i\mathsf{g}B(e)})]$. Expanding the ordered product to the second order in g, the term of first order of $\chi(\prod_e e^{i\mathsf{g}B(e)})$ is $i\mathsf{g}(\dim\pi)^{-1} \operatorname{Tr} \pi_*(\sum_e B(e))$, which vanishes because $\operatorname{Tr} \circ \pi_*$ vanishes on $\mathfrak{g}$ (Lemma 13.6 below), the term of second order has the mean $-\frac{1}{2}\mathsf{g}^2 C \operatorname{Var} \mathsf{X}^1(C)$ with $\mathsf{X}^1(C) \coloneqq \sum_{e \in C} B^1(e)$, the terms of odd order have mean zero, and the remainder of fourth order has a finite moment. Hence the pair terms are $O(\mathsf{g}^4)$ and $\mathfrak{Y}(W) = \mathsf{g}^2 C(c_3 - c_2) + O(\mathsf{g}^4)$ with $c_k \coloneqq \operatorname{Cov}(\mathsf{X}^1(I_0), \mathsf{X}^1(I_k))$, which for the free covariance is positive by Lemma 13.5 below. These fields share with Yang–Mills theory the feature on which the theorem rests.

13.3 The Normalisations Drop Out

We next show that the limits exist on the eleven diagrams and that $\mathfrak{Y}(W)$ is the limit of the same number computed from the expectations $N_K(\mathcal{D}) \coloneqq \mathbb{E}_K[W_{\mathcal{D}}]$, in which no normalisation appears.

Proposition 13.4 (The Normalisations Drop Out)**.** *(i) Let $K > j_q$. If $N_K(\mathcal{D}) \neq 0$ on the six diagrams $R_1, \ldots, R_4, P_2, P_3$, then each of $\mathfrak{Y}$, $\mathfrak{y}_S$, $\mathfrak{y}_2$, and $\mathfrak{y}_3$ takes the same value on W_K and on N_K.*

(ii) There is $\bar g_1' > 0$ *such that for* $0 < \bar g \le \bar g_1'$ *the limits* $W(\mathcal{D})$ *exist on the eleven diagrams,* $|W_K(\mathcal{D})| \ge 1/2$ *for* $K \ge K_0(\mathcal{D})$, $|W(\mathcal{D})| \ge 1/2$, $W_K(I_a) = W_K(I_0)$ *and* $W_K(R_n^{(1)}) = W_K(R_n)$ *for* $K > j_q$, *and* $W(I_a) = W(I_0)$ *and* $W(R_n^{(1)}) = W(R_n)$. *For these* $\bar g$, $\mathfrak{Y}^{\mathrm{sym}}(W) = \mathfrak{Y}(W) = \lim_K \mathfrak{Y}(N_K)$, $\mathfrak{y}_S(W) = \lim_K \mathfrak{y}_S(N_K)$, *and* $\mathfrak{y}_k(W) = \lim_K \mathfrak{y}_k(N_K)$.

(iii) Let $\bar g \le \bar g_1'$ *and* $q' \ge q$. *If the limits* $W^{(q')}$ *of the renormalised expectations with the normalisation of the level* q' *exist and do not vanish on the six diagrams, then* $\mathfrak{Y}(W^{(q')}) = \mathfrak{Y}(W)$.

Proof. (i) By Proposition 7.13, $\log|\kappa_K^{(q)}(C,\chi)| = \sum_v N_v^{(0)}(C)\,\mathrm{Re}\, m_{v,K} + \sum_{c\in\mathcal{K}(C)} \mathrm{Re}\, z_{v_c,w_c,K}$, where $N_v^{(0)}(C) = N_v(C)/\varepsilon_K$ is the number of bonds of $\mathbb{T}_K$ of orientation v on C and the coefficients do not depend on C. Hence $\sum_{C\in\mathcal{D}} \log|\kappa_K^{(q)}(C,\chi)|$ is a linear function of the numbers $N_v(\mathcal{D})$ and $K_{v,w}(\mathcal{D})$, and by Lemma 13.1(ii) its weighted sums with the weights of $\mathfrak{Y}$ and of $\mathfrak{y}_S$ vanish, and its value on P_k is twice its value on R_1. Since $\log|W_K(\mathcal{D})| = \log|N_K(\mathcal{D})| - \sum_{C\in\mathcal{D}} \log|\kappa_K^{(q)}(C,\chi)|$, this is (i).

(ii) Each of the eleven diagrams is an ensemble at level q of at most two rectangles with the representation π. A rectangle with sides $a \ge b$ has $s(C) = b$, $\delta(C) = b$, and $P(C) = 2(a+b)$, so that $\Lambda(C) \le \max(2, 1 + a/b)$ by Lemma 2.5, $r(C) = \min(b/2, b, (a+b)/4) = b/2$, and $P(C)/r(C) = 4(1 + a/b)$; for the nine rectangles $a/b \le \max(s/t, 4t/s)$, so that they are admissible and $P(C)/r(C) \le 4\Lambda_*$, and $P_q(C) \le 2(4t+s)L^q/u$. The left side of (12.19) is therefore at most $\ell_{\mathrm{w}} := 8(4t+s)^2 L^{2q}/u^2 + 8\Lambda_*$ for each diagram. Let $\bar g_1$ and ℓ be those of Proposition 12.5 at the level q, and let $\bar g_1' \le \bar g_1$ be so small that $\ell(c_\pi, \bar g) \ge \ell_{\mathrm{w}}$ for $\bar g \le \bar g_1'$, which is possible because $\ell(c_\pi, \bar g) \to \infty$ as $\bar g \to 0$. Proposition 12.5 then gives $q \ge q_\sharp(c_\pi, \bar g)$, the limits, and the lower bounds. The diagrams I_a and $R_n^{(1)}$ are the images of I_0 and R_n under the translations by ate_μ and te_μ, which belong to $\mathfrak{G}_q$ because $t \in uL^{-q}\mathbb{N}$, so that Corollary 1.6(ii) gives the equalities for W_K and for W. By the lower bounds $N_K(\mathcal{D}) \ne 0$ for K large, and (i) together with the continuity of $\log|\cdot|$ on $\mathbb{C}\setminus\{0\}$ gives the three limits. Finally $\mathfrak{Y}^{\mathrm{sym}}(W) = \mathfrak{Y}(W)$ by Lemma 13.1(iii) applied to $f = \log|W|$.

(iii) By (1.12) and Proposition 7.13 at the level q', $\log|\kappa_K^{(q')}(C,\chi)|$ is again a linear function of the lengths and the corner numbers of C with coefficients independent of C, so that (i) holds with $W_K^{(q')}$ in place of W_K. Since $W_K^{(q')}(\mathcal{D}) \to W^{(q')}(\mathcal{D}) \ne 0$, also $N_K(\mathcal{D}) \ne 0$ for K large, and $\mathfrak{Y}(W^{(q')}) = \lim_K \mathfrak{Y}(N_K) = \mathfrak{Y}(W)$ by (ii). *q.e.d.*

In other words, $\mathfrak{Y}(W)$ does not depend on the renormalisation prescription within the class of perimeter and corner normalisations, and in particular not on the level at which the diagrams are read; any change of the masses $m_{v,K}$, of the corner constants, or of a constant factor per loop leaves it unchanged.

13.4 The Free Field

The lower bound on the shape term rests on an explicit computation for the free field. At the stopping scale the field is close to a flat field, whose holonomies around the cycles of the torus need not be trivial, and in the components of the Lie algebra on which these holonomies act diagonally the free field acquires twisted periodicity conditions. We therefore compute the covariance of the fluxes for every twist, and we need a lower bound which is uniform in it.

Let $\eta > 0$ and $\mathsf{T} \in \eta\mathbb{N}$, let $\theta \in [0,1)^4$ be a *twist*, and let $\mathbb{T}^*_{\eta,\theta}$ be the set of the $(\mathsf{T}/\eta)^4$ momenta $p = (2\pi/\mathsf{T})(n+\theta)$, $n \in \mathbb{Z}^4$, taken modulo $(2\pi/\eta)\mathbb{Z}^4$, with the momentum $p = 0$

removed if $\theta = 0$. Put $\hat{p}_\alpha := (2/\eta)\sin(\eta p_\alpha/2)$ and $\hat{p}^2 := \sum_\alpha \hat{p}_\alpha^2$, which is positive on $\mathbb{T}^*_{\eta,\theta}$, and define for $z \in \eta\mathbb{Z}^4$ the kernel

$$k_\theta(z) := \frac{1}{\mathsf{T}^4} \sum_{p \in \mathbb{T}^*_{\eta,\theta}} e^{ip\cdot z}\, \frac{\hat{p}_\mu^2 + \hat{p}_\nu^2}{\hat{p}^2} \,. \tag{13.6}$$

For a rectangle I of plaquettes of the lattice $\eta\mathbb{Z}^4$ in the plane of e_μ and e_ν, which we identify with the set of the base points of its plaquettes, and for $\mathsf{w} \in \eta\mathbb{Z}$, put

$$c^\theta(\mathsf{w}) := \eta^4 \sum_{x \in \mathsf{I}} \sum_{y \in \mathsf{I}+\mathsf{w}e_\mu} \operatorname{Re} k_\theta(x-y) \,. \tag{13.7}$$

At $\theta = 0$ this is the covariance of the fluxes $\mathsf{X}(\mathsf{I}) = \oint_{\partial \mathsf{I}} \mathsf{B}$ and $\mathsf{X}(\mathsf{I}+\mathsf{w}e_\mu)$ of the free field, the centred Gaussian 1-form on the periodic lattice of spacing η and period T with $\mathbb{E}[\mathsf{B}_\alpha(x)\mathsf{B}_\beta(y)] = \delta_{\alpha\beta}\mathsf{T}^{-4}\sum_p e^{ip\cdot(x-y)}/\hat{p}^2$: by the lattice Stokes formula $\mathsf{X}(\mathsf{I}) = \eta^2 \sum_{x\in\mathsf{I}} \mathsf{F}(x)$ with the field strength $\mathsf{F} = \partial_\mu \mathsf{B}_\nu - \partial_\nu \mathsf{B}_\mu$, and $\mathbb{E}[\mathsf{F}(x)\mathsf{F}(y)] = k_0(x-y)$ because $|e^{i\eta p_\alpha} - 1|^2 = \eta^2 \hat{p}_\alpha^2$.

Lemma 13.5 (The Free Field's Third Difference)**.** *Let I have width t along e_μ and height s along e_ν, with $t, s \in \eta\mathbb{N}$, $12t \le \mathsf{T}$, $s \le \mathsf{T}/2$, and $\eta \le \mathsf{T}/4$. Then for every twist θ,*

$$c^\theta(3t) - c^\theta(2t) \ge \frac{s^2 t^3}{112\,\mathsf{T}^5} \,. \tag{13.8}$$

Proof. We split the multiplier of (13.6) as $(\hat{p}_\mu^2 + \hat{p}_\nu^2)/\hat{p}^2 = 1 - \sum_{\alpha \notin \{\mu,\nu\}} \hat{p}_\alpha^2/\hat{p}^2$ and treat the two parts separately.

The constant part. For $z \in \eta\mathbb{Z}^4$ the sum of $e^{ip\cdot z}$ over all the $(\mathsf{T}/\eta)^4$ momenta $(2\pi/\mathsf{T})(n+\theta)$ factorises over the coordinates, and in each coordinate it equals $(\mathsf{T}/\eta)e^{2\pi i\theta_\alpha z_\alpha/\mathsf{T}}$ if $z_\alpha \in \mathsf{T}\mathbb{Z}$ and 0 otherwise. We take lifts in which $\mathsf{I}+\mathsf{w}e_\mu$ is the translate of I by $\mathsf{w}e_\mu$, with $t \le \mathsf{w} \le \mathsf{T}-t$. For $x \in \mathsf{I}$ and $y \in \mathsf{I}+\mathsf{w}e_\mu$ we then have $0 < y_\mu - x_\mu < \mathsf{w} + t \le \mathsf{T}$, so that $x - y \notin \mathsf{T}\mathbb{Z}^4$, and the constant part contributes 0 to $k_\theta(x-y)$ if $\theta \ne 0$, and $-\mathsf{T}^{-4}$, the removed momentum, if $\theta = 0$. Either way its contribution to $c^\theta(\mathsf{w})$ does not depend on w in this range, which contains $2t$ and $3t$ because $12t \le \mathsf{T}$.

The resolvent in one direction. In the second part we fix $p_\perp := (p_\alpha)_{\alpha \ne \mu}$ with $\sum_{\alpha\notin\{\mu,\nu\}} \hat{p}_\alpha^2 > 0$, the other terms vanishing, put $\mathsf{m}^2 := \sum_{\alpha\ne\mu} \hat{p}_\alpha^2 > 0$, and let $\vartheta > 0$ with $\cosh\vartheta = 1 + \eta^2\mathsf{m}^2/2$. The sum over p_μ is then

$$\mathsf{h}(z) := \frac{1}{\mathsf{T}} \sum_{p_\mu} \frac{e^{ip_\mu z}}{\hat{p}_\mu^2 + \mathsf{m}^2} = \frac{\eta}{2\sinh\vartheta} \sum_{n\in\mathbb{Z}} e^{2\pi i\theta_\mu n} e^{-\vartheta|z-n\mathsf{T}|/\eta} \,, \qquad z \in \eta\mathbb{Z} \,, \tag{13.9}$$

the sum over $p_\mu \in (2\pi/\mathsf{T})(\mathbb{Z}+\theta_\mu)$ modulo $2\pi/\eta$. Indeed, both sides satisfy $h(z+\mathsf{T}) = e^{2\pi i\theta_\mu} h(z)$ and solve $\eta^{-2}(2h(z) - h(z+\eta) - h(z-\eta)) + \mathsf{m}^2 h(z) = \eta^{-1}\sum_n e^{2\pi i\theta_\mu n}\delta_{z,n\mathsf{T}}$, whose solution in the space of such h is unique because the operator has the eigenvalues $\hat{p}_\mu^2 + \mathsf{m}^2 > 0$ there. For the left side this is the sum of $e^{ip_\mu z}$ over the momenta, computed above in one coordinate. For the right side, $g(z) := (\eta/(2\sinh\vartheta))e^{-\vartheta|z|/\eta}$ solves the equation with the source $\eta^{-1}\delta_{z,0}$: at $z \ne 0$ because $2\cosh\vartheta = 2 + \eta^2\mathsf{m}^2$, and at $z = 0$ because $2 - 2e^{-\vartheta} + \eta^2\mathsf{m}^2 = 2\cosh\vartheta - 2e^{-\vartheta} = 2\sinh\vartheta$; the sum $\sum_n e^{2\pi i\theta_\mu n} g(z - n\mathsf{T})$ converges absolutely and has the periodicity and the sources. Since g is real and even, $\mathsf{h}(-z) = \overline{\mathsf{h}(z)}$, and $\operatorname{Re}\mathsf{h}$ is even.

Monotonicity. For $0 \le z \le \mathsf{T}$ put $y := \vartheta z/\eta$, $\mathsf{N} := \vartheta\mathsf{T}/\eta$, $\mathsf{q} := e^{-\mathsf{N}}$, and $\psi := 2\pi\theta_\mu$. Summing the two geometric series over $n \le 0$ and $n \ge 1$,

$$\frac{2\sinh\vartheta}{\eta} \operatorname{Re}\mathsf{h}(z) = \frac{S(y)}{|1 - \mathsf{q}e^{i\psi}|^2} \,, \qquad S(y) := e^{-y}(1 - \mathsf{q}\cos\psi) + \mathsf{q}e^{y}(\cos\psi - \mathsf{q}) \,. \tag{13.10}$$

We claim that $-S'(y) \ge (1-\mathsf{q})e^{-y}(1-e^{2y-\mathsf{N}})$ for $0 \le y \le \mathsf{N}/2$. Indeed $-S'(y) = e^{-y}(1-\mathsf{q}\cos\psi) - \mathsf{q}e^{y}(\cos\psi - \mathsf{q})$. If $\cos\psi \le \mathsf{q}$, the second term is nonnegative and the claim follows from $1-\mathsf{q}\cos\psi \ge 1-\mathsf{q}$. Otherwise $\cos\psi - \mathsf{q} \le 1-\mathsf{q}\cos\psi$, the difference being $-(1-\cos\psi)(1+\mathsf{q})$, and $\mathsf{q}e^{y} = e^{2y-\mathsf{N}}e^{-y}$, so that $-S'(y) \ge e^{-y}(1-\mathsf{q}\cos\psi)(1-e^{2y-\mathsf{N}})$, which gives the claim. Since $|1-\mathsf{q}e^{i\psi}|^2 \le 4$, for $0 \le z < z' < \mathsf{T}/2$,

$$\operatorname{Re}\mathsf{h}(z) - \operatorname{Re}\mathsf{h}(z') \ge \frac{\eta(1-\mathsf{q})}{8\sinh\vartheta} \int_{\vartheta z/\eta}^{\vartheta z'/\eta} e^{-y}(1-e^{2y-\mathsf{N}})\,\mathrm{d}y > 0\,, \tag{13.11}$$

so that $\operatorname{Re}\mathsf{h}$ is strictly decreasing on $[0,\mathsf{T}/2)$, for every twist.

The difference. For $x \in \mathsf{l}$ and $y \in \mathsf{l} + \mathsf{w}e_\mu$ the coordinates other than x_μ and x_ν agree, and the two rectangles have the same rows. The second part of $c^\theta(\mathsf{w})$ is therefore $-\mathsf{T}^{-3}\sum_{p_\perp}(\sum_{\alpha\notin\{\mu,\nu\}}\hat p_\alpha^2)|\eta\sum_{x_\nu}e^{ip_\nu x_\nu}|^2\eta^2\sum_{x_\mu,y_\mu}\operatorname{Re}\mathsf{h}(x_\mu-y_\mu-\mathsf{w})$, with x_ν over the rows and x_μ, y_μ over the columns of l, and by the first step

$$c^\theta(3t)-c^\theta(2t) = \frac{1}{\mathsf{T}^3}\sum_{p_\perp}\Big(\sum_{\alpha\notin\{\mu,\nu\}}\hat p_\alpha^2\Big)\Big|\eta\sum_{x_\nu}e^{ip_\nu x_\nu}\Big|^2\eta^2\sum_{x_\mu,y_\mu}\left[\operatorname{Re}\mathsf{h}(z+2t)-\operatorname{Re}\mathsf{h}(z+3t)\right], \tag{13.12}$$

with $z \coloneqq y_\mu - x_\mu \in (-t,t)$, where we used that $\operatorname{Re}\mathsf{h}$ is even. Since $t < z+2t < z+3t < 4t \le \mathsf{T}/3$, every bracket is positive by (13.11), and every term of (13.12) is nonnegative. We keep one of them. Let α and α' be the directions other than μ and ν, and choose p_ν, p_α, $p_{\alpha'}$ in their classes with $|p_\nu|, |p_{\alpha'}| \le \pi/\mathsf{T}$ and $\pi/\mathsf{T} \le |p_\alpha| \le 2\pi/\mathsf{T}$, which is possible because every class $(2\pi/\mathsf{T})(\mathbb{Z}+\theta_\beta)$ meets both $[-\pi/\mathsf{T},\pi/\mathsf{T}]$ and $\{\pi/\mathsf{T} \le |p| \le 2\pi/\mathsf{T}\}$. Since $\eta \le \mathsf{T}/4$, these momenta have $|\eta p/2| \le \pi/4$, and $\sin y \ge (2\sqrt2/\pi)y$ on $[0,\pi/4]$ gives $\hat p_\alpha^2 \ge 8/\mathsf{T}^2$, $\mathsf{m}^2 \le 6\pi^2/\mathsf{T}^2$ because $\hat p_\beta \le |p_\beta|$, and, with $s \le \mathsf{T}/2$,

$$\Big|\eta\sum_{x_\nu}e^{ip_\nu x_\nu}\Big| = \eta\left|\frac{\sin(sp_\nu/2)}{\sin(\eta p_\nu/2)}\right| \ge \frac{2\sqrt2}{\pi}\,s\,, \tag{13.13}$$

the left side being s if $p_\nu = 0$.

The bracket. Now $\eta\mathsf{m} \le \sqrt6\pi/4 < 2$, so that $\vartheta = 2\operatorname{arsinh}(\eta\mathsf{m}/2) \in [\eta\mathsf{m}/2, \eta\mathsf{m}]$, because $y/2 \le \operatorname{arsinh} y \le y$ on $[0,1]$, and $\sinh\vartheta = \eta\mathsf{m}(1+\eta^2\mathsf{m}^2/4)^{1/2} \le \sqrt2\,\eta\mathsf{m}$. In (13.11) with $z+2t$ and $z+3t$ the interval of integration starts at most at $3\vartheta t/\eta \le 3\mathsf{m}t$ and has length $\vartheta t/\eta \ge \mathsf{m}t/2$; on it $2y-\mathsf{N} \le (\vartheta/\eta)(8t-\mathsf{T}) \le -\vartheta\mathsf{T}/(3\eta) \le -\mathsf{m}\mathsf{T}/6$; and $1-\mathsf{q} \ge 1-e^{-\mathsf{m}\mathsf{T}/2}$. With $1-e^{-x} \ge x(1-x/2)$ for $x \ge 0$, the bracket is at least

$$\frac{1}{8\sqrt2\,\mathsf{m}}(1-e^{-\mathsf{m}\mathsf{T}/2})(1-e^{-\mathsf{m}\mathsf{T}/6})e^{-3\mathsf{m}t}\,\frac{\mathsf{m}t}{2}\Big(1-\frac{\mathsf{m}t}{4}\Big) \ge \frac{t}{512\sqrt2} \ge \frac{t}{725}\,, \tag{13.14}$$

because $2\sqrt2 \le \mathsf{m}\mathsf{T} \le \sqrt6\pi$ and $\mathsf{m}t \le \sqrt6\pi/12$ give $e^{-\mathsf{m}\mathsf{T}/2} \le e^{-\sqrt2} \le 1/4$, $e^{-\mathsf{m}\mathsf{T}/6} \le e^{-\sqrt2/3} \le 5/8$, $e^{-3\mathsf{m}t} \ge e^{-2} \ge 2/15$, and $1-\mathsf{m}t/4 \ge 5/6$, and $(3/4)(3/8)(2/15)(5/6) = 1/32$. The kept term of (13.12) is therefore at least $\mathsf{T}^{-3}(8/\mathsf{T}^2)(8s^2/\pi^2)\,t^2\,t/725 = 64s^2t^3/(725\pi^2\mathsf{T}^5)$, since the sum over x_μ and y_μ has $(t/\eta)^2$ terms, and $64/(725\pi^2) \ge 1/112$. *q.e.d.*

For the free field with the charge g the fluxes $\mathsf{X}_a \coloneqq \oint_{I_a}\mathsf{B}$ form a Gaussian charge model whose covariance is invariant under the translation by te_μ, with $c^{\mathrm{G}}_k = c^0(kt)$. By (13.5) its number $\mathfrak{Y}$ vanishes, while its shape term is $\mathsf{g}^2(c^0(3t)-c^0(2t)) > 0$ by the lemma: in the abelian theory the positive shape term is cancelled exactly by the pair terms.

13.5 The Pair Terms

We now show that the pair terms $\mathfrak{y}_k(W) = \log|W(P_k)/(W(I_0)W(I_k))|$ are of the fourth order in the coupling, up to logarithms. Reflection positivity gives only their sign, and only for some t: if $(k+1)t/2 \in uL^{-q}\mathbb{Z}$, the reflection $\theta := \theta_{\mu,o_\mu+(k+1)t/2}$ is of level q and maps I_0 onto I_k with the opposite orientation, so that Corollaries 1.6(iii) and 1.7(ii) give $W(P_k) = W(I_0 \cup \theta\bar{I}_0) \geq |W(I_0)|^2$ and $\mathfrak{y}_k(W) \geq 0$. An upper bound needs more, namely that the two rectangles are nearly independent under the measure of the stopping scale.

The argument has two parts. At the stopping step the normalised dressed loop of P_k on the term ℓ_0 is, up to terms which decay super-polynomially in $\bar{g}$, the product of those of I_0 and I_k, because the dressed loop of an ensemble is built from objects which see only the loops that carry them, and the objects which reach from one rectangle to the other are paid by their size; this is Lemma 13.9. The pair term is then governed by the covariance, under the probability measure $\langle\cdot\rangle_{\ell_0}$, of the deviations from one of the two normalised dressed loops. Each deviation consists of the deviation of a character of the background holonomy, which is quadratic in the distance of the holonomy from 1 by Lemma 13.6 below, and of the insertions, which are of the order $\bar{g}^2$; its L^2-norm is therefore of the order $\bar{g}^2$, and the covariance of the order $\bar{g}^4$ (Lemma 13.7 and Theorem 13.10).

Lemma 13.6 (Characters near One)**.** *Let π be an irreducible unitary representation of G of dimension n_π with the character χ. Then $\operatorname{Tr}\pi_*(Y) = 0$ for every $Y \in \mathfrak{g}$ and $\det\pi(U) = 1$ for every $U \in G$. If $U \in G$ and $h := \|\pi(U) - 1\| \leq 2/n_\pi$, then*

$$|\chi(U) - 1| \leq \frac{3}{5}h^2 \,. \tag{13.15}$$

Proof. The differential $\mathrm{d}\pi$ is a homomorphism of Lie algebras from $\operatorname{Lie} G$ into the Lie algebra of $U(n_\pi)$, and the trace is a homomorphism from the latter into the abelian Lie algebra $i\mathbb{R}$; so $\operatorname{Tr}\circ\mathrm{d}\pi$ vanishes on the commutators, which span $\operatorname{Lie} G$ because G is simple. Since $\pi_*(Y) = -i\,\mathrm{d}\pi(iY)$ (Section 8.2), $\operatorname{Tr}\pi_*(Y) = 0$, and $\det\pi(e^{iY}) = e^{i\operatorname{Tr}\pi_*(Y)} = 1$. The set of the $U \in G$ with $\det\pi(U) = 1$ is therefore a subgroup which contains the image of the exponential map, a neighbourhood of 1, and since a connected group is generated by every neighbourhood of 1, it is G.

If $n_\pi = 1$, then $\pi = \det\pi = 1$ and $\chi = 1$. Let $n_\pi \geq 2$, so that $h \leq 1$, and let $e^{i\omega_j}$, $\omega_j \in (-\pi,\pi]$, be the eigenvalues of the unitary matrix $\pi(U)$. Each satisfies $2|\sin(\omega_j/2)| = |e^{i\omega_j} - 1| \leq h$, and since $\arcsin$ is convex on $[0,\frac{1}{2}]$ with $\arcsin\frac{1}{2} < 0.524$, we have $\arcsin y \leq 1.05\,y$ there, so that $|\omega_j| = 2\arcsin(|e^{i\omega_j} - 1|/2) \leq 1.05\,h$. Hence $|\sum_j \omega_j| \leq 1.05\,n_\pi h \leq 2.1$, while $e^{i\sum_j \omega_j} = \det\pi(U) = 1$, so that $\sum_j \omega_j = 0$. Therefore

$$\chi(U) - 1 = \frac{1}{n_\pi}\sum_{j=1}^{n_\pi}(e^{i\omega_j} - 1 - i\omega_j) \,, \tag{13.16}$$

and $|e^{i\omega} - 1 - i\omega| \leq \omega^2/2$ gives $|\chi(U) - 1| \leq (1.05\,h)^2/2 \leq \frac{3}{5}h^2$. *q.e.d.*

The objects at the stopping step. Let $\mathcal{D}$ be one of the eleven diagrams. By the proof of Proposition 13.4(ii) its loops are admissible, and for $\bar{g} \leq \bar{g}'_1$ we have $q \geq q_\sharp(c_\pi,\bar{g})$, so that the results of Section 8 apply to it. The index $K_0(\mathcal{D})$ of (8.1) depends on $\mathcal{D}$ only through the largest P_q of its loops, and we put $K_{\mathrm{w}} := \max(K_0(R_4), j_q + 2)$, which is at least $K_0(\mathcal{D})$ for the eleven diagrams. Let $K \geq K_{\mathrm{w}}$, so that the stopping step $K' = K - j_q \geq 2$ is common to the eleven diagrams. The density $\varrho_{K'}$, its labelled terms, the weights w_ℓ, and the expectations

$\langle\cdot\rangle_\ell$ of Theorem 8.1 do not involve the ensemble, and we write $w \coloneqq w_{\ell_0}$ and $\langle\cdot\rangle \coloneqq \langle\cdot\rangle_{\ell_0}$, a probability measure on $G^{\mathcal{B}}$. On ℓ_0 we write $E(\mathcal{D}) \coloneqq E_{\mathrm{const},K'}$ for the constant of the dressed loop of $\mathcal{D}$, and

$$S(\mathcal{D}) \coloneqq \prod_{C\in\mathcal{D}} f_{K'}^{(C)} + \sum_{\mathfrak{B}\neq\emptyset} Q_{K'}(\mathfrak{B})\, w_{K'}(\mathfrak{B}) \tag{13.17}$$

for its bracket, a function of V, and we put $\Lambda(\mathcal{D}) \coloneqq \sum_{\ell\neq\ell_0\,|\,Z_\ell>0} w_\ell \langle\widehat{\mathcal{W}}_\ell\rangle_\ell$, computed with the dressed loops of $\mathcal{D}$. Then (8.5) reads

$$W_K(\mathcal{D}) = w\, e^{E(\mathcal{D})}\langle S(\mathcal{D})\rangle + \Lambda(\mathcal{D})\,. \tag{13.18}$$

For a single rectangle C we write $\xi_C \coloneqq S(C) - 1$ and $I(C) \coloneqq S(C) - f_{K'}^{(C)}$, so that $|I(C)| \le e^{I_*(C)} - 1$ by Lemma 8.6. Finally let

$$\bar{C} \coloneqq C_{\mathrm{fs}} N_q^4 \Big(1 + \frac{1}{2}\log \hat{g}_{K'}^{-2}\Big)\,, \qquad T_{\mathrm{ng}} \coloneqq 2\bar{C} + 4\log\bar{g}^{-2}\,, \tag{13.19}$$

which do not depend on K, and, for each of the nine rectangles C,

$$\bar{h}(C) \coloneqq 4.8\, c_\pi c_h B_3 P_{K'}(C)^2 (2NT_{\mathrm{ng}})^{1/2}(1+\beta_0)\bar{g}\,, \qquad \sigma(C) \coloneqq \frac{3}{5}\bar{h}(C)^2 + e^{I_*(C)} - 1 + 3\bar{g}^4\,. \tag{13.20}$$

Lemma 13.7 (Sizes at the Stopping Step)**.** *For $\bar{g}$ sufficiently small, depending on q, t/u, s/u, T/u, π, and the constants of the construction, the following hold for every $K \ge K_{\mathrm{w}}$ and every one of the eleven diagrams $\mathcal{D}$.*

(i) $|E(\mathcal{D})| \le \bar{g}^{10}$, $1 - w \le \bar{g}^{10}$, $|\Lambda(\mathcal{D})| \le \bar{g}^{10}$, *and* $I_*(\mathcal{D}) \le \bar{g}$.

(ii) The event $\mathcal{V} \coloneqq \{V \in \operatorname{supp}\varrho_{\ell_0} \,|\, A(U_{K'}(V)) \le T_{\mathrm{ng}} g_{K'}^2\}$, with A the Wilson action, has $\langle \mathbf{1}_{\mathcal{V}^c}\rangle \le \bar{g}^8$, and on $\mathcal{V}$, for each of the nine rectangles C, $\|\pi(U_{K'}(V)(C)) - 1\| \le \bar{h}(C)$ and $|f_{K'}^{(C)} - 1| \le \frac{3}{5}\bar{h}(C)^2$.

(iii) For each of the nine rectangles C, $|\xi_C| \le 3$ on the support of ϱ_{ℓ_0} and $\langle|\xi_C|^2\rangle^{1/2} \le \sigma(C)$.

(iv) There are C_σ and m_σ, depending on the same quantities, with $\sigma(C) \le C_\sigma \bar{g}^2 (\log\bar{g}^{-2})^{m_\sigma}$ for each of the nine rectangles.

Proof. We first collect the sizes of the parameters of the stopping step. By (3.3), $\hat{g}_{K'}^{-2} = \bar{g}^{-2} + b' j_q + C_W$, and j_q grows like $\log\log\bar{g}^{-2}$ by (3.44), so that $\hat{g}_{K'} \le \bar{g}$ and $\log\bar{g}^{-2} \le \log\hat{g}_{K'}^{-2} \le 2\log\bar{g}^{-2}$ for $\bar{g}$ small, as in the proof of Proposition 12.5. Hence $(\log\bar{g}^{-2})^r \le R_{K'} \le R_{K'-1} \le 2L(\log\hat{g}_{K'-1}^{-2})^r \le C(\log\bar{g}^{-2})^r$ by Lemma 3.4, with constants C of the kind of the statement here and below. Consequently $P_{K'}(C) = P_q(C) L^3 M R_{K'-1} \le C(\log\bar{g}^{-2})^r$ for each of the nine rectangles, $N_q = 2L^{m_T+q}L^3 M R_{K'-1} \le C(\log\bar{g}^{-2})^r$, and $\bar{C} + T_{\mathrm{ng}} \le C(\log\bar{g}^{-2})^{4r+1}$. In units of the step K' the corners of the rectangles lie in $a_{K'}\mathbb{Z}^4$ with $a_{K'} = L^{j_q - q} \ge L^3 M R_{K'}$ by Lemma 3.14, and their shortest segments, the distances between their opposite sides, and the distance $(k-1)t/(uL^{-j_q})$ of I_0 from I_k are at least $a_{K'}$.

(i) By Definition 5.2(iv), $|E(\mathcal{D})| \le \sum_{j\le K'} \epsilon_j(\mathcal{D}, T)$, and we sum the parts of (7.31) as in the proof of Corollary 7.11, keeping the exponentials. For a loop C of $\mathcal{D}$ put $x \coloneqq L r_{K'}(C) \ge L a_{K'}/2 \ge L^4 M R_{K'}/2$. The corner part is at most $C_E^{\mathcal{K}} \bar{\mathfrak{a}}(P(C)/r(C))(x+1)(1+1/(c_E^T x)) e^{-2c_E^T x}$ by Lemma 2.11 with $c = 2c_E^T$ and $|\mathcal{K}(C)| \le P(C)/(2r(C))$, and $2c_E^T x \ge \nu L^3 R_{K'}/(64 A_{\mathrm{env}}^*)$. The torus part is at most $9\bar{C}_Z \bar{\mathfrak{a}} P_q(C) L \iota_{T,K'} e^{-c^Z \iota_{T,K'}}$, by the proof of Corollary 7.11, and since $y \mapsto y e^{-c^Z y}$ decreases for $c^Z y \ge 1$ and $\iota_{T,K'} \ge 96 L^3 M R_{K'}$ by (7.12), it is at most $9\bar{C}_Z \bar{\mathfrak{a}} P_q(C) L \cdot$

$96L^3MR_{K'}e^{-3\nu L^3R_{K'}/(8A^*_{\rm env})}$. The cross part is at most $3C_{\rm cr}\bar{\mathfrak{a}}P_q(C)L\cdot L^3MR_{K'}e^{-\nu_\times L^3R_{K'}}$ in the same way, with $a_{K'}\ge L^3MR_{K'}$. Each of these three is at most a constant times $(\log\bar g^{-2})^r e^{-c_\flat(\log\bar g^{-2})^r}$ with $c_\flat:=\min(\nu/(64A^*_{\rm env}),\nu_\times)L^3$, and since $r=2$ this is at most $\bar g^{11}$ for $\bar g$ small. The **R**-part is at most $C_E^R(P(C)/r(C)+P_q(C))(\frac12+2/(ec_E^R))S^{\mathbf R}(\bar g)$ by the proof of Corollary 7.11, and the cross part of the **R**-steps, present for the pairs, is at most $6C_E^{\mathbf R\times}P_q(\mathcal D)S^{\mathbf R}(\bar g)/(e\kappa)$ by the same proof. To see that $S^{\mathbf R}(\bar g)$ decays faster than every power of $\bar g$, put $y_n:=\frac23(\bar g^{-2}+b'n)$ and $a:=(A_0/4)(\log y_0)^{p_0-1}$, which tends to infinity as $\bar g\to0$ because $p_0>1$. The term $j=K-n$ of $S^{\mathbf R}(\bar g)$ is at most $4L^2M^2(\log g_j^{-2})^{2r}\exp(-(A_0/4)(\log y_n)^{p_0})$, by that proof, and $(\log y_n)^{p_0}\ge(\log y_0)^{p_0-1}\log y_n$, while $g_j^{-2}\le\hat g_j^{-2}\le 3y_n$ by Lemma 3.2(i) and (3.3); for $\bar g$ small, $(\log 3y)^{2r}\le y$ for $y\ge y_0$ and $a\ge3$, so that

$$S^{\mathbf R}(\bar g)\le 4L^2M^2\sum_{n\ge0}y_n^{1-a}\le 4L^2M^2y_0^{1-a}\Big(1+\frac{3y_0}{2b'(a-2)}\Big)\le 8L^2M^2y_0^{2-a}\,, \tag{13.21}$$

the sum being bounded by its first term and the integral $\int_0^\infty y_u^{1-a}\,\mathrm du$, which is at most $\bar g^{11}$ for $\bar g$ small. Adding the parts over the at most two loops of $\mathcal D$ gives $|E(\mathcal D)|\le\bar g^{10}$. For the large-field terms, $1-w\le\bar\delta_{\rm LF}$ by Proposition 8.4, and $|\Lambda(\mathcal D)|\le e^{B_E(\mathcal D)+I_*(\mathcal D)}\bar\delta_{\rm LF}$ by the proof of Theorem 8.7; here $\bar\delta_{\rm LF}\le e^{2\bar C}(e^{\bar s}-1)$ with $\bar s:=2N_q^4\exp(-c_{\rm LF}A_0(\log((1+\beta_0)^{-2}\bar g^{-2}))^{p_0})$, whose logarithm is at most $C(\log\bar g^{-2})^{4r+1}-c(\log\bar g^{-2})^{p_0}$ with $p_0\ge5r>4r+1$, so that $\bar\delta_{\rm LF}\le\bar g^{11}$ for $\bar g$ small. Finally $B_E(\mathcal D)\le1$ and $I_*(\mathcal D)\le\bar g$ for $\bar g$ small, by (12.26) and by (iv) below, whose proof does not use (i).

(ii) We first bound the probability of $\mathcal V^c$. In the proof of Proposition 8.4, every term is divided by the common factor $e^{-E^{\ell_0}}$, after which the density of ℓ_0 is at most $\exp(-g_{K'}^{-2}A(U_{K'}(V))+C_{\mathcal E}N_q^4)$, since $\chi_{K'}(\mathbb T)\le1$ and the regular part, the **R**-part, and the boundary part are bounded by $C_{\mathcal E}\sum_n|\Gamma_n|=C_{\mathcal E}N_q^4$ on ℓ_0, and its integral is at least e^{-C_K} by (8.28). Since $C_{\mathcal E}N_q^4\le C_K\le\bar C$, because $C_{\rm fs}\ge C_{\mathcal E}$ and $\log g_{K'}^{-1}\le\frac12\log\hat g_{K'}^{-2}$, the normalised Haar measure gives

$$\langle\mathbf 1_{\mathcal V^c}\rangle\le e^{C_K}\int_{\mathcal V^c}e^{-g_{K'}^{-2}A(U_{K'}(V))+C_{\mathcal E}N_q^4}\,\mathrm dV\le e^{2\bar C-T_{\rm ng}}=\bar g^8\,. \tag{13.22}$$

On $\mathcal V$ we apply Supplement, Lemma 17.6, with $T=T_{\rm ng}$ and $\varepsilon^{\rm act}:=3(2NT_{\rm ng})^{1/2}g_{K'}$. Its conditions hold for $\bar g$ small: $\varepsilon_{K'}$ is small; $2B_3\varepsilon_{K'}\le a_0$; $\varepsilon^{\rm act}\le a_1$; and $\varepsilon^{\rm act}\le2\varepsilon_{K'}=2\hat g_{K'}p_0(\hat g_{K'})$, because $g_{K'}\le\rho^{1/2}\hat g_{K'}$ by Lemma 3.2(i), so that it suffices that $9\cdot2NT_{\rm ng}\rho\le4A_0^2(\log\hat g_{K'}^{-2})^{2p_0}$, whose left side is at most $C(\log\bar g^{-2})^{4r+1}$ while $2p_0>4r+1$. The lemma gives $|U(\partial p)-1|<B_3\varepsilon^{\rm act}L^{-2K'}$ on the fine plaquettes, $U:=U_{K'}(V)$, and hence $\|U(\partial p)-1\|\le t:=c_hB_3\varepsilon^{\rm act}L^{-2K'}$. In fine units a rectangle C of the nine is a closed path of $P:=P_{K'}(C)L^{K'}$ bonds in a box of side at most $P/2$, which embeds in the torus because $3\,\mathrm{diam}(C)<T$. Hence $3(P/2)Pt\le\frac32c_hB_3P_{K'}(C)^2\varepsilon^{\rm act}$, which is at most $\min(1/10,\log(1+r_G))$ for $\bar g$ small, and Lemma 12.6 gives $\|\pi(U(C))-1\|\le1.06\cdot\frac32c_\pi c_hB_3P_{K'}(C)^2\varepsilon^{\rm act}\le\bar h(C)$, using $g_{K'}\le(1+\beta_0)\bar g$. For $\bar g$ small, $\bar h(C)\le2/n_\pi$, and Lemma 13.6 gives $|f_{K'}^{(C)}-1|=|\chi(U(C))-1|\le\frac35\bar h(C)^2$.

(iii) On the support of ϱ_{ℓ_0}, $|\xi_C|\le|f_{K'}^{(C)}-1|+|I(C)|\le2+e^{I_*(C)}-1\le3$ by (i), and on $\mathcal V$, $|\xi_C|\le\frac35\bar h(C)^2+e^{I_*(C)}-1$ by (ii). Splitting $\xi_C=\xi_C\mathbf 1_{\mathcal V}+\xi_C\mathbf 1_{\mathcal V^c}$, Minkowski's inequality and (13.22) give $\langle|\xi_C|^2\rangle^{1/2}\le\frac35\bar h(C)^2+e^{I_*(C)}-1+3\langle\mathbf 1_{\mathcal V^c}\rangle^{1/2}\le\sigma(C)$.

(iv) By the sizes collected above, $\bar h(C)^2\le C\bar g^2P_{K'}(C)^4T_{\rm ng}\le C\bar g^2(\log\bar g^{-2})^{8r+1}$. In (8.37), $\bar\rho\le C\bar g^2(\log\bar g^{-2})^{4p_0}$ by (8.36), $\varepsilon_{K'}/\min(\alpha_{0,K'},\alpha_{1,K'})$ is a power of $\log\hat g_{K'}^{-2}$ by Definition 3.11, $S_{\rm wr}(b)\le3$ for $b\ge1$ by Lemma 2.11, and $P_{K'}(C)\le C(\log\bar g^{-2})^r$, so that $I_*(C)\le C\bar g^2(\log\bar g^{-2})^{m'}$ with an exponent m' of the constants of stage 0, and $e^{I_*(C)}-1\le2I_*(C)$. With $3\bar g^4\le3\bar g^2$ this gives (iv) with $m_\sigma:=\max(8r+1,m')$. *q.e.d.*

We next make precise the statement that a dressed loop sees only the loops which carry its objects. The *network* of a group is the union of its domains, of all scales, its intervals, and the networks of the clusters merged into it; it is connected, contains the networks of its old members, and has diameter at most $M\sigma(W)$ in units of the step at which it was made, by the proof of Lemma 7.3. For an ensemble $\mathcal{C}$ with a loop C of the character χ_C we write $\mathcal{C}_C := ((C, \chi_C))$.

Lemma 13.8 (Locality between Ensembles)**.** *Let $\mathcal{C}$ be an ensemble as in Theorem 1.5, C one of its loops, and $K \geq K_0(\mathcal{C})$. On the trivial term, at every step $1 \leq k \leq K'$, the following hold.*

(i) Every group of $\mathcal{C}$ which is not a cross group and whose intervals lie on C is a group of $\mathcal{C}_C$, with the same domains, intervals, network, and insertion, the insertion being the same function of the block field. Two such groups are compatible for $\mathcal{C}$ if and only if they are compatible for $\mathcal{C}_C$.

(ii) Every group of $\mathcal{C}_C$ whose network is at distance greater than 2 from the other loops of $\mathcal{C}$, in units of the step k, is a group of $\mathcal{C}$ of the kind (i).

(iii) A cluster of the reference gas of $\mathcal{C}$ of the step k all of whose groups are of the kind (i) is a cluster of the reference gas of $\mathcal{C}_C$ with the same weight, and a cluster of $\mathcal{C}_C$ all of whose groups satisfy the hypothesis of (ii) is a cluster of $\mathcal{C}$ with the same weight.

Proof. A group, with its insertion, is determined by its constituents. At the step 1 these are its roots and its cluster occurrences, and a group is the only group of the term formed by them [1, Definition 10.1, Lemma 10.2(a)]. At a step $k+1 \geq 2$ they are the polymers of the step merged into it, its old members, which are groups of the step k, the slots on its arcs, and the clusters of the reference gas of the step k merged into it (Section 6.7 and the merging rule of Section 7.1), and at an **R**-step also the polymers of the **R**-operation linked to it (Supplement, Proposition S.10). Its domains are those of its constituents, its intervals are the smallest arcs of its loops which contain the intervals of its members and the traversals of its slots and merged clusters, and its insertion is the Gaussian integral of the step of the product of the activities of its polymers, the insertions of its old members, its slots, and the weights of its merged clusters, less its flat part (7.16). Compatibility is a relation between the domains, the arcs, and the merged clusters of two groups, and a cluster of the reference gas is a multiset of groups whose weight is the Ursell function of its incompatibility graph times the flat values of its groups. None of these rules involves a loop which carries no interval of the group: the polymers and their activities are those of Bałaban's expansion, which does not involve the ensemble; the slots and the transports are attached to the traversals of the arcs, along the same background field; and the smallest arc of C which contains a set of traversals of C is determined by C. We argue by induction on k.

The step 1. By [1, Lemma 10.2(c)], applied to C and to each other loop of $\mathcal{C}$ as in the proof of Theorem 5.4, the groups of $\mathcal{C}$ with intervals on C only are the groups of $\mathcal{C}_C$ none of whose roots has a contact on another loop, with the same intervals, weights, and compatibility. A root with a contact on a loop C' contains a bond of the dependence set $\mathcal{F}_r$ of a link r of C' [1, (3.87), Lemma 3.5], whose bonds have both ends in the two blocks of the first step at the ends of the bond of C' which contains r, so that it lies within distance 1 of C' in units of the step 1. This gives (i) and (ii) at the step 1, and (iii) follows from the description of the clusters.

The step $k \to k+1$. Let W be a group of $\mathcal{C}$ of the step $k+1$ which is not a cross group and has its intervals on C. Its old members and the groups of its merged clusters have their

intervals on C, since the arcs of W contain theirs; in particular no merged cluster contains groups of two loops, which would put arcs on both into W. By (i) and (iii) at the step k they are objects of $\mathcal{C}_C$ with the same insertions and weights, and the rules above assemble from the same constituents a group of $\mathcal{C}_C$ with the same domains, intervals, network, and insertion. Conversely, let W be a group of $\mathcal{C}_C$ of the step $k+1$ whose network is at distance greater than 2 from the other loops, in units of the step $k+1$. The networks of its old members and merged clusters lie in that of W, at distance greater than $2L$ from the other loops in units of the step k, so that by (ii) and (iii) at the step k they are objects of $\mathcal{C}$ of the kind (i); none of its polymers meets a traversal of another loop; and the rules assemble from these constituents a group of $\mathcal{C}$ which is not a cross group. Compatibility is the same in both ensembles because it depends only on the domains, the arcs, and the merged clusters, and (iii) follows from (i) and (ii) by the description of the clusters. *q.e.d.*

The groups of $\mathcal{C}_C$ which do not satisfy the hypothesis of (ii) have no counterpart in $\mathcal{C}$, in general: their constituents may meet objects of another loop, with which they form a cross group of $\mathcal{C}$. These groups, the cross groups, and the clusters which contain groups of two loops are the objects which distinguish the pair from the two rectangles, and the next lemma shows that they are negligible.

Lemma 13.9 (The Pair at the Stopping Step)**.** *Let $k \in \{2,3\}$, $K \geq K_{\mathrm{w}}$, and let $D := (k-1)tL^{j_q}/u$ be the distance of I_0 from I_k in units of the step K'. On the support of ϱ_{ℓ_0},*

$$|S(P_k) - S(I_0)S(I_k)| \leq 3e^{I_*(P_k)}(e^{N_{\mathrm{ng}}} - 1) \,, \tag{13.23}$$

where, with $I_{\mathrm{wr}}(P_k) := P_{K'}(P_k)C^2\bar{\rho}\, S_{\mathrm{wr}}(c_{\mathrm{att}}N_q/M)$ the second part of (8.37) *and $A^\times(P_k)$ its last part, the cross total* (8.38) *of the pair, for which $d_\times = D$,*

$$N_{\mathrm{ng}} := A^\times(P_k) + e^{-\nu D/(3M)}I_*(P_k) + I_{\mathrm{wr}}(P_k) + LP_{K'}(P_k)C_{S'}\mathfrak{a}_{K'}e^{-\nu D/(96A^*_{\mathrm{env}}M)} \,. \tag{13.24}$$

Proof. Let $\mathcal{D}$ be one of P_k, I_0, I_k. We call a group of $\mathcal{D}$ at the step K' on ℓ_0 *far* if it is not a cross group and its network lies within distance $D/3$ of the loop C which carries its intervals, in units of the step K', and a cluster of the reference gas of the step K' *far* if all its groups are far and carried by the same loop. The other groups and clusters are *near*. We put

$$F(\mathcal{D}) := \sum_{\mathfrak{B}\ \mathrm{far}} Q^{\mathrm{far}}(\mathfrak{B})\, w_{K'}(\mathfrak{B}) \,, \qquad Q^{\mathrm{far}}(\mathfrak{B}) := \exp\Big(- \sum_{\Gamma\ \mathrm{far},\ \Gamma \not\perp \mathfrak{B}} \Phi^\circ(\Gamma)\Big) \,, \tag{13.25}$$

the first sum over the families of far groups, the empty family included, and the second over the far clusters which contain a group incompatible with a member of $\mathfrak{B}$.

The far parts factorise. Let $C \in \{I_0, I_k\}$ and let C' be the other rectangle. A far group of P_k carried by C has its network at distance at least $D - D/3 > 2$ from C', so by Lemma 13.8(i) it is a group of $\mathcal{C}_C = C$ with the same network, hence far, and with the same insertion; conversely a far group of C is a group of P_k of the kind (i) by Lemma 13.8(ii), with the same network. By Lemma 13.8(iii) the same holds for the far clusters, with the same weights. Two incompatible groups have networks at distance at most M in units of the step K', by the proof of Lemma 7.3, while the networks of far groups carried by I_0 and by I_k are at distance at least $D/3 > M$; so these groups are compatible, and so are a far cluster carried by one rectangle and a far group carried by the other. Hence the families of far groups of P_k are the unions $\mathfrak{B}_0 \cup \mathfrak{B}_k$ of families of far groups of I_0 and of I_k, and $Q^{\mathrm{far}}(\mathfrak{B}_0 \cup \mathfrak{B}_k)$ is the product of the two factors of I_0 and of I_k. The insertions of far groups act on the space of one loop, so that the operator of which $w_{K'}(\mathfrak{B}_0 \cup \mathfrak{B}_k)$ is the normalised trace is the tensor

product of those of $\mathfrak{B}_0$ and $\mathfrak{B}_k$, along the same background field, and the normalised trace of a tensor product is the product of the normalised traces. Therefore

$$F(P_k) = F(I_0)\,F(I_k)\ . \tag{13.26}$$

The near parts are small. Let $\mathfrak{M} := \sum_W e^{\theta(W)}\|Y_W\|$, the sum over all groups of $\mathcal{D}$, $\mathfrak{N}_{\rm g} := \sum_{W\ \rm near} e^{\theta(W)}\|Y_W\|$, and $\mathfrak{N}_{\rm c} := \sum_{\Gamma\ \rm near}|\Phi^\circ(\Gamma)|$. By (7.18) and the bound after it, $Q_{K'}(\mathfrak{B}) = Q^{\rm far}(\mathfrak{B})\exp(-\sum_{\Gamma\ \rm near,\Gamma\not\subset\mathfrak{B}}\Phi^\circ(\Gamma))$ for a family of far groups, and $|Q^{\rm far}(\mathfrak{B})| \le \exp(\sum_{\Gamma\not\subset\mathfrak{B}}|\Phi^\circ(\Gamma)|) \le \prod_{W\in\mathfrak{B}} e^{\theta(W)}$, while $|w_{K'}(\mathfrak{B})| \le \prod_{W\in\mathfrak{B}}\|Y_W\|$ as in the proof of Lemma 8.6. Hence, separating in $S(\mathcal{D}) - F(\mathcal{D})$ the families of far groups from those with a near member, and bounding sums over families by sums over sets of groups,

$$|S(\mathcal{D})-F(\mathcal{D})| \le e^{\mathfrak{M}}(e^{\mathfrak{N}_{\rm c}}-1)+\prod_{W\ \rm far}(1+e^{\theta(W)}\|Y_W\|)\Big(\prod_{W\ \rm near}(1+e^{\theta(W)}\|Y_W\|)-1\Big) \le e^{\mathfrak{M}}(e^{\mathfrak{N}_{\rm g}+\mathfrak{N}_{\rm c}}-1)\ , \tag{13.27}$$

where we used $(e^a-1)+(e^b-1) \le e^{a+b}-1$ for $a,b\ge 0$. In the same way $|S(\mathcal{D})| \le e^{\mathfrak{M}}$ and $|F(\mathcal{D})| \le e^{\mathfrak{M}}$, and $\mathfrak{M} \le I_*(\mathcal{D})$ by the proof of Lemma 8.6, since there are no attached groups on ℓ_0.

We bound the near masses. The groups of the cross species of P_k have the total at most $A^\times(P_k)$ by Lemma 8.5, with $d_\times = D$, and a single rectangle has none. A near group which is not a cross group has a network which contains a traversal of its loop and a point at distance at least $D/3$ from it; if it was made at the step $j \le K'$, its network has diameter at least $L^{K'-j}D/3$ in units of the step j, so that $\sigma(W) \ge D/(3M)$. If it is of the deviation species, the weighted bound of Definition 5.2(ii) gives for these groups the total $e^{-\nu D/(3M)}\sum_{\mathfrak{n}} C_\chi\tilde\rho_{K'}\delta^*_{\mathfrak{n}}(U_{K'}) \le e^{-\nu D/(3M)}I_*(\mathcal{D})$, by the proof of Lemma 8.6. On ℓ_0 there are no attached groups, so that a group which is neither of the cross species nor of the deviation species is one of the groups whose domains do not lift to the universal cover (Section 6.16), and the total of these is at most $I_{\rm wr}(\mathcal{D})$ by the proof of Lemma 8.6. Thus $\mathfrak{N}_{\rm g} \le A^\times(\mathcal{D}) + e^{-\nu D/(3M)}I_*(\mathcal{D}) + I_{\rm wr}(\mathcal{D})$. A near cluster of nonzero weight contains no cross group, whose flat value is 0; it contains either groups of both rectangles, which are joined by a chain of incompatible groups, or a group whose network reaches distance $D/3$ from its loop, and in both cases its network has diameter at least $D/3$ in units of the step K', that is $LD/3$ in units of the step $K'-1$, and contains a traversal of its groups' arcs. By (7.6) at the step K', summed over the $LP_{K'}(\mathcal{D})$ traversals of the step $K'-1$, $\mathfrak{N}_{\rm c} \le LP_{K'}(\mathcal{D})C_{S'}\mathfrak{a}_{K'}e^{-2c_E^T LD/3}$, and $2c_E^T LD/3 = \nu D/(96A^*_{\rm env}M)$. Since $A^\times$, I_*, $I_{\rm wr}$, and $P_{K'}$ do not decrease when a loop is added, $\mathfrak{N}_{\rm g}+\mathfrak{N}_{\rm c} \le N_{\rm ng}$ for each of the three diagrams.

Conclusion. By (13.26),

$$S(P_k)-S(I_0)S(I_k) = (S(P_k)-F(P_k)) - F(I_0)(S(I_k)-F(I_k)) - S(I_k)(S(I_0)-F(I_0))\ , \tag{13.28}$$

and (13.27) with $I_*(I_0)+I_*(I_k) \le I_*(P_k)$ gives (13.23). *q.e.d.*

Theorem 13.10 (The Pair Terms)**.** *For $\bar g$ sufficiently small, depending on q, t/u, s/u, T/u, π, and the constants of the construction, and for $k \in \{2,3\}$,*

$$|\mathfrak{y}_k(W)| \le 16\,\sigma(I_0)^2 \le 16C_\sigma^2\bar g^4(\log\bar g^{-2})^{2m_\sigma}\ . \tag{13.29}$$

Proof. Let $K \ge K_{\rm w}$, and write $X_{\mathcal{D}} := e^{E(\mathcal{D})}S(\mathcal{D})$, so that $W_K(\mathcal{D}) = w\langle X_{\mathcal{D}}\rangle + \Lambda(\mathcal{D})$ by (13.18), and abbreviate the indices P_k, I_0, I_k by P, 0, k. Expanding the product $W_K(I_0)W_K(I_k)$ and adding and subtracting $w\langle X_0X_k\rangle$ and $w\langle X_0\rangle\langle X_k\rangle$,

$$\begin{aligned}\Delta := W_K(P_k) - W_K(I_0)W_K(I_k) &= w(\langle X_0X_k\rangle - \langle X_0\rangle\langle X_k\rangle) + w\langle X_P - X_0X_k\rangle \\ &\quad + w(1-w)\langle X_0\rangle\langle X_k\rangle + \Lambda(P_k) - w\Lambda(I_0)\langle X_k\rangle - w\Lambda(I_k)\langle X_0\rangle - \Lambda(I_0)\Lambda(I_k)\ .\end{aligned} \tag{13.30}$$

The first term is a covariance, in which the constant parts of $S = 1+\xi$ drop out: $\langle X_0 X_k\rangle - \langle X_0\rangle\langle X_k\rangle = e^{E(I_0)+E(I_k)}\langle(\xi_{I_0} - \langle\xi_{I_0}\rangle)(\xi_{I_k} - \langle\xi_{I_k}\rangle)\rangle$. By the Cauchy–Schwarz inequality, $\langle|\xi - \langle\xi\rangle|^2\rangle \le \langle|\xi|^2\rangle$, and Lemma 13.7(i), (iii), its modulus is at most $e^{2\bar g^{10}}\sigma(I_0)\sigma(I_k)$, and $\sigma(I_k) = \sigma(I_0)$ because I_k is a translate of I_0, with the same $P_{K'}$ and I_*.

The other terms are small. On the support of ϱ_{ℓ_0}, $X_P - X_0X_k = e^{E(P_k)}(S(P_k) - S(I_0)S(I_k)) + (e^{E(P_k)} - e^{E(I_0)+E(I_k)})S(I_0)S(I_k)$. By Lemma 13.7(i), $|E(\mathcal{D})| \le \bar g^{10}$ and $I_*(\mathcal{D}) \le \bar g \le 1/10$, and $N_{\mathrm{ng}} \le \bar g^{10}$ for $\bar g$ small, because each of its four terms carries one of the factors $e^{-\nu_\times D/M}$ and $S^{\mathbf{R}}(\bar g)$ (in (8.38), with $d_\times = D$; the second is smaller than every power of $\bar g$ by (13.21)), $e^{-\nu D/(3M)}$, $e^{-2c_{\mathrm{att}}L^3R_{K'}}$ (from $S_{\mathrm{wr}}(b) \le 3e^{-b}$ for $b \ge 1$ and $N_q/M \ge 2L^3R_{K'}$), and $e^{-\nu D/(96A^*_{\mathrm{env}}M)}$, with $D \ge L^3MR_{K'}$ and $R_{K'} \ge (\log\bar g^{-2})^2$, against a power of $\log\bar g^{-2}$. Lemma 13.9, $|S(\mathcal{D})| \le e^{I_*(\mathcal{D})}$, and $|e^a - e^b| \le e^{\max(|a|,|b|)}|a-b|$ give $|X_P - X_0X_k| \le 14\bar g^{10}$, and $|\langle X_{\mathcal{D}}\rangle| \le 2$. With Lemma 13.7(i) for $1-w$ and Λ, the sum of the moduli of the terms of (13.30) other than the first is at most $(14+4+1+4+1)\bar g^{10} \le \bar g^9$ for $\bar g \le 1/24$. Since $\sigma(I_0) \ge 3\bar g^4$, we have $\bar g^9 \le \frac12\sigma(I_0)^2$, and $e^{2\bar g^{10}} \le \frac32$, so that

$$|\Delta| \le 2\,\sigma(I_0)^2\ . \tag{13.31}$$

Finally we take the logarithm. By Proposition 13.4(ii), $W_K(R_1) = W_K(I_0) = W_K(I_k)$ and $|W_K(I_0)| \ge \frac12$, so that $W_K(P_k) = W_K(I_0)W_K(I_k)(1+z)$ with $z := \Delta/(W_K(I_0)W_K(I_k))$, $|z| \le 8\sigma(I_0)^2 \le \frac12$ for $\bar g$ small by Lemma 13.7(iv), and

$$|\mathfrak{y}_k(W_K)| = |\log|1+z|| \le 2|z| \le 16\,\sigma(I_0)^2\ , \tag{13.32}$$

because $\log(1+x) \le x$ and $\log(1-x) \ge -2x$ for $0 \le x \le \frac12$. The right side does not depend on K, and $\mathfrak{y}_k(W_K) \to \mathfrak{y}_k(W)$ as $K \to \infty$ by the continuity of $\log|\cdot|$ on $\mathbb{C}\setminus\{0\}$, since the limits on the eleven diagrams exist and do not vanish (Proposition 13.4(ii)). The second inequality of (13.29) is Lemma 13.7(iv). *q.e.d.*

13.6 The Shape Term

The shape term $\mathfrak{y}_S$ is a combination of the four rectangles $R_1, \dots, R_4$ alone, and we reduce it, in three steps, to the second moments of the fluxes of the background field. At the stopping step its logarithms are, up to terms of the order $\bar g^4$, the real parts of the expectations of the deviations ξ_{R_n}, and each deviation is the sum of the deviation of the leading word and of the insertions. The insertions cancel in the shape term. This is the mechanism which made the coefficients of the normalisation independent of the loop: the expectation of an insertion is a number attached to a traversal, which, by the invariance of the construction under the translations that preserve its partitions, depends only on the position of the traversal relative to the nearest corner, so that its sum over a rectangle is a linear function of the lengths of the sides and of the number of corners of each type, and the weights of the shape term annihilate such functions (Proposition 13.12). The leading words, finally, are expanded to the second order in the logarithms of the holonomies of the rectangles I_a, and the identity which writes the boundary of R_n as a product of loops around $I_0, \dots, I_{n-1}$ turns the shape term into a second difference of the covariances of these logarithms (Lemma 13.13).

Proposition 13.11 (The Shape Term at the Stopping Step)**.** *For $\bar g$ sufficiently small, depending on q, t/u, s/u, T/u, π, and the constants of the construction, and for $K \ge K_{\mathrm{w}}$,*

$$\Big|\mathfrak{y}_S(W_K) - \sum_{n=1}^{4} n_{R_n}\,\mathrm{Re}\langle\xi_{R_n}\rangle\Big| \le \bar g^9 + 8\max_n \sigma(R_n)^2\ . \tag{13.33}$$

Proof. By (13.18), $W_K(R) = we^{E(R)}\langle S(R)\rangle(1+z_R)$ with $z_R := \Lambda(R)/(we^{E(R)}\langle S(R)\rangle)$, for each of the four rectangles R. By Lemma 13.7, $|\langle S(R)\rangle - 1| = |\langle \xi_R\rangle| \le \sigma(R) \le \frac{1}{2}$, $w \ge 1 - \bar g^{10}$, $|E(R)| \le \bar g^{10}$, and $|\Lambda(R)| \le \bar g^{10}$, so that $|z_R| \le 3\bar g^{10} \le \frac{1}{2}$ and $|\log|1+z_R|| \le 6\bar g^{10}$. Taking logarithms,

$$\log|W_K(R)| = \log w + \operatorname{Re} E(R) + \log|\langle S(R)\rangle| + \log|1+z_R| \,. \tag{13.34}$$

The weights $n_{R_n} = 1, -3, 3, -1$ sum to zero, so that $\log w$ drops out of $\mathfrak{y}_S(W_K)$, and their moduli sum to 8, so that the second and the fourth terms contribute at most $8\cdot 7\bar g^{10} \le \bar g^9$. For the third term, $|\log(1+x) - x| \le |x|^2$ for complex x with $|x| \le \frac{1}{2}$, by the series of the logarithm, and $\log|1+x| = \operatorname{Re}\log(1+x)$; with $x = \langle \xi_R\rangle$ this gives $|\log|\langle S(R)\rangle| - \operatorname{Re}\langle \xi_R\rangle| \le \sigma(R)^2$, and (13.33) follows. *q.e.d.*

Since $\xi_R = (f^{(R)}_{K'} - 1) + I(R)$, the sum in (13.33) splits into the contribution of the leading words and that of the insertions. We treat the insertions first. The objects of the construction which we use are those of the stopping step on the term ℓ_0; the partitions of the steps $1, \dots, K'$ are invariant under the translations by the vectors of $\Pi\mathbb{Z}^4$, in units of the step K', where $\Pi := L^{-1}\Pi_{K'}$ with the number $\Pi_{K'}$ of Lemma 7.1, a power of L. By that lemma and (3.44), $M \le \Pi \le LMR_{K'-1} = L^{-2}a_{K'}$, so that Π divides $a_{K'}$, and the corners of the four rectangles, their sides $t_{\mathfrak b} := tL^{j_q}/u$ and $s_{\mathfrak b} := sL^{j_q}/u$ in units of the step K', and the differences of their corners lie in $\Pi\mathbb{Z}^4$ and $\Pi\mathbb{Z}$, with $\min(t_{\mathfrak b}, s_{\mathfrak b}) \ge a_{K'} \ge L^2\Pi$. We put

$$\mathsf{D} := \Pi\Big\lfloor \frac{\min(t_{\mathfrak b}, s_{\mathfrak b})}{3\Pi} \Big\rfloor \,, \qquad \text{so that} \qquad \frac{2}{9}\min(t_{\mathfrak b}, s_{\mathfrak b}) \le \mathsf{D} \le \frac{1}{3}\min(t_{\mathfrak b}, s_{\mathfrak b}) \,, \tag{13.35}$$

because $\min(t_{\mathfrak b}, s_{\mathfrak b})/(3\Pi) \ge L^2/3 \ge 3$. Every side of each of the four rectangles has a length $N_v \ge 3\mathsf{D}$, and $\mathsf{D} \ge \frac{2}{9}L^3MR_{K'}$.

Proposition 13.12 (The Insertions Cancel)**.** *For $\bar g$ sufficiently small, depending on the quantities of Proposition 13.11, and for $K \ge K_{\mathrm w}$,*

$$\Big|\sum_{n=1}^{4} n_{R_n}\langle I(R_n)\rangle\Big| \le 8\max_n \mathsf{e}(R_n) \,, \tag{13.36}$$

where, for a rectangle C,

$$\mathsf{e}(C) := \frac{1}{2}I_*(C)^2 e^{I_*(C)} + I_*(C)(\bar h(C) + 2\bar g^8) + 2e^{-\nu(\mathsf{D}/4 - M)/(64M)}I_*(C) + I_{\mathrm{wr}}(C) \,, \tag{13.37}$$

and $8\max_n \mathsf{e}(R_n) \le C\bar g^3(\log \bar g^{-2})^{m'_\sigma}$ with constants of the same kind.

Proof. Let C be one of the four rectangles. On ℓ_0, $I(C) = \sum_{\mathfrak B \ne \emptyset} Q_{K'}(\mathfrak B) w_{K'}(\mathfrak B)$, the members of a family having pairwise disjoint arcs on C. For a traversal $\mathfrak n$ of the step K' and a point V of the support of ϱ_{ℓ_0} let $\bar\nu_{\mathfrak n} := \sum_{W \ni \mathfrak n} e^{\theta(W)}\|Y_W\|$, a function of V, so that $\sum_{\mathfrak n}\bar\nu_{\mathfrak n} \le I_*(C)$ at every such V, by the proof of Lemma 8.6. The bounds of the class are bounds on these sums at each V, and not on the sums of the suprema of their terms; accordingly every estimate below is made at a fixed V and then integrated, the sums over the groups and the integral being exchanged by Tonelli's theorem under these absolute bounds.

Families of two or more members. The members of a family have distinct first traversals. Bounding each member through its first traversal, $|Q_{K'}(\mathfrak B)w_{K'}(\mathfrak B)| \le \prod_{W\in\mathfrak B} e^{\theta(W)}\|Y_W\|$ shows that the families of m members contribute at most the elementary symmetric sum $e_m((\bar\nu_{\mathfrak n})_{\mathfrak n}) \le I_*(C)^m/m!$, and those with $m \ge 2$ at most $e^{I_*(C)} - 1 - I_*(C) \le \frac{1}{2}I_*(C)^2 e^{I_*(C)}$.

One member. For $\mathfrak{B} = \{W\}$, let J be the arc of W, $\mathsf{t}_W \coloneqq \pi(U_{K'}(J))$ the transport along it, and $\mathsf{H}_W \coloneqq \pi(U_{K'}(C_J))$, with C_J the loop C read from the initial site of J. Then $w_{K'}(\{W\}) = \mathrm{tr}(Y_W \mathsf{t}_W^{-1} \mathsf{H}_W)$, and H_W is unitarily conjugate to $\pi(U_{K'}(C))$, so that $\|\mathsf{H}_W - 1\| = h_C \coloneqq \|\pi(U_{K'}(C)) - 1\|$ and $|w_{K'}(\{W\}) - \mathrm{tr}(Y_W \mathsf{t}_W^{-1})| \le \|Y_W\| h_C$. Since $h_C \le \bar{h}(C)$ on $\mathcal{V}$, $h_C \le 2$ everywhere, and $\langle \mathbf{1}_{\mathcal{V}^c} \rangle \le \bar{g}^8$ (Lemma 13.7(ii)), and $|Q_{K'}(\{W\})| \le e^{\theta(W)}$,

$$\Big| \langle I(C) \rangle - \sum_W \mathsf{z}_W^0 \Big| \le \frac{1}{2} I_*(C)^2 e^{I_*(C)} + I_*(C)(\bar{h}(C) + 2\bar{g}^8) \,, \qquad \mathsf{z}_W^0 \coloneqq Q_{K'}(\{W\}) \langle \mathrm{tr}(Y_W \mathsf{t}_W^{-1}) \rangle \,, \tag{13.38}$$

the sum over all groups of C, which converges absolutely, since $\sum_W |\mathsf{z}_W^0| \le I_*(C)$.

Small and large groups. We call a group W *small* if it is of the deviation species and $M\sigma(W) + 1 \le \mathsf{D}/4$, and *large* otherwise. A large group of the deviation species has $\sigma(W) \ge (\mathsf{D}/4 - 1)/M$, so that by the weighted bound of Definition 5.2(ii) these groups contribute at most $e^{-\nu(\mathsf{D}/4-1)/M} I_*(C)$ to $\sum_W |\mathsf{z}_W^0|$, and the groups of the other species at most $I_{\mathrm{wr}}(C)$, as in the proof of Lemma 13.9.

The interaction, localised. For a small W let $Q^{\mathsf{D}}(W) \coloneqq \exp(-\sum' \Phi^\circ(\Gamma))$, the sum over the clusters of the reference gas of the step K' which contain a group incompatible with W and whose networks lie within distance $\mathsf{D}/4$ of the network of W, and $\mathsf{z}_W \coloneqq Q^{\mathsf{D}}(W) \langle \mathrm{tr}(Y_W \mathsf{t}_W^{-1}) \rangle$. A cluster which contains a group incompatible with W and is left out of this sum has a group within distance M of the network of W and a network of diameter at least $\mathsf{D}/4 - M$, in units of the step K', so that $a^\circ(\Gamma) \ge \nu(\mathsf{D}/4 - M)/(64M)$ by (7.8). The theorem of Kotecký and Preiss in the form [15, Theorem 5.4], with the condition (7.7), gives $\sum_{\Gamma \not\perp W} |\Phi^\circ(\Gamma)| e^{a^\circ(\Gamma)} \le a^\circ(W) = \theta(W)$, so that the clusters left out have $\sum |\Phi^\circ(\Gamma)| \le \theta(W) e^{-\nu(\mathsf{D}/4-M)/(64M)}$. Since $Q_{K'}(\{W\}) = Q^{\mathsf{D}}(W) e^{-z}$ with z the sum of $\Phi^\circ(\Gamma)$ over these clusters, $|Q^{\mathsf{D}}(W)| \le e^{\theta(W)}$, $|z| \le \theta(W)$, and $|e^{-z} - 1| \le |z| e^{|z|}$,

$$|Q_{K'}(\{W\}) - Q^{\mathsf{D}}(W)| \le \theta(W) e^{2\theta(W)} e^{-\nu(\mathsf{D}/4-M)/(64M)} \le e^{\theta(W) + \nu\sigma(W)} e^{-\nu(\mathsf{D}/4-M)/(64M)} \,, \tag{13.39}$$

because $\theta(W) e^{\theta(W)} \le e^{2\theta(W)}$ and $\theta(W) \le \nu\sigma(W)/16$. By the weighted bound, $\sum_{W \text{ small}} |\mathsf{z}_W^0 - \mathsf{z}_W| \le e^{-\nu(\mathsf{D}/4-M)/(64M)} I_*(C)$, and $\sum_{W \text{ small}} |\mathsf{z}_W| \le I_*(C)$.

Agreement. Let C and C' be two of the four rectangles, possibly equal, τ a translation by a vector of $\Pi\mathbb{Z}^4$, W a small group of C whose arc has the first traversal $\mathfrak{p}$, and $\mathbb{B}$ the open ball of radius D about the initial site of $\mathfrak{p}$. Suppose that τ maps the bonds of C with initial site in $\mathbb{B}$ onto the bonds of C' with initial site in $\tau\mathbb{B}$, with their orientations. We claim that τW is a small group of C' with the first traversal $\tau\mathfrak{p}$ and that $\mathsf{z}_{\tau W}(C') = \mathsf{z}_W(C)$. Indeed, the network of W lies within $\mathsf{D}/4$ of the initial site of $\mathfrak{p}$, and the networks of the clusters of $\sum'$ within $\mathsf{D}/2$; so the arcs of W and of the groups of these clusters are paths of consecutive bonds of C with initial sites in $\mathbb{B}$, hence also of $\tau^{-1}C'$, since both loops visit no site twice, and each has fewer than $2\mathsf{D} \le P_{K'}(C)/6$ traversals. By the argument of Lemma 7.4(i), these groups, their flat values, the hard core between them, the weights Φ°, and the insertion Y_W as a function of V, depend on the loop only through these arcs; and by the argument of Lemma 7.4(ii), which applies to the objects of a step as functions of the field, τ carries them to groups, clusters, and weights of C' with the same sizes and species, with $\Phi^\circ(\tau\Gamma; C') = \Phi^\circ(\Gamma; C)$, $Y_{\tau W}(C')(\tau V) = Y_W(C)(V)$, and $\mathsf{t}_{\tau W}(\tau V) = \mathsf{t}_W(V)$, where $(\tau V)(b) \coloneqq V(\tau^{-1} b)$. The same argument with τ^{-1} shows that $\Gamma \mapsto \tau\Gamma$ maps the clusters of $\sum'$ for W onto those for τW, so that $Q^{\mathsf{D}}(\tau W) = Q^{\mathsf{D}}(W)$. Finally the density ϱ_{ℓ_0} is composed of the same operations, so that $\varrho_{\ell_0}(\tau V) = \varrho_{\ell_0}(V)$, and the Haar measure on $G^{\mathcal{B}}$ is invariant under τ, the periods of the torus being multiples of Π; hence $\langle F \circ \tau^{-1} \rangle = \langle F \rangle$ for bounded measurable F, and with $F(V) \coloneqq \mathrm{tr}(Y_W(C)(V) \mathsf{t}_W(V)^{-1})$ this gives the claim.

Corners and sides. For a traversal $\mathfrak{p}$ of C let $\mathsf{y}_C(\mathfrak{p}) \coloneqq \sum \mathsf{z}_W(C)$, the sum over the small groups W with the first traversal $\mathfrak{p}$. The initial site of $\mathfrak{p}$ lies on a side of C of orientation v, at a distance $\mathsf{d}(\mathfrak{p}) \in [0, N_v)$ from the initial corner of that side. We call $\mathfrak{p}$ a corner traversal of that corner if $\mathsf{d}(\mathfrak{p}) < \mathsf{D}$, a corner traversal of the terminal corner of the side if $\mathsf{d}(\mathfrak{p}) \ge N_v - \mathsf{D}$, and a middle traversal otherwise; since $N_v \ge 3\mathsf{D}$, these classes partition the traversals. For a middle traversal $\mathfrak{p}$ the bonds of C with initial site at distance less than D from that of $\mathfrak{p}$ are the bonds of its own side at distance less than D along it, because the adjacent sides are at distance at least D and the opposite side at distance at least $3\mathsf{D}$. For $0 \le \varrho < \Pi$ let $\mathfrak{p}_v(\varrho)$ be the traversal of the side of R_1 of orientation v with $\mathsf{d}(\mathfrak{p}_v(\varrho)) = \mathsf{D} + \varrho$, a middle traversal, since $\varrho < \Pi \le \mathsf{D} \le N_v(R_1) - 2\mathsf{D}$. If $\mathsf{d}(\mathfrak{p}) \equiv \mathsf{D} + \varrho$ modulo Π, the translation τ with $\tau\mathfrak{p}_v(\varrho) = \mathfrak{p}$ has its vector in $\Pi\mathbb{Z}^4$, and it maps the bonds of R_1 near $\mathfrak{p}_v(\varrho)$ onto those of C near $\mathfrak{p}$; the agreement, applied with τ and τ^{-1}, gives $\mathsf{y}_C(\mathfrak{p}) = \mathsf{y}_{R_1}(\mathfrak{p}_v(\varrho))$. For a corner traversal $\mathfrak{p}$ of a corner $\mathfrak{c}$, the bonds of C with initial site at distance less than D from that of $\mathfrak{p}$ lie on the two sides at $\mathfrak{c}$, within $2\mathsf{D}$ of $\mathfrak{c}$. The translation which takes the corner of R_1 of the type of $\mathfrak{c}$ to $\mathfrak{c}$ has its vector in $\Pi\mathbb{Z}^4$, and it maps the corner traversals of that corner of R_1 onto those of $\mathfrak{c}$, together with the bonds near each; so the corner total $\mathsf{Z}_\mathfrak{c} \coloneqq \sum_\mathfrak{p} \mathsf{y}_C(\mathfrak{p})$, over the corner traversals of $\mathfrak{c}$, depends only on the type of $\mathfrak{c}$.

The sum. The middle traversals of the side of orientation v have $\mathsf{d}(\mathfrak{p}) = \mathsf{D}, \dots, N_v - \mathsf{D} - 1$; there are $N_v - 2\mathsf{D} \in \Pi\mathbb{Z}$ of them, and each residue of $\mathsf{d}(\mathfrak{p}) - \mathsf{D}$ modulo Π occurs equally often. Hence, the rearrangement being justified by the absolute convergence,

$$\sum_{W \text{ small}} \mathsf{z}_W(C) = \sum_\mathfrak{c} \mathsf{Z}_\mathfrak{c} + \sum_{v \in \mathcal{V}} \frac{N_v(C) - 2\mathsf{D}}{\Pi}\, \mathsf{S}_v \,, \qquad \mathsf{S}_v \coloneqq \sum_{\varrho=0}^{\Pi-1} \mathsf{y}_{R_1}(\mathfrak{p}_v(\varrho)) \,, \tag{13.40}$$

the first sum over the four corners of C and the terms with $N_v(C) = 0$ absent. Each rectangle has one corner of each of the four types and one side of each of the orientations $\pm e_\mu$, $\pm e_\nu$ (Lemma 13.1(i)), and the weights n_{R_n} satisfy $\sum_n n_{R_n} = 0$ and $\sum_n n_{R_n} N_v(R_n) = 0$ for every v, by the proof of Lemma 13.1(ii). Therefore $\sum_n n_{R_n} \sum_{W \text{ small}} \mathsf{z}_W(R_n) = 0$.

Conclusion. By (13.38), the bound on the large groups, (13.39), and the vanishing of the weighted sum of the small groups, $|\langle I(C)\rangle - \sum_{W \text{ small}} \mathsf{z}_W(C)| \le \mathsf{e}(C)$, where we used $e^{-\nu(\mathsf{D}/4-1)/M} \le e^{-\nu(\mathsf{D}/4-M)/(64M)}$; the moduli of the weights sum to 8, which gives (13.36). For the order, $I_*(C) \le C\bar{g}^2(\log \bar{g}^{-2})^{m'}$ and $\bar{h}(C) \le C\bar{g}(\log \bar{g}^{-2})^{4r+1/2}$ by the proof of Lemma 13.7(iv), while $\mathsf{D}/(4M) \ge L^3(\log \bar{g}^{-2})^r/18$ and $I_{\mathrm{wr}}(C)$ carries the factor $e^{-2c_{\mathrm{att}}L^3 R_{K'}}$, so that the last two terms of (13.37) decay faster than every power of $\bar{g}$, since $r = 2$. *q.e.d.*

It remains to treat the leading words. On $\mathcal{V}$ we write $U \coloneqq U_{K'}(V)$, $o_a \coloneqq o + ate_\mu$ for the initial corner of I_a, and, for $0 \le a \le b \le 3$, $\mathsf{L}_{a,b}$ for the closed path which runs from o_a along the bottom side of R_4 to o_b, around I_b, and back to o_a, so that $\mathsf{L}_{a,a} = I_a$. By the proof of Lemma 13.7(ii), applied to the defining representation of G, for which the number (1.7) is 1, every I_b satisfies $\|U(I_b) - 1\| \le h^G \coloneqq 1.06 \cdot \frac{3}{2} c_h B_3 P_{K'}(I_0)^2 \varepsilon^{\mathrm{act}}$ on $\mathcal{V}$, and $U(\mathsf{L}_{a,b})$ is unitarily conjugate to $U(I_b)$. If $h^G \le r_G$, the principal logarithm $X_{a,b} \coloneqq -i \log U(\mathsf{L}_{a,b})$ therefore lies in $\mathfrak{g}$, and $\|X_{a,b}\| \le 1.05\, h^G$, by the argument of the proof of Lemma 13.6 applied to the eigenvalues. We put $X_a \coloneqq X_{a,a}$,

$$\mathsf{K}_{a,b} \coloneqq \mathrm{tr}(\pi_*(X_a)\, \pi_*(X_{a,b})) \,, \qquad \Psi_k \coloneqq \langle \mathbf{1}_\mathcal{V} \mathsf{K}_{0,k} \rangle \,, \tag{13.41}$$

which are real, since both factors are Hermitian, and $\mathsf{I} \coloneqq 1.05\, c_\pi h^G$.

Lemma 13.13 (The Lasso)**.** *For $\bar g$ sufficiently small, depending on the quantities of Proposition 13.11, and for $K \geq K_{\rm w}$, $h^G \leq r_G$, $\mathsf{l} \leq 1/16$, and*

$$\sum_{n=1}^{4} n_{R_n} \langle f_{K'}^{(R_n)} - 1 \rangle = \Psi_3 - \Psi_2 + r_H \;, \qquad |r_H| \leq 512\, \mathsf{l}^3 + 16 \bar g^8 \;. \tag{13.42}$$

Proof. The conditions on h^G and l hold for $\bar g$ small, since $h^G \leq \bar h(I_0)/c_\pi$, which tends to zero with $\bar g$ by Lemma 13.7(iv).

The lasso. As closed paths from o, $\partial R_1 = \mathsf{L}_{0,0}$, and ∂R_{n+1} is $\mathsf{L}_{0,n}$ followed by ∂R_n: in the product the return of $\mathsf{L}_{0,n}$ along the bottom side cancels against the first part of ∂R_n, and the left side of I_n against the right side of R_n. Hence $U(\partial R_n) = U(\mathsf{L}_{0,n-1}) \cdots U(\mathsf{L}_{0,0})$. For $a \leq b$, the paths $\mathsf{L}_{0,a}$ and $\mathsf{L}_{0,b}$ both run first from o to o_a and finally back, so that $U(\mathsf{L}_{0,a}) = \mathsf{t} U(I_a) \mathsf{t}^{-1}$ and $U(\mathsf{L}_{0,b}) = \mathsf{t} U(\mathsf{L}_{a,b}) \mathsf{t}^{-1}$ with the same transport $\mathsf{t} \in G$ from o to o_a. The principal logarithm commutes with unitary conjugation, so $X_{0,a} = \mathsf{t} X_a \mathsf{t}^{-1}$ and $X_{0,b} = \mathsf{t} X_{a,b} \mathsf{t}^{-1}$, and since $\pi_*(gXg^{-1}) = \pi(g) \pi_*(X) \pi(g)^{-1}$, the numbers $\mathrm{tr}(\pi_*(X_{0,a}) \pi_*(X_{0,b}))$ equal $\mathsf{K}_{a,b}$.

The second order. Put $Z_a := \pi_*(X_{0,a})$, a Hermitian matrix with $\pi(U(\mathsf{L}_{0,a})) = e^{iZ_a}$ and $\|Z_a\| \leq c_\pi \|X_{0,a}\| \leq \mathsf{l}$, and $S := \sum_{a<n} \|Z_a\| \leq 4\mathsf{l} \leq \frac14$. Write $e^{iZ_a} = 1 + \mathsf{k}_a$ with $\mathsf{k}_a = iZ_a + \mathsf{j}_a$ and $\mathsf{j}_a = -\frac12 Z_a^2 + \mathsf{E}_a$; the exponential series gives $\|\mathsf{E}_a\| \leq \frac14 \|Z_a\|^3$, $\|\mathsf{j}_a\| \leq 0.57 \|Z_a\|^2$, and $\|\mathsf{k}_a\| \leq 1.3 \|Z_a\|$ for $\|Z_a\| \leq \frac14$. In the expansion of the ordered product of the $1 + \mathsf{k}_a$, the terms with three or more factors k have total norm at most $(1.3S)^3 e^{1.3S}/6 \leq 0.51 S^3$; the terms with two factors are $-\sum Z_a Z_b$, over the ordered pairs of the product, up to $0.66 \sum_{a<b} \|Z_a\| \|Z_b\| (\|Z_a\| + \|Z_b\|) \leq 0.22 S^3$, because the last sum is at most $S^3/3$; and $\sum_a \|\mathsf{E}_a\| \leq S^3/4$. So

$$\pi(U(\partial R_n)) = 1 + i \sum_{a<n} Z_a - \frac12 \sum_{a<n} Z_a^2 - \sum Z_a Z_b + R_n \;, \qquad \|R_n\| \leq S^3 \;. \tag{13.43}$$

By Lemma 13.6, $\mathrm{tr}\, Z_a = 0$, and $\mathrm{tr}(Z_a Z_b) = \mathrm{tr}(Z_b Z_a)$, so that $f_{K'}^{(R_n)} - 1 = -\frac12 \mathrm{tr}(\sum_{a<n} Z_a)^2 + r_n$ with $|r_n| \leq S^3 \leq 64\, \mathsf{l}^3$. With $\mathsf{v}(m) := \sum_{n>m} n_{R_n}$, so that $\mathsf{v}(0), \dots, \mathsf{v}(3) = 0, -1, 2, -1$,

$$\sum_{n=1}^{4} n_{R_n} \mathrm{tr}\Big(\sum_{a<n} Z_a\Big)^2 = \sum_{a,b\leq 3} \mathsf{v}(\max(a,b)) \,\mathrm{tr}(Z_a Z_b) = \sum_a \mathsf{v}(a) \mathsf{K}_{a,a} + 2 \sum_{a<b} \mathsf{v}(b) \mathsf{K}_{a,b} \;. \tag{13.44}$$

Stationarity. Let τ be the translation by $t_{\mathfrak b} e_\mu$, whose vector lies in $\Pi \mathbb{Z}^4$, acting on V as in the proof of Proposition 13.12. By the argument of Lemma 7.4(ii), $U_{K'}(\tau V) = \tau U_{K'}(V)$, so that $U_{K'}(\tau V)(\mathsf{L}_{a,b}) = U_{K'}(V)(\mathsf{L}_{a-1,b-1})$ for $a \geq 1$, and the event $\mathcal{V}$ is invariant under τ, since $A(U_{K'}(\cdot))$ and the support of ϱ_{ℓ_0} are. With the invariance of $\langle \cdot \rangle$ under τ, applied a times, $\langle \mathbf{1}_{\mathcal{V}} \mathsf{K}_{a,b} \rangle = \Psi_{b-a}$.

Conclusion. Hence $\langle \mathbf{1}_{\mathcal{V}} \sum_n n_{R_n} \mathrm{tr}(\sum_{a<n} Z_a)^2 \rangle = \Psi_0 \sum_a \mathsf{v}(a) + 2 \sum_{a<b} \mathsf{v}(b) \Psi_{b-a} = 2(\Psi_2 - \Psi_3)$, because $\sum_a \mathsf{v}(a) = 0$ and the coefficients of Ψ_1, Ψ_2, Ψ_3 in the second sum are $-1+2-1 = 0$, $2-1 = 1$, and -1. On $\mathcal{V}^c$ we use $|f_{K'}^{(R_n)} - 1| \leq 2$, the moduli of the weights summing to 8, and $\langle \mathbf{1}_{\mathcal{V}^c} \rangle \leq \bar g^8$. Together, $\sum_n n_{R_n} \langle f_{K'}^{(R_n)} - 1 \rangle = \Psi_3 - \Psi_2 + r_H$ with $|r_H| \leq 8 \cdot 64\, \mathsf{l}^3 + 16 \bar g^8$. *q.e.d.*

13.7 The Coarse Covariance at Flat Fields

The second moments Ψ_k are expectations under the measure $\langle \cdot \rangle$ of the stopping step, which at small coupling is concentrated near the fields V whose minimal configurations are nearly

flat. The integration by parts of Section 13.8 will express them, to the order $\bar{g}^2$, through a quadratic form evaluated at such fields, and in this section we compute that form at the fields which are exactly flat. There it is the covariance of the fluxes of a Gaussian field on $\mathbb{T}_{j_q}$, the block field of the free field, and the question is how far it is from the covariance of the free field itself, which Lemma 13.5 bounds below. The two differ by the covariance of the fluxes given the block averages (Lemma 13.15). This is the covariance of a constrained Gaussian field of Bałaban's, whose kernel decays exponentially on the scale of the blocks, so that it contributes almost nothing to the separated rectangles I_0 and I_k (Lemma 13.17). The integration by parts needs, besides, an inverse of the Hessian of the action, which has a kernel at a flat field; we regularise it by a small number λ, and the price, a term proportional to λ, is controlled by a coercivity estimate for the block field (Lemmata 13.18 and 13.19).

The setting. We use the units of the step K' of Section 8: the block field V lives on the bonds $\mathcal{B}$ of $\mathbb{T}_{j_q}$, which has the spacing 1 and N_q sites per side, and the fine lattice $\mathbb{T}_K$ has the spacing $\eta \coloneqq L^{-K'}$. In these units the rectangles have the sides $t_{\mathfrak{b}} = tL^{j_q}/u$ and $s_{\mathfrak{b}} = sL^{j_q}/u$, integers with $12t_{\mathfrak{b}} < N_q$ and $3s_{\mathfrak{b}} < N_q$, and we write l_a for the set of the plaquettes of $\mathbb{T}_K$ in the plane of e_μ and e_ν inside the rectangle bounded by I_a, so that $|\mathsf{l}_a| \coloneqq \eta^2 \# \mathsf{l}_a = s_{\mathfrak{b}} t_{\mathfrak{b}}$. For a field V which satisfies [7, (7), p. 278], the minimal configurations U with $M^{K'}U = V$ form one orbit of the gauge transformations of $\mathbb{T}_K$ which equal 1 at the sites of $\mathbb{T}_{j_q}$ [7, (4), p. 278; Theorem 1, p. 279], and $U_{K'}(V)$ denotes any of them. Since the corner o of the rectangles is a site of $\mathbb{T}_q \subset \mathbb{T}_{j_q}$, the holonomy of a closed path based at o is the same for all of them, so that the logarithms $X_{a,b}$ of Section 13.6 are functions of V; and $\mathsf{A}(V) \coloneqq A(U_{K'}(V))$, the Wilson action of the minimal configuration, is a function of V which is invariant under the gauge transformations of $\mathbb{T}_{j_q}$. On $\mathfrak{g}$ we use the inner product $X \cdot Y \coloneqq \operatorname{tr}(XY)$, which is invariant under Ad and is the product of [5, p. 392], with $|X|_2 \coloneqq (X \cdot X)^{1/2}$, so that $|X|_2 \le \|X\| \le N^{1/2}|X|_2$ because tr is normalised. We put $d_{\mathfrak{g}} \coloneqq \dim G$ and fix an orthonormal basis $(t_\alpha)_{\alpha \le d_{\mathfrak{g}}}$ of $\mathfrak{g}$.

The Casimir number. For the representation π of the eleven diagrams, with the normalised trace tr of its space, we define $C_2(\pi)$ and record a consequence of Schur's lemma:

$$\sum_{\alpha=1}^{d_{\mathfrak{g}}} \pi_*(t_\alpha)^2 = C_2(\pi) \cdot 1 \,, \qquad \operatorname{tr}(\pi_*(Y)\,\pi_*(Z)) = \frac{C_2(\pi)}{d_{\mathfrak{g}}}\, Y \cdot Z \quad (Y, Z \in \mathfrak{g}) \,. \tag{13.45}$$

Indeed, the sum does not depend on the orthonormal basis, since an orthogonal change of basis leaves $\sum_\alpha t_\alpha \otimes t_\alpha$ invariant, and $\pi(g)\pi_*(t_\alpha)\pi(g)^{-1} = \pi_*(\mathsf{Ad}(g)t_\alpha)$ maps it to the sum for the basis $(\mathsf{Ad}(g)t_\alpha)$; so it commutes with $\pi(G)$ and is a scalar, π being irreducible. The form $(Y, Z) \mapsto \operatorname{tr}(\pi_*(Y)\pi_*(Z))$ is real, symmetric, and Ad-invariant, so it equals $(SY) \cdot Z$ for a symmetric operator S on $\mathfrak{g}$ which commutes with $\mathsf{Ad}(G)$; an eigenspace of S is then invariant under $\mathsf{Ad}(G)$, hence an ideal of $\mathfrak{g}$, hence $\mathfrak{g}$ itself because G is simple, and the trace of the first identity gives $S = C_2(\pi)/d_{\mathfrak{g}}$. Finally $C_2(\pi) > 0$: the number $n_\pi C_2(\pi) = \sum_\alpha \operatorname{Tr}(\pi_*(t_\alpha)^2)$ is a sum of squared Hilbert–Schmidt norms of Hermitian matrices, which vanishes only if $\pi_* = 0$, and then $\pi = 1$ on the exponentials, which generate the connected group G, while π is not trivial.

The fluxes and their derivatives. Let $\mathcal{V}_3 \coloneqq \{V \in \operatorname{supp} \varrho_{\ell_0} \,|\, \mathsf{A}(V) < 3T_{\mathrm{ng}} g_{K'}^2\}$, an open neighbourhood of $\mathcal{V}$ in the support. The proof of Lemma 13.7(ii) with $3T_{\mathrm{ng}}$ in place of T_{ng}, whose conditions hold for $\bar{g}$ small by the same argument, multiplies $\varepsilon^{\mathrm{act}}$ by $\sqrt{3}$ and gives $\|U(I_b) - 1\| \le \sqrt{3} h^G$ on $\mathcal{V}_3$, so that the logarithms $X_{a,b}$ of Section 13.6 are defined on $\mathcal{V}_3$ and

have $\|X_{a,b}\| \le 1.05\sqrt{3}h^G \le 2h^G$ once $\sqrt{3}h^G \le r_G$. We put

$$X^\alpha := t_\alpha \cdot X_{0,0} \;, \qquad \Xi_\alpha := t_\alpha \cdot (X_{0,3} - X_{0,2}) \;, \qquad \text{so that} \qquad \mathsf{K}_{0,3} - \mathsf{K}_{0,2} = \frac{C_2(\pi)}{d_\mathfrak{g}} \sum_\alpha X^\alpha \, \Xi_\alpha \tag{13.46}$$

by (13.45). For $u \in \mathfrak{g}^\mathcal{B}$ we write $e^{iu}V$ for the field $(e^{iu_b}V(b))_b$, $\partial_u F(V) := \frac{\mathrm{d}}{\mathrm{d}s} F(e^{isu}V)|_{s=0}$ for a function F, and $\nabla F(V) \in \mathfrak{g}^\mathcal{B}$ for its gradient, with $\langle u, \nabla F\rangle = \partial_u F$ for the inner product $\langle u, u'\rangle := \sum_b u_b \cdot u'_b$; we put $\|u\| := \langle u, u\rangle^{1/2}$ and $\|u\|_\infty := \max_b \|u_b\| \le N^{1/2}\|u\|$, and we let $\Delta(V)$ be the symmetric operator on $\mathfrak{g}^\mathcal{B}$ with $\langle u, \Delta(V)u\rangle = \frac{\mathrm{d}^2}{\mathrm{d}s^2}\mathsf{A}(e^{isu}V)|_{s=0}$. For $\lambda > 0$ such that $\Delta(V) + \lambda$ is invertible we define

$$c_\lambda(V) := \sum_{\alpha=1}^{d_\mathfrak{g}} \langle \nabla X^\alpha, (\Delta(V) + \lambda)^{-1} \nabla \Xi_\alpha \rangle \;. \tag{13.47}$$

This number does not depend on the orthonormal basis, and it is invariant under the gauge transformations ω of $\mathbb{T}_{j_q}$: these act on $\mathfrak{g}^\mathcal{B}$ by the orthogonal map $u_b \mapsto \mathsf{Ad}(\omega(b_-))u_b$, under which ∇ and Δ are covariant, and on the logarithms based at o by $\mathsf{Ad}(\omega(o))$, which changes the basis. The functions X^α, Ξ_α, and A are real analytic on $\mathcal{V}_3$, by [7, Proposition 9, p. 309], as Lemma 13.21 below makes precise.

The linearisation at a flat field. We call V_0 *flat* if $V_0(\partial p) = 1$ for every plaquette p of $\mathbb{T}_{j_q}$. On the $\mathfrak{g}$-valued 1-forms and 2-forms of $\mathbb{T}_K$ we use the products $\langle A, A'\rangle := \eta^4 \sum_b A_b \cdot A'_b$ and $\langle F, F'\rangle := \eta^4 \sum_p F_p \cdot F'_p$, over the bonds and the positively oriented plaquettes, and for a configuration U we write D for the covariant exterior derivative of 1-forms at U, $(DA)(p)$ being the coefficient of $i\eta^2$ in $U'(\partial p)U(\partial p)^{-1}$ to first order for $U' = e^{i\eta A}U$, and ∂_U for the covariant derivative of 0-forms, as in [5, (3.8)–(3.10), p. 392], which writes D for both. Let Q be the linearisation at U of the average of order K', $M^{K'}(e^{i\eta A}U) = e^{i(QA+O(A^2))}M^{K'}(U)$ on each bond of $\mathcal{B}$, and Q^{dr} the linear part of the dressed average of Supplement, Lemma 17.7, at U. For a 2-form χ and $t \in \mathfrak{g}$ let $\chi \otimes t$ be the 2-form $p \mapsto \chi(p)t$, and let $\chi_a := \eta^{-2}\mathbf{1}_{\mathsf{l}_a}$, the plaquettes being oriented from e_μ to e_ν, so that $\langle \chi_a \otimes t, F\rangle = \eta^2 \sum_{p \in \mathsf{l}_a} t \cdot F(p)$, the flux of $t \cdot F$ through I_a.

Lemma 13.14 (Flat Fields)**.** *Let V_0 be flat, and let $\sigma_1, \dots, \sigma_4$ be integers such that each hyperplane $\{x_\beta \equiv \sigma_\beta + \frac{1}{2}\}$ has distance at least 3 from the rectangle bounded by R_4.*

(i) There are commuting $h_1, \dots, h_4 \in G$ and a gauge transformation ω of $\mathbb{T}_{j_q}$ with $\omega(o) = 1$ such that V_0^ω equals h_β on the bonds $\langle y, y + e_\beta\rangle$ with $y_\beta \equiv \sigma_\beta$ and 1 on the other bonds, and $U_{K'}(V_0^\omega)$ may be taken to be the flat configuration U_s which equals h_β on the bonds $\langle x, x + \eta e_\beta\rangle$ of $\mathbb{T}_K$ with $x_\beta \equiv \sigma_\beta + \frac{1}{2}(1 - \eta)$ and 1 on the other bonds.

(ii) Let $(E_r)_{r \le d_\mathfrak{g}}$ be an orthonormal basis of $\mathfrak{g}^\mathbb{C}$, for the Hermitian extension of the product, with $\mathrm{Ad}(h_\beta)E_r = e^{2\pi i\theta_{r,\beta}}E_r$ and $\theta_r \in [0,1)^4$. In the gauge of (i), the components $\mathrm{tr}(E_r^* B)$ of a $\mathfrak{g}$-valued form B on $\mathbb{T}_K$ or on $\mathbb{T}_{j_q}$, read in the box $\prod_\beta(\sigma_\beta + \frac{1}{2}, \sigma_\beta + \frac{1}{2} + N_q)$ and extended by $B(x + N_q e_\beta) = e^{2\pi i\theta_{r,\beta}}B(x)$, are scalar forms with the twisted periodicity θ_r; the operators D, ∂_U, Q, Q^{dr} at U_s, and the translations of $\mathbb{T}_{j_q}$ by whole blocks, act on each component as the corresponding operators at $U = 1$ on such forms, Q^{dr} as the average $Q_{K'}$ of [2, (1.18), p. 20] with centred blocks.

(iii) Let H be a linear map from the forms on $\mathcal{B}$ to the 1-forms on $\mathbb{T}_K$ such that $QHu = u$ and $\|DHu\| = \min\{\|DA\| \,|\, QA = u\}$, and $K := (DH)^* DH$. The operator DH does not depend on the choice of H and is the same for Q^{dr} in place of Q, and so is the subspace $D(\ker Q)$.

(iv) In the gauge of (i), $\Delta(V_0) = K$, $\nabla X^\alpha(V_0) = H^* D^*(\chi_0 \otimes t_\alpha)$, *and* $\nabla \Xi_\alpha(V_0) = H^* D^*((\chi_3 - \chi_2) \otimes t_\alpha)$.

Proof. (i) The torus cut along the four hyperplanes $\{x_\beta \equiv \sigma_\beta + \frac{1}{2}\}$ is a box, and the transport of V_0 from o along paths in the box does not depend on the path, V_0 being flat; with $\omega(y)$ the inverse of the transport from o to y, $V_0^\omega = 1$ on the bonds of the box, and on a bond which crosses the hyperplane of direction β it equals the holonomy h_β of a closed path from o which winds once in that direction, independently of the bond by flatness. Two such holonomies commute, since the closed path around the cycles β and γ and back is contractible. The configuration U_s is flat, since a plaquette which meets the fine bonds of a hyperplane contains two of them with opposite orientations, or four at the intersection of two hyperplanes, where the h_β commute. Near the hyperplane of direction β it is the gauge transform of 1 by the transformation equal to 1 on one side and to h_β^{-1} on the other, which is 1 or h_β^{-1} at the sites of $\mathbb{T}_{j_q}$, and the average [8, (0.4), p. 253] is covariant under such a transformation through its values at the endpoints of each bond [8, (0.6), p. 253]; so all the averages of U_s are configurations of the same form, and $M^{K'} U_\mathrm{s} = V_0^\omega$. Its action is 0, so it is a minimum, and it lies on the unique critical orbit [7, Theorem 1, p. 279].

(ii) The gauge transformations which cross the hyperplanes act on $\mathfrak{g}$ by $\mathsf{Ad}(h_\beta^{\pm 1})$, which multiplies the component r by $e^{\pm 2\pi i \theta_{r,\beta}}$; off the hyperplanes $U_\mathrm{s} = 1$, and the operators are covariant (Supplement, Lemma 17.7(a), and [5, (3.32), p. 395]), so in the extended components they are the operators at $U = 1$. At $U = 1$ each step of the dressed average is the straight average with centred blocks (Supplement, proof of Lemma 17.7), whose composite is [2, (1.18), p. 20].

(iii) The set $\mathcal{F}(u) := \{DA \,|\, QA = u\}$ is an affine subspace, so it has a unique point of least norm, which is DHu for every admissible H. At U_s the gauge directions satisfy $D\partial_U = 0$, and differentiating the covariance [8, (0.6), p. 253] gives $Q\partial_U \tilde{\omega} = \bar{D}(\tilde{\omega}|_{\mathbb{T}_{j_q}})$, with $\bar{D}$ the covariant derivative of 0-forms of $\mathbb{T}_{j_q}$ at V_0^ω; so $\mathcal{F}(u) = \mathcal{F}(u + \bar{D}\omega')$ for every 0-form ω' of $\mathbb{T}_{j_q}$, the restriction to the sites being onto. The linearisation of [8, (0.4)] at 1 averages A over the paths $\Gamma \cup [x, x'] \cup (-\Gamma')$, with Γ from c_- to x and Γ' from c_+ to x' inside the blocks, since the factors $(-c)$ and $\mathbf{U}(c)$ cancel to first order; the parts along Γ and Γ' are the values at c_- and c_+ of a 0-form ψ_A of $\mathbb{T}_{j_q}$, so that one step is the straight average minus $\bar{D}\psi_A$. Composing over the steps, with the identity $Q_{K'} \partial \tilde{\omega} = \bar{D} Q'_{K'} \tilde{\omega}$ of [2, (1.20), p. 20], $Q - Q^{\mathrm{dr}}$ takes values in the range of $\bar{D}$ in each component by (ii), so $\mathcal{F}(u)$ is the same for Q^{dr}, and DH is the same. Finally $D(\ker Q)$ is the orthogonal complement of $D(\operatorname{ran} H)$ in $\operatorname{ran} D$ by Lemma 13.15(ii) below, for Q and for Q^{dr}.

(iv) Let $u \in \mathfrak{g}^{\mathcal{B}}$. The map $A \mapsto \frac{1}{i} \log(M^{K'}(e^{i\eta A} U_\mathrm{s})(V_0^\omega)^{-1})$ is analytic near 0 with the derivative Q, which has the right inverse H, so the implicit function theorem gives an analytic family $A_\mathrm{c}(s) = sHu + O(s^2)$ with $M^{K'}(e^{i\eta A_\mathrm{c}(s)} U_\mathrm{s}) = e^{isu} V_0^\omega$. At the flat U_s the expansion [5, (3.12), p. 392] has $J = 0$ and $\Delta = D^* D$, since the form Δ' of [5, (3.10)] vanishes, so that $\mathsf{A}(e^{isu} V_0^\omega) \leq A(e^{i\eta A_\mathrm{c}(s)} U_\mathrm{s}) = \frac{1}{2} s^2 \|DHu\|^2 + O(s^3)$. Conversely, by [7, Proposition 9, p. 309], $U_{K'}(e^{isu} V_0^\omega) = (e^{i\eta A(s)} U_\mathrm{s})^{w(s)}$ with A and the gauge transformation w analytic, $A(0) = 0$, $w(0) = 1$, and by the covariance $M^{K'}(e^{i\eta A(s)} U_\mathrm{s}) = (e^{isu} V_0^\omega)^{w(s)^{-1}}$ with w read at the sites, so that $QA'(0) = u + \bar{D}\omega'$ for a 0-form ω'; hence $DA'(0) \in \mathcal{F}(u)$ and $\mathsf{A}(e^{isu} V_0^\omega) = \frac{1}{2} s^2 \|DA'(0)\|^2 + O(s^3) \geq \frac{1}{2} s^2 \|DHu\|^2 + O(s^3)$. Therefore $\langle u, \Delta(V_0) u \rangle = \|DHu\|^2$ and $DA'(0) = DHu$. The loops $\mathsf{L}_{0,k}$ lie at distance at least 3 from the hyperplanes, where $U_\mathrm{s} = 1$, so by the lattice Stokes formula the holonomy of $\mathsf{L}_{0,k}$ at $e^{i\eta A(s)} U_\mathrm{s}$ is $1 + is\eta^2 \sum_{p \in \mathsf{l}_k} DA'(0)(p) + O(s^2)$, the conjugation by the transport along the bottom side of R_4 and by $w(s)(o)$ changing only terms of the second order because the holonomy at $s = 0$ is 1. Hence $\partial_u X^\alpha = \langle \chi_0 \otimes t_\alpha, DHu \rangle$ and

$\partial_u \Xi_\alpha = \langle (\chi_3 - \chi_2) \otimes t_\alpha, DHu \rangle$, which is the claim. *q.e.d.*

At a flat field the problem has become linear, and in each component it is the problem of the free field with a twist. We first split the fluxes, in an abstract setting, into the part which the block averages determine and the rest.

Lemma 13.15 (Flux Splitting)**.** *Let E, F, E' be finite-dimensional real or complex inner product spaces, $D\colon E \to F$ and $Q\colon E \to E'$ linear, Q onto.*

(i) There is a linear $H\colon E' \to E$ with $QH = 1$ such that HB minimises $\|DA\|$ on $\{QA = B\}$ for every B, and every such H has $D^*DH(E') \subset \operatorname{ran} Q^*$.

(ii) $\operatorname{ran} D = D(\operatorname{ran} H) \oplus D(\ker Q)$*, the two summands being orthogonal.*

(iii) $\Pi_{\operatorname{ran} D} = \Pi_{D \operatorname{ran} H} + \Pi_{D \ker Q}$*, and* $\Pi_{D \operatorname{ran} H} = DH(H^*D^*DH)^+H^*D^*$*, where Π denotes the orthogonal projection and $^+$ the Moore–Penrose inverse.*

Proof. (i) Let ι be the inclusion of $\ker Q$. Since Q is onto, $QQ^+ = 1$ and $\{QA = B\} = Q^+B + \ker Q$. Put $H := (1 - \iota(D\iota)^+D)Q^+$, so that $QH = 1$ because $Q\iota = 0$. Then $DHB = (1 - D\iota(D\iota)^+)DQ^+B$ is DQ^+B minus its orthogonal projection onto $D(\ker Q) = \operatorname{ran} D\iota$, that is the point of least norm of $DQ^+B + D(\ker Q)$, so HB is a minimiser. At every minimiser HB the first variation gives $\langle DHB, Dz \rangle = 0$ for $z \in \ker Q$, so $D^*DHB \in (\ker Q)^\perp = \operatorname{ran} Q^*$.

(ii) For $z \in \ker Q$ and $B \in E'$, by (i), $\langle DHB, Dz \rangle = \langle Q^*y, z \rangle = \langle y, Qz \rangle = 0$ for some y; and every A is $HQA + (A - HQA)$ with $Q(A - HQA) = 0$.

(iii) The first identity follows from (ii), and the second is $T(T^*T)^+T^* = \Pi_{\operatorname{ran} T}$ for $T := DH$. *q.e.d.*

We apply the lemma with E the $\mathfrak{g}$-valued 1-forms on $\mathbb{T}_K$, F the 2-forms, E' the forms on $\mathcal{B}$, and D, Q, H of Lemma 13.14 at a flat field. In the component r the projection $\Pi_{\operatorname{ran} D}$ is diagonal in the momenta $p = (2\pi/N_q)(n + \theta_r)$, and its entry on the plaquettes of the plane of e_μ and e_ν is $(\hat p_\mu^2 + \hat p_\nu^2)/\hat p^2$, the momentum $p = 0$ being absent if $\theta_r = 0$: the range of D at the momentum p consists of the forms $a \wedge b$ with $a_\beta = \eta^{-1}(e^{i\eta p_\beta} - 1)$, $|a_\beta|^2 = \hat p_\beta^2$, and $|\iota_{\bar a}(e_\mu \wedge e_\nu)|^2/|a|^2$ is that entry. Hence, with the kernel (13.6) at $\mathsf{T} = N_q$ and the number (13.7) for $\mathsf{I} = \mathsf{I}_0$,

$$\operatorname{Re}\langle \chi_0, \Pi_{\operatorname{ran} D, r} \chi_k \rangle = c^{\theta_r}(kt_{\mathfrak{b}}) \,, \qquad \mathfrak{f}_r := \operatorname{Re}\langle \chi_0, \Pi_{D \ker Q, r}(\chi_3 - \chi_2) \rangle \,, \tag{13.48}$$

where the index r marks the operator on the component r; $\mathfrak{f}_r$ is the *fluctuation part* of the third difference. In the language of Gaussian fields, Lemma 13.15(iii) says that the fluxes of the free field, with covariance $\Pi_{\operatorname{ran} D}$, are the sum of the fluxes of the block field, with covariance $\Pi_{D \operatorname{ran} H}$, and of an independent field with the covariance given the block averages.

To bound $\mathfrak{f}_r$ we need pointwise values of a field which is known through an equation and through local L^2-norms. The next lemma provides them; it is a discrete interior estimate, uniform in the spacing.

Lemma 13.16 (An Interior Bound)**.** *On $\varepsilon\mathbb{Z}^4$, $\varepsilon \le 1/16$, let $\nabla_\kappa f(x) := \varepsilon^{-1}(f(x + \varepsilon e_\kappa) - f(x))$, ∇_κ^* its adjoint for $\langle f, h \rangle := \varepsilon^4 \sum_x \bar f(x) h(x)$, $\Delta := \sum_\kappa \nabla_\kappa^* \nabla_\kappa$, B_ρ the Euclidean ball of radius ρ about a point x_0, and $\|f\|_{B_\rho} := \|f \mathbf{1}_{B_\rho}\|$. There is C_{LR}, a number, such that if f and $g_1, \dots, g_4$ are complex functions with $\Delta f = \sum_\kappa c_\kappa \nabla_\kappa g_\kappa$ on B_1, for constants $|c_\kappa| \le 1$, then $|f(x)| \le C_{\mathrm{LR}}(\|f\|_{B_1} + \sup_{B_1} |g|)$ for $x \in B_{1/2}$.*

Proof. Let G be the Green's function of the simple random walk on $\mathbb{Z}^4$, with $\mathcal{L}G = -\mathbf{1}_{\{0\}}$ for the generator $\mathcal{L}$, and $\mathcal{G}(z) := \frac{1}{8}\varepsilon^{-2}G(z/\varepsilon)$. Since Δ acts on functions of y/ε as $-8\varepsilon^{-2}\mathcal{L}$, $\Delta_y\mathcal{G}(x-y) = \varepsilon^{-4}\mathbf{1}_{\{y=x\}}$, and $h(x) = \varepsilon^4\sum_y \mathcal{G}(x-y)(\Delta h)(y)$ for finitely supported h. By [16, Theorem 4.3.1], $G(z) = a_4|z|^{-2} + O(|z|^{-4})$, so that $|G(z)| \le c|z|^{-2}$ and, by the mean value theorem for the leading term, $|G(z+e_\kappa) - G(z)| \le c|z|^{-3}$ for $z \neq 0$, while $G(0)$ is finite; hence $|\mathcal{G}(z)| \le c(|z| \vee \varepsilon)^{-2}$ and $|\nabla_\kappa\mathcal{G}(z)|, |\nabla^*_\kappa\mathcal{G}(z)| \le c(|z| \vee \varepsilon)^{-3}$, with numbers c. Extend f and g by 0 off B_1, let $\psi \in C^\infty(\mathbb{R}^4)$ with $\psi = 1$ on $B_{3/4}$ and $\operatorname{supp}\psi \subset B_{7/8}$, and let $x \in B_{1/2}$. Then $f(x) = (\psi f)(x)$ is the sum of $\varepsilon^4\sum_y \mathcal{G}(x-y)\psi(y)(\Delta f)(y)$ and $\varepsilon^4\sum_y \mathcal{G}(x-y)([\Delta,\psi]f)(y)$, in which only values in $B_{7/8+2\varepsilon} \subset B_1$ enter. Summing by parts, the first sum is $\sum_\kappa c_\kappa\varepsilon^4\sum_y \nabla^*_\kappa(\psi\mathcal{G}(x-\cdot))(y)g_\kappa(y)$, and $|\nabla^*_\kappa(\psi\mathcal{G}(x-\cdot))(y)| \le c'(|x-y| \vee \varepsilon)^{-3}$ on B_1, whose ε^4-sum over B_1 is at most a number c''; so the first sum is at most $4c''\sup_{B_1}|g|$. In the second, $[\Delta,\psi]h(y) = -\varepsilon^{-2}\sum_\kappa\sum_\pm(\psi(y\pm\varepsilon e_\kappa)-\psi(y))h(y\pm\varepsilon e_\kappa)$ vanishes unless $\frac{3}{4}-\varepsilon \le |y-x_0| < \frac{7}{8}+\varepsilon$, so that $|x-y| \ge \frac{1}{8}$ there, and $[\Delta,\psi]^* = -[\Delta,\psi]$. With $\psi(y\pm\varepsilon e_\kappa) - \psi(y) = \pm\varepsilon\partial_\kappa\psi(y) + r_\pm$, $|r_\pm| \le \frac{1}{2}\varepsilon^2\|\partial^2\psi\|_\infty$, we get $|[\Delta,\psi]\mathcal{G}(x-\cdot)(y)| \le c'''$, a number, so the second sum is at most $c'''\varepsilon^4\sum_{B_1}|f| \le c'''|B_2|^{1/2}\|f\|_{B_1}$ by the Cauchy–Schwarz inequality. *q.e.d.*

The fluctuation part is a covariance of Bałaban's constrained Gaussian field, and its decay is inherited from his bounds on the propagators.

Lemma 13.17 (The Fluctuation Part)**.** *There are C_7 and $\delta_2 > 0$, depending on d, L, and G, such that at every flat field and for every component r, $|\mathfrak{f}_r| \le C_7(s/t)^2e^{-\delta_2 t_b/2}$.*

Proof. We work at U_s in the gauge of Lemma 13.14(i), with the sequence $\Omega_j = \mathbb{T}$ of the term ℓ_0 and the dressed average, so that $Q = Q^{\mathrm{dr}}$ by Lemma 13.14(iii); distances are in units of the blocks of $\mathbb{T}_{j_q}$, where the distance of [3, (2.46), p. 231] is at least the ℓ^1-distance of the blocks.

Bałaban's constrained covariance. The cubes of the class [5, (3.35), p. 396] have side $O(1)M < N_q$, so on each of them U_s is a gauge transform of 1, and it satisfies [5, (3.35)–(3.36)] with $A = 0$. By Supplement, Lemma 17.7(c), Bałaban's operators and [5, Theorems 3.1, 3.3, 3.13] are available for it with constants of d, L, and G, and Supplement, Lemma 17.8, gives [5, (3.132), p. 422]. Since U_s is flat, the form Δ' and the current J of [5, (3.10)–(3.11), p. 392] vanish, hence so do Δ'_π, $\Delta^{(2)}$, and $\Delta^{(2)}_\pi$ by [5, (3.120), (3.134), (3.135), pp. 419–422]; by [5, (3.119)] and $D\partial_U = 0$, $\Delta_\pi = D^*D$; and by [5, (3.122), (3.128)], with $a = 1$, $G_1 = G = (D^*D + \partial_U R\partial_U^* + Q^*Q)^{-1}$, where R is Bałaban's projection on the 0-forms. The covariance $\mathfrak{G}$ of the Gaussian field constrained by $QA = 0$ and $R\partial_U^*A = 0$ [5, (3.148), p. 425] is then

$$\mathfrak{G} = G - G\partial_U R\partial_U^* G - GQ^*(QGQ^*)^{-1}QG \tag{13.49}$$

by [5, (3.153), p. 426], and it has the bounds of [5, Theorem 3.3] except the one with the covariant Laplacian [5, Theorem 3.13, p. 426].

The flux covariance. For a 1-form λ let $v := \mathfrak{G}\lambda$ and $w := (QGQ^*)^{-1}QG\lambda$. Then $v = G(\lambda - \partial_U R\partial_U^* G\lambda - Q^*w)$, and $Qv = 0$, $R\partial_U^* v = 0$, because $\mathfrak{G} = \mathfrak{P}G_1$ with $\mathfrak{P}$ the projection onto $S := \{QA = 0, R\partial_U^*A = 0\}$ [5, (3.153); p. 425]. Applying G^{-1} we obtain

$$D^*Dv = \lambda - \partial_U R\partial_U^* G\lambda - Q^*w\,, \tag{13.50}$$

and for $z \in S$ it gives $\langle D^*Dv - \lambda, z\rangle = -\langle\partial_U^*G\lambda, R\partial_U^*z\rangle - \langle w, Qz\rangle = 0$, R being an orthogonal projection. Moreover $D(S) = D(\ker Q)$: for $z \in \ker Q$, the form $z' := z - \partial_U G'R\partial_U^* z$ has $Qz' = 0$, because $Q\partial_U G'R = \bar{D}Q'G'R = 0$ [5, p. 426, l. 17], $R\partial_U^*z' = 0$, because z' is z

transformed to the gauge $R\partial_U^* A = 0$ [5, p. 419, ll. 26–27], and $Dz' = Dz$. For $\lambda \coloneqq D^*\phi$ we therefore have $Dv \in D(\ker Q)$ and $Dv - \phi \perp D(\ker Q)$, that is,

$$D\mathfrak{G}D^* = \Pi_{D\ker Q} \,, \qquad \mathfrak{f}_r = \mathrm{Re}\langle \chi_0, D\mathfrak{G}D^*(\chi_3 - \chi_2)\rangle_r \,, \tag{13.51}$$

all these operators commuting with the decomposition into components.

The local equation. Let x_0 be the base point of a plaquette of $\mathsf{l}_0 \cup \mathsf{l}_2 \cup \mathsf{l}_3$, B_ρ the ball of radius ρ about it, and ϕ a 2-form which vanishes on B_2. On B_3 we have $U_\mathrm{s} = 1$, so $D = d$ and $\partial_U = \nabla$ there. For $F \coloneqq D\mathfrak{G}D^*\phi$, $dF = 0$ on B_2, and (13.50) with $\lambda = D^*\phi = 0$ on $B_{3/2}$ gives $d^*F = -Q^*w + \nabla(\cdot)$ there, hence $dd^*F = -dQ^*w$. On 2-forms $dd^* + d^*d = \sum_\kappa \nabla_\kappa^* \nabla_\kappa$ componentwise, because the forward and backward differences commute, so $\Delta F_{\mu\nu} = -\nabla_\mu (Q^*w)_\nu + \nabla_\nu (Q^*w)_\mu$ on B_1, and Lemma 13.16, for each of the six components of each component r, gives

$$|F(x_0)| \le \sqrt{6}\, C_{\mathrm{LR}} \Big(\|F\|_{B_1} + \sup_{B_1} |Q^*w| \Big) \,. \tag{13.52}$$

The multipliers. The average Q is a mean of line integrals over one block length [2, (1.18), p. 20], so for a bond b of $\mathcal{B}$ and the unit form e_b at b, $|Q^*e_b| \le 1$, Q^*e_b is supported on the fine bonds between the blocks of b_- and b_+, $|QA(b)| \le \sup|A|$, and $|QA(b)| \le \sqrt{2}\|A\|_{\tilde\Delta(b)}$, with $\tilde\Delta(b)$ the two blocks which it reads. On 1-forms the components of the covariant gradient ∇_U of [5, (3.39), p. 397] are $D_\mu A_\nu$ and $(DA)_{\mu\nu} = D_\mu A_\nu - D_\nu A_\mu$, so $|DA| \le 2|\nabla_U A|$ pointwise. Let $\delta_2 \coloneqq \frac{1}{2}\min(\delta_0, \delta_1)$ with δ_0 of [5, Theorem 3.1] and δ_1 of (13.53) below, and $C_\delta \coloneqq (2/(1 - e^{-\delta_2}))^4 \ge \sum_{y \in \mathbb{Z}^4} e^{-\delta_2 |y|_1}$. At one scale [5, (3.132)] reads

$$|(QGQ^*)^{-1}(b, b')| \le O(1) e^{-\delta_1 d(b,b')} \,. \tag{13.53}$$

For $\phi = \chi_k$ we have $QGD^*\phi(b) = \langle \chi_k, DGQ^* e_b \rangle$, and (13.53) together with [5, (3.42)] for G [5, Theorem 3.3] gives $|w(y)| \le C_2 |\mathsf{l}_k| e^{-\delta_2 d(y, \mathsf{l}_k)}$, with $d(y, \mathsf{l}_k)$ the distance of y from the blocks which meet l_k, the convolution of the two exponentials being bounded by $e^{-\delta_2 d} C_\delta$. For ϕ supported in $B_1(x)$ with $\|\phi\| \le 1$ the same argument with [5, (3.46)] for G and $|QA(b)| \le \sqrt{2}\|A\|_{\tilde\Delta(b)}$ gives $|w_\phi(y)| \le C_4 e^{-\delta_2 d(y, B_1(x))}$; here and below the C_i depend on d, L, and G.

The bound. Fix $k \in \{2, 3\}$, the base point x of a plaquette of l_0, and $F \coloneqq D\mathfrak{G}D^*\chi_k$. Then $\|F\|_{B_1(x)}$ is the supremum of $|\langle \phi, F\rangle|$ over the ϕ supported in $B_1(x)$ with $\|\phi\| \le 1$, and $\langle \phi, F\rangle = \langle D\mathfrak{G}D^*\phi, \chi_k\rangle = \eta^2 \sum_{p \in \mathsf{l}_k} F_\phi(p)$ with $F_\phi \coloneqq D\mathfrak{G}D^*\phi$, $\mathfrak{G}$ being symmetric. The blocks which meet $B_1(x)$ or $B_1(x')$, $x' \in \mathsf{l}_k$, and their neighbours are at distance at least $(k-1)t_\mathrm{b} - 5 \ge t_\mathrm{b} - 5$; so (13.52) at x', with [5, (3.46)] for $\mathfrak{G}$ and the bound on w_ϕ, gives $|F_\phi(x')| \le C_5 e^{-\delta_2(t_\mathrm{b}-5)}$, hence $\|F\|_{B_1(x)} \le C_5 |\mathsf{l}_k| e^{-\delta_2(t_\mathrm{b}-5)}$. Then (13.52) at x, with the bound on w, gives $|F(x)| \le C_6 |\mathsf{l}_k| e^{-\delta_2(t_\mathrm{b}-5)}$, and $|\langle \chi_0, F\rangle| \le C_6 |\mathsf{l}_0| |\mathsf{l}_k| e^{-\delta_2(t_\mathrm{b}-5)}$. Since $|\mathsf{l}_a| = s_\mathrm{b} t_\mathrm{b} = (s/t) t_\mathrm{b}^2$ and $y^4 e^{-\delta_2 y/2} \le (8/(e\delta_2))^4$, the two values of k give the claim with $C_7 \coloneqq 2C_6 e^{5\delta_2} (8/(e\delta_2))^4$. *q.e.d.*

The coarse form K has a kernel, and the following estimate controls how far a coarse field orthogonal to the gauge directions can vary without costing action.

Lemma 13.18 (Coercivity of the Block Field)**.** *At a flat field, in a component r, let $\bar D$ be the covariant derivative of the 0-forms of $\mathbb{T}_{j_q}$ and ∇_μ the difference of a form on $\mathcal{B}$ in the direction μ, acting on each of its components.*

(i) $\operatorname{ran} K \subset \ker \bar D^*$.

(ii) Every form u on $\mathcal{B}$ with $\bar{D}^*u = 0$ has $\|\nabla_\mu u\| \le 4\|DHu\|$.

Proof. By Lemma 13.14(ii) and (iii) we may lift the component to quasi-periodic forms, on which $D = d$, $Q = Q_{K'}$ with centred blocks, and the products are taken over one period. (i) For a 0-form ω of $\mathbb{T}_{j_q}$ let $\tilde{\omega}$ be constant on each block with $Q'_{K'}\tilde{\omega} = \omega$. By [2, (1.20), p. 20], $Q\partial\tilde{\omega} = \bar{D}\omega$, and $d\partial\tilde{\omega} = 0$; since the curvature of a minimiser is unique, $DH\bar{D}\omega = 0$. Hence $K\bar{D} = 0$ and $\bar{D}^*K = 0$.

(ii) Put $A := Hu$ and $F := dA$, so that $u = QA$. By [2, (1.18)], $u_b = \sum_{x\in B(b_-)} \eta^4 \int_{[x,x+e_\nu]} A$ for $b = \langle b_-, b_- + e_\nu\rangle$, the line integral being η times the sum over the fine bonds. For $\nu \neq \mu$ the lattice Stokes formula on the square S_x spanned by $[x, x+e_\nu]$ and $[x, x+e_\mu]$ gives $\int_{[x,x+e_\nu]+e_\mu} A - \int_{[x,x+e_\nu]} A = \psi(x+e_\nu) - \psi(x) + \Phi_F(S_x)$, with $\psi(z) := \int_{[z,z+e_\mu]} A$ and $\Phi_F(S) := \eta^2 \sum_{q\subset S} F_{\mu\nu}(q)$; for $\nu = \mu$ it holds with $\Phi_F := 0$. The translation by e_μ maps $B(b_-)$ onto $B(b_- + e_\mu)$ and that by e_ν maps it onto $B(b_+)$, so averaging gives

$$\nabla_\mu u = \bar{D}(Q'_{K'}\psi) + \mathcal{S}F\,, \qquad (\mathcal{S}F)_b := \sum_{x\in B(b_-)} \eta^4 \Phi_F(S_x)\,. \tag{13.54}$$

If $\bar{D}^*u = 0$, then $\bar{D}^*\nabla_\mu u = \nabla_\mu \bar{D}^* u = 0$, so $\|\nabla_\mu u\|^2 = \langle \nabla_\mu u, \mathcal{S}F\rangle$ and $\|\nabla_\mu u\| \le \|\mathcal{S}F\|$. A fine plaquette of the plane of e_μ and e_ν lies in S_x for at most η^{-2} points x of $B(b_-)$, and all the S_x lie in $R_b := B(b_-) + [0,1]e_\mu + [0,1]e_\nu$, which contains at most $4\eta^{-4}$ such plaquettes; so $|(\mathcal{S}F)_b| \le \eta^4 \sum_{q\subset R_b} |F_{\mu\nu}(q)| \le 2(\eta^4 \sum_{q\subset R_b} |F_{\mu\nu}(q)|^2)^{1/2}$ by the Cauchy–Schwarz inequality. Each plaquette lies in R_b for at most four bonds b of each direction, so $\|\mathcal{S}F\|^2 \le 16\|F\|^2$. *q.e.d.*

Lemma 13.19 (The Flux of the Block Field)**.** *At a flat field, in each component, $\|\Pi_{D\,\mathrm{ran}\,H}\chi_0\|^2 \le C_V(s_\mathrm{b}+2)(t_\mathrm{b}+2)$, with C_V depending on d, L, and G.*

Proof. The norm is the supremum of $|\langle\chi_0, F\rangle|$ over the $F \in D(\mathrm{ran}\,H)$ with $\|F\| = 1$. Let $F = DHu$. By Lemma 13.15(i), $D^*F = Q^*\sigma$ for some σ, and applying H^*, $\sigma = H^*D^*DHu = Ku$. Near a base point x_0 of a plaquette of l_0 we have $D = d$, so $dF = 0$, $d^*F = Q^*Ku$, and $\Delta F = dQ^*Ku$ on $B_1(x_0)$, and Lemma 13.16 gives $|F(x_0)| \le \sqrt{6}\,C_{\mathrm{LR}}(\|F\|_{B_1(x_0)} + \sup_{B_1(x_0)}|Q^*Ku|)$. By [5, (3.133), p. 422], with the factor of [5, p. 422, ll. 32–33] for the variational H, which is 1 at one scale, and $|DA| \le 2|\nabla_U A|$, $|DHe_b(x)| \le C_H e^{-\delta_H|y(x)-b_-|_1}$, with $\delta_H := \frac{1}{2}\delta_1$ and $y(x)$ the block of x. With the weighted norms $\|F\|^2_{w,y} := \eta^4 \sum_x e^{-\delta_H|y(x)-y|_1}|F(x)|^2$ and $C'_\delta := (2/(1-e^{-\delta_H}))^4$, the Cauchy–Schwarz inequality gives $|(Ku)_b| = |\langle DHe_b, F\rangle| \le C_H C'^{1/2}_\delta \|F\|_{w,b_-}$. A fine bond lies in the supports of at most two of the Q^*e_b, $B_1(x_0)$ meets these supports only for $|b_- - y(x_0)|_1 \le 5$, and it lies in the blocks with $|y - y(x_0)|_1 \le 4$, so that $|F(x_0)| \le C_8\|F\|_{w,y(x_0)}$ with $C_8 := \sqrt{6}\,C_{\mathrm{LR}}(e^{2\delta_H} + 2C_H C'^{1/2}_\delta e^{5\delta_H})$. Now $\langle\chi_0, F\rangle = \eta^2 \sum_{p\in\mathsf{l}_0} F_{\mu\nu}(p)$, and l_0 meets at most $(s_\mathrm{b}+2)(t_\mathrm{b}+2)$ blocks, in each of which it has at most η^{-2} plaquettes; by the Cauchy–Schwarz inequality over these blocks and $\sum_y e^{-\delta_H|y(x)-y|_1} \le C'_\delta$, $|\langle\chi_0, F\rangle| \le C_8 \sum_y \|F\|_{w,y} \le C_8 C'^{1/2}_\delta((s_\mathrm{b}+2)(t_\mathrm{b}+2))^{1/2}\|F\|$, which gives the claim with $C_V := C_8^2 C'_\delta$. *q.e.d.*

We can now compute the number (13.47) at a flat field. The regularisation enters through the operator P below, which is positive and is small on the fluxes of the block field because these vary slowly along the translations by whole blocks.

Proposition 13.20 (The Coarse Covariance at a Flat Field)**.** *Let V_0 be flat, $0 < \lambda \le 1$, and, in the gauge of Lemma 13.14(i), $P := \Pi_{D\,\mathrm{ran}\,H} - DH(K+\lambda)^{-1}H^*D^*$. Then:*

(i) with C_7, δ_2 of Lemma 13.17 and C_V of Lemma 13.19,

$$c_\lambda(V_0) \ge d_{\mathfrak{g}}\Big(\frac{s^2t^3}{112\,T^5} - C_7\Big(\frac{s}{t}\Big)^2 e^{-\delta_2 t_{\mathrm{b}}/2} - 10\pi^2 C_V \lambda t_{\mathrm{b}}^2 (s_{\mathrm{b}}+2)(t_{\mathrm{b}}+2)\Big)\ ; \tag{13.55}$$

(ii) $DH\lambda(K+\lambda)^{-1}u = PDHu$ for every form u on $\mathcal{B}$, and in every component r and for every $F \in D(\operatorname{ran} H)$,

$$|\langle \chi_3 - \chi_2, P_r F\rangle| \le 2\pi t_{\mathrm{b}} (\lambda C_V (s_{\mathrm{b}}+2)(t_{\mathrm{b}}+2))^{1/2} \|F\|\ . \tag{13.56}$$

Proof. The exact form. By Lemma 13.14(iv) and the gauge invariance of c_λ, $c_\lambda(V_0) = \sum_\alpha \langle \chi_0 \otimes t_\alpha, \Pi_\lambda((\chi_3-\chi_2)\otimes t_\alpha)\rangle$ with $\Pi_\lambda \coloneqq DH(K+\lambda)^{-1}H^*D^*$. The sum over the orthonormal basis is a partial trace over $\mathfrak{g}$, which is the same in the basis (E_r), and D, Q, H, K commute with the decomposition into components, so $c_\lambda(V_0) = \sum_r \operatorname{Re}\langle \chi_0, \Pi_{\lambda,r}(\chi_3-\chi_2)\rangle$, the total being real. By Lemma 13.15(iii), $\Pi_{D\operatorname{ran} H} = DHK^+H^*D^*$, and H^*D^* maps into $(\ker DH)^\perp = (\ker K)^\perp$, so that $P = DHg(K)H^*D^*$ with $g(x) \coloneqq \lambda x^{-1}(x+\lambda)^{-1}$ on $(\ker K)^\perp$ and $g \coloneqq 0$ on $\ker K$; in particular $0 \le P \le \Pi_{D\operatorname{ran} H}$. By Lemma 13.15(iii) and (13.48),

$$c_\lambda(V_0) = \sum_r (c^{\theta_r}(3t_{\mathrm{b}}) - c^{\theta_r}(2t_{\mathrm{b}}) - \mathfrak{f}_r - \mathfrak{l}_r)\ , \qquad \mathfrak{l}_r \coloneqq \operatorname{Re}\langle \chi_0, P_r(\chi_3-\chi_2)\rangle\ . \tag{13.57}$$

The translation. Let $\mathcal{T}$ be the translation by one block in the direction μ, acting on the forms of the component r with the phase of Lemma 13.14(ii) across the hyperplane. It is unitary and commutes with D and Q, hence with DH, whose value at u is the point of least norm of $\{DA \,|\, QA = u\}$, and with K; and $\chi_k = \mathcal{T}^{kt_{\mathrm{b}}}\chi_0$, the rectangles not meeting the hyperplanes. With $J \coloneqq H^*D^*\chi_0$ and $v \coloneqq g(K)^{1/2}J$ we get $\langle \chi_0, P_r\mathcal{T}^{kt_{\mathrm{b}}}\chi_0\rangle = \langle v, \mathcal{T}^{kt_{\mathrm{b}}}v\rangle = \int e^{ikt_{\mathrm{b}}p}\,\mathrm{d}\nu(p)$, with $\nu \ge 0$ the spectral measure of $\mathcal{T}$ in v on $(-\pi,\pi]$, so that

$$\mathfrak{l}_r = \int (\cos 3t_{\mathrm{b}}p - \cos 2t_{\mathrm{b}}p)\,\mathrm{d}\nu(p)\ . \tag{13.58}$$

Since $\cos 3x - \cos 2x = -2\sin\frac{5x}{2}\sin\frac{x}{2}$, $|\cos 3x - \cos 2x| \le \frac{5}{2}x^2$; and $p^2 \le \frac{\pi^2}{4}|1-e^{ip}|^2$ on $(-\pi,\pi]$. Hence $|\mathfrak{l}_r| \le \frac{5\pi^2}{8}t_{\mathrm{b}}^2\|(1-\mathcal{T})v\|^2 = \frac{5\pi^2}{8}t_{\mathrm{b}}^2\|\nabla_\mu v\|^2$.

The coercivity. Since $g(K)$ vanishes on $\ker K$, $v \in \operatorname{ran} K \subset \ker \bar{D}^*$ by Lemma 13.18(i), and Lemma 13.18(ii) gives $\|\nabla_\mu v\|^2 \le 16\langle v, Kv\rangle = 16\langle J, Kg(K)J\rangle \le 16\lambda\langle J, K^+J\rangle = 16\lambda\|\Pi_{D\operatorname{ran} H}\chi_0\|^2$, because $Kg(K) = \lambda(K+\lambda)^{-1} \le \lambda K^+$ on $(\ker K)^\perp$, and $\langle J, K^+J\rangle = \langle \chi_0, \Pi_{D\operatorname{ran} H}\chi_0\rangle$. With Lemma 13.19, $|\mathfrak{l}_r| \le 10\pi^2 C_V \lambda t_{\mathrm{b}}^2(s_{\mathrm{b}}+2)(t_{\mathrm{b}}+2)$.

Conclusion of (i). Lemma 13.5 with $\mathsf{T} = N_q$, $\eta = L^{-K'}$, $t = t_{\mathrm{b}}$, and $s = s_{\mathrm{b}}$, whose conditions hold because $12t_{\mathrm{b}} < N_q$ and $3s_{\mathrm{b}} < N_q$, gives $c^{\theta_r}(3t_{\mathrm{b}}) - c^{\theta_r}(2t_{\mathrm{b}}) \ge s_{\mathrm{b}}^2 t_{\mathrm{b}}^3/(112N_q^5) = s^2t^3/(112T^5)$ for every twist, the number being invariant under the common rescaling of the lengths; Lemma 13.17 bounds $\mathfrak{f}_r$, and there are $d_{\mathfrak{g}}$ components.

(ii) On $(\ker K)^\perp$, $\lambda(K+\lambda)^{-1} = g(K)K$, and DH vanishes on $\ker K$, which $\lambda(K+\lambda)^{-1}$ maps into itself; so $DH\lambda(K+\lambda)^{-1}u = DHg(K)Ku = DHg(K)H^*D^*DHu = PDHu$. Since $0 \le P \le 1$, the Cauchy–Schwarz inequality for the form $\langle \cdot, P\cdot\rangle$ gives $|\langle \chi_3-\chi_2, P_rF\rangle| \le \langle \chi_3-\chi_2, P_r(\chi_3-\chi_2)\rangle^{1/2}\|F\|$, and as in the translation step $\langle \chi_3-\chi_2, P_r(\chi_3-\chi_2)\rangle = \int|e^{it_{\mathrm{b}}p}-1|^2\,\mathrm{d}\nu(p) \le t_{\mathrm{b}}^2\int p^2\,\mathrm{d}\nu(p) \le \frac{\pi^2}{4}t_{\mathrm{b}}^2\|\nabla_\mu v\|^2$, which the coercivity step bounds by $4\pi^2 t_{\mathrm{b}}^2\lambda C_V(s_{\mathrm{b}}+2)(t_{\mathrm{b}}+2)$. *q.e.d.*

13.8 The Second Moments of the Fluxes

We now compute $\Psi_3 - \Psi_2$ to the order $\bar g^2$. Near the fields of small action the measure $\langle\cdot\rangle$ has the density $\exp(-g_{K'}^{-2}\mathsf{A} + \mathcal{A}')$, whose second part is of the order one per block, and an integration by parts against the Haar measure turns $\langle X^\alpha \partial_w \mathsf{A}\rangle$ into $g_{K'}^2\langle \partial_w X^\alpha\rangle$, up to terms with the derivatives of w and of $\mathcal{A}'$. If w solves $(\Delta+\lambda)w = \nabla\Xi_\alpha$, then $\sum_\alpha \partial_w X^\alpha$ is the number c_λ of Section 13.7; but such a w satisfies $\partial_w\mathsf{A} = \Xi_\alpha$ only approximately, and since X^α is of the order $\bar g$, the defect $\partial_w\mathsf{A} - \Xi_\alpha$ has to be small compared with $\bar g$ on all the fields of small action. At a flat field both $\nabla\mathsf{A}$ and Ξ_α vanish, and at the field $e^{iu}W_0$ near a flat W_0 the defect is, to first order in u, the number $-\lambda\langle (K+\lambda)^{-1}\nabla\Xi_\alpha(W_0), u\rangle$, which the regularisation keeps small. The difficulty is that a field of action of the order $g_{K'}^2$ need not lie within a distance of the order $g_{K'}$ of the flat fields: these form a variety with singular points, the holonomies with large centralisers, near which the Hessian $K = \Delta(W_0)$ has small eigenvalues in the directions which leave the variety. We meet it with a Łojasiewicz inequality for commuting holonomies and a stratification of the variety by the dimension of the centraliser, which give at every field of small action a flat field and a lower bound on the Hessian there that are matched to each other: the field lies close to the flat field on the scale which the bound sets (Proposition 13.27). The defect is then followed along a segment from a field which differs from the flat one only in the directions of the kernel of K (Proposition 13.28).

The cutoff and the regularisation. We keep the units and the notation of Section 13.7, and we write $g \coloneqq g_{K'}$. The data parameter ε_1 of stage 5, at which [7, Theorem 1, p. 279] is applied, gives the constants $r_V \coloneqq C_1\varepsilon_1/(2N^{1/2})$ and $\varepsilon_V \coloneqq c_h B_5 \varepsilon_1$, with C_1 and B_5 of [7, (172)–(173), p. 305], and we write $\mathcal{R}$ for the set of the $V \in G^{\mathcal{B}}$ with $|V(\partial p') - 1| < \varepsilon_1$ for every plaquette p' of $\mathbb{T}_{j_q}$. Let $\varepsilon^\zeta \coloneqq 3(8NT_{\mathrm{ng}})^{1/2}g$ and $\mathcal{O} \coloneqq \{V \in G^{\mathcal{B}} \,|\, |V(\partial p') - 1| < \varepsilon^\zeta \text{ for every } p'\}$, an open subset of $\mathcal{R}$ for $\bar g$ small. On $\mathcal{O}$, [7, Theorem 1] with its ε_1 equal to ε^ζ gives the minimal configurations, with $|U_{K'}(V)(\partial p) - 1| < B_3\varepsilon^\zeta\eta^2$ on the fine plaquettes. Since $B_3\varepsilon^\zeta < \varepsilon_{K'}$ for $\bar g$ small, by $T_{\mathrm{ng}} \le C(\log\bar g^{-2})^{4r+1}$, $g \le \rho^{1/2}\hat g_{K'}$, and $p_0 > 2r + \frac12$, every test function of the term ℓ_0 passes on $\mathcal{O}$, as in the lower bound of the proof of Proposition 8.4; hence $\chi_{K'}(\mathbb{T}) = 1$ and $\varrho_{\ell_0} = \exp A_{K'}(U_{K'}(V))$ on $\mathcal{O}$, and $\mathcal{O} \subset \operatorname{supp}\varrho_{\ell_0}$. We fix $\vartheta \in C^\infty(\mathbb{R})$ with $0 \le \vartheta \le 1$, $\vartheta = 1$ on $(-\infty, 1]$, and $\vartheta = 0$ on $[2,\infty)$, and we put $\zeta(V) \coloneqq \vartheta(\mathsf{A}(V)/(T_{\mathrm{ng}}g^2))$ for $V \in \operatorname{supp}\varrho_{\ell_0}$ and $\zeta(V) \coloneqq 0$ otherwise. By Supplement, Lemma 17.6, with $T = 2T_{\mathrm{ng}}$, every $V \in \operatorname{supp}\varrho_{\ell_0}$ with $\mathsf{A}(V) \le 2T_{\mathrm{ng}}g^2$ has $|V(\partial p') - 1| \le 2(4NT_{\mathrm{ng}})^{1/2}g < \varepsilon^\zeta$ for every p'. So ζ vanishes outside the compact set $\mathcal{K}$ of the fields with $|V(\partial p') - 1| \le 2(4NT_{\mathrm{ng}})^{1/2}g$ for every p', which lies in $\mathcal{O}$; it is smooth on $G^{\mathcal{B}}$, because A is real analytic on $\mathcal{O}$ [7, Proposition 9, p. 309]; $\zeta = 1$ on $\mathcal{V}$; and $\operatorname{supp}\zeta \subset \mathcal{V}_3$. On $\operatorname{supp}\zeta$ the plaquettes satisfy $\|V(\partial p') - 1\| \le \varepsilon_\zeta \coloneqq 2c_h(4NT_{\mathrm{ng}})^{1/2}g$, and $\|X_{0,b}\| \le 2h^G$ by Section 13.7.

We fix an exponent p_λ with $14r + 2 < p_\lambda < q_0 - 11r - \frac12$, an interval which is not empty because $q_0 > 25r + \frac52$ (Appendix A), and we put

$$\begin{aligned} \lambda &\coloneqq N_q^{-2}(\log\bar g^{-2})^{-p_\lambda}\,, \qquad F_\alpha \coloneqq \zeta X^\alpha\,, \\ w_\alpha &\coloneqq (\Delta(V) + \lambda)^{-1}\nabla\Xi_\alpha\,, \qquad \mathfrak{e}_\alpha \coloneqq \langle w_\alpha, \nabla\mathsf{A}\rangle - \Xi_\alpha\,, \end{aligned} \tag{13.59}$$

the last two at the fields at which $\Delta(V) + \lambda$ is invertible. The sizes are those collected in the proof of Lemma 13.7: for $\bar g$ small and $K \ge K_{\mathrm{w}}$, N_q and $P \coloneqq P_{K'}(R_4) = 2(4t_{\mathrm{b}} + s_{\mathrm{b}}) \le 2N_q$ are at most $C(\log\bar g^{-2})^r$, $T_{\mathrm{ng}} \le C(\log\bar g^{-2})^{4r+1}$, and $h^G \le C\bar g(\log\bar g^{-2})^{4r+1/2}$; moreover $\bar g^2/2 \le \hat g_{K'}^2 \le g^2 \le (1+\beta_0)^2\bar g^2$, because $\hat g_{K'}^{-2} = \bar g^{-2} + b' j_q + C_W$, $t_{\mathrm{b}} \ge a_{K'} \ge (\log\bar g^{-2})^r$, and the reduced radii $r_{i,K'} \coloneqq \alpha_{i,K'}/(8B_3)$ of Supplement, Section 15, satisfy $r_{i,K'} \ge c\,\bar g(\log\bar g^{-2})^{q_0}$ by Definition 3.3. Here and below C, c, and m are positive constants which depend on q, t/u, s/u, T/u, π, p_λ, and the constants of the construction, and we write $X \preceq \bar g^a$ if $|X| \le C\bar g^a(\log\bar g^{-2})^m$; if $X \preceq \bar g^{a+b}$ with $b > 0$, then $|X| \le \bar g^{a+b/2}$ for $\bar g$ small.

The minimal configuration depends analytically on its data, and we need this in a quantitative form, at complex data and with the gauge transformation which returns it to Bałaban's representative.

Lemma 13.21 (The Moved Minimiser)**.** *Let $V_1 \in \mathcal{R}$, and let U_1 be a minimal configuration for V_1.*

(i) There are maps $B \mapsto \mathcal{H}(B)$, into the $\mathfrak{g}^{\mathbb{C}}$-valued 1-forms on $\mathbb{T}_K$, and $B \mapsto w_B$, into the $G^{\mathbb{C}}$-valued functions on the sites of $\mathbb{T}_K$, analytic on the set of the $\mathfrak{g}^{\mathbb{C}}$-valued forms B on $\mathcal{B}$ with $\|B\|_\infty < r_V$, with $\mathcal{H}(0) = 0$, $w_0 = 1$,

$$\|\mathcal{H}(B)_b\| < \varepsilon_V \ , \qquad \|(\nabla^\eta \mathcal{H}(B))(b)\| < \varepsilon_V \ , \qquad \|\log w_B(x)\| \le 72d\,\varepsilon_V \ , \tag{13.60}$$

with ∇^η the covariant lattice derivative at U_1 of [7, (19), p. 281] in units of the blocks, with the same bound ε_V for the second derivative $D^{\eta*}_{U_1} D^{\eta}_{U_1} \mathcal{H}(B)$ of [7, (19)], which [7, Proposition 9, p. 309] bounds together with the first two, and such that, for real B, $(e^{i\eta\mathcal{H}(B)}U_1)^{w_B}$ is a minimal configuration for the data $e^{iB}V_1$. Consequently, for real B, $\mathsf{A}(e^{iB}V_1) = A(e^{i\eta\mathcal{H}(B)}U_1)$, and the holonomy at a site y of $\mathbb{T}_{j_q}$ of a closed path of $\mathbb{T}_K$ based at y is $w_B(y)$ times its holonomy at $e^{i\eta\mathcal{H}(B)}U_1$ times $w_B(y)^{-1}$.

(ii) For $\|B\|_\infty < r_V$, $\|\mathcal{H}(B)_b\|, \|(\nabla^\eta\mathcal{H}(B))(b)\|, \|(D^{\eta*}_{U_1} D^{\eta}_{U_1}\mathcal{H}(B))(b)\| \le \varepsilon_V \|B\|_\infty / r_V$ and $\|\log w_B(x)\| \le 72d\,\varepsilon_V \|B\|_\infty / r_V$. For $\|B\|_\infty \le r_V/2$, every $\mathfrak{g}^{\mathbb{C}}$-valued form v on $\mathcal{B}$, and $m \ge 1$, the derivatives $\partial_v^m f(B) := (\mathrm{d}/\mathrm{d}z)^m f(B + zv)|_{z=0}$ satisfy $\|\partial_v^m \mathcal{H}(B)_b\|, \|\partial_v^m(\nabla^\eta\mathcal{H})(B)(b)\| \le m!\,\varepsilon_V (2\|v\|_\infty / r_V)^m$ and $\|\partial_v^m \log w_B(x)\| \le 72d\,m!\,\varepsilon_V(2\|v\|_\infty/r_V)^m$.

Proof. (i) The data V_1 satisfy [7, (7), p. 278] with the parameter ε_1, and $\|B\|_\infty < r_V$ gives $|B| \le N^{1/2}\|B\| < C_1\varepsilon_1/2$, hence $|e^{iB} - 1| \le e^{|B|} - 1 < C_1\varepsilon_1$, by [4, (20), p. 21] and the submultiplicativity of $|\cdot|$; so $V' := e^{iB}$ satisfies [7, (172), p. 305]. By [7, Proposition 9, p. 309] and [7, pp. 305–307], the minimal configuration for $V'V_1$, transformed to the Landau gauge relative to U_1, is $e^{i\eta\mathcal{H}(B)}U_1$ with $\mathcal{H}$ analytic in the $\mathfrak{g}^{\mathbb{C}}$-valued B satisfying [7, (172)] [7, p. 307, ll. 1–2], and $\mathcal{H}$ satisfies [7, (19)–(21), p. 281] with $\varepsilon_2 = B_5\varepsilon_1$. On the term ℓ_0 every set Ω_j of the determining sequence is the torus, so the conditions [7, (19)] for $j = K'$, where $L^{K'}\eta = 1$, read $|\mathcal{H}| < B_5\varepsilon_1$ and $|\nabla^\eta\mathcal{H}| < B_5\varepsilon_1$, which give the first two bounds after multiplication by c_h. The configuration in the Landau gauge is carried back to the axial gauge by a gauge transformation which is analytic in $\mathcal{H}$, hence in B [7, p. 307, ll. 2–5], and which is normalised by the condition that its averaged transformation be 1 on the sets Λ_j [7, p. 281, ll. 1–3]; by the uniqueness in Supplement, Lemma 17.14, applied at the G-valued base $\mathbf{U} = U_1$, with $A' = 0$ and $X = \mathcal{H}(B)$, so that $\varepsilon_A = 0$ and $\varepsilon_X = B_5\varepsilon_1$, this is the transformation w_B of that lemma, and $|\log w_B| \le 72dB_5\varepsilon_1$. For real B the minimal configurations for $e^{iB}V_1$ form one orbit, the action is gauge invariant, and a gauge transformation w changes the holonomy at y of a closed path based at y by the conjugation with $w(y)$.

(ii) The function $z \mapsto \mathcal{H}(zB/\|B\|_\infty)_b$ is analytic on $|z| < r_V$, bounded by ε_V, and vanishes at $z = 0$, so Schwarz's lemma, applied to its pairings with the functionals of norm one, gives $\|\mathcal{H}(B)_b\| \le \varepsilon_V\|B\|_\infty/r_V$; the same argument applies to $(\nabla^\eta\mathcal{H}(B))(b)$ and to $\log w_B(x)$. For $\|B\|_\infty \le r_V/2$ and $|z| < r_V/(2\|v\|_\infty)$ we have $\|B + zv\|_\infty < r_V$, and Cauchy's estimate on this disc gives the bounds on the derivatives. *q.e.d.*

The fluxes are logarithms of holonomies of the moved minimiser, and Cauchy's estimate in the data bounds their derivatives; the same argument applied to the action bounds

the Hessian and its variation. We write $\partial_w\partial_u F(V) := \partial_\sigma\partial_s F(e^{isu}e^{i\sigma w}V)|_{s=\sigma=0}$, so that $\langle u, \partial_w\nabla F(V)\rangle = \partial_w\partial_u F(V)$ with $\partial_w\nabla F(V) := \frac{\mathrm{d}}{\mathrm{d}\sigma}\nabla F(e^{i\sigma w}V)|_{\sigma=0}$, and similarly $\partial_w\Delta(V) := \frac{\mathrm{d}}{\mathrm{d}\sigma}\Delta(e^{i\sigma w}V)|_{\sigma=0}$. Let $\mathcal{R}'$ be the set of the $V_1 \in \mathcal{R}$ with $\|U_{K'}(V_1)(I_b) - 1\| \le r_G/2$ for $b \in \{0,2,3\}$, a condition on the holonomy at the site o_b of $\mathbb{T}_{j_q}$ which does not depend on the choice of the minimal configuration.

Lemma 13.22 (Derivatives of the Fluxes and of the Action)**.** *There are constants C_Ξ and C_Δ of the construction such that the following hold for all real forms u, w on $\mathcal{B}$.*

(i) At every $V_1 \in \mathcal{R}'$ the functions $X_{0,k}$, $k \in \{0,2,3\}$, are real analytic, and $\|\partial_u X_{0,k}(V_1)\| \le C_\Xi P\|u\|_\infty$, $\|\nabla X^\alpha(V_1)\|, \|\nabla\Xi_\alpha(V_1)\| \le C_\Xi P$, and $\|\partial_w\nabla X^\alpha(V_1)\|, \|\partial_w\nabla\Xi_\alpha(V_1)\| \le C_\Xi P^2\|w\|_\infty$.

(ii) At every $V_1 \in \mathcal{R}$, $\|\Delta(V_1)\| \le C_\Delta N_q^4$ and $\|\partial_w\Delta(V_1)\| \le C_\Delta N_q^4\|w\|_\infty$.

(iii) If $V_1 \in \mathcal{R}$ and $e^{-itw}V_1 \in \mathcal{R}$ for $0 \le t \le (2\mathsf{A}(V_1)/(C_\Delta N_q^4))^{1/2}$ and every w with $\|w\| = 1$, then $\|\nabla\mathsf{A}(V_1)\| \le (2C_\Delta N_q^4\mathsf{A}(V_1))^{1/2}$.

Proof. (i) Let U be a minimal configuration for V_1, and for complex B with $\|B\|_\infty < r_V$ let $H_k(B)$ be the holonomy of $\mathsf{L}_{0,k}$ at $e^{i\eta\mathcal{H}(B)}U$, with $\mathcal{H}$ of Lemma 13.21 at the base V_1. The path $\mathsf{L}_{0,k}$ has $2kt_\mathsf{b} + 2t_\mathsf{b} + 2s_\mathsf{b} \le P$ blocks, hence at most P/η fine bonds, and each of its factors $e^{\pm i\eta\mathcal{H}_b}$ differs from 1 by at most $e^{\eta\|\mathcal{H}_b\|} - 1 \le 2\eta\varepsilon_V\|B\|_\infty/r_V$, by Lemma 13.21(ii). Expanding the product of the factors $(1+a_l)U(b_l)^{\pm1}$ as in (8.11), with the unitary $U(b_l)$, gives $\|H_k(B) - H_k(0)\| \le e^{2P\varepsilon_V\|B\|_\infty/r_V} - 1$. Let $\rho_P := \min(r_V/2, r_V/(50P\varepsilon_V))$, so that $1/\rho_P \le c_P P$ with $c_P := (2 + 50\varepsilon_V)/r_V$. Since $H_k(0)$ is unitarily conjugate to $U(I_k)$ and $r_G \le 1/10$, for $\|B\|_\infty < \rho_P$ we get $\|H_k(B) - 1\| \le \frac{1}{20} + e^{1/25} - 1 \le \frac{1}{10}$, so that $-i\log H_k(B)$ is analytic, with norm at most $\frac{1}{9}$; and $\|\mathsf{Ad}(w_B(o))\| \le e^{144d\varepsilon_V} \le 2$ for ε_1 small. By Lemma 13.21(i), $X_{0,k}(e^{iB}V_1) = \mathsf{Ad}(w_B(o))(-i\log H_k(B))$ for real B, and the right side is an analytic extension to $\|B\|_\infty < \rho_P$, of norm at most $\frac{1}{4}$. Hence X^α and Ξ_α are real analytic near V_1, with analytic extensions bounded by $\frac{1}{4}$ and $\frac{1}{2}$, since $|t_\alpha \cdot Y| \le |Y|_2 \le \|Y\|$ for complex Y. Cauchy's estimate in z on $|z| < \rho_P/\|u\|_\infty$, for the field $e^{izu}V_1$, gives $\|\partial_u X_{0,k}(V_1)\| \le \|u\|_\infty/(4\rho_P)$, and with $\|u\|_\infty \le N^{1/2}\|u\|$ the bounds on ∇X^α and $\nabla\Xi_\alpha$. For the second derivatives, $e^{isu}e^{i\sigma w}V_1 = e^{iB(s,\sigma)}V_1$ with $B(s,\sigma)_b := -i\log(e^{isu_b}e^{i\sigma w_b})$, which is analytic and has $\|B(s,\sigma)\|_\infty \le 2(|s|\|u\|_\infty + |\sigma|\|w\|_\infty)$ for $|s|\|u\|_\infty + |\sigma|\|w\|_\infty \le \frac{1}{4}$, by the series of the exponential and the logarithm; so Cauchy's estimate on the polydisc $|s|\|u\|_\infty, |\sigma|\|w\|_\infty \le \rho_P/5$ gives $|\partial_w\partial_u\Xi_\alpha(V_1)| \le \frac{25}{2}\|u\|_\infty\|w\|_\infty/\rho_P^2$, and the same with $\frac{25}{4}$ for X^α. The constant $C_\Xi := 13N^{1/2}\max(c_P, c_P^2)$ covers all the bounds.

(ii) For real s and σ let $F(s,\sigma) := \mathsf{A}(e^{isu}e^{i\sigma w}V_1)$, so that $\langle u, \Delta(V_1)u\rangle = \partial_s^2 F(0,0)$ and $\langle u, \partial_w\Delta(V_1)u\rangle = \partial_\sigma\partial_s^2 F(0,0)$. By Lemma 13.21(i), $F(s,\sigma) = A^{\mathbb{C}}(e^{i\eta\mathcal{H}(B(s,\sigma))}U)$ with $A^{\mathbb{C}}(U') := \sum_p[1 - \frac{1}{2}(\operatorname{tr}U'(\partial p) + \operatorname{tr}U'(\partial p)^{-1})]$, which equals the action at unitary configurations and is analytic. Let p be a fine plaquette. Moving the transports of U to the right, $(e^{i\eta\mathcal{H}}U)(\partial p)$ is the product of four exponentials $e^{i\eta A'_l}$, with A'_l the transported values $\pm\mathcal{H}_{b_l}$, times $U(\partial p)$; by the Baker–Campbell–Hausdorff formula the four exponentials combine to e^{iZ} with Z equal to $\eta\sum_l A'_l$ up to $O(\eta^2\varepsilon_V^2)$, and $\eta\sum_l A'_l$ is η^2 times a combination of the covariant derivatives $\nabla^\eta\mathcal{H}$, up to a term bounded by $\|U(\partial p) - 1\|\varepsilon_V\eta$. With (13.60) and $|U(\partial p) - 1| < B_3\varepsilon_1\eta^2$ [7, Theorem 1], every plaquette of $e^{i\eta\mathcal{H}(B)}U$ is within $C_{\mathrm{pl}}\eta^2$ of 1, $C_{\mathrm{pl}} := 4\varepsilon_V + c_h B_3\varepsilon_1$, for ε_1 small. Its logarithm Y_p has $\|Y_p\| \le 2C_{\mathrm{pl}}\eta^2 \le 1$, and the term of p in $A^{\mathbb{C}}$ is $\sum_{n\ge1}(-1)^{n+1}\operatorname{tr}(Y_p^{2n})/(2n)!$, of modulus at most $\cosh\|Y_p\| - 1 \le \|Y_p\|^2$. Summing over the $6N_q^4\eta^{-4}$ fine plaquettes, $|F| \le C_A N_q^4$ with $C_A := 24C_{\mathrm{pl}}^2$ on the polydisc $|s|\|u\|_\infty, |\sigma|\|w\|_\infty \le r_V/5$, and Cauchy's estimate there gives $|\langle u, \Delta(V_1)u\rangle| \le$

$50C_AN_q^4\|u\|_\infty^2/r_V^2$ and $|\langle u, \partial_w \Delta(V_1)u\rangle| \le 250C_AN_q^4\|u\|_\infty^2\|w\|_\infty/r_V^3$. The operators are symmetric and $\|u\|_\infty \le N^{1/2}\|u\|$, which gives (ii) with $C_\Delta := 250NC_A \max(r_V^{-2}, r_V^{-3})$.

(iii) If $\mathsf{A}(V_1) = 0$, then V_1 is a minimum of $\mathsf{A} \ge 0$ and $\nabla\mathsf{A}(V_1) = 0$. Otherwise let $\|w\| = 1$ and $t_0 := (2\mathsf{A}(V_1)/(C_\Delta N_q^4))^{1/2}$. The second derivative of $t \mapsto \mathsf{A}(e^{-itw}V_1)$ is $\langle w, \Delta(e^{-itw}V_1)w\rangle \le C_\Delta N_q^4$ by (ii), so that $0 \le \mathsf{A}(e^{-it_0w}V_1) \le \mathsf{A}(V_1) - t_0\langle w, \nabla\mathsf{A}(V_1)\rangle + \frac{1}{2}t_0^2C_\Delta N_q^4$, that is, $\langle w, \nabla\mathsf{A}(V_1)\rangle \le 2\mathsf{A}(V_1)/t_0 = (2C_\Delta N_q^4\mathsf{A}(V_1))^{1/2}$. *q.e.d.*

The integration by parts itself is exact. On $\mathcal{O}$ we write $\mathcal{A}'(V) := A_{K'}(U_{K'}(V)) + g^{-2}\mathsf{A}(V)$ for the logarithm of the density of ℓ_0 without the Wilson action, and for $b \in \mathcal{B}$ and $\alpha \le d_{\mathfrak{g}}$ we let $e_{b,\alpha}$ be the form equal to t_α at b and to 0 elsewhere, an orthonormal basis of $\mathfrak{g}^{\mathcal{B}}$.

Lemma 13.23 (Integration by Parts)**.** *Let F be a C^1 function on $G^{\mathcal{B}}$ whose support lies in $\mathcal{O}$, and let v be a C^1 map from a neighbourhood of* $\operatorname{supp} F$ *in $\mathcal{O}$ to $\mathfrak{g}^{\mathcal{B}}$, with the divergence* $\operatorname{div} v := \sum_{b,\alpha} \partial_{e_{b,\alpha}}(t_\alpha \cdot v_b)$*. Then*

$$\langle F\langle v, \nabla\mathsf{A}\rangle\rangle = g^2\langle\langle v, \nabla F\rangle\rangle + g^2\langle F \operatorname{div} v\rangle + g^2\langle F\langle v, \nabla\mathcal{A}'\rangle\rangle \,. \tag{13.61}$$

Proof. For a C^1 function h on $G^{\mathcal{B}}$, $\int \partial_{e_{b,\alpha}} h(V)\,\mathrm{d}V = \frac{\mathrm{d}}{\mathrm{d}s}\int h(e^{ise_{b,\alpha}}V)\,\mathrm{d}V|_{s=0} = 0$, the Haar measure of the factor of b being invariant under the left multiplication by e^{ist_α}, and the derivative passing under the integral over the compact group. We apply this to $h := (t_\alpha \cdot v_b)F\varrho_{\ell_0}$, which is C^1 on $G^{\mathcal{B}}$ when extended by 0, because $\varrho_{\ell_0} = \exp(-g^{-2}\mathsf{A} + \mathcal{A}')$ is smooth on $\mathcal{O}$. Since ∂_u is linear in u, $\sum_{b,\alpha}(t_\alpha \cdot v_b)\partial_{e_{b,\alpha}} f = \langle v, \nabla f\rangle$, and summing over b and α gives $0 = \int \varrho_{\ell_0}[F \operatorname{div} v + \langle v, \nabla F\rangle + F\langle v, -g^{-2}\nabla\mathsf{A} + \nabla\mathcal{A}'\rangle]\,\mathrm{d}V$. Dividing by $g^{-2}Z_{\ell_0}$ gives (13.61). *q.e.d.*

The last term of (13.61) contains the derivative of the effective action beyond the Wilson action. Its terms are of the order one per block, and their derivatives are bounded by Cauchy's estimate on the radius of analyticity of Bałaban's terms, which is larger than the size of the fields by a power of $\log \bar{g}^{-2}$.

Lemma 13.24 (The Rest of the Density)**.** *Let* $r_E := \min(r_V/2, r_V \min(r_{0,K'}, r_{1,K'})/(8C_{Q^*}N^{1/2}\varepsilon_V))$*, with the constant $C_{Q^*} = 2(1 + (2d+1)L^{d-1}) \ge 1$ of Supplement, Section 15, and with ε_V a common bound of $\mathcal{H}(B)$, of its first differences, and of $D^*D\mathcal{H}(B)$ on $\|B\|_\infty < r_V$, in the units of [7, (19), p. 281]. There is a constant M_E of the construction such that for $\bar{g}$ small, every $V \in \mathcal{K}$, and every real form w on $\mathcal{B}$,*

$$|\langle w, \nabla\mathcal{A}'(V)\rangle| \le \frac{M_EN_q^4\|w\|_\infty}{r_E} \,. \tag{13.62}$$

Proof. Let $U := U_{K'}(V)$, and for complex t with $|t| < r_E/\|w\|_\infty$ let $U_t := e^{i\eta\mathcal{H}(tw)}U$, with $\mathcal{H}$ of Lemma 13.21 at the base V. On ℓ_0 the function $\mathcal{A}'$ is, by [9, (2.23), p. 258] and Section 8.1, with the coupling function $g_{K'}^2(x)$ of [9, (2.24)] equal to g^2 because $\phi_j = 1$ on the torus, the sum of the regular part $\mathbf{E}_{K'}$, the $\mathbf{R}$-part $\mathbf{R}_{K'}$, and the boundary part, each a gauge invariant function of the configuration, and of constants which do not depend on V. For real t the configuration U_t is a gauge transform of a minimal configuration for $e^{itw}V$, by Lemma 13.21(i), so that $\mathcal{A}'(e^{itw}V)$ is the sum of the three parts at U_t and of the constants, and the parts at U_t are analytic in t.

The class. We show that U_t lies in the regular subclass $\mathcal{S}_z$ of Supplement, Section 15, for every point z, at the reduced radii $r_{0,K'}$, $r_{1,K'}$ of the layer K', the only layer on ℓ_0.

The configuration U is G-valued with $|U(\partial p) - 1| < B_3\varepsilon^\zeta\eta^2 \le \frac{1}{2}r_{0,K'}\eta^2$, and its current is at most $B_3\varepsilon^\zeta$ in units of the blocks by [7, (2), (170), pp. 278, 305], for $\bar g$ small, because $\varepsilon^\zeta \preceq \bar g$ while $r_{0,K'} \ge c\bar g(\log \bar g^{-2})^{q_0}$. The field $A' \coloneqq \mathcal{H}(tw)$ is $\mathfrak{g}^{\mathbb{C}}$-valued with $|A'|, |\nabla^\eta A'| \le N^{1/2}\varepsilon_V|t|\|w\|_\infty/r_V \le \frac{1}{8}\min(r_{0,K'}, r_{1,K'})/C_{Q^*}$ by Lemma 13.21(ii), and the same bound holds for D^*DA', by the same Schwarz estimate applied to $D^*D\mathcal{H}$, which [7, (19)] bounds together with $\mathcal{H}$ and its first differences [7, Proposition 9, p. 309], so that, by the argument of the proof of Lemma 13.22(ii), the plaquettes of U_t are within $\frac{1}{2}r_{0,K'}\eta^2 + \frac{1}{2}r_{0,K'}\eta^2$ of 1, and the current of U_t, which is the current of U plus D^*DA' plus terms of the second order in A', is at most $B_3\varepsilon^\zeta + \frac{1}{8}r_{0,K'}/C_{Q^*} + \frac{1}{8}r_{0,K'}/C_{Q^*}$, which is below $r_{0,K'}/C_{Q^*}$ for $\bar g$ small because $\varepsilon^\zeta \preceq \bar g$. For real t the restriction of a minimal configuration to the near domain $\square_0$ of a cube is, in a gauge on $\square_0$, the local minimiser of its own block data [8, (3.18), p. 273]; both sides of this identity, transported to U_t by the gauge transformation of Lemma 13.21(i), are analytic in t, so it holds for complex t by the identity theorem, in a complex gauge on $\square_0$.

The parts. On ℓ_0 every Λ_j is the torus, so $\Lambda_j^0 = \mathbb{T}^{(j)}$, the function ϕ_j of [9, (2.25), p. 259] is 1, and the tents h_z of [9, p. 283] sum to 1; by [9, (2.25)–(2.26)],

$$\mathbf{E}_{K'}(U_t) = \sum_{j=1}^{K'} \sum_{z\in\mathbb{T}^{(j)}} \Big[\mathbf{E}^{(j)}(\mathbb{T}, U_t, z) - \mathbf{E}^{(j)}(\mathbb{T}, 1, z) - \beta_j(g_{j-1})A(h_z, U_t)\Big] . \tag{13.63}$$

Supplement, Proposition S.15(a), for the single family $\mathbf{E}^{(j)}$, that is with the second family 0, $d = 1$ by [8, (1.18), p. 263], and $N_j = 0$, bounds the bracket by $C_\alpha E_0 (L^{j-K'})^{4+\beta'}$, since every z lies in $\Omega_{K'} \setminus \Omega_{K'+1}$. The lattice $\mathbb{T}^{(j)}$ has $L^{4(K'-j)}N_q^4$ points, and $L^{\beta'} \ge 2$, so $|\mathbf{E}_{K'}(U_t)| \le C_\alpha E_0 N_q^4 \sum_{m\ge0} L^{-\beta' m} \le 2C_\alpha E_0 N_q^4$. In the same way Supplement, Proposition S.15(b), with the class norm 1 by [9, (2.31), p. 260], bounds the terms of [9, (2.30)] at a point of the scale j by $C_\alpha g_j^{\kappa_0}(L^{j-K'})^4$, so that $|\mathbf{R}_{K'}(U_t)| \le C_\alpha N_q^4 \sum_{j\le K'} g_j^{\kappa_0} \le C_\alpha N_q^4$, by $g_j^{-2} \ge g_{K'}^{-2} + b(K'-j) - C_W$ (3.4), $\kappa_0 \ge 13$, and $\bar g$ small. The terms of the boundary part are analytic on the spaces of [9, pp. 261–262], which contain U_t, with the bound [9, (2.42)], and [9, (2.47)–(2.48), p. 264] bound their sum by $2B_\partial \sum_n |\Gamma_n| = 2B_\partial N_q^4$. So $t \mapsto \mathcal{A}'(e^{itw}V)$ extends analytically to $|t| < r_E/\|w\|_\infty$, where it differs from a constant by at most $M_E N_q^4$, $M_E \coloneqq 2C_\alpha E_0 + C_\alpha + 2B_\partial$, and Cauchy's estimate at $t = 0$ gives (13.62). *q.e.d.*

We turn to the geometry of the fields of small action. Their plaquettes are close to 1, so in a suitable gauge they are close to configurations which are constant off four hyperplanes, with values $h_1, \dots, h_4$ that nearly commute; how close the h_β are to a commuting quadruple is a question about the real algebraic set of commuting quadruples, which Łojasiewicz's inequality answers with an unknown exponent. The same inequality, applied to the eigenvalues of the operator $M(k)$ below on the strata of that set, bounds from below the twists of the components of a flat field which do not lie in the centraliser, and with them the spectral gap of K.

Lemma 13.25 (Łojasiewicz Inequalities for Commuting Holonomies)**.** *Let $Z \subset G^4$ be the set of the commuting quadruples. For $k \in G^4$ let $\Phi(k) \coloneqq \max_{\beta<\gamma}\|k_\beta k_\gamma k_\beta^{-1}k_\gamma^{-1} - 1\|$, let $M(k) \coloneqq \sum_\beta (\mathrm{Ad}(k_\beta) - 1)^*(\mathrm{Ad}(k_\beta) - 1)$ on $\mathfrak{g}$ with the product $X \cdot Y$, with the eigenvalues $0 \le \mu_1(k) \le \dots \le \mu_{d_\mathfrak{g}}(k)$, and for a compact $S \subset G^4$ let $d_\infty(k, S) \coloneqq \min_{z\in S}\max_\beta\|k_\beta - z_\beta\|$. Put $Z_{\ge0} \coloneqq Z$ and $Z_{\ge j} \coloneqq \{k \in Z \,|\, \mu_j(k) = 0\}$ for $1 \le j \le d_\mathfrak{g}$. There are constants $C_{\mathrm{Lo}} > 0$, $\alpha_{\mathrm{Lo}} \in (0, 1]$, $\mathsf{c}_j > 0$, and $\varpi_j \ge 1$, depending on G only, such that*

(i) $d_\infty(k, Z) \le C_{\mathrm{Lo}}\Phi(k)^{\alpha_{\mathrm{Lo}}}$ for every $k \in G^4$;

(ii) $\mu_{j+1}(k) \ge \mathsf{c}_j d_\infty(k, Z_{\ge j+1})^{\varpi_j}$ for $0 \le j < d_\mathfrak{g}$ and every $k \in Z_{\ge j}$.

Proof. We use Łojasiewicz's inequality in the form of [13, Theorem 6.4, p. 34]: if K is a subset of a real analytic manifold and $f, h\colon K \to \mathbb{R}$ are subanalytic functions with compact graphs and $f^{-1}(0) \subset h^{-1}(0)$, then $|f| \geq c|h|^\rho$ on K for some $c, \rho > 0$. The manifold is $M_N(\mathbb{C})^4$, a real vector space. The group G is a closed subgroup of $U(N)$, hence a closed real analytic submanifold, so that G^4 and $Z = G^4 \cap \{k_\beta k_\gamma = k_\gamma k_\beta\}$ are compact and semianalytic. By [13, Remarks 3.11(2), p. 19], $x \mapsto \min_{t\in T} f(x,t)$ is subanalytic if T is compact and subanalytic and f is continuous and subanalytic, and a maximum is $-\min(-\cdot)$. Now $\Phi(k)$ is the maximum, over $\beta < \gamma$ and over the pairs of unit vectors $x, y \in \mathbb{C}^N$, of the polynomial $\mathrm{Re}\langle y, (k_\beta k_\gamma k_\beta^* k_\gamma^* - 1)x\rangle$; in the same way $(k, z) \mapsto \max_\beta \|k_\beta - z_\beta\|$ is subanalytic, and so is $d_\infty(\cdot, S)$ for compact subanalytic S. By the Courant–Fischer principle, $\mu_i(k)$ is the minimum over the orthonormal i-frames $(f_1, \dots, f_i)$ of $\mathfrak{g}$ of the maximum over the unit vectors $a \in \mathbb{R}^i$ of $\langle \sum_l a_l f_l, M(k) \sum_l a_l f_l\rangle$, a polynomial, so μ_i is continuous and subanalytic, and $Z_{\geq j} = Z \cap \mu_j^{-1}(0)$ is compact and subanalytic. All the functions are continuous on compact sets, so their graphs are compact.

(i) We take $K = G^4$, $f = \Phi$, and $h = d_\infty(\cdot, Z)$, which vanish exactly on Z. Since $\Phi \leq 2$, the bound $d_\infty(k, Z) \leq (\Phi(k)/c)^{1/\rho}$ gives (i) with $\alpha_{\mathrm{Lo}} := \min(1, 1/\rho)$ and $C_{\mathrm{Lo}} := c^{-1/\rho} 2^{\max(0, 1/\rho - 1)}$.

(ii) We take $K = Z_{\geq j}$, $f = \mu_{j+1}$, and $h = d_\infty(\cdot, Z_{\geq j+1})$. On $Z_{\geq j}$ the function μ_{j+1} vanishes exactly on $Z_{\geq j+1}$, a closed set which contains $(1,1,1,1)$, at which $M = 0$, and on which h vanishes exactly. The theorem gives $c_j, \rho_j > 0$ with $\mu_{j+1} \geq c_j h^{\rho_j}$ on $Z_{\geq j}$, and since $h \leq 2$, (ii) holds with $\varpi_j := \max(1, \rho_j)$ and $\mathsf{c}_j := c_j 2^{\min(0, \rho_j - 1)}$. *q.e.d.*

We fix the hyperplanes once and for all. Let $\sigma := o - 4(1,1,1,1)$, so that each hyperplane $\{x_\beta \equiv \sigma_\beta + \frac{1}{2}\}$ has distance at least 3 from the rectangle bounded by R_4, as Lemma 13.14 requires, because $N_q > 12 t_{\mathrm{b}}$ and $N_q > 3 s_{\mathrm{b}}$; let $x_0 := \sigma + (1,1,1,1)$, and let $\mathcal{T}_{\mathrm{tr}}$ be the set of the bonds $\langle x, x + e_\beta\rangle$ of $\mathbb{T}_{j_q}$ with $x_\gamma \equiv x_{0,\gamma}$ for $\gamma < \beta$ and $x_\beta \not\equiv \sigma_\beta$, a spanning tree, in which γ_x denotes the path from x_0 to the site x. For a bond $b = \langle y, y + e_\beta\rangle$ we put $n(b) := e_\beta$ if $y_\beta \equiv \sigma_\beta$ and $n(b) := 0$ otherwise, and c_β is the closed path of N_q bonds from x_0 in the direction e_β.

Lemma 13.26 (A Tree Gauge)**.** *Let $V \in G^{\mathcal{B}}$ with $\|V(\partial p') - 1\| \leq \varepsilon$ for every plaquette p' of $\mathbb{T}_{j_q}$. There is a gauge transformation ω of $\mathbb{T}_{j_q}$ with $V^\omega = 1$ on $\mathcal{T}_{\mathrm{tr}}$ such that, with $h_\beta := V^\omega(c_\beta)$, $h^{e_\beta} := h_\beta$, and $h^0 := 1$, every bond has $\|V^\omega(b) - h^{n(b)}\| \leq 50 N_q^2 \varepsilon$, and $\Phi(h) \leq 8 N_q^2 \varepsilon$.*

Proof. Since $\mathcal{T}_{\mathrm{tr}}$ is a spanning tree, there is a unique gauge transformation ω with $\omega(x_0) = 1$ and $V^\omega = 1$ on its bonds, obtained by fixing ω along the paths γ_x outwards from x_0. Then $V^\omega(b)$ is the holonomy at V^ω of the closed path $\ell_b := \gamma_{b_-} b \gamma_{b_+}^{-1}$, all its other bonds being bonds of the tree, and it equals the holonomy of ℓ_b at V, because a gauge transformation conjugates the holonomy of a closed path by its value at the base point; in the same way h_β is the value of V^ω on the last bond of c_β, the others being bonds of the tree. A closed path of length m in $\mathbb{Z}^4$ reduces to the empty path by at most $m^2/2$ moves, each a transposition of two consecutive orthogonal steps or the cancellation of two consecutive opposite steps. Indeed, if a direction e_β occurs, the signed steps in this direction sum to zero, so two of them which are consecutive among the steps of this direction have opposite signs; the steps between them are orthogonal to e_β, so at most $m - 2$ transpositions make the two adjacent, and one cancellation removes them, which gives at most $\frac{m}{2}(m-1)$ moves in all. A transposition of two orthogonal steps replaces the holonomy PXS by $PX'S$, where X and X' are the transports along the two sides of the plaquette which the steps span and $X'^{-1}X$ is a holonomy of that plaquette, so that it changes the holonomy by $\|X - X'\| = \|X'^{-1}X - 1\| \leq \varepsilon$, by the unitary

invariance of the norm; a cancellation leaves it unchanged. The tree paths do not cross the hyperplanes, so $\ell_b c_\beta^{-1}$ for $n(b) = e_\beta$, and ℓ_b for $n(b) = 0$, wind around no cycle of the torus and lift to closed paths of $\mathbb{Z}^4$ of length at most $9N_q$, whose moves project to moves on the torus; and $c_\beta c_\gamma c_\beta^{-1} c_\gamma^{-1}$ lifts to a closed path of length $4N_q$. This gives the two bounds, since $(9N_q)^2/2 \le 50N_q^2$ and $(4N_q)^2/2 = 8N_q^2$. *q.e.d.*

In the gauge of Lemma 13.14(i) for the hyperplanes above, a flat field W' with the commuting holonomies $\tilde h_\beta$ equals $\tilde h^{n(b)}$ on every bond b. We write $\mathfrak{g}^{\tilde h} := \{Y \in \mathfrak{g} \,|\, \mathsf{Ad}(\tilde h_\beta)Y = Y \text{ for every } \beta\}$, $\mathcal{Z}$ for the space of the real forms c on $\mathcal{B}$ with values in $\mathfrak{g}^{\tilde h}$ which are constant in each direction, $c_b = c_\beta$ for the bonds of the direction β, with $|c| := \max_\beta \|c_\beta\|$, $\Pi_{\mathcal{Z}}$ for the orthogonal projection onto $\mathcal{Z}$, and $\bar D$ for the covariant derivative at W' of the 0-forms of $\mathbb{T}_{j_q}$. Let $\varpi := \max_j \varpi_j$,

$$\varsigma_j := \frac{\alpha_{\mathrm{Lo}}}{2}(4\varpi)^{-j} \quad (0 \le j \le d_{\mathfrak{g}} + 1)\,, \qquad \varsigma_* := \varsigma_{d_{\mathfrak{g}}+1}\,, \qquad \mathsf{r}_j := \bar g^{\varsigma_j}\,, \tag{13.64}$$

so that $\frac12 \ge \varsigma_0 > \varsigma_1 > \cdots > \varsigma_* > 0$ and $\varpi_j \varsigma_{j+1} \le \varsigma_j/4$, and let $c_\varkappa := \frac14 \min(1, \min_i \mathsf{c}_i 2^{-\varpi_i}/(4\pi^2))$.

Proposition 13.27 (Fields Near the Flat Ones)**.** *For $\bar g$ small and every $V \in \operatorname{supp}\zeta$ there are an index $j \in \{0, \dots, d_{\mathfrak{g}}\}$, a gauge transformation ω' of $\mathbb{T}_{j_q}$, commuting $\tilde h_1, \dots, \tilde h_4 \in G$ with the flat field W' as above, a form $c \in \mathcal{Z}$, and a form $u_{\mathrm{H}} \in \mathfrak{g}^{\mathcal{B}}$ such that, with $B := e^{ic}W'$ and $V_s := e^{isu_{\mathrm{H}}}B$ for $s \in [0,1]$, $V^{\omega'} = V_1$ and the following hold, where $K := \Delta(W')$ and $\delta_* := 2(|c| + \|u_{\mathrm{H}}\|_\infty)$.*

(i) The number $\varkappa := \frac14 N_q^{-2} \min(1, \mathsf{c}_j(\mathsf{r}_{j+1}/2)^{\varpi_j}/(4\pi^2))$, the second entry being omitted if $j = d_{\mathfrak{g}}$, satisfies $\varkappa \ge c_\varkappa \mathsf{r}_j^{1/4} N_q^{-2}$; moreover $\ker K = \operatorname{ran}\bar D \oplus \mathcal{Z}$, $K \ge \varkappa$ on $(\ker K)^\perp$, and $\bar D^* \bar D \ge 64\varkappa$ on $(\ker \bar D)^\perp$.

(ii) $\bar D^* u_{\mathrm{H}} = 0$ and $\Pi_{\mathcal{Z}} u_{\mathrm{H}} = 0$, so that $u_{\mathrm{H}} \perp \ker K$.

(iii) $\mathsf{A}(B) + \frac38 \varkappa \|u_{\mathrm{H}}\|^2 \le \mathsf{A}(V)$, $\langle u_{\mathrm{H}}, K u_{\mathrm{H}}\rangle \le 3\mathsf{A}(V)$, and $\mathsf{A}(V_s) \le 3\mathsf{A}(V)$ for $s \in [0,1]$. Moreover $V_s = e^{iu'_s}W'$ with real forms u'_s, $\|u'_s\|_\infty \le \delta_*$, the fields $e^{i\sigma u'_s}W'$, $\sigma \in [0,1]$, lie in $\mathcal{R}'$, and for all these fields V', $\|\Delta(V') - K\| \le C_\Delta N_q^4 \delta_* \le \min(\varkappa/4, \lambda/2)$ and $\|(\Delta(V') + \lambda)^{-1}\| \le 2/\lambda$.

(iv) $\nabla\mathsf{A}(B) \in \mathcal{Z}$, and $(\Delta(B) + \lambda)^{-1}$ maps $\mathcal{Z}$ and $\mathcal{Z}^\perp$ into themselves.

(v) $|c| \preceq \bar g^{\varsigma_j}$, $\|u_{\mathrm{H}}\|_\infty \le N^{1/2}\|u_{\mathrm{H}}\| \preceq \bar g^{1-\varsigma_j/8}$, and $\delta_* \preceq \bar g^{\varsigma_j}$.

Proof. The flat field. By Lemma 13.26 with $\varepsilon = \varepsilon_\zeta$ there is ω with $V^\omega = 1$ on the tree and holonomies h with $\Phi(h) \le 8N_q^2\varepsilon_\zeta$. Let $\delta_i := d_\infty(h, Z_{\ge i})$ for $i \le d_{\mathfrak{g}}$ and $\delta_{d_{\mathfrak{g}}+1} := \infty$. By Lemma 13.25(i), $\delta_0 \le C_{\mathrm{Lo}}(8N_q^2\varepsilon_\zeta)^{\alpha_{\mathrm{Lo}}} \preceq \bar g^{\alpha_{\mathrm{Lo}}}$, so $\delta_0 \le \mathsf{r}_0$ for $\bar g$ small, and there is a least $j \le d_{\mathfrak{g}}$ with $\delta_{j+1} > \mathsf{r}_{j+1}$; then $\delta_j \le \mathsf{r}_j$. Let $\tilde h \in Z_{\ge j}$ with $d_\infty(h, \tilde h) = \delta_j$, and let W' be the flat field with the holonomies $\tilde h$. Then $\|V^\omega(b) - W'(b)\| \le 50N_q^2\varepsilon_\zeta + \mathsf{r}_j \le 2\mathsf{r}_j$ for every bond, so $V^\omega = e^{iu}W'$ bondwise with $u_b \in \mathfrak{g}$ and $\|u\|_\infty \le 4\mathsf{r}_j$, because $2\mathsf{r}_j \le r_G$ and $\|\log(1+x)\| \le 2\|x\|$ for $\|x\| \le \frac12$.

The components. The commuting unitary operators $\mathsf{Ad}(\tilde h_\beta)$ have a common orthonormal eigenbasis (E_r) of $\mathfrak{g}^{\mathbb{C}}$, $\mathsf{Ad}(\tilde h_\beta)E_r = e^{2\pi i\theta_{r,\beta}}E_r$ with $\theta_r \in [0,1)^4$, in which $M(\tilde h)E_r = \sum_\beta |e^{2\pi i\theta_{r,\beta}} - 1|^2 E_r$. So $\ker M(\tilde h) = \mathfrak{g}^{\tilde h}$ is spanned by the E_r with $\theta_r = 0$, and every r with $\theta_r \ne 0$ has $4\pi^2 \sum_\beta \|\theta_{r,\beta}\|^2 \ge \mu_{j'+1}(\tilde h)$, $\|\cdot\|$ the distance to $\mathbb{Z}$ and $j' := \dim \mathfrak{g}^{\tilde h}$, because

$|e^{2\pi ix}-1| \le 2\pi\|x\|$. If $j < d_{\mathfrak{g}}$, the triangle inequality gives $d_\infty(\tilde h, Z_{\ge j+1}) \ge \delta_{j+1}-\delta_j > \mathsf{r}_{j+1}-\mathsf{r}_j \ge \mathsf{r}_{j+1}/2$ for $\bar g$ small, so $\tilde h \notin Z_{\ge j+1}$ and $j' = j$, and Lemma 13.25(ii) gives $\mu_{j+1}(\tilde h) \ge \mathsf{c}_j(\mathsf{r}_{j+1}/2)^{\varpi_j}$. Hence every component with $\theta_r \ne 0$ has $\sum_\beta \|\theta_{r,\beta}\|^2 \ge 4N_q^2\varkappa$; if $j = d_{\mathfrak{g}}$ there is no such component.

(i). Since $\mathsf{r}_{j+1}^{\varpi_j} = \bar g^{\varpi_j\varsigma_{j+1}} \ge \mathsf{r}_j^{1/4}$, $\varkappa \ge c_\varkappa \mathsf{r}_j^{1/4} N_q^{-2}$. On the component r the translations $\mathcal{T}_\mu$ by one block, $(\mathcal{T}_\mu f)(x) \coloneqq f(x-e_\mu)$ with the phase of Lemma 13.14(ii) across the hyperplane, as in the proof of Proposition 13.20, commute and have the joint eigenvalues $(e^{-2\pi i(n_\mu+\theta_{r,\mu})/N_q})_\mu$, $n \in \mathbb{Z}^4$, with the eigenvectors $x \mapsto e^{2\pi i(n+\theta_r)\cdot x/N_q}$ on the forms of each degree and, for 1-forms, of each direction. Since $|e^{i\phi}-1| \ge (4/\pi)\operatorname{dist}(\phi/2, \pi\mathbb{Z})$, $|e^{2\pi i(n+\theta)/N_q}-1|^2 \ge 16\|\theta\|^2/N_q^2$, and this is at least $16/N_q^2$ if $\theta = 0$ and $n \notin N_q\mathbb{Z}$. Hence $\sum_\mu \|(\mathcal{T}_\mu - 1)u\|^2 \ge 64\varkappa\|u\|^2$ for the forms u, of degree 0 or 1, in a component with $\theta_r \ne 0$, and for those orthogonal to the constants in a component with $\theta_r = 0$. On 0-forms $\bar D^*\bar D = \sum_\mu (\mathcal{T}_\mu - 1)^*(\mathcal{T}_\mu - 1)$, whose kernel consists of the constants of the components with $\theta_r = 0$, which gives the last claim. For $c \in \mathcal{Z}$ let A_c be the fine 1-form equal to c_β on the fine bonds of the direction β. Since $\mathsf{Ad}(\tilde h_\beta)$ fixes the values of c, A_c has the constant components of the components with $\theta_r = 0$, so $DA_c = 0$ and $Q^{\mathrm{dr}}A_c = c$ by Lemma 13.14(ii), the straight average of a constant being that constant; by Lemma 13.14(iii), DHc is the point of least norm of $\{DA \,|\, Q^{\mathrm{dr}}A = c\}$, which is 0, so that $\mathcal{Z} \subset \ker DH = \ker K$. Next, $\operatorname{ran}\bar D \subset \ker K$, because $K\bar D = 0$ by the proof of Lemma 13.18(i). Finally $\mathcal{Z} \perp \operatorname{ran}\bar D$: for a 0-form ω and $c \in \mathcal{Z}$, $(\mathsf{Ad}(W'(b))\omega(b_+)) \cdot c_b = \omega(b_+)\cdot c_b$, so $\langle \bar D\omega, c\rangle = \sum_\beta \sum_b (\omega(b_+)-\omega(b_-))\cdot c_\beta$, the inner sum over the bonds of the direction β, which vanishes because every site is the endpoint b_+ of exactly one of them and the endpoint b_- of exactly one; thus $\bar D^* c = 0$. For $u \perp \operatorname{ran}\bar D \oplus \mathcal{Z}$, that is $\bar D^* u = 0$ and $\Pi_{\mathcal{Z}} u = 0$, Lemma 13.18(ii) in each direction gives $64\langle u, Ku\rangle = 64\|DHu\|^2 \ge \sum_\mu \|\nabla_\mu u\|^2 \ge 64\varkappa\|u\|^2$. So $\ker K = \operatorname{ran}\bar D \oplus \mathcal{Z}$ and $K \ge \varkappa$ on its orthogonal complement.

(ii), the slice. For a 0-form $\omega_1 \perp \ker\bar D$ and $c \in \mathcal{Z}$ we define u_{H} by $e^{iu_{\mathrm{H},b}} \coloneqq e^{i\omega_1(b_-)}e^{iu_b}e^{-i\mathsf{Ad}(W'(b))\omega_1(b_+)}e^{-ic_b}$, so that $e^{iu_{\mathrm{H},b}}e^{ic_b}W'(b) = e^{i\omega_1(b_-)}V^\omega(b)e^{-i\omega_1(b_+)}$, that is $(V^\omega)^{e^{i\omega_1}} = e^{iu_{\mathrm{H}}}e^{ic}W'$, and $\omega' \coloneqq e^{i\omega_1}\omega$. Then $u_{\mathrm{H}} = u - \bar D\omega_1 - c + \mathcal{N}(\omega_1, c)$, with $(\bar D\omega_1)_b = \mathsf{Ad}(W'(b))\omega_1(b_+) - \omega_1(b_-)$, and by the Baker–Campbell–Hausdorff series, which converges because the four exponents have norms at most $\frac{1}{10}$ on the set below for $\bar g$ small, with a constant C_{BCH} of N, the remainder satisfies $\|\mathcal{N}_b\| \le C_{\mathrm{BCH}}(\|\omega_1(b_-)\| + \|\omega_1(b_+)\| + \|c_b\|)(\|\omega_1(b_-)\| + \|\omega_1(b_+)\| + \|c_b\| + \|u_b\|)$, every product of two of the exponents containing an exponent other than u_b, together with the corresponding Lipschitz bound. Since $\bar D^* c = 0$ and $\Pi_{\mathcal{Z}}\bar D\omega_1 = 0$, the conditions $\bar D^* u_{\mathrm{H}} = 0$ and $\Pi_{\mathcal{Z}} u_{\mathrm{H}} = 0$ say that (ω_1, c) is a fixed point of

$$\mathcal{T}_{\mathrm{sl}}(\omega_1, c) \coloneqq \Big((\bar D^*\bar D)^+ \bar D^*(u + \mathcal{N}(\omega_1, c)),\, \Pi_{\mathcal{Z}}(u + \mathcal{N}(\omega_1, c)) \Big) . \tag{13.65}$$

By (i), $\|(\bar D^*\bar D)^+\bar D^*\| \le (64\varkappa)^{-1/2}$. Let $R_u \coloneqq 8N_q^2\|u\|_\infty$, which is at least $4\|u\|$ because $\mathcal{B}$ has $4N_q^4$ bonds and $|\cdot|_2 \le \|\cdot\|$, and $R_\omega \coloneqq R_u(64\varkappa)^{-1/2} \ge R_u$. On the set of the pairs with $\|\omega_1\| \le R_\omega$ and $\|c\| \le R_u$, every site lying on eight bonds and $\|\cdot\|_\infty \le N^{1/2}\|\cdot\|$, the bound on $\mathcal{N}_b$ gives $\|\mathcal{N}\| \le 90NC_{\mathrm{BCH}}R_\omega^2$ and the Lipschitz constant $90NC_{\mathrm{BCH}}R_\omega$. If $C_{\mathrm{sl}}N_q^2\mathsf{r}_j \le \varkappa$ with $C_{\mathrm{sl}} \coloneqq 180NC_{\mathrm{BCH}}$, then $90NC_{\mathrm{BCH}}R_\omega^2 \le \frac{3}{4}R_u$ and $2(64\varkappa)^{-1/2}\cdot 90NC_{\mathrm{BCH}}R_\omega \le \frac{1}{2}$, so $\mathcal{T}_{\mathrm{sl}}$ maps the set into itself and is a contraction there, and it has a fixed point. The condition holds for $\bar g$ small, because $N_q^2\mathsf{r}_j/\varkappa \le c_\varkappa^{-1}N_q^4\bar g^{3\varsigma_j/4}$. At the fixed point, c is constant, so $|c| \le N^{1/2}R_u/N_q^2 = 8N^{1/2}\|u\|_\infty \le 32N^{1/2}\mathsf{r}_j$; and $\|\omega_1\|_\infty \le N^{1/2}R_\omega$ gives $\|u_{\mathrm{H}}\|_\infty \le C'_{\mathrm{sl}}N_q^2\mathsf{r}_j\varkappa^{-1/2}$ with a constant C'_{sl} of N. Finally $u_{\mathrm{H}} \perp \operatorname{ran}\bar D \oplus \mathcal{Z} = \ker K$.

(iii), the coercivity. By the series of the exponential and the logarithm, $V_s = e^{iu'_s}W'$ with $u'_{s,b} \coloneqq -i\log(e^{isu_{\mathrm{H},b}}e^{ic_b})$ and $\|u'_s\|_\infty \le 2(\|u_{\mathrm{H}}\|_\infty + |c|) = \delta_*$. The fields $V' = e^{i\sigma u'_s}W'$,

$\sigma \in [0,1]$, lie in $\mathcal{R}$: every bond value is within δ_* of that of the flat W', so their plaquettes satisfy $|V'(\partial p') - 1| \le N^{1/2}\|V'(\partial p') - 1\| \le 4N^{1/2}\delta_* < \varepsilon_1$. They lie in $\mathcal{R}'$, because the holonomies of W' around the I_b are 1, and by Lemma 13.21(i) at the base W' and the bound on $\|H_k(B) - H_k(0)\|$ in the proof of Lemma 13.22(i), with $B = \sigma u'_s$, those of V' differ from 1 by at most $e^{2P\varepsilon_V \delta_*/r_V} - 1 \le r_G/2$. Since $\frac{\mathrm{d}}{\mathrm{d}\sigma}\Delta(e^{i\sigma u'_s}W') = \partial_{u'_s}\Delta(e^{i\sigma u'_s}W')$, Lemma 13.22(ii) gives $\|\Delta(V') - K\| \le C_\Delta N_q^4 \delta_*$, with $K = \Delta(W')$ by Lemma 13.14(iv). With the slice bounds, $N_q^4\delta_*/\varkappa \preceq \bar g^{5\varsigma_j/8}$ and $N_q^4\delta_*/\lambda \preceq \bar g^{7\varsigma_j/8}$, so $C_\Delta N_q^4\delta_* \le \min(\varkappa/4, \lambda/2)$ for $\bar g$ small; since $K \ge 0$, this gives $\Delta(V') + \lambda \ge \lambda/2$. Now $e^{i(s+h)u_{\mathrm{H}}}B = e^{ihu_{\mathrm{H}}}V_s$, so the function $s \mapsto \mathsf{A}(V_s)$, which is real analytic because $V_s \in \mathcal{R}$, has the derivatives $\langle u_{\mathrm{H}}, \nabla\mathsf{A}(V_s)\rangle$ and $\langle u_{\mathrm{H}}, \Delta(V_s)u_{\mathrm{H}}\rangle$; since A is invariant under the gauge transformations of $\mathbb{T}_{j_q}$ (see (iv)), Taylor's formula gives

$$\mathsf{A}(V) = \mathsf{A}(V_1) = \mathsf{A}(B) + \langle u_{\mathrm{H}}, \nabla\mathsf{A}(B)\rangle + \int_0^1 (1-s)\langle u_{\mathrm{H}}, \Delta(V_s)u_{\mathrm{H}}\rangle \,\mathrm{d}s \,. \tag{13.66}$$

The middle term vanishes, by (iv) below and $\Pi_{\mathcal{Z}} u_{\mathrm{H}} = 0$. By (i) and (ii), $\langle u_{\mathrm{H}}, Ku_{\mathrm{H}}\rangle \ge \varkappa\|u_{\mathrm{H}}\|^2$, so that $\langle u_{\mathrm{H}}, \Delta(V_s)u_{\mathrm{H}}\rangle$ lies between $\langle u_{\mathrm{H}}, Ku_{\mathrm{H}}\rangle \mp \frac14\varkappa\|u_{\mathrm{H}}\|^2$ and is at least $\frac34\varkappa\|u_{\mathrm{H}}\|^2$. Hence (13.66) gives $\mathsf{A}(V) - \mathsf{A}(B) \ge \frac38\varkappa\|u_{\mathrm{H}}\|^2$, and also $\frac12\langle u_{\mathrm{H}}, Ku_{\mathrm{H}}\rangle \le \mathsf{A}(V) - \mathsf{A}(B) + \frac18\varkappa\|u_{\mathrm{H}}\|^2 \le \frac43(\mathsf{A}(V) - \mathsf{A}(B))$, so that $\langle u_{\mathrm{H}}, Ku_{\mathrm{H}}\rangle \le \frac83\mathsf{A}(V)$. Taylor's formula on $[0,s]$ with the same upper bound for the integrand gives $\mathsf{A}(V_s) \le \mathsf{A}(B) + \frac12(\langle u_{\mathrm{H}}, Ku_{\mathrm{H}}\rangle + \frac14\varkappa\|u_{\mathrm{H}}\|^2) \le \mathsf{A}(B) + \frac53(\mathsf{A}(V) - \mathsf{A}(B)) \le 3\mathsf{A}(V)$, using $\mathsf{A}(B) \ge 0$.

(iv), the symmetry. Let $\mathcal{T}$ be the translation of $\mathbb{T}_{j_q}$ by one block in a direction ν, $(\mathcal{T}V)(b) := V(b - e_\nu)$. With the convention $V^\upsilon(b) = \upsilon(b_-)V(b)\upsilon(b_+)^{-1}$, under which the gauge transformations act on $\mathfrak{g}^{\mathcal{B}}$ as in Section 13.7, $\mathcal{T}W' = W'^{g_\nu}$, with g_ν equal to $\tilde h_\nu$ on the sites x with $x_\nu \equiv \sigma_\nu + 1$ and to 1 elsewhere: the bonds of the direction ν which enter this layer get the value $\tilde h_\nu \tilde h_\nu^{-1} = 1$, those which leave it the value $\tilde h_\nu$, and on the bonds inside the layer g_ν conjugates values in $\{1, \tilde h_\beta\}$, which commute with $\tilde h_\nu$. Since $\mathcal{T}c = c$ and $\mathsf{Ad}(\tilde h_\nu)c_\beta = c_\beta$, also $\mathcal{T}B = B^{g_\nu}$. The function A is invariant under the gauge transformations of $\mathbb{T}_{j_q}$, because $U_{K'}(V^\upsilon) = U_{K'}(V)^{\bar\upsilon}$ [7, (181), p. 307] and the action is gauge invariant, and under the translations by whole blocks, because these commute with $M^{K'}$ and leave the action invariant, so that $\mathcal{T}U_{K'}(V)$ is minimal for $\mathcal{T}V$. With the orthogonal map $S_\nu u := \mathsf{Ad}(g_\nu)^{-1}\mathcal{T}u$, acting on the values at b by $\mathsf{Ad}(g_\nu(b_-))^{-1}$, we get $(\mathcal{T}(e^{iu}B))^{g_\nu^{-1}} = e^{iS_\nu u}B$, hence $\mathsf{A}(e^{iS_\nu u}B) = \mathsf{A}(e^{iu}B)$ for all u. Differentiating, $S_\nu\nabla\mathsf{A}(B) = \nabla\mathsf{A}(B)$ and S_ν commutes with $\Delta(B)$. On the component r, S_ν is the twisted translation $\mathcal{T}_\nu$ of (i), so the joint fixed vectors of $S_1, \dots, S_4$ are the constants of the components with $\theta_r = 0$, and the real ones form $\mathcal{Z}$. This gives (iv).

(v), the orders. By (iii) and $\mathsf{A}(V) \le 2T_{\mathrm{ng}}g^2$, $\|u_{\mathrm{H}}\|^2 \le \frac{16}{3}T_{\mathrm{ng}}g^2/\varkappa \preceq \bar g^{2-\varsigma_j/4}$, by (i); and $|c| \le 32N^{1/2}\bar g^{\varsigma_j}$, so $\delta_* \preceq \bar g^{\varsigma_j}$, because $\varsigma_j \le \frac12 < 1 - \varsigma_j/8$. *q.e.d.*

With the decomposition in hand we can bound the defect $\mathfrak{e}_\alpha$. We follow it along the segment $s \mapsto V_s$ from the field B, which differs from the flat W' only in the directions of $\mathcal{Z}$. At B the gradient of the action lies in $\mathcal{Z}$, so only the component of $\nabla\Xi_\alpha$ in $\mathcal{Z}$ enters, and it is of the second order in c because the fluxes vanish to the second order along $\mathcal{Z}$; the flux difference $\Xi_\alpha(B)$ itself is a commutator of a small transport with the flux of I_0. Along the segment the derivative of the defect is $-\lambda\langle u_{\mathrm{H}}, w_\alpha\rangle$ up to terms of a higher order, and at a flat field this is the term which Proposition 13.20(ii) controls.

Proposition 13.28 (The Defect of the Integration by Parts)**.** *There is a constant C_E of the kind of Section 13.8 such that for $\bar g$ small, $K \ge K_{\mathrm{w}}$, and every $V \in \operatorname{supp}\zeta$, the operator $\Delta(V) + \lambda$ is at least $\lambda/2$, and*

$$\Big|\sum_{\alpha=1}^{d_{\mathfrak{g}}} X^\alpha(V)\,\mathfrak{e}_\alpha(V)\Big| \le C_E\bar g^2(\log\bar g^{-2})^{7r+1-p_\lambda/2} + \bar g^{2+\varsigma_*/2} \,. \tag{13.67}$$

Proof. Reduction. Under a gauge transformation ω of $\mathbb{T}_{j_q}$, the gradients and Δ are covariant under the orthogonal map $u_b \mapsto \mathsf{Ad}(\omega(b_-))u_b$, A is invariant, and the logarithms based at o change by $\mathsf{Ad}(\omega(o))$; so, as for c_λ in Section 13.7, the sum in (13.67) at V^ω is the same sum at V for the orthonormal basis $(\mathsf{Ad}(\omega(o))^{-1}t_\alpha)$, and it does not depend on the basis, being the contraction of a bilinear expression in t_α with $\sum_\alpha t_\alpha \otimes t_\alpha$. The spectrum of Δ is invariant as well. We may therefore replace V by $V_1 = V^{\omega'}$ of Proposition 13.27, whose notation we use; there $\Delta(V_1) + \lambda \geq \lambda/2$ by Proposition 13.27(iii). Since $|X_{0,0}(V_1)|_2 \leq \|X_{0,0}(V_1)\| \leq 2h^G$, the Cauchy–Schwarz inequality over α gives

$$\Big|\sum_\alpha X^\alpha \mathfrak{e}_\alpha\Big| \leq 2d_{\mathfrak{g}}^{1/2} h^G \max_\alpha |\mathfrak{e}_\alpha(V_1)| \,, \tag{13.68}$$

and it remains to bound $\mathfrak{e}_\alpha(V_1)$ for a fixed α. For the fields V_s, $s \in [0,1]$, among which $V_0 = B$, we have $\mathsf{A}(V_s) \leq 3\mathsf{A}(V) \leq 6T_{\mathrm{ng}} g^2$, and $e^{-itw}V_s \in \mathcal{R}$ for $\|w\| = 1$ and $0 \leq t \leq (2\mathsf{A}(V_s)/(C_\Delta N_q^4))^{1/2}$, a number $\preceq \bar{g}$, because the plaquettes of V_s are within $4N^{1/2}\delta_*$ of 1; so Lemma 13.22(iii) gives $\|\nabla \mathsf{A}(V_s)\| \leq \gamma_A := (12 C_\Delta N_q^4 T_{\mathrm{ng}})^{1/2} g$. On all the fields V' of Proposition 13.27(iii), Lemma 13.22(i) and the bound on the resolvent give $\|w_\alpha(V')\| \leq 2C_\Xi P/\lambda$.

The flux at the fields of $\mathcal{Z}$. For a real $z \in \mathcal{Z}$ with $|z| < r_V$ let $B_z := e^{iz}W'$, so that $B = B_c$, and let $\mathsf{t}_k(z) \in G$ be the transport from o to o_k along the bottom side of R_4 at a minimal configuration for B_z, which does not depend on its choice because both ends are sites of $\mathbb{T}_{j_q}$. By the proof of Lemma 13.13, $X_{0,k} = \mathsf{Ad}(\mathsf{t}_k)X_{k,k}$. We claim that $X_{k,k}(B_z) = X_{0,0}(B_z)$ for $k \leq 3$. The argument of Proposition 13.27(iv) for the translation $\mathcal{T}^{-1}$ in the direction μ gives $\mathcal{T}^{-1}B_z = B_z^{g'}$, with g' equal to $\tilde{h}_\mu^{-1}$ on the layer $x_\mu \equiv \sigma_\mu$ and to 1 elsewhere, and by induction $\mathcal{T}^{-n}B_z = B_z^{G_n}$, where G_n equals 1 off the layers $x_\mu \equiv \sigma_\mu, \sigma_\mu - 1, \dots, \sigma_\mu - n + 1$. For $n = k t_{\mathrm{b}} \leq 3t_{\mathrm{b}}$ these layers do not contain o, because $o_\mu = \sigma_\mu + 4$ and $N_q > 3t_{\mathrm{b}} + 4$. Since $\mathcal{T}^{-n}$ commutes with $M^{K'}$, the holonomy of I_0 at o for $\mathcal{T}^{-n}B_z$ is that of I_k at o_k for B_z, and by [7, (181), p. 307] it is also the holonomy of I_0 for B_z conjugated by $G_n(o) = 1$; this is the claim. Hence

$$\hat{\Xi}(z) := \Xi_\alpha(B_z) = t_\alpha \cdot (\mathsf{Ad}(\mathsf{t}_3(z)) - \mathsf{Ad}(\mathsf{t}_2(z)))X_{0,0}(B_z) \,. \tag{13.69}$$

At $z = 0$ the minimal configuration is U_{s} (Lemma 13.14(i)), which is 1 near R_4, so $\mathsf{t}_k(0) = 1$ and $X_{0,0}(B_0) = 0$; moreover $\frac{\mathrm{d}}{\mathrm{d}s} X^\beta(B_{sz})|_{s=0} = \langle \chi_0 \otimes t_\beta, DHz \rangle = 0$ by Lemma 13.14(iv) and $\mathcal{Z} \subset \ker DH$. By Lemma 13.21(i) at the base W', the transports $\mathsf{t}_k(z)$ are real analytic in z, so in (13.69) the first factor vanishes to the first order at $z = 0$ and the second to the second order, and the Taylor polynomial of degree two of $\hat{\Xi}$ at 0 vanishes. By the proof of Lemma 13.22(i) at the base $W' \in \mathcal{R}'$, $\hat{\Xi}$ extends analytically to the complex $z \in \mathcal{Z}^{\mathbb{C}}$ with $|z| < \rho_P$, with modulus at most $\frac{1}{2}$, and Schwarz's lemma for the function $\zeta \mapsto \hat{\Xi}(\zeta z/|z|)$, which vanishes to the third order at $\zeta = 0$, gives $|\hat{\Xi}(z)| \leq \frac{1}{2}(|z|/\rho_P)^3 \leq \frac{1}{2}c_P^3 P^3 |z|^3$.

The defect at B. Since $\nabla \mathsf{A}(B) \in \mathcal{Z}$ and $(\Delta(B) + \lambda)^{-1}$ maps $\mathcal{Z}$ into itself (Proposition 13.27(iv)), $\langle w_\alpha(B), \nabla \mathsf{A}(B) \rangle = \langle \Pi_{\mathcal{Z}} \nabla \Xi_\alpha(B), (\Delta(B) + \lambda)^{-1} \nabla \mathsf{A}(B) \rangle$. Let $m := |c| + \bar{g}$ and $z' \in \mathcal{Z}$ with $|z'| = 1$. For complex τ with $|\tau| \leq m$ the form $Z(\tau)$ with $e^{iZ(\tau)_{\mathfrak{b}}} = e^{i\tau z'_{\mathfrak{b}}} e^{ic_{\mathfrak{b}}}$ lies in $\mathcal{Z}^{\mathbb{C}}$, because $\mathfrak{g}^{\tilde{h}}$ is a Lie algebra, and $|Z(\tau)| \leq 2(e^m - 1 + |c|) \leq 5m < \rho_P$ for $\bar{g}$ small, by the series of the logarithm; so Cauchy's estimate for $\tau \mapsto \hat{\Xi}(Z(\tau))$ on $|\tau| \leq m$, whose derivative at 0 is $\partial_{z'} \Xi_\alpha(B)$, gives $|\partial_{z'} \Xi_\alpha(B)| \leq \frac{1}{2} c_P^3 P^3 (5m)^3/m \leq 63 c_P^3 P^3 m^2$. Since $\|z'\| \geq N_q^2 N^{-1/2}$ for such z', $\|\Pi_{\mathcal{Z}} \nabla \Xi_\alpha(B)\| \leq \hat{C}_\Xi N_q m^2$ with $\hat{C}_\Xi := 504 N^{1/2} c_P^3$, using $P \leq 2N_q$. For $\Xi_\alpha(B)$ itself we use (13.69) at $z = c$: by unitarity, $\|(\mathsf{Ad}(\mathsf{t}_3) - \mathsf{Ad}(\mathsf{t}_2))Y\| \leq 2\|\mathsf{t}_3 - \mathsf{t}_2\|\|Y\|$ and $\|\mathsf{t}_3 - \mathsf{t}_2\|$ is the distance from 1 of the transport from o_2 to o_3. By Lemma 13.21 at the base W', this transport is $w_c(o_2)\mathsf{t}' w_c(o_3)^{-1}$, up to the order of the factors, with t' the transport of $e^{i\eta \mathcal{H}(c)} U_{\mathrm{s}}$ along t_{b}/η fine bonds on which $U_{\mathrm{s}} = 1$, so that, all the factors being unitary, its distance

from 1 is at most $(t_{\mathrm{b}}+144d)\varepsilon_V|c|/r_V$. Moreover $\|X_{0,0}(B)\| \le \|X_{0,0}(V_1)\| + C_\Xi P\|u_{\mathrm{H}}\|_\infty \le 2h^G + C_\Xi P\|u_{\mathrm{H}}\|_\infty$, by Lemma 13.22(i) along the segment, whose fields lie in $\mathcal{R}'$. Together, with $\|(\Delta(B)+\lambda)^{-1}\| \le 2/\lambda$,

$$|\mathfrak{e}_\alpha(B)| \le \frac{2}{\lambda}\hat{C}_\Xi N_q m^2 \gamma_A + \frac{2(P+144d)\varepsilon_V}{r_V}|c|(2h^G + C_\Xi P\|u_{\mathrm{H}}\|_\infty)\ . \tag{13.70}$$

The segment. Let $w_s := w_\alpha(V_s)$ and $\phi(s) := \mathfrak{e}_\alpha(V_s)$. Since $V_{s+h} = e^{ihu_{\mathrm{H}}}V_s$, we have $\frac{\mathrm{d}}{\mathrm{d}s}\Xi_\alpha(V_s) = \langle u_{\mathrm{H}}, \nabla\Xi_\alpha(V_s)\rangle = \langle u_{\mathrm{H}}, (\Delta(V_s)+\lambda)w_s\rangle$, and $\frac{\mathrm{d}}{\mathrm{d}s}\langle w, \nabla\mathsf{A}(V_s)\rangle = \partial_\sigma\partial_t \mathsf{A}(e^{itw}e^{i\sigma u_{\mathrm{H}}}V_s)|_{t=\sigma=0}$ for a fixed w. By the Baker–Campbell–Hausdorff formula, $e^{itw}e^{i\sigma u_{\mathrm{H}}} = e^{i(tw+\sigma u_{\mathrm{H}}+\frac{1}{2}t\sigma\, i[w,u_{\mathrm{H}}])+O_3}$ bondwise, where $i[w,u_{\mathrm{H}}]$ is real and O_3 collects the terms of order three in (t,σ), so that this mixed derivative is $\langle w, \Delta(V_s)u_{\mathrm{H}}\rangle + \frac{1}{2}\langle i[w,u_{\mathrm{H}}], \nabla\mathsf{A}(V_s)\rangle$. Therefore

$$\phi'(s) = \langle \partial_s w_s, \nabla\mathsf{A}(V_s)\rangle - \lambda\langle u_{\mathrm{H}}, w_s\rangle + \frac{1}{2}\langle i[w_s,u_{\mathrm{H}}], \nabla\mathsf{A}(V_s)\rangle\ , \tag{13.71}$$

the terms with $\Delta(V_s)$ cancelling. We bound the three terms.

The first and the third term. Differentiating the resolvent, $\partial_s w_s = -R_s(\partial_{u_{\mathrm{H}}}\Delta(V_s))R_s\nabla\Xi_\alpha(V_s) + R_s\partial_{u_{\mathrm{H}}}\nabla\Xi_\alpha(V_s)$ with $R_s := (\Delta(V_s)+\lambda)^{-1}$, so Lemma 13.22(i), (ii) and $\|R_s\| \le 2/\lambda$ give $\|\partial_s w_s\| \le C_\Xi P\|u_{\mathrm{H}}\|_\infty(4C_\Delta N_q^4/\lambda^2 + 2P/\lambda)$. Since $|[a,b]|_2 \le 2\|b\||a|_2$, $\|i[w_s,u_{\mathrm{H}}]\| \le 2\|u_{\mathrm{H}}\|_\infty\|w_s\|$. Hence the first and the third term are at most

$$\gamma_A C_\Xi P\|u_{\mathrm{H}}\|_\infty\Big(\frac{4C_\Delta N_q^4}{\lambda^2} + \frac{4P}{\lambda}\Big) \preceq \bar{g}^{2-\varsigma_j/8}\ , \tag{13.72}$$

by Proposition 13.27(v), $\gamma_A \preceq \bar{g}$, and $1/\lambda \preceq 1$.

The term of the regularisation. We compare it with its value at the flat field. Writing $R_s - (K+\lambda)^{-1} = -(K+\lambda)^{-1}(\Delta(V_s)-K)R_s$,

$$\begin{aligned}\lambda\langle u_{\mathrm{H}}, w_s\rangle = {}& \lambda\langle (K+\lambda)^{-1}u_{\mathrm{H}}, \nabla\Xi_\alpha(W')\rangle - \lambda\langle (K+\lambda)^{-1}u_{\mathrm{H}}, (\Delta(V_s)-K)w_s\rangle \\ & + \lambda\langle (K+\lambda)^{-1}u_{\mathrm{H}}, \nabla\Xi_\alpha(V_s) - \nabla\Xi_\alpha(W')\rangle\ .\end{aligned} \tag{13.73}$$

Since $u_{\mathrm{H}} \perp \ker K$ and $K \ge \varkappa$ there, $\|\lambda(K+\lambda)^{-1}u_{\mathrm{H}}\| \le (\lambda/\varkappa)\|u_{\mathrm{H}}\|$. By Proposition 13.27(iii), $\|\Delta(V_s)-K\| \le C_\Delta N_q^4\delta_*$, and Lemma 13.22(i) along the fields $e^{i\sigma u'_s}W'$, which lie in $\mathcal{R}'$, gives $\|\nabla\Xi_\alpha(V_s) - \nabla\Xi_\alpha(W')\| \le C_\Xi P^2\delta_*$; so the last two terms of (13.73) are at most

$$\frac{\|u_{\mathrm{H}}\|\delta_*}{\varkappa}(2C_\Delta C_\Xi P N_q^4 + C_\Xi P^2) \preceq \bar{g}^{1+5\varsigma_j/8}\ , \tag{13.74}$$

by Proposition 13.27(i) and (v). In the first term we use $\nabla\Xi_\alpha(W') = H^*D^*((\chi_3-\chi_2)\otimes t_\alpha)$ (Lemma 13.14(iv)) and Proposition 13.20(ii), which give $\lambda\langle (K+\lambda)^{-1}u_{\mathrm{H}}, \nabla\Xi_\alpha(W')\rangle = \langle P_\lambda DHu_{\mathrm{H}}, (\chi_3-\chi_2)\otimes t_\alpha\rangle$, where P_λ denotes the operator P of that proposition. We expand $t_\alpha = \sum_r a_r E_r$ in the basis of the components, with $\sum_r |a_r|^2 = 1$; the operators D, H, P_λ act on each component, so (13.56) and the Cauchy–Schwarz inequality over r give

$$\begin{aligned}|\langle P_\lambda DHu_{\mathrm{H}}, (\chi_3-\chi_2)\otimes t_\alpha\rangle| &\le 2\pi t_{\mathrm{b}}(\lambda C_V(s_{\mathrm{b}}+2)(t_{\mathrm{b}}+2))^{1/2}\|DHu_{\mathrm{H}}\| \\ &\le 2\pi(6C_V T_{\mathrm{ng}})^{1/2}N_q g(\log\bar{g}^{-2})^{-p_\lambda/2}\ ,\end{aligned} \tag{13.75}$$

because $\|DHu_{\mathrm{H}}\|^2 = \langle u_{\mathrm{H}}, Ku_{\mathrm{H}}\rangle \le 3\mathsf{A}(V) \le 6T_{\mathrm{ng}}g^2$ by Proposition 13.27(iii), $t_{\mathrm{b}} \le N_q$, $(s_{\mathrm{b}}+2)(t_{\mathrm{b}}+2) \le N_q^2$, and $\lambda^{1/2} = N_q^{-1}(\log\bar{g}^{-2})^{-p_\lambda/2}$.

Conclusion. The right side of (13.75) is at most $C\bar{g}(\log\bar{g}^{-2})^{3r+1/2-p_\lambda/2}$, by the sizes of Section 13.8. By Proposition 13.27(v), $|c|$, $m \preceq \bar{g}^{\varsigma_j}$ and $\|u_{\mathrm{H}}\|_\infty \preceq \bar{g}^{1-\varsigma_j/8}$, so the two terms of

(13.70) are $\preceq \bar g^{1+2\varsigma_j}$ and $\preceq \bar g^{1+7\varsigma_j/8}$. With (13.72), (13.74), and $2-\varsigma_j/8 > 1+5\varsigma_j/8$, we obtain from $\mathfrak{e}_\alpha(V_1) = \phi(0) + \int_0^1 \phi'(s)\,\mathrm{d}s$ and $\phi(0) = \mathfrak{e}_\alpha(B)$ that $|\mathfrak{e}_\alpha(V_1)| \le C\bar g(\log \bar g^{-2})^{3r+1/2-p_\lambda/2} + \mathsf{E}$ with $\mathsf{E} \preceq \bar g^{1+5\varsigma_j/8}$. Since $h^G \le C\bar g(\log \bar g^{-2})^{4r+1/2}$ and $\varsigma_j \ge \varsigma_*$, (13.68) gives (13.67), the term with E being at most $\bar g^{2+\varsigma_*/2}$ for $\bar g$ small. *q.e.d.*

We can now assemble the second moments. The integration by parts of Lemma 13.23 with the vector fields w_α turns $\sum_\alpha \langle F_\alpha \Xi_\alpha\rangle$ into $g^2\langle \zeta c_\lambda\rangle$, up to the defect of Proposition 13.28 and to three terms which carry a factor g^2 besides the factor h^G of F_α; and at a field near a flat one the number c_λ is close to its value at the flat field, which Proposition 13.20(i) bounds below.

Proposition 13.29 (The Second Moments of the Fluxes)**.** *For $\bar g$ sufficiently small, depending on q, t/u, s/u, T/u, π, and the constants of the construction, and for every $K \ge K_{\mathrm w}$, $\Psi_3 - \Psi_2 \ge 3c_O\bar g^2$, with c_O of Theorem 1.8.*

Proof. The cutoff. By (13.41) and (13.46), $\Psi_3 - \Psi_2 = (C_2(\pi)/d_{\mathfrak g})\langle \mathbf{1}_{\mathcal V} \sum_\alpha X^\alpha \Xi_\alpha\rangle$. The function ζ equals 1 on $\mathcal V$ and vanishes outside $\mathcal V_3$, on which $|\sum_\alpha X^\alpha \Xi_\alpha| \le |X_{0,0}|_2 |X_{0,3} - X_{0,2}|_2 \le 8(h^G)^2$, and $\langle \mathbf{1}_{\mathcal V^{\mathrm c}}\rangle \le \bar g^8$ by Lemma 13.7(ii); hence

$$\Big| \frac{d_{\mathfrak g}}{C_2(\pi)}(\Psi_3 - \Psi_2) - \sum_\alpha \langle F_\alpha \Xi_\alpha\rangle \Big| \le 8(h^G)^2 \bar g^8 \,. \tag{13.76}$$

The integration by parts. By Proposition 13.28, $\Delta + \lambda \ge \lambda/2$ on the compact set $\operatorname{supp}\zeta$, on which the holonomies of the I_b are within $\sqrt{3}h^G < r_G/2$ of 1; hence there is an open neighbourhood of $\operatorname{supp}\zeta$ contained in $\mathcal R'$ on which $\Delta + \lambda$ is invertible, and w_α is real analytic there, by Lemma 13.22. On $\operatorname{supp}\zeta$ we write $\Xi_\alpha = \langle w_\alpha, \nabla \mathsf A\rangle - \mathfrak{e}_\alpha$, and Lemma 13.23 with $F = F_\alpha$, whose support lies in $\mathcal K \subset \mathcal O$, and $v = w_\alpha$ gives, after summation over α and with $\sum_\alpha \langle w_\alpha, \nabla X^\alpha\rangle = c_\lambda$,

$$\sum_\alpha \langle F_\alpha \Xi_\alpha\rangle = g^2\langle \zeta c_\lambda\rangle + g^2 \sum_\alpha \Big(\langle X^\alpha \langle w_\alpha, \nabla\zeta\rangle\rangle + \langle F_\alpha \operatorname{div} w_\alpha\rangle + \langle F_\alpha \langle w_\alpha, \nabla \mathcal A'\rangle\rangle \Big) - \Big\langle \zeta \sum_\alpha X^\alpha \mathfrak{e}_\alpha \Big\rangle \,. \tag{13.77}$$

On $\operatorname{supp}\zeta$ we have $|X^\alpha| \le 2h^G$, $\|w_\alpha\| \le 2C_\Xi P/\lambda$, and $\|\nabla \mathsf A\| \le (4C_\Delta N_q^4 T_{\mathrm{ng}})^{1/2} g$, by Lemma 13.22(i) and (iii), whose condition holds there because the plaquettes are within ε_ζ of 1.

The four error terms. The gradient $\nabla\zeta = \vartheta'(\mathsf A/(T_{\mathrm{ng}}g^2))\nabla \mathsf A/(T_{\mathrm{ng}}g^2)$ vanishes on $\mathcal V$, so the first is at most $g^2 d_{\mathfrak g} 2h^G (2C_\Xi P/\lambda)\|\vartheta'\|_\infty (4C_\Delta N_q^4/T_{\mathrm{ng}})^{1/2} g^{-1}\bar g^8 \preceq \bar g^{10}$. For the second, $\operatorname{div} w_\alpha = \sum_{b,\beta} \langle e_{b,\beta}, \partial_{e_{b,\beta}} w_\alpha\rangle$, and differentiating the resolvent as in the proof of Proposition 13.28, with $\|e_{b,\beta}\|_\infty \le N^{1/2}$,

$$|\operatorname{div} w_\alpha| \le 4N_q^4 d_{\mathfrak g} N^{1/2} C_\Xi \Big(\frac{2P^2}{\lambda} + \frac{4C_\Delta N_q^4 P}{\lambda^2} \Big) \,, \tag{13.78}$$

so that the second term is $\preceq \bar g^3$. For the third, Lemma 13.24 and $\|w_\alpha\|_\infty \le N^{1/2}\|w_\alpha\|$ bound it by $g^2 d_{\mathfrak g} 2h^G M_E N_q^4 N^{1/2}(2C_\Xi P/\lambda)/r_E$, and $r_E \ge c\bar g(\log \bar g^{-2})^{q_0}$ by the bound on $r_{i,K'}$; with the sizes of Section 13.8 it is at most $C\bar g^2(\log \bar g^{-2})^{11r+1/2+p_\lambda - q_0}$. The last term of (13.77) is bounded by Proposition 13.28, since $0 \le \zeta \le 1$.

The main term. On $\operatorname{supp}\zeta$, in the gauge of Proposition 13.27, $c_\lambda(V) = c_\lambda(V_1)$ and $V_1 = e^{iu_1'} W'$, and we follow c_λ along the fields $e^{i\sigma u_1'} W'$, which lie in $\mathcal R'$ and satisfy $\|(\Delta +$

$\lambda)^{-1}\| \le 2/\lambda$ by Proposition 13.27(iii). Differentiating each of the three factors of (13.47) by Lemma 13.22(i), (ii),

$$|c_\lambda(V) - c_\lambda(W')| \le d_{\mathfrak{g}} C_\Xi^2 \Big(\frac{4P^3}{\lambda} + \frac{4C_\Delta N_q^4 P^2}{\lambda^2} \Big) \delta_* \preceq \bar{g}^{\varsigma_*} \ . \tag{13.79}$$

By Proposition 13.20(i), whose last two terms tend to 0 with $\bar{g}$ because $t_{\mathfrak{b}} \ge (\log \bar{g}^{-2})^r$ and $\lambda t_{\mathfrak{b}}^2 (s_{\mathfrak{b}}+2)(t_{\mathfrak{b}}+2) \le N_q^2 (\log \bar{g}^{-2})^{-p_\lambda} \preceq (\log \bar{g}^{-2})^{2r-p_\lambda}$, we get $c_\lambda \ge d_{\mathfrak{g}} s^2 t^3/(112 e^{1/2} T^5) > 0$ on $\operatorname{supp} \zeta$ for $\bar{g}$ small. Since $\zeta \ge \mathbf{1}_{\mathcal{V}}$ and $\langle \mathbf{1}_{\mathcal{V}} \rangle \ge 1 - \bar{g}^8$, $g^2 \langle \zeta c_\lambda \rangle \ge (1-\bar{g}^8) g^2 d_{\mathfrak{g}} s^2 t^3/(112 e^{1/2} T^5)$.

Conclusion. The exponents $7r + 1 - p_\lambda/2$ and $11r + \frac{1}{2} + p_\lambda - q_0$ are negative by the choice of p_λ, so all the error terms in (13.76) and (13.77) are $o(\bar{g}^2)$ as $\bar{g} \to 0$, uniformly in $K \ge K_{\mathrm{w}}$. With $g^2 \ge \bar{g}^2/2$,

$$\Psi_3 - \Psi_2 \ge \frac{C_2(\pi)}{d_{\mathfrak{g}}} \Big((1-\bar{g}^8) \frac{\bar{g}^2}{2} \cdot \frac{d_{\mathfrak{g}} s^2 t^3}{112 e^{1/2} T^5} - o(\bar{g}^2) \Big) \ge \frac{C_2(\pi) s^2 t^3}{224 e^2 T^5} \bar{g}^2 = 3 c_O \bar{g}^2 \tag{13.80}$$

for $\bar{g}$ small. *q.e.d.*

13.9 Proof of Theorem 1.8

We now collect the pieces. The shape term is reduced to $\Psi_3 - \Psi_2$ by Propositions 13.11 and 13.12 and Lemma 13.13, with errors of the orders $\bar{g}^3$ and $\bar{g}^4$ up to logarithms, Proposition 13.29 bounds $\Psi_3 - \Psi_2$ below, and Theorem 13.10 bounds the pair terms.

Proof of Theorem 1.8. Let $\bar{g}$ be so small that all the results of this section hold, in particular $\bar{g} \le \bar{g}_1'$.

(i) This is Proposition 13.4(ii).

(ii) Let $K \ge K_{\mathrm{w}}$. By Proposition 13.11, the decomposition $\xi_{R_n} = (f_{K'}^{(R_n)} - 1) + I(R_n)$, Proposition 13.12, and Lemma 13.13, in which $\Psi_3 - \Psi_2$ is real,

$$\mathfrak{y}_S(W_K) \ge \Psi_3 - \Psi_2 - |r_H| - 8 \max_n \mathsf{e}(R_n) - \bar{g}^9 - 8 \max_n \sigma(R_n)^2 \ . \tag{13.81}$$

Here $|r_H| \le 512\, \mathsf{l}^3 + 16 \bar{g}^8$ with $\mathsf{l} = 1.05 c_\pi h^G \le C \bar{g} (\log \bar{g}^{-2})^{4r+1/2}$, $8 \max_n \mathsf{e}(R_n) \le C \bar{g}^3 (\log \bar{g}^{-2})^{m_\sigma'}$, and $\sigma(R_n) \le C_\sigma \bar{g}^2 (\log \bar{g}^{-2})^{m_\sigma}$ by Lemma 13.7(iv), so that with Proposition 13.29 the right side is at least $2 c_O \bar{g}^2$ for $\bar{g}$ small, for every $K \ge K_{\mathrm{w}}$. By Proposition 13.4(i) and (ii), $\mathfrak{y}_S(W) = \lim_K \mathfrak{y}_S(N_K) = \lim_K \mathfrak{y}_S(W_K) \ge 2 c_O \bar{g}^2$. By Theorem 13.10, $|\mathfrak{y}_k(W)| \le 16 C_\sigma^2 \bar{g}^4 (\log \bar{g}^{-2})^{2m_\sigma} \le \frac{1}{2} c_O \bar{g}^2$ for $\bar{g}$ small, and (13.1) gives $\mathfrak{Y}(W) = \mathfrak{y}_S(W) - \mathfrak{y}_2(W) + \mathfrak{y}_3(W) \ge c_O \bar{g}^2 \ge \frac{1}{2} c_O \bar{g}^2$.

For a level $q' \ge q$ the eleven diagrams are ensembles at the level q', with admissible loops, and $q' \ge q \ge q_\sharp(c_\pi, \bar{g})$, so that Theorem 1.5 gives the limits $W^{(q')}(\mathcal{D})$. By (1.12), $W_K^{(q')}(\mathcal{D}) / W_K(\mathcal{D})$ is a product of the $j_{q'} - j_q$ step factors $\kappa_k(C, \chi)$, $K - j_{q'} < k \le K - j_q$, of the at most two loops of $\mathcal{D}$. By Definition 7.8, the exponent of such a factor is bounded by $\sum_v N_v^{(k-1)}(C) |\kappa_v^{(k)} / \Pi_k + \mathbf{1}_{\mathbf{R}}(k) \mu_{k,v}^{\mathbf{R}}| + |\mathcal{K}(C)| \max_{v,w} |\zeta_{v,w}^{(k)} + \mathbf{1}_{\mathbf{R}}(k) \zeta_{k,v,w}^{\mathbf{R}}|$, where the coefficients are bounded by $C_{S'} \bar{\mathfrak{a}} + C_E^R$ and $C_\zeta \bar{\mathfrak{a}} + C_E^R$, by the proof of Proposition 7.13, and $N_v^{(k-1)}(C) \le P_q(C) a_{K'} L^{K'-k+1} \le P_q(C) a_{K'} L^{j_{q'} - j_q}$, which does not depend on K. So $|W_K^{(q')}(\mathcal{D})| \ge e^{-C'} |W_K(\mathcal{D})|$ with C' independent of K, and $W^{(q')}(\mathcal{D}) \ne 0$ by (i). Proposition 13.4(iii) gives $\mathfrak{Y}(W^{(q')}) = \mathfrak{Y}(W)$.

(iii) By Proposition 13.4(ii) and part (ii), $\mathfrak{Y}^{\mathrm{sym}}(W) = \mathfrak{Y}(W) > 0$, and W does not vanish on the eleven diagrams by (i). Proposition 13.2, on the other hand, gives $\mathfrak{Y}^{\mathrm{sym}} = 0$ for the function of every Gaussian charge model and for every limit of such functions which does not vanish on the eleven diagrams, so W is neither. *q.e.d.*

14 Notes on the Papers

We record the places at which Bałaban's papers sketch or defer an argument which we use, the places at which we read a printed formula otherwise than it is printed, and the conventions in which our notation differs from his.

Deferred and sketched arguments. The asymptotic behaviour of the couplings, [8, Theorem 2, (0.31), p. 259], is stated with the remark that "a proof of this theorem, based on perturbative calculations, will be given in a separate paper"; the bounds [9, (2.6), (2.7), p. 255] on the running couplings are stated to "follow from the renormalization group equations (0.20) [I]". We do not use these statements: Section 4 proves what we need for the construction of this paper. The bounds on the walk factors of the propagators are given "rather briefly" [5, p. 409], and the bounds on the factors of the walk expansions are stated up to powers of the constants [5, p. 413]; we use them as printed, and the comparison of two fine lattices in Section 9 is proved in Supplement, Proposition S.12. Bałaban sketches a second proof of [9, Theorem 2] by gauge invariance [9, pp. 283–284]; we use the first. That the configurations which satisfy pointwise conditions belong to the spaces $U_j^c(X, \alpha_0, \alpha_1)$ of the regular terms is stated in [8, p. 263]: for the configurations which satisfy the conditions (i)–(iii) of the definition of these spaces with smaller constants α_0', α_1', the bounds [8, (1.17)] on the configurations built from their averages are obtained "from Proposition 9 [15]", which is [7, Proposition 9, p. 309], and "for α_0' sufficiently small the above estimates imply the condition (iv)". The application of that proposition to the averages of a complex configuration is not carried out there. We use the statement as it is printed: it is the reason for the constant c_F in (5.17), and Lemma 6.19 depends on it. The expansion of the fluctuation integral of a step on a term with large-field regions is described in [9, pp. 276–278] by reference to [10]: the localisation is repeated "with the only change that the random walk expansions we use now are constructed for the sequence $\{\Omega_j\}$", the terms of the action whose localisation domains are not contained in Λ_{k+1} are stated to satisfy the conditions of the lemmata of that paper on the spaces [9, (3.43)], and the polymer expansion is "the exponentiated cluster expansion of Sect. 7 [I], or rather its appropriate generalization", of which it is said that "the differences, like the restriction of the integration to Λ_{k+1}, or the larger class of localization domains, do not matter". The generalisation is not written out. We use it for the activities of the vacuum polymers of such a term, in Sections 6.12 and 6.14. For the same terms the localisation of the propagators by parameters is said to be repeated "with the only change that the random walk expansions we use now are constructed for the sequence $\{\Omega_j\}$", with cubes "in proper scales", and "the remaining considerations and bounds are the same" [9, p. 276]; and the statement that the analyticity domains, which become smaller at each step by the powers of $\frac{1}{2}$ in [9, (2.34)–(2.39)], still contain the configurations of the next step is given without a computation: "the localization of the fluctuation field, and the exponential decay of the minimizing function $\mathbf{H}_k$ imply that this difference is still much greater than bounds on this function" [9, p. 277]. We use both in Section 6.14; what the outer fields add to the second, the rotation by the gauge transformation of the step, is computed there and in Supplement, Proposition S.9(vi).

Formulae read otherwise than printed. At a few places a printed reference or formula cannot be meant as it stands, because it contradicts another printed line or the computation into which it enters. At each of them exactly one reading agrees with the rest of the text, and we use that reading; for each we give the printed text, the lines which exclude it, and the place at which our argument depends on it.

References to displays. Three references point to displays which do not exist or which say

something else, and in each case the context names the display that is meant. Theorem 1 of [9, p. 262] assumes that the couplings satisfy "the inequalities (I.0.33)", and the second choice of the threshold of the first step replaces g_0 in the logarithmic factor by a bound $\gamma_0 \geq g_0$ "coming from the inequality (0.33) [I]" [9, p. 246]. The first paper of the series has no display (0.33), since the numbered displays of its Section 0 end with (0.32) on p. 260. The only one of them which bounds the sequence of the couplings is the two-sided bound (0.31) of [8, Theorem 2, p. 259], whose left inequality at $k = 0$ gives $g_0 \leq \gamma_0$ once $g \leq \gamma$, and Bałaban refers to it elsewhere as "the basic inequality (0.31)", "the result of Theorem 2" [12, p. 355], there with the second paper of the series printed as its source. We therefore read both references as (0.31). Theorem 3.6 is quoted with this hypothesis, and Supplement, Proposition S.1, verifies it for our construction in the form of the inequalities (2.6)–(2.9) of [9, pp. 255–256], which the proof uses in its place [9, p. 255]. Next, the estimates on the propagators in a background field use "the weighted distance $d(y, y')$ defined by (2.36) in [4]" [5, p. 397], where [4] is [3]. The display (2.36) of that paper, on p. 229, is the partition of unity $\sum_\square h_\square^2 = 1$, whereas the weighted distance of two points of the set $\mathfrak{B}$ of the blocks of all scales, which is the set in which [5] takes its points, is defined in (2.46), p. 231. We read (2.46); this is the distance in the bounds on the walks of Section 9.2 and in Supplement, Lemma 17.3. Finally, the gauge transformation u_1 of [7, (124), p. 296], which is determined by U_1 through the conditions $\overline{R_0 u_1}^j = 1$ on Λ_j, "is given by the formula (106) [5]" [7, pp. 296–297], where [5] is [5]. That paper numbers all its displays in the form (3.n), and its (3.106), p. 414, is the Neumann series of a propagator, while the display (106) of [7, p. 294] itself is an identity between quadratic forms. The formula meant is (106) of [4, p. 33], the reference [4] of the same paper, which writes the gauge transformation determined by U_1 explicitly through the averages of U_1 [4, pp. 32–33]. We read it so in Supplement, Lemma 17.14, and in its application in Section 13.8.

The sections of [I]. The paper on the complete model cites the cluster expansion of the fluctuation integral as "Sect. 7 [I]", with the formulae (I.7.1)–(I.7.13), and the localisation of the action as "Sect. 6 [I]", with (I.6.34) and (I.6.41) [9, pp. 275–278], although its reference [I] is the first paper of the series [8], which has no such sections, and it lists the second paper [10] separately. The material is that of [10]: its formulae (2.1), the Mayer expansion, (2.11)–(2.13), the polymer expansion and its exponentiated form, (1.34), the space of analyticity of the form of [9, (3.43), p. 276], and (1.41), the localised action, are the formulae cited as (I.7.1), (I.7.11)–(I.7.13), (I.6.34), and (I.6.41), and the bound of [10, (2.14)] is printed there as a bound of "(7.14)" [10, p. 15]. We read a reference (I.7.n) as [10, (2.n)] and a reference (I.6.n) as [10, (1.n)]; Section 6.4 depends on this reading.

Thresholds of scaled fields. Bałaban's fluctuation fields are scaled by the coupling and carry Gaussian weights of order one: the field A of the first step enters the configuration as $\exp ig_0 A$ [9, (1.19), p. 250], and the field A_k of the step $k+1$ as $\exp i[g_k C A_k - h\tilde{D}(g_k C A_k)]$ [9, (3.15), p. 268]. A threshold δ for configurations is therefore the threshold $g^{-1}\delta$ for the field, and the first step prints it so, pairing $|A(b)| < g_0^{-1}\delta_0$ with $|\exp ig_0 A(b)\, U_1(b) U_{1,\square'}^{-1}(b) - 1| < 2\delta_0$ [9, (1.18), (1.19), p. 250]. The decomposition of unity of the general step [9, (3.16), p. 268] prints $|A_k(b)| < \delta_k$ instead, although $\delta_k = g_k(A_1/A_0)p_0(g_k)$ [9, p. 265] is a threshold for configurations, with which [9, (3.19), p. 268] compares $|V^{(k)}(b)V_{\square'}^{(k)}(b)^{-1} - 1|$. As printed, the large-field function $\chi^{(k)c}(R_{k+1})$ of (3.16) would select every field above a positive power of g_k, where the Gaussian weight is close to 1, whereas [12, p. 380] draws from it the small factor $\exp(-\gamma_0 g_j^{-2}\delta_j^2 (LM_2R_j)^{-d}|R_j|)$, which is the Gaussian weight at the field $g_j^{-1}\delta_j$. We therefore read the threshold of (3.16) as $g_k^{-1}\delta_k$, which is our ϑ_k/g_k, and Definition 3.11(ii) states it as the condition $|g_k A_k(b)| < \vartheta_k$ on the unscaled field. The field B_j of the **R**-operation is in the same position. It carries the Gaussian weight $\exp(-\frac{1}{2}\langle B_j, C^*\Delta^{(j)} C B_j\rangle)$ and enters the

configuration as $\exp i[g_j C B_j - h\tilde{D}(g_j C B_j)]$ [11, (1.60), pp. 188–189], while $\delta'_j = g_j A_1 p_1(g_j)$ [11, p. 183] is a threshold for configurations [11, (1.27), p. 182]. The restriction $|B_j(b)| < \delta'_j$ printed in [11, (1.55), p. 188] would confine the field to a ball whose radius is a positive power of g_j, so that the small-field function $\chi'^{(j)}$ would carry almost none of the Gaussian mass; the restriction of the field A_j in the same paper is printed as $|A_j(b)| < g_j^{-1}\delta_j$ [11, (1.9), p. 178], the condition paired with (1.55) is $|\exp i g_j B_j(b) - 1| < 2\delta'_j$ [11, (1.59), p. 188], and [12, p. 383] bounds a scaled field below by $g_j^{-1}\delta'_j = A_1 p_1(g_j)$. We read the threshold of (1.55) as $g_j^{-1}\delta'_j$, which Definition 3.11(ii) states as $|g_j B_j(b)| < \delta'_j$. The value of δ_0 itself is printed as the chain $\delta_0 = g_0 A_1 p_0(g_0) = (A_1/A_0)\varepsilon_0$ [9, p. 247], whose two members differ by the factor A_0 in the convention $p_0(g) = A_0(\log g^{-2})^{p_0}$ of [9, p. 246] and agree in the convention of [12, pp. 380–383] discussed below. The second member is the one with which the text goes on: the next sentence assumes that A_0 is much larger than A_1, the general step defines $\delta_k = g_k(A_1/A_0)p_0(g_k) = (A_1/A_0)\varepsilon_k$ [9, p. 265], and the estimates (3.7) and (3.9) of [9, p. 266] replace δ_k by $(A_1/A_0)\varepsilon_k$ and use that A_1/A_0 is small. We take $\delta_0 = (A_1/A_0)\varepsilon_0$, which is ϑ_0 of Definition 3.3(ii) and fixes the radius ϑ_0/g_0 of the fluctuation integral of the first step in Section 5.4.

Two exponents. In the displays of [12, pp. 380–383] the symbol $p_0(g)$ stands for $(\log g^{-2})^{p_0}$, without the factor A_0 of (3.1), in the same way as $p_1(g) = (\log g^{-2})^{p_1}$ [11, p. 183]. The constants are written out there: the factor of a component of a large-field region is $\exp(-\gamma_0 \min\{\frac{1}{2}B_3^{-2}A_0^2, 2A_1^2, A_1^2\}p_0^2(g_j)(d'_j + 1))$ [12, p. 381], in which A_0 comes from the plaquette thresholds ε_j and A_1 from the thresholds δ_j of the factors on p. 380, and the bound of $\exp(-\frac{1}{4}g_m^{-2}B_3^{-2}\frac{1}{2}\varepsilon_m^2)$ by $\exp(-\frac{1}{8}B_3^{-2}(1+\beta_0)^{-2}A_0^2 p_0^2(g_j))$ on p. 381 follows from $g_m^{-2}\varepsilon_m^2 = A_0^2(\log g_m^{-2})^{2p_0}$ and [9, (2.7), p. 255] in this reading only, since in the convention of (3.1) its right side would carry the factor A_0^4 in the exponent and its left side the factor A_0^2. Supplement, Proposition S.1, takes its conditions on A_1 and A_0 from these displays in this reading. The second exponent is printed r_0. It occurs twice, in the assumptions $\nu \geq 2p_0 + dr_0$ [12, p. 363] and $2p_1 - (d+5)r_0 > p_0$ [12, p. 383], and it is defined nowhere in the series. In both places it is the exponent of the cube sizes. On p. 383 the assumption serves to bound the factor $\exp(-R_j^{-d-5}p_1^2(g_j))$ by $\exp(-p_0(g_j))$, and since $(\log g_j^{-2})^r \leq R_j < L(\log g_j^{-2})^r$ by [9, (2.5), p. 255], this bound holds for small g_j when $2p_1 - (d+5)r > p_0$ and fails when $2p_1 - (d+5)r < p_0$; on p. 363 the factor R_k^{d-1} of the bound is to be absorbed by $L^{-(d-1)N} \leq (\log g_k^{-2})^{-(d-1)\nu}$. We read r_0 as r, in the condition on p_1 of Appendix A and in the condition on the depth of Section 3.4.

Iterated averages. The averaging operation of order k enters [8, (1.1), p. 260] as $\bar{U}^k = M^k(U)$, while the first paper of the series defines only the average M of order one [8, (0.4), p. 253]. The averages of higher order are defined in [4, (43), p. 24] recursively, the average of order $k+1$ being the average of order one of the average of order k, so that M^k is the k-fold iterate of M, and we use it in this sense.

Two averages of one step. The linearised averages in the operators of [5] are the linear parts of the average $\bar{\bar{U}}'$ of the fluctuation relative to the background [5, (3.13), p. 392; (3.113), p. 418], [4, (121)–(124), p. 36], while the constraints of the minimisers are those of the average M itself [7, (3), p. 278]. The two linear parts differ by a gauge transformation of the coarse sites, which does not vanish at a flat background, and the papers pass from one to the other without a remark, which is harmless for gauge-invariant quantities. Where both occur we distinguish them, as Q_1 and Q_1^{pl} in Section 9.3 and as Q^{dr} and Q in Section 13.7; the relation between them is Supplement, Proposition S.12(j).

The spaces of the regular terms. The first paper of the series proves its inductive description on spaces $U_j^c(X, \alpha_0, \alpha_1)$ with numbers α_0, α_1 which do not depend on the coupling

[8, Theorem 3, p. 264], and the paper on the complete model states the properties of the regular terms on the smaller spaces with the radii [9, (2.28), p. 259], which are proportional to the coupling up to logarithms. On the smaller spaces a second derivative at the trivial configuration is not bounded uniformly in the coupling by the bound of the term, and the coefficients of the flow are second derivatives. We use Theorem 3 of [8] for the part of the regular terms which does not involve the **R**-terms, along the couplings of a history, which need not satisfy the equation [8, (0.20), p. 256] of that paper; its proof treats one step at a time and uses of the couplings only that they are small (Supplement, Lemma 3.5).

Free choices. The papers centre the cubes of the families π_j on a lattice of spacing M [8, p. 257] and leave the position of the other partitions free; we centre all of them at the origin (Section 7.1), which is compatible with the blocks of the averages [8, (0.3), p. 252] because all sides are odd, and which makes every partition invariant under the reflections $x_\mu \mapsto -x_\mu$. Every object of a step which the papers define by reference to a coordinate system, the axial contours of the fluctuation fields, the gauge-fixing graphs, and the designated bonds of the averages, is attached to its block or cube through the centre and the fixed orientation of the axes, so that a translation which preserves the partitions maps the object of a cube to the object of its image. This is the rule under which Lemma 7.4(ii) holds; the reflections θ_μ of Section 12 map the objects of a cube to those of its image because the averages use the contours of all orders [8, p. 252], and the axial contours, which start at the centre of the block and change the coordinates in a fixed order [2, (1.7), p. 18], [8, p. 275], are mapped to the axial contours of the image block by a reflection of one coordinate, which does not change that order. A permutation of the axes changes that order, which is why the covariance of the normalisations is proved for the reflections and not for the exchange of two axes (Section 12.3).

Notation. We write ϑ_k for the threshold of the fluctuation field which Bałaban writes δ_k [9, p. 265], because δ_k is the difference of the couplings of the two runs in Sections 9–11. The constant of the boundary terms which [9, (2.47), p. 264] prints B_1 is our B_∂. The exponent of the age bound (3.31) is written $\beta < 1$ in [9, Theorem 2, p. 263], and its proof gives it as $1-\beta$ with an arbitrary $\beta > 0$, "with $1-\beta$, $\beta > 0$, instead of $\beta < 1$" [9, p. 283]; we write the exponent as $1-\beta_E$, with a small $\beta_E > 0$, because β is also Bałaban's constant of the spaces of the boundary terms. In [9, (2.5), p. 255] the letter r denotes both the exponent of the powers of L among which R_j is chosen and the fixed exponent of $\log g_j^{-2}$; we keep r for the second. Our couplings, thresholds, cube sizes, and depths are those of the tube of a target (Definition 3.11), evaluated at $\hat{g}_j$, where Bałaban's are evaluated at g_j; Proposition 3.12 states that his bounds hold for them.

Appendix A The Constants in the Order of Choices

The estimates of the paper contain many constants, and their consistency rests on the order in which they are chosen: a smallness condition can be met only if the quantity which it constrains is chosen after the constants on which the condition depends. We fix the constants in seven stages, numbered from 0 to 6, so that a constant of a stage depends only on the constants of the earlier stages. Every condition of the paper is then of one of two kinds, a lower bound on a quantity of its own stage in terms of earlier ones, or an upper bound on a quantity of a later stage in terms of earlier ones, and conditions of both kinds are met by taking each quantity large or small enough when its stage comes. The *constants of the construction* are those of the stages 0 to 5, together with the constants computed from them. The last stage chooses the bound γ of Section 3.1, and with it the coupling $\bar{g}$ and the target $(K, \bar{g})$; every requirement that $\bar{g}$ be sufficiently small which involves no representation is a

condition of this stage.

Stage 0*: the dimension, the group, and the exponents.* These numbers are fixed once and for all.

$d = 4$, $G \subset U(N)$	the dimension and the gauge group, with the numbers κ of (4.2), c of (4.3), and $\dim G$, the radius r_G of the logarithmic chart, the constant c_G of the Haar measure of balls (Section 8.3), and the constant c_h which compares the two norms of Appendix B
$r = 2$	the exponent of the cube sizes R_j, for which [9, p. 245] requires $r \geq 2$
p_0	the exponent of the thresholds, $p_0 \geq 5r$ [9, p. 246]
p_1	the exponent of the thresholds of the **R**-operation, $p_1 < p_0$ [11, p. 183], $p_0 - p_1 - 1 \geq \lceil r \rceil/2$ [11, (1.94), p. 198], and $2p_1 - (d+5)r > p_0$ [12, p. 383], where the exponent printed r_0 is read as r, the only exponent of the cube sizes in the construction
$q_0 = q_1$	the exponents of the radii of analyticity, integers greater than 1 [9, (2.28), p. 259], with $q_0 > p_0 + 3r$ and $q_0 > 25r + \frac{5}{2}$ (Section 13.8)
ν_L	$p_0 + q_1 + 7r + 1$, the exponent of the depth (Definition 3.9)
$\beta_0 = 1/7$	the constant of Bałaban's inequalities (2.6), which "can be chosen arbitrarily small, if g is sufficiently small" [9, p. 255]; it satisfies $\beta_0 \leq \sqrt{2} - 1$ (Section 3.1) and $(1 + \beta_0)^8 \leq 3 < L/4$ (Section 6.9)
$\beta = 1/8$	the constant of Bałaban's spaces of the boundary terms [9, (2.34)–(2.39), p. 261], a small positive number "but not too small" [9, p. 261]; with it $\kappa_s = \beta/30 = 1/240$ in (3.37)
ε_0	$0 < \varepsilon_0 < 1/125$, the loss in the exponents of the comparison of the two runs (Proposition 10.6)
β_H, β_E, β', θ_V	$\beta_H \geq 19/20$, $\beta_E \leq 1/20$, and β', θ_V of (11.8)
Hölder exponent	a number in $[\frac{3}{4}, 1)$, written β in Section 9 only, with γ_w of (9.21) and γ_H of (11.18)
Λ_*	a number $\Lambda_* \geq 2$, the bound on the chord–arc constants of the admissible loops (Definition 1.4)

Stage 1*: the blocking factor.* The integer L is odd and satisfies (C.1), which bounds it from below in terms of d, ε_0, and Λ_*. The constant $C_{\text{dep}} = 16L^4(L+1)^2$ of Lemma 3.10 is computed here.

Stage 2*: the constants computed from d, L, and G.* These are Bałaban's constants which depend on d and L only: the decay rate δ_0 and the constants B_0, $B_0(\beta)$ of the propagators [5, Theorem 3.1, (3.42), (3.43), p. 397], the constant c_0 of [3, (2.58)], and the constant B_3 of the minimisers [7, Theorem 1, p. 279]. With them come the constant C_W of Theorem 4.4, the numbers $b_* = c \ln L$ and $b^* = 3c \ln L$ of (4.4), the constant C_M of Theorem 10.8, which counts the ingredients of a term per point, and

$$C_P \coloneqq \max(2, 2C_M) \,, \tag{A.1}$$

the constant of Proposition 8.4; it satisfies the two requirements $C_P \geq 2$ and $C_P \geq 2C_M$ which Sections 8.3 and 11.7 place on it.

Stage 3*: the decay rates and the sizes.* First Bałaban's decay rate κ of the inductive bounds, large at fixed β [9, p. 260], and the exponent $\kappa_0 \geq 13$ of (3.29), which "can be chosen

arbitrarily large, similarly as κ, if the other parameters are fixed" [9, p. 260]; the comparison of two cutoffs uses two of its powers to pay the radius of the spaces of the tube (Section 11.4) and the others for the summability in K; then the rate κ_1 of the decoupling parameters of [10], which is bounded below in terms of κ and L. Then the sizes, each bounded below in terms of the earlier constants and the sizes before it, in the order: the size R_0M_0 of [5, p. 408], enlarged beyond Bałaban's smallest choice, which leaves his results valid because "in [4] we have proved all theorems under the assumption that R, M are sufficiently large" [5, p. 408]; the integer K and the block size KR_0M_0 of [5]; the size M_1 of the cubes for which the theorems of [5, 7, 6] hold, with $M_2 = L^2M_1$ [9, p. 245]; the separation R of the walk problems [3, (2.2)]; and the cube size $M = L^m$, which is "sufficiently large" [9, p. 245].

Stage 4*: the regularity parameters of the propagators.* The parameters α_0 and α_1 of the domains of [5, (3.35)–(3.37), p. 396], with $\alpha_J := KR_0M_0\alpha_0$ and the radius $\alpha'_* = \min(\alpha_1, \alpha_J/\hat{x}_J)$ of Supplement, Proposition S.12(g), bounded above in terms of the stages 0–3.

Stage 5*: the parameters of the variational problem.* The parameter ε_3 of [7, Proposition 4, p. 292], and then the data parameter ε_1 of [7, Theorem 1, p. 279] at which its theorems are applied, bounded above in terms of the stages 0–4. In Sections 10 and 11 the letter ε_1 denotes the data parameter (10.11) of a history, a function of $\bar{g}$ which lies below this fixed value once $\bar{g}$ is small.

Stage 6*: the coupling.* First Bałaban's large constants: A_1 and then A_0 of the thresholds [9, p. 246, p. 265], with A_1/A_0 small [9, p. 247], and C_0^α, C_1^α of the radii [9, (2.28), p. 259], each bounded below in terms of the stages 0–5. Then the bound γ, small in terms of everything before it: the conditions (3.7) and (3.38), $\gamma \le \gamma_1$ of Proposition 4.3, the conditions (T1) of Appendix C, and the requirements "$\bar{g}$ sufficiently small" of the statements which involve no representation, where $\bar{g} \le \gamma$ by (3.2). The requirements which involve the representations are upper bounds on the number γ_q of (3.46), and they are met by the choice of the level q of the ensemble, which comes after the coupling. The integer K and the tube are chosen last. The thresholds ε_j, ϑ_j, δ'_j, the radii $\alpha_{0,j}$, $\alpha_{1,j}$, the cube sizes R_j, and the depths N_k of Definition 3.3 and Definition 3.9 are functions of the tube.

The derived constants. Every other constant of the paper is computed from those above and is introduced where it first occurs. They fall into five groups, which differ in what they may depend on.

(a) The constants of Bałaban's theorems, E_0, E_1, R_1, the constant B_0 of the boundary terms, and B_∂ (Section 3.2), depend on the constants of the construction and not on the tube, the step, K, or the torus (Proposition 3.12). The constants $E_\pm$ of the ultraviolet stability (3.32) are not of this kind: they are independent of the lattice spacing and of the torus, and they depend on the tube and the step through the coupling g_k, as in Bałaban's statement. The constants C_Λ, C_3, C_{rem}, C_R, and γ_1 of Proposition 4.3 are of the same kind.

(b) The constants of the inductive class (Definition 5.2): the constants C_τ, c_τ, C_ρ, C_t of (5.6), the box constant $C_{\text{box}} = 2916$, the reserve ν of the first step [1], taken with $\nu \le \eta(\mathcal{C})$ of the proof of Theorem 5.4, the unit constant $u_\star := (2 \cdot 3^d + 2^{d+4} + 17)13^d\Lambda_*$ of the sizes (Supplement, Proposition S.10(viii)), the extent rate $\nu_e := \nu/(2C_{\text{box}}\Lambda_*)$ and the rate $\eta_w := 5\nu/(16\Lambda_*) = \frac{5}{8}\nu_e C_{\text{box}}$ of the tail weights (5.9), the cross rate $\nu_\times \le \nu_e$, the rate $c_{\text{att}} \le \nu_e/8$ of the attached and of the wrapping insertions, the rates $c'_* \le \nu_e/4$, κ_e, c_{th}, the constants C_{coll} of (6.87) and C'_w, c_G of (6.90), the constants C_ν, C_K and the rate c_δ of the carrying of the attached insertions (Supplement, Section 7), and the constants C_χ, C_{new}, C_{rel}, $C_{S'}$ of (6.49), and $C_{\text{cr}} \ge C_{S'}$. For ensembles with admissible loops they

depend on the constants of the construction, among them Λ_*, and on the representations through $c_{\mathcal{C}}$, as nondecreasing functions of it, and on nothing else (Definition 5.2(v)).

(c) The constants of the stopping scale, $C_{\mathcal{E}}$, C_∂, C_λ, C_{fs}, C_A, C_e, and C_{ent} of Section 8, which are of the kind (a), and the constant c_π of (1.7), which depends on the representation. The numbers $B_E(\mathcal{C})$ and $I_*(\mathcal{C})$ of Section 8.4 are not constants of a stage: they are bounds attached to an ensemble, which depend on $\bar{g}$ and not on K.

(d) The constants of the comparison of two cutoffs: $C_{\mathrm{df}}(\beta)$ and $C_\square(\beta)$ of Section 9.3, C_w of Theorem 9.5, $C_G(\varepsilon_0)$ and C_U of Proposition 10.6, C_f, C_{sh}, C_h of Section 10, E_1', E_V, C_α of Proposition 11.5, C_β of Lemma 11.6, and $C_{\mathcal{T}}$, C_s, C_S'', C_a of Section 11.5. They do not depend on K, the step, the lattice spacing, or the torus; some of them depend on $\bar{g}$ through the stopping index j_q and through the bound $\bar{\varepsilon}_1$ of (10.11), as stated where they are introduced.

(e) The rates of the sums over K: $\tilde{\theta}$ of (11.22), which depends on γ through ε^*, and ℓ_K, μ_K, w_K of (11.23).

Appendix B Conventions

Lattices and units. The lattice of the scale j of a run with K steps has the spacing $a_j = L^j\varepsilon$, with $\varepsilon = uL^{-K}$ the finest spacing, and a length "in units of the scale j" is a length divided by a_j. Distances are ℓ^∞-distances unless stated otherwise. The bonds and the plaquettes of a set are those which intersect it [8, p. 251].

Cubes and partitions. For a cube $\square$ of the family π_j, of side M, the cube $\tilde{\square}^n$ is the cube of side $(1+2n)M$ with the same centre [8, p. 257]. The operation $\sim$ applied to a set adds one layer of the cubes of the family in terms of which the set is defined [9, p. 245], a cube being added when it has a point in common with the set, and $X^{\sim -1} \coloneqq ((X^c)^\sim)^c$ [9, p. 247]. Partitions are compatible, every cube of a coarser partition being a union of cubes of a finer one [9, p. 245], and every partition consists of cubes centred at the points of $s\mathbb{Z}^4$, s its side (Section 7.1).

Norms. On matrices $\|\cdot\|$ is the operator norm (Section 8.2), which is invariant under multiplication by unitary matrices; on the insertions of the cross groups, which are tensors with one factor for each interval, it is the projective norm (5.5) built from it; and $|\cdot|$ is the norm in which Bałaban states the regularity of the fields; the constant $c_h \geq 1$ of Section 8.2 compares the two. The trace $\mathrm{tr} = N^{-1}\,\mathrm{Tr}$ is normalised.

Letters with several meanings. A few letters keep the meanings which they have in the papers that we cite, and the context separates them.

(i) β is Bałaban's constant $1/8$ of the spaces of the boundary terms in Sections 3–8 and in the coefficients $1-\beta(1-2^{-(k-m)})$ of the radii; the exponent $\beta < 1$ of the age bound (3.31), which is $1-\beta_E$, in Section 3 and in Lemma 11.4; and the Hölder exponent of stage 0 in Section 9.

(ii) M is the cube size L^m of [9, p. 245] and [8, p. 257], except in Section 9.2 and in the bounds (9.6), (9.20), where it is the block size KR_0M_0 of [5, p. 408]; K is the number of steps, except in KR_0M_0.

(iii) κ is the number of Lemma 4.2 in Section 4 and Bałaban's decay rate elsewhere.

(iv) B_0 is the constant of the boundary terms [9, (2.42)] in Sections 3 and 10, and the constant of the propagators [5, (3.42)] in Section 9.

(v) δ' without an index is the deviation of the coefficients in Theorem 4.1 and Proposition 4.3; δ'_j is the threshold of the **R**-operation.

(vi) r is the exponent of stage 0; $r(C)$ and $r_0(C)$ are the lengths of a loop of Section 2, r_k the radius of the fluctuation cutoff, and r_G the radius of the logarithmic chart.

Appendix C The Conditions and Their Sources

We list the conditions on the constants by the stage of Appendix A to which they belong, with the place where each is used and, where it is Bałaban's, where he imposes it. Each of them bounds a quantity of its stage from below, or a quantity of a later stage from above, in terms of the earlier stages; so all of them hold once the stages are chosen in order, and none of them involves the number of loops, their sizes, or K.

Stage 1. The blocking factor L is an odd integer, as in [8, p. 251], with

$$L \ge L_0(\Lambda_*)\,, \qquad L_0(\Lambda) \coloneqq \max(2^{52},\, 4\Lambda^2 L_0^\times)\,, \qquad L_0^\times \coloneqq 1.5\cdot 10^6(2\cdot 3^d + 2^{d+4} + 17)13^d\,, \tag{C.1}$$

so that $L_0^\times = 1.86360525\cdot 10^{13}$ at $d=4$. The two terms have different origins. The term $2^{52} = 4^{26}$ gives the inequality $L^{5(1-\varepsilon_0)/128} \ge 4$, because $128/(5(1-\varepsilon_0)) < 128\cdot 125/620 < 26$ for $\varepsilon_0 < 1/125$; this inequality is used in Supplement, Proposition S.14, for the variation of the terms of the effective action between the scales of a young history, where a factor 2^n from the radii of the analyticity domains has to be beaten by the decay $L^{-5(1-\varepsilon_0)n/128}$. The term $4\Lambda^2 L_0^\times$ is the second condition of (6.43), $L \ge 6\cdot 10^6 \bar\Lambda u_\star$, for a loop with the chord–arc constant Λ and with Λ in the place of Λ_* in the unit constant $u_\star$ of Appendix A, since then $6\cdot 10^6 \Lambda u_\star = 4\Lambda^2 L_0^\times$. The chord–arc constant enters twice because it enters the counts of the step, through the number of the traversals of a loop in a cube (Lemma 5.5), and the constant $u_\star$, which bounds the cubes and the bonds of a group per unit of its size: a group made by the **R**-operation occupies on its loop an interval which may be longer than the diameters of its polymers by the factor 4Λ (Supplement, Proposition S.10(viii)). This bound contains the lower bounds 5000Λ, $3(1+\Lambda)$, 20Λ, and 81Λ of the same counts. Since L_0 increases and every admissible loop has $\Lambda(C) \le \Lambda_*$, the condition (C.1) covers all of them at once. It covers as well the other lower bounds on L of the paper and of the supplement, all of which are below 2^{52}: $L > 11$ [8, p. 251], $L \ge 3$ (Lemma 3.10), $L \ge 5$ in (7.10), $(1+\beta_0)^8 \le L/4$ (Section 6.9), $L \ge 21$ (Lemma 8.2), $L \ge 195$ (Lemma 7.5), and $L \ge 12d+6$, $L^2 > 9d$, $L \ge 128(1+\beta_0)$ (Supplement, Propositions S.12–S.14).

Stage 3. The rate κ_1 satisfies $\frac12(\kappa_1 - 1) \ge 2L\kappa$ [10, p. 20]. The size R_0M_0 satisfies the two conditions of Lemma 9.4, $\frac16\delta_0 R_0 M_0 > 8\log c_0(\frac16) + 1$ and $4\delta_0 R_0 M_0 \ge 3(1+\beta)\log L$, with the Hölder exponent β, and the conditions of the same form $c\delta R_0 M_0 \ge C\log L + C'$, with a rate δ of stage 2 and numbers c, C, C' depending on d and on the Hölder exponent, of Supplement, Proposition S.12; Bałaban's results hold for every such enlargement of R_0M_0 [5, p. 408]. The integer K of [5, p. 408] is a power of L with $K \ge 4$, and the separation of the walk problems is $R \ge 12dL$. The cube sizes satisfy Bałaban's lower bounds $M \ge M(\kappa)$ [8, Theorem 3, p. 264], $\delta_0 M \ge \kappa$ [10, p. 21], $\delta_1 M \ge \kappa_1$ [10, p. 5], and $\delta(M/M_1) \ge 2$ [11, p. 186], and the lower bounds on M_1 and M of Supplement, Propositions S.12–S.14, each of which has the form $c\delta M_1 \ge C\log L + C'$ or $M \ge C(d,L)M_1$.

Stage 4. The parameters α_0 and α_1 are so small that "$O(1)M\alpha_0$ is still a sufficiently

small number" [5, p. 396], with M the block size KR_0M_0, and that the upper bounds of Supplement, Proposition S.12, hold; these involve the constants of the stages 0–3 only.

Stage 5. The parameters of the variational problem satisfy $\varepsilon_3 \le a_3$ [7, Proposition 4, p. 292] and $\varepsilon_1 \le a_1$, and every cube of the parameter M_V at which [7, Theorem 1] is applied has $M_V \le M(\varepsilon_1) = R_1M_1a_1/\varepsilon_1$ [7, p. 279], an upper bound on ε_1; the further upper bounds on ε_3 and ε_1 in terms of α_0, α_1, and the earlier constants are those of Supplement, Propositions S.12 and S.13.

Stage 6. The constants of the thresholds satisfy $A_1 \ge 48(15/8)^{1/2}$ and $A_0^2 \ge \frac{15}{8}A_1^2 \max(8(1+\beta_0)^2B_3^2, 4O(1)B_3M^2)$, with the constant $O(1)$ of [12, p. 381], which keep the comparisons of [12, pp. 381–382] valid when the coefficient of the action is reduced by the factor 15/8 (Supplement, Proposition S.1). The bound γ satisfies (3.7) and (3.38), $\gamma \le \gamma_1$ of Proposition 4.3, the bound $\delta' \le (2/5)c \ln L$ of Theorem 4.1, which holds for γ small because $\delta' \to 0$ with γ, and $C_W\gamma^2 \le 11/40$ (Supplement, Lemma 2.1). Moreover γ satisfies the following conditions, which we call (T1): for every $g \le (1+\beta_0)\gamma$,

$$\varepsilon^*(g)C_{\mathcal{T}} \le \frac{1}{4}\,, \qquad 4\hat{\varepsilon}^*(g)N'\Big(\frac{g}{1+\beta_0}\Big) \le 1\,, \qquad 4C_{\mathrm{ov}}\varepsilon^{*B}(g)N_0^+(g)W^\sharp \le 1\,. \tag{C.2}$$

Here $\varepsilon^*(g)$ is the coefficient ε_k^* of (11.19) as a function of the coupling, $C_{\mathcal{T}} \ge (1-L^{-\beta'})^{-1}$ the constant of Supplement, Proposition S.16(e), $\hat{\varepsilon}^*(g) := \sup_{g' \le 2g} \varepsilon^*(g')$, and $N'(g) := \log_L(MR(g)) + 3$, with $R(g)$ the smallest power of L which is at least $(\log g^{-2})^r$, the bound on the number of steps through which a recent **R**-term is carried without contraction. In the third condition $\varepsilon^{*B}(g)$ is the coefficient of the recursion of the boundary terms, a constant times $gp_0(g)$, $N_0^+(g) := 1 + 2(\log(32L(L+1)) + 2\log\log g^{-2})/\log L$ bounds the depth, $C_{\mathrm{ov}} = 2$ counts the generations of created boundary terms at a point, and $W^\sharp$ is a constant of L and d (Supplement, Proposition S.14). Each condition holds for γ small: $\varepsilon^*(g)$ decays as a negative power of $\log g^{-2}$, $\varepsilon^{*B}(g)$ as g times a power of it, while N' and N_0^+ grow as double logarithms; the running couplings of the histories which the conditions concern are at most $(1+\beta_0)\bar{g} \le (1+\beta_0)\gamma$ by (3.9). None of the conditions on γ involves a loop or a representation, so that the construction, and with it the step factors of the normalisation, is the same for all the ensembles.

Every statement of the paper which requires the coupling to be sufficiently small imposes one of two kinds of conditions (Section 1.3). If it involves no representation, it is an upper bound on $\bar{g} \le \gamma'$ in terms of the constants of the construction and of Λ_*, and, where it concerns a given ensemble, of the data of the ensemble which the statement names. If it involves the representations of an ensemble $\mathcal{C}$, it concerns the steps $k \le K'$ at which the loops are carried, and it is an upper bound on the number γ_q of (3.46) in terms of the constants of the construction, of Λ_*, and of $c_{\mathcal{C}}$: these are the conditions of Theorem 5.4, of Sections 6.9 and 6.15, of Lemma 7.3, Proposition 7.6, and Corollary 7.11, the condition (8.42), and the condition (11.26) on the coefficient $\varepsilon^Q(g)$ of the old vertices in the recursion of the insertions (Supplement, Proposition S.16(f)), together with those of Supplement, Propositions S.9 and S.10. They are not conditions on the constants of a stage: they are met by the choice of the level q in Theorem 1.5.

References

[1] , *A Wilson Loop after the First Renormalisation Step*, Manuscript, 2026, 2026.

[2] Tadeusz Bałaban, *Propagators and Renormalization Transformations for Lattice Gauge Theories. I*, Comm. Math. Phys. **95** (1984), 17–40.

[3] ———, *Propagators and Renormalization Transformations for Lattice Gauge Theories. II*, Comm. Math. Phys. **96** (1984), 223–250.

[4] ———, *Averaging Operations for Lattice Gauge Theories*, Comm. Math. Phys. **98** (1985), 17–51.

[5] ———, *Propagators for Lattice Gauge Theories in a Background Field*, Comm. Math. Phys. **99** (1985), 389–434.

[6] ———, *Spaces of Regular Gauge Field Configurations on a Lattice and Gauge Fixing Conditions*, Comm. Math. Phys. **99** (1985), 75–102.

[7] ———, *The Variational Problem and Background Fields in Renormalization Group Method for Lattice Gauge Theories*, Comm. Math. Phys. **102** (1985), 277–309.

[8] ———, *Renormalization Group Approach to Lattice Gauge Field Theories. I. Generation of Effective Actions in a Small Field Approximation and a Coupling Constant Renormalization in Four Dimensions*, Comm. Math. Phys. **109** (1987), 249–301.

[9] ———, *Convergent Renormalization Expansions for Lattice Gauge Theories*, Comm. Math. Phys. **119** (1988), 243–285.

[10] ———, *Renormalization Group Approach to Lattice Gauge Field Theories. II. Cluster Expansions*, Comm. Math. Phys. **116** (1988), 1–22.

[11] ———, *Large Field Renormalization. I. The Basic Step of the* **R** *Operation*, Comm. Math. Phys. **122** (1989), 175–202.

[12] ———, *Large Field Renormalization. II. Localization, Exponentiation, and Bounds for the* **R** *Operation*, Comm. Math. Phys. **122** (1989), 355–392.

[13] Edward Bierstone and Pierre D. Milman, *Semianalytic and Subanalytic Sets*, Inst. Hautes Études Sci. Publ. Math. **67** (1988), 5–42.

[14] Richard A. Brandt, Filippo Neri, and Masa-aki Sato, *Renormalization of Loop Functions for All Loops*, Phys. Rev. D **24** (1981), 879–902.

[15] Sacha Friedli and Yvan Velenik, *Statistical Mechanics of Lattice Systems: A Concrete Mathematical Introduction*, Cambridge University Press, Cambridge, 2017.

[16] Gregory F. Lawler and Vlada Limic, *Random Walk: A Modern Introduction*, Cambridge Studies in Advanced Mathematics, vol. 123, Cambridge University Press, Cambridge, 2010.

[17] Konrad Osterwalder and Erhard Seiler, *Gauge Field Theories on a Lattice*, Ann. Physics **110** (1978), 440–471.

[18] Alexander M. Polyakov, *Gauge Fields as Rings of Glue*, Nuclear Phys. B **164** (1980), 171–188.