Subsequential Infinite-Volume Limits of the Wilson Loop Functions

Abstract

The limits of the renormalised Wilson loop expectations constructed in the paper live on a four-dimensional torus, and the bound on which their existence rests grows exponentially with the volume of the torus. We show that a chessboard estimate, which uses nothing but the reflection positivity of the lattice measures, replaces this bound by one which does not depend on the period of the torus. As a consequence the limits converge along a subsequence of periods, for every ensemble of admissible loops in $\mathbb{R}^4$ at once, and the limiting functions are bounded, reflection positive, and covariant under the symmetries which the paper establishes on the torus. Their average over the permutations of the axes is, in addition, covariant under the whole hypercubic group, and it remains bounded and reflection positive.

1 The Result

Everything in the paper takes place on the torus $(\mathbb{R}/T\mathbb{Z})^4$ of the period $T = 2L^{m_T}u$, and we ask what happens as $m_T \to \infty$. Most of the construction does not see the period. The bare coupling $g_0(K, \bar{g})$ of Paper, Theorem 4.1, is determined by the coefficients β_{k+1} of Paper, (4.6), which are defined on the infinite lattice [1, (1.20)–(1.22), p. 264], so that the map from the bare coupling to the running couplings, and with it the bare coupling for a target $(K, \bar{g})$, may be chosen the same for all periods; we make this choice. The step factors, and therefore the normalisations, depend neither on the loop nor on the torus (Paper, Proposition 7.13), and the threshold γ_0 and the function $q_\sharp$ of Paper, Theorem 1.5, depend only on the constants of the construction and on Λ_*. Of the three quantities which enter the uniform bound of Paper, Theorem 8.6, the sum $B_E(\mathcal{C})$ of the remainders does not involve the torus, and the insertion total $I_*(\mathcal{C})$ decreases as the torus grows. The third one grows, however: the bound $\bar{\delta}_{\mathrm{LF}}$ on the weight of the large-field terms (Paper, Proposition 8.4) contains the factor e^{2C_K} with $C_K = C_{\mathrm{fs}}N_q^4(1 + \log g_K^{-1})$, where N_q is the number of sites per side of the lattice at the stopping scale (Paper, (8.2)). This factor is the price of comparing whole integrals over that lattice, and it makes the bound of Paper, Theorem 8.6, exponential in the volume.

Reflection positivity turns a cost which is extensive in the volume into a cost per block. If the torus is cut into $|\Lambda|$ congruent blocks and an ensemble is placed in one of them, the chessboard estimate bounds the $|\Lambda|$-th power of its expectation by the expectation of the ensemble obtained by reflecting it into every block. The uniform bound of the paper, applied to this larger ensemble, contains the factor e^{2C_K} only once, so that its $|\Lambda|$-th root is a number of the block alone, while the costs of the loops, which are sums over the loops, are divided back to those of one copy. The chessboard estimate is due to Fröhlich, Israel, Lieb, and Simon [2]; we prove the form which we need, from the reflection positivity of the lattice measures in Paper, Lemma 12.2, in Section 2.

Throughout, L, u, Λ_*, γ_0, and $q_\sharp$ are as in Paper, Theorem 1.5, and $\bar g \in (0,\gamma_0]$ is fixed. For an integer $s \geq 0$ we write $T_s := 2L^s u$ and $\mathbb{T}^{(s)} := (\mathbb{R}/T_s\mathbb{Z})^4$, which is the torus of the paper with $m_T = s$, and a superscript (s) marks the objects of the paper on it, such as $\mathbb{E}_K^{(s)}$, $W_K^{(s)}$, and $W^{(s)}$. The limit in the volume concerns loops in $\mathbb{R}^4$, which we now define.

Definition 1.1 (Ensembles in $\mathbb{R}^4$)**.** Let $q \geq 0$ be an integer. A *loop at level q in $\mathbb{R}^4$* is a closed path of bonds of the lattice $uL^{-q}\mathbb{Z}^4$ which visits no site twice, with its orientation, and a *loop ensemble at level q in $\mathbb{R}^4$* is a finite family $\mathcal{C} = ((C_1,\chi_1),\dots,(C_n,\chi_n))$ of pairwise vertex-disjoint loops at level q in $\mathbb{R}^4$, each with the character χ_i of an irreducible unitary representation of G. The *extent* $E(\mathcal{C})$ is the largest of the four numbers $\max x_\mu - \min x_\mu$, the maximum and the minimum being taken over the sites x of the loops of $\mathcal{C}$. We put $b_q(\mathcal{C}) := uL^{n_q(\mathcal{C})}$, where $n_q(\mathcal{C})$ is the least integer $n \geq -q$ with $uL^n \geq E(\mathcal{C}) + 2uL^{-q}$, and $s_q(\mathcal{C}) := \max(0, n_q(\mathcal{C}) + 1)$. Finally $\mathfrak{E}$ denotes the set of the pairs $(\mathcal{C}, q)$ of an integer $q \geq 0$ and a loop ensemble $\mathcal{C}$ at level q in $\mathbb{R}^4$ whose loops are admissible in the sense of Paper, Definition 1.4, and for which $q \geq q_\sharp(c_{\mathcal{C}}, \bar g)$.

Let $(\mathcal{C}, q) \in \mathfrak{E}$ and $s \geq s_q(\mathcal{C})$, and let $\pi_s \colon \mathbb{R}^4 \to \mathbb{T}^{(s)}$ be the covering map. The sites of $\mathcal{C}$ lie in a closed box of side $E(\mathcal{C}) < b_q(\mathcal{C})$, and π_s is injective on every box of side less than $T_s = 2uL^s \geq 2Lb_q(\mathcal{C})$. The family $\pi_s\mathcal{C}$ of the images, with the same characters, is therefore a loop ensemble at level q on $\mathbb{T}^{(s)}$ in the sense of Paper, Definition 1.1, the condition $3\,\mathrm{diam}(\pi_s C_i) < T_s$ holding because $\mathrm{diam}(C_i) \leq E(\mathcal{C}) < b_q(\mathcal{C})$; and its loops have the chord–arc constants, the ratios P/r, and the numbers P_q of the loops of $\mathcal{C}$, since these are computed in a lift. We write $W^{(s)}(\mathcal{C})$ for the limit of Paper, Theorem 1.5, of the ensemble $\pi_s\mathcal{C}$ with the normalisation of the level q.

Two numbers enter the bound. The first is the number $B_E(\mathcal{C})$ of Paper, (7.32), which is defined without reference to the torus and bounds the sum of the remainders on every torus. The second replaces the insertion total of Paper, (8.37), which involves the torus only through its last factor $S_{\mathrm{wr}}(c_{\mathrm{att}}N_q/M)$; since $S_{\mathrm{wr}}(b) = \sum_{i\geq 0} L^i e^{-bL^i}$ decreases in b and $N_q = 2L^{m_T+j_q} \geq 2$, the insertion total is at most

$$I_*^\flat(\mathcal{C}) := P_{K'}(\mathcal{C})\Big[\frac{2c_\sigma C_\chi(1+2\kappa_s)B_3 S_w\,\bar\rho\,\varepsilon_{K'}}{\min(\alpha_{0,K'},\alpha_{1,K'})} + C^2\bar\rho\,S_{\mathrm{wr}}\Big(\frac{2c_{\mathrm{att}}}{M}\Big)\Big]\,, \tag{1.1}$$

with the constants of Paper, (8.37). For a length $b > 0$ we put, with the constants of Paper, Proposition 8.4, and with $\hat g_{K'}^{-2} = \bar g^{-2} + b' j_q + C_W$ (Paper, (3.3)),

$$\Gamma_q(b,\bar g) := \Big(\frac{bL^{j_q}}{u}\Big)^4\Big[2C_{\mathrm{fs}}\Big(1+\frac{1}{2}\log\hat g_{K'}^{-2}\Big) + \frac{2}{(MR_{K'})^4}\exp(-c_{\mathrm{LF}}\,p_0((1+\beta_0)\bar g))\Big]\,. \tag{1.2}$$

The numbers $P_{K'}(\mathcal{C}) = P_q(\mathcal{C})L^3MR_{K'-1}$, $\varepsilon_{K'}$, $\alpha_{0,K'}$, $\alpha_{1,K'}$, $R_{K'}$, and $\hat g_{K'}$ are functions of q, $\bar g$, and the constants of the construction (Paper, Section 8), and the constants in (1.1) and (1.2) do not depend on the torus (Paper, Definition 5.1, uniformity, and (8.29)). Hence neither $I_*^\flat(\mathcal{C})$ nor $\Gamma_q(b,\bar g)$ depends on the period or on K; the first is a sum over the loops of $\mathcal{C}$, and so is $B_E(\mathcal{C})$.

The theorem says that the limits on the tori converge along one subsequence of periods for all ensembles at once, and that the limits keep the bound and the structure which the paper proves on each torus.

Theorem 1.2 (Subsequential Infinite-Volume Limits)**.** *There is a strictly increasing sequence $(s_k)_{k\geq 1}$ of integers such that, for every $(\mathcal{C}, q) \in \mathfrak{E}$, the limit*

$$W^\infty(\mathcal{C}) := \lim_{k\to\infty} W^{(s_k)}(\mathcal{C}) \tag{1.3}$$

exists, the level being that of the pair. For every level q and all ensembles $\mathcal{C}$, $g\mathcal{C}$, $h\mathcal{C}$, $\mathcal{D}_a$ below which form pairs of $\mathfrak{E}$ with this level, the limits have the following properties.

(i) $|W^\infty(\mathcal{C})| \le \exp(B_E(\mathcal{C}) + I^\flat_*(\mathcal{C}) + \Gamma_q(b_q(\mathcal{C}), \bar g))$.

(ii) $W^\infty(\mathcal{C})$ *does not depend on the order of the loops of* $\mathcal{C}$; $W^\infty(g\mathcal{C}) = W^\infty(\mathcal{C})$ *for every map* $g(x) = Ox + v$ *of* $\mathbb{R}^4$ *with a diagonal matrix* O *with entries* ± 1 *and* $v \in uL^{-q}\mathbb{Z}^4$; $W^\infty(\bar{\mathcal{C}}) = \overline{W^\infty(\mathcal{C})}$, *and* $W^\infty(\mathcal{C})$ *does not change if a loop* C_i *is replaced by* C_i^{-1} *and its character by* $\bar\chi_i$; *and* $W^\infty(h\mathcal{C}) = W^\infty(\mathcal{C})$ *for every map* $h(x) = Ox + v$ *with a signed permutation matrix* O *and* $v \in uL^{-q}\mathbb{Z}^4$ *for which there are maps* $g_1, \dots, g_n$ *of the previous kind with* $hC_i = g_i C_i$ *for every* i.

(iii) Let $\mu \in \{0, \dots, 3\}$ *and* $y \in uL^{-q}\mathbb{Z}$, *let* θ *be the reflection of* $\mathbb{R}^4$ *which replaces* x_μ *by* $2y - x_\mu$, *and let* $\mathcal{D}_1, \dots, \mathcal{D}_m$ *be loop ensembles at level* q *in* $\mathbb{R}^4$, *possibly empty, whose loops are admissible and lie in* $\{x_\mu > y\}$, *with* $q \ge q_\sharp(c, \bar g)$ *for the largest number* c *of the form* c_π *among their representations. Then, with* $\mathcal{D}_0 := \emptyset$ *and* $W^\infty(\emptyset) := 1$, *the matrices*

$$(W^\infty(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b))_{1\le a,b\le m} \qquad \text{and} \qquad (W^\infty(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b) - W^\infty(\mathcal{D}_a)\overline{W^\infty(\mathcal{D}_b)})_{1\le a,b\le m} \tag{1.4}$$

are Hermitian and positive semidefinite, and $W^\infty(\theta\bar{\mathcal{D}}_a) = \overline{W^\infty(\mathcal{D}_a)}$ *for every* a.

The bound (i) is the one which the compactness argument needs, a number of the ensemble which does not depend on the period. It is not a product over the loops, since it contains the fourth power of the extent of the whole ensemble through $b_q(\mathcal{C})$, and Remark 4.1 says what the theorem leaves open.

The covariance in (ii) under a map which exchanges two axes is restricted, although the lattice measures are invariant under every such map (Paper, Lemma 12.1): the normalisations are not known to be invariant under the exchange of two axes (Paper, Section 12.3). The full hypercubic group is recovered by averaging over the permutations of the axes. For a permutation σ of $\{0, \dots, 3\}$ let P_σ be the linear map of $\mathbb{R}^4$ with $(P_\sigma x)_{\sigma(\mu)} = x_\mu$, so that $P_\sigma e_\mu = e_{\sigma(\mu)}$ and $P_\sigma P_\tau = P_{\sigma\tau}$. It maps $uL^{-q}\mathbb{Z}^4$ onto itself and preserves the distance $|\cdot|_\infty$, the segments, and the corners, so that $P_\sigma\mathcal{C}$, the family of the images with the same characters, has the chord–arc constants, ratios P/r, numbers P_q, extent, and number c of $\mathcal{C}$, and $(P_\sigma\mathcal{C}, q) \in \mathfrak{E}$ whenever $(\mathcal{C}, q) \in \mathfrak{E}$. We put

$$W^{\mathrm{hc}}(\mathcal{C}) := \frac{1}{24}\sum_\sigma W^\infty(P_\sigma\mathcal{C})\,. \tag{1.5}$$

On each torus $\mathbb{T}^{(s)}$ the map with the formula of P_σ is a symmetry of level q, so that $\mathbb{E}^{(s)}_K[W_{\pi_s P_\sigma \mathcal{C}}] = \mathbb{E}^{(s)}_K[W_{\pi_s\mathcal{C}}]$ by Paper, Lemma 12.1. Hence $W^{\mathrm{hc}}(\mathcal{C})$ is the limit, as $K \to \infty$ and then along (s_k), of $\mathbb{E}^{(s)}_K[W_{\pi_s\mathcal{C}}]$ multiplied by $\frac{1}{24}\sum_\sigma \prod_i \kappa^{(q)}_K(P_\sigma C_i, \chi_i)^{-1}$: the expectation divided by a normalisation which is invariant under the whole hypercubic group, but which is an average of products over the loops rather than a product. If the normalisations are invariant under the exchanges of axes, then $W^{\mathrm{hc}} = W^\infty$.

The translations in Theorem 1.2(ii) are restricted to the vectors of $uL^{-q}\mathbb{Z}^4$ only because the loops are paths in this lattice, whereas neither the tori nor the measures on them single out an origin, and we remove the restriction by a convention. A *translate* of a loop ensemble $\mathcal{C}$ at level q in $\mathbb{R}^4$ is the family $\mathcal{C} + w$ of its loops shifted by one vector $w \in \mathbb{R}^4$, with the same characters, and for $(\mathcal{C}, q) \in \mathfrak{E}$ and $w \in \mathbb{R}^4$ we put

$$W^\infty(\mathcal{C} + w) := W^\infty(\mathcal{C})\,, \qquad W^{\mathrm{hc}}(\mathcal{C} + w) := W^{\mathrm{hc}}(\mathcal{C})\,, \tag{1.6}$$

the level being q. These numbers depend only on the family $\mathcal{C}+w$ and on q. Indeed, let $\mathcal{C}+w=\mathcal{C}'+w'$ with a second loop ensemble $\mathcal{C}'$ at level q, so that $\mathcal{C}'=\mathcal{C}+t$ with $t:=w-w'$. Since a loop is a closed path without self-intersections, it is not contained in a line and has a corner; the corners of $\mathcal{C}$ and of $\mathcal{C}'$ are sites of $uL^{-q}\mathbb{Z}^4$, and the translation by t maps the former to the latter, so that $t\in uL^{-q}\mathbb{Z}^4$. Theorem 1.2(ii) therefore gives $W^\infty(\mathcal{C}')=W^\infty(\mathcal{C})$ and, applied to $P_\sigma\mathcal{C}'=P_\sigma\mathcal{C}+P_\sigma t$ for every σ, $W^{\mathrm{hc}}(\mathcal{C}')=W^{\mathrm{hc}}(\mathcal{C})$. The convention is the result of carrying out the construction on the lattices $w+uL^{-K}\mathbb{Z}^4$ in place of $uL^{-K}\mathbb{Z}^4$, since the translation by w is an isometry of every torus $\mathbb{T}^{(s)}$ which carries the lattices, the measures, and the loops of the one construction to those of the other.

Corollary 1.3 (Hypercubic Covariance)**.** *For every level q and all ensembles $\mathcal{C}$, $\mathcal{D}_a$ below which form pairs of $\mathfrak{E}$ with this level, the following hold.*

(i) $|W^{\mathrm{hc}}(\mathcal{C})|\le\exp(B_E(\mathcal{C})+I^{\flat}_*(\mathcal{C})+\Gamma_q(b_q(\mathcal{C}),\bar g))$.

(ii) $W^{\mathrm{hc}}(\mathcal{C})$ *does not depend on the order of the loops of* $\mathcal{C}$; $W^{\mathrm{hc}}(h\mathcal{C}')=W^{\mathrm{hc}}(\mathcal{C}')$ *for every translate* $\mathcal{C}'$ *of* $\mathcal{C}$ *and every map* $h(x)=Ox+v$ *of* $\mathbb{R}^4$ *with a signed permutation matrix* O *and* $v\in\mathbb{R}^4$*, with the convention* (1.6)*;* $W^{\mathrm{hc}}(\bar{\mathcal{C}})=\overline{W^{\mathrm{hc}}(\mathcal{C})}$*; and* $W^{\mathrm{hc}}(\mathcal{C})$ *does not change if a loop* C_i *is replaced by* C_i^{-1} *and its character by* $\bar\chi_i$.

(iii) With μ, y, θ, and $\mathcal{D}_1,\dots,\mathcal{D}_m$ as in Theorem 1.2(iii), the two matrices (1.4) *with* W^{hc} *in place of* W^∞ *are Hermitian and positive semidefinite, and* $W^{\mathrm{hc}}(\theta\bar{\mathcal{D}}_a)=\overline{W^{\mathrm{hc}}(\mathcal{D}_a)}$ *for every a.*

With the convention (1.6) the reflection in (iii) may be taken in any hyperplane $\{x_\mu=y\}$, $y\in\mathbb{R}$, for the translates $\mathcal{D}_a+w$ by one vector w with $y-w_\mu\in uL^{-q}\mathbb{Z}$: writing θ_0 for the reflection which replaces x_μ by $2(y-w_\mu)-x_\mu$, we have $\theta(x+w)=\theta_0x+w$, so that $(\mathcal{D}_a+w)\cup\theta(\bar{\mathcal{D}}_b+w)=(\mathcal{D}_a\cup\theta_0\bar{\mathcal{D}}_b)+w$, and the matrices of the translated family are those of the family $\mathcal{D}_1,\dots,\mathcal{D}_m$ for the reflection θ_0.

2 The Chessboard Estimate

The estimate rests on one Cauchy–Schwarz inequality for each reflection, and on an argument which shows that these inequalities, applied repeatedly, spread one function over all the blocks. We first set up the blocks and the maps which move a function from one block to another.

Let $s\ge0$, $q\ge0$, and $K\ge q$ be integers, let n be an integer with $-q\le n\le s$, and put $b:=uL^n$ and $m:=L^{s-n}$, so that $T_s=2mb$ and $b\in uL^{-q}\mathbb{Z}$. Let $\Lambda:=(\mathbb{Z}/2m\mathbb{Z})^4$. For $j\in\mathbb{Z}/2m\mathbb{Z}$ and $y\in\mathbb{R}/T_s\mathbb{Z}$ put $\psi_j(y):=y+jb$ if j is even and $\psi_j(y):=(j+1)b-y$ if j is odd; this is well defined, because $T_s=2mb$ and the parity of j is defined modulo $2m$. For $t\in\Lambda$ let $\varphi_t(x):=(\psi_{t_0}(x_0),\dots,\psi_{t_3}(x_3))$ and $|t|:=t_0+\dots+t_3$ modulo 2. The map φ_t is the composition of a translation by a vector of $b\mathbb{Z}^4\subset uL^{-q}\mathbb{Z}^4$ with reflections θ_μ, so it belongs to the group $\mathfrak{G}_q$ of Paper, Section 1.3, and it maps the open block $B_0^\circ:=(0,b)^4$ onto the open block $B_t^\circ:=\prod_\mu(t_\mu b,(t_\mu+1)b)$.

For $\mu\in\{0,\dots,3\}$ and $k\in\mathbb{Z}$ the reflection $\theta_{\mu,kb}$ is a reflection of level q, since $kb\in uL^{-q}\mathbb{Z}$. For $t\in\Lambda$ let $\hat\theta_k t$ be obtained from t by replacing t_μ by $2k-1-t_\mu$. Then

$$\theta_{\mu,kb}\circ\varphi_t=\varphi_{\hat\theta_k t}\,,\qquad |\hat\theta_k t|\ne|t|\,. \tag{2.1}$$

Indeed, only the coordinate μ changes, and $\theta_{\mu,kb}$ replaces y by $2kb-y$: for even $j=t_\mu$ we get $2kb-jb-y=(2k-j)b-y=\psi_{2k-1-j}(y)$, since $2k-1-j$ is odd, and for odd j we get $2kb-(j+1)b+y=\psi_{2k-1-j}(y)$, since $2k-1-j$ is even.

The maps act on functions of the gauge field on $\mathbb{T}_K^{(s)}$ as in Paper, Section 12.1: $(F\circ h)(U)=F(hU)$ with $(hU)(b)=U(hb)$. Since $(h(h'U))(b)=(h'U)(hb)=U(h'hb)$, we have $h(h'U)=(h'h)U$ and

$$(F\circ h)\circ h'=F\circ(h'h)\,, \tag{2.2}$$

and if F depends only on the values of U on the bonds with both ends in a set X of sites, then $F\circ h$ depends only on those with both ends in hX. For a bounded measurable function F and $t\in\Lambda$ we put $\vartheta_tF:=F\circ\varphi_t$ if $|t|$ is even and $\vartheta_tF:=\overline{F\circ\varphi_t}$ if $|t|$ is odd. Since $(\Theta_{\mu,kb}F)(U)=\overline{F(\theta_{\mu,kb}U)}$, so that $\Theta_{\mu,kb}F=\overline{F\circ\theta_{\mu,kb}}$, the rules (2.2) and (2.1) give

$$\Theta_{\mu,kb}\vartheta_tF=\vartheta_{\hat\theta_k t}F\ : \tag{2.3}$$

for even $|t|$ the left side is $\overline{F\circ(\theta_{\mu,kb}\varphi_t)}=\overline{F\circ\varphi_{\hat\theta_k t}}$, with $|\hat\theta_k t|$ odd, and for odd $|t|$ it is $F\circ\varphi_{\hat\theta_k t}$, with $|\hat\theta_k t|$ even.

We first spread a function over the slabs of one direction. For $\mu\in\{0,\dots,3\}$ and $j\in\mathbb{Z}/2m\mathbb{Z}$ we write $\vartheta_j^\mu:=\vartheta_{je_\mu}$, which maps the open slab $\{0<x_\mu<b\}$ onto the open slab $\{jb<x_\mu<(j+1)b\}$.

Lemma 2.1 (One Direction)**.** *Let $\mu\in\{0,\dots,3\}$, and let G be a bounded measurable function of the gauge field on $\mathbb{T}_K^{(s)}$ which depends only on the values of U on the bonds with both ends in the open slab $\{0<x_\mu<b\}$. Then*

$$\mathbb{E}_K^{(s)}\Big[\prod_{j\in\mathbb{Z}/2m\mathbb{Z}}\vartheta_j^\mu G\Big]\ge 0\qquad\textit{and}\qquad |\mathbb{E}_K^{(s)}[G]|^{2m}\le\mathbb{E}_K^{(s)}\Big[\prod_{j\in\mathbb{Z}/2m\mathbb{Z}}\vartheta_j^\mu G\Big]\,. \tag{2.4}$$

Proof. For $S\subset\mathbb{Z}/2m\mathbb{Z}$ put $Z(S):=\mathbb{E}_K^{(s)}[\prod_{j\in S}\vartheta_j^\mu G]$, so that $Z(\emptyset)=1$ and $Z(\{0\})=\mathbb{E}_K^{(s)}[G]$. For $k\in\mathbb{Z}/2m\mathbb{Z}$ let $H_k^+:=\{k,k+1,\dots,k+m-1\}$, let H_k^- be its complement, and let θ_k also denote the involution $j\mapsto 2k-1-j$ of $\mathbb{Z}/2m\mathbb{Z}$, which maps H_k^+ onto H_k^-.

We first show that, for every S and every k,

$$|Z(S)|^2\le Z(S_k^+)\,Z(S_k^-)\,,\qquad S_k^\pm:=(S\cap H_k^\pm)\cup\theta_k(S\cap H_k^\pm)\,. \tag{2.5}$$

The open half-torus $H_{\mu,kb}=\{kb<x_\mu<kb+mb\}$ of the reflection $\theta_{\mu,kb}$ contains the slabs $\{jb<x_\mu<(j+1)b\}$ with $j\in H_k^+$, because $T_s/2=mb$. The functions

$$A:=\prod_{j\in S\cap H_k^+}\vartheta_j^\mu G\,,\qquad B:=\prod_{j\in\theta_k(S\cap H_k^-)}\vartheta_j^\mu G \tag{2.6}$$

therefore depend only on the values of U on bonds with both ends in $H_{\mu,kb}$, which belong to the set E_+ of Paper, Section 12.1, so that A and B lie in $\mathcal{A}_+$. Since $\Theta_{\mu,kb}$ is multiplicative, (2.3) gives $\Theta_{\mu,kb}B=\prod_{j\in S\cap H_k^-}\vartheta_j^\mu G$ and $\Theta_{\mu,kb}A=\prod_{j\in\theta_k(S\cap H_k^+)}\vartheta_j^\mu G$, so that

$$Z(S)=\mathbb{E}_K^{(s)}[A\,\Theta_{\mu,kb}B]\,,\qquad Z(S_k^+)=\mathbb{E}_K^{(s)}[A\,\Theta_{\mu,kb}A]\,,\qquad Z(S_k^-)=\mathbb{E}_K^{(s)}[B\,\Theta_{\mu,kb}B]\,. \tag{2.7}$$

By Paper, Lemma 12.2, $\langle F,F'\rangle:=\mathbb{E}_K^{(s)}[F\,\Theta_{\mu,kb}F']$ is a positive semidefinite sesquilinear form on $\mathcal{A}_+$, and (2.5) is its Cauchy–Schwarz inequality $|\langle A,B\rangle|^2\le\langle A,A\rangle\langle B,B\rangle$. In particular

$Z(S_k^+) = \langle A, A\rangle \geq 0$ for every S, and for $S = \mathbb{Z}/2m\mathbb{Z}$, for which $S_k^+ = \mathbb{Z}/2m\mathbb{Z}$, this is the first assertion of (2.4).

For the second assertion we fix $\epsilon > 0$, put $\zeta := Z(\mathbb{Z}/2m\mathbb{Z}) + \epsilon > 0$, and let $\rho(S) := |Z(S)|\zeta^{-|S|/(2m)}$. Since $|S_k^+| + |S_k^-| = 2|S|$, the inequality (2.5) gives

$$\rho(S)^2 \leq \rho(S_k^+)\,\rho(S_k^-) \tag{2.8}$$

for every S and every k. Let ρ_* be the largest of the finitely many numbers $\rho(S)$, so that $\rho_* \geq \rho(\emptyset) = 1$. If $\rho(S) = \rho_*$, then (2.8) and $\rho(S_k^\pm) \leq \rho_*$ force $\rho(S_k^+) = \rho(S_k^-) = \rho_*$ for every k. Among the sets with $\rho(S) = \rho_*$ we choose one, S, of the largest cardinality. Then $|S_k^\pm| \leq |S|$ for every k, and since the two cardinalities add up to $2|S|$, both are equal to $|S|$, that is, $|S \cap H_k^+| = |S|/2$ for every k. Comparing this for k and $k+1$ gives $\mathbf{1}_S(k+m) = \mathbf{1}_S(k)$, so that S is invariant under the shift $j \mapsto j+m$, which maps H_k^+ onto H_k^-. The set S_k^+ maximises ρ and has $|S|$ elements as well, so it is invariant under the same shift; since $S_k^+ \cap H_k^+ = S \cap H_k^+$ and $S_k^+ \cap H_k^- = \theta_k(S \cap H_k^+)$, this means that $(S \cap H_k^+) + m = \theta_k(S \cap H_k^+)$, while the invariance of S gives $(S \cap H_k^+) + m = S \cap H_k^-$. Hence $\theta_k(S \cap H_k^+) = S \cap H_k^-$, that is, $\theta_k S = S$, for every k. The involutions θ_k generate a group which acts transitively on $\mathbb{Z}/2m\mathbb{Z}$, because $\theta_{k+1}\theta_k$ is the shift by 2 and θ_0 maps 0 to -1; so S is either empty or all of $\mathbb{Z}/2m\mathbb{Z}$, and $\rho_* = \rho(S)$ is either 1 or $Z(\mathbb{Z}/2m\mathbb{Z})/\zeta < 1$. Thus $\rho(\{0\}) \leq 1$, that is, $|\mathbb{E}_K^{(s)}[G]| \leq (Z(\mathbb{Z}/2m\mathbb{Z}) + \epsilon)^{1/(2m)}$, and letting $\epsilon \to 0$ gives the second assertion. *q.e.d.*

Applied in the four directions in turn, the lemma spreads a function of one block over all the blocks.

Lemma 2.2 (Chessboard Estimate)**.** *Let F be a bounded measurable function of the gauge field on $\mathbb{T}_K^{(s)}$ which depends only on the values of U on the bonds with both ends in B_0°. Then*

$$\mathbb{E}_K^{(s)}\Big[\prod_{t\in\Lambda} \vartheta_t F\Big] \geq 0 \qquad \textit{and} \qquad |\mathbb{E}_K^{(s)}[F]|^{|\Lambda|} \leq \mathbb{E}_K^{(s)}\Big[\prod_{t\in\Lambda} \vartheta_t F\Big]\,. \tag{2.9}$$

Proof. Put $G_0 := F$ and $G_{\nu+1} := \prod_{j\in\mathbb{Z}/2m\mathbb{Z}} \vartheta_j^\nu G_\nu$ for $\nu = 0, 1, 2, 3$, and let Λ_ν be the set of the $t \in \Lambda$ with $t_{\nu'} = 0$ for every $\nu' \geq \nu$. We claim that $G_\nu = \prod_{t\in\Lambda_\nu} \vartheta_t F$. This holds for $\nu = 0$. If it holds for ν, then, since ϑ_j^ν is multiplicative and commutes with complex conjugation, it suffices to see that $\vartheta_j^\nu \vartheta_t F = \vartheta_{t+je_\nu} F$ for $t \in \Lambda_\nu$; and indeed $(F \circ \varphi_t) \circ \varphi_{je_\nu} = F \circ (\varphi_{je_\nu}\varphi_t)$ by (2.2), where $\varphi_{je_\nu}\varphi_t = \varphi_{t+je_\nu}$ because the two maps change different coordinates and ψ_0 is the identity, while the numbers of complex conjugations add up to $|t| + j = |t + je_\nu|$ modulo 2. The maps φ_t with $t \in \Lambda_\nu$ change only the coordinates $x_{\nu'}$ with $\nu' < \nu$ and map B_0° into $\{0 < x_\nu < b\}$, so that G_ν depends only on the values of U on the bonds with both ends in this slab, and Lemma 2.1 with $\mu = \nu$ and $G = G_\nu$ gives $\mathbb{E}_K^{(s)}[G_{\nu+1}] \geq 0$ and $|\mathbb{E}_K^{(s)}[G_\nu]|^{2m} \leq \mathbb{E}_K^{(s)}[G_{\nu+1}]$. Since $G_4 = \prod_{t\in\Lambda} \vartheta_t F$ and $|\Lambda| = (2m)^4$, the chain

$$|\mathbb{E}_K^{(s)}[F]|^{(2m)^4} \leq \mathbb{E}_K^{(s)}[G_1]^{(2m)^3} \leq \mathbb{E}_K^{(s)}[G_2]^{(2m)^2} \leq \mathbb{E}_K^{(s)}[G_3]^{2m} \leq \mathbb{E}_K^{(s)}[G_4] \tag{2.10}$$

proves (2.9). *q.e.d.*

3 The Bound Uniform in the Volume

We now apply the chessboard estimate to a Wilson loop insertion. The reflected copies of an ensemble form an ensemble of the same kind, its normalisation is the product of the normalisations of the copies up to complex conjugation, and the uniform bound of the paper applies to it.

Proposition 3.1 (Bound Uniform in the Volume)**.** *Let $s \geq 0$, let $\mathcal{C}$ be a loop ensemble at level q on $\mathbb{T}^{(s)}$ whose loops are admissible, with $\bar{g}$ and q as in Paper, Theorem 1.5, for $\mathcal{C}$, let n be an integer with $-q \leq n \leq s$, and put $b \coloneqq uL^n$. Assume that there is $v \in uL^{-q}\mathbb{Z}^4$ such that every site of $\mathcal{C}$ lies in the image in $\mathbb{T}^{(s)}$ of the open box $v + (0,b)^4$. Then*

$$|W^{(s)}(\mathcal{C})| \leq \exp(B_E(\mathcal{C}) + I_*^\flat(\mathcal{C}) + \Gamma_q(b,\bar{g})) \,. \tag{3.1}$$

Proof. By Paper, Corollary 1.6(ii), which holds for $W_K^{(s)}$ at every $K > j_q$, the renormalised expectations do not change when $\mathcal{C}$ is translated by $-v \in uL^{-q}\mathbb{Z}^4$, and the translate has the same numbers B_E and $I_*^\flat$; we may therefore assume $v = 0$. We use the notation of Section 2 with these s, q, n, and $K > j_q$.

The reflected ensemble. For $t \in \Lambda$ let $\vartheta_t\mathcal{C}$ be the ensemble $\varphi_t\mathcal{C}$ of the images of the loops of $\mathcal{C}$ with the image orientations, with the characters of $\mathcal{C}$ if $|t|$ is even and with the conjugate characters if $|t|$ is odd. Since $\varphi_t \in \mathfrak{G}_q$, each $\vartheta_t\mathcal{C}$ is an ensemble at level q whose loops have the chord–arc constants, ratios P/r, and numbers P_q of those of $\mathcal{C}$, and $c_{\bar{\pi}} = c_\pi$ for every representation. The loops of $\vartheta_t\mathcal{C}$ lie in the open block B_t°, and these blocks are disjoint, so that the concatenation $\mathcal{D}$ of the ensembles $\vartheta_t\mathcal{C}$, $t \in \Lambda$, is a loop ensemble at level q with $|\Lambda|n$ admissible loops and $c_\mathcal{D} = c_\mathcal{C}$; hence $\bar{g}$ and q are as in Paper, Theorem 1.5, for $\mathcal{D}$ as well, the function $q_\sharp$ depending on the ensemble only through $c_\mathcal{D}$.

The chessboard estimate. The observable $F \coloneqq W_\mathcal{C}$ depends only on the values of U on the bonds of the loops of $\mathcal{C}$, which have both ends in the convex set B_0°. By Paper, Section 12.1, $W_\chi(C) \circ \varphi_t = W_\chi(\varphi_t C)$, and $\overline{W_\chi(C')} = W_{\bar{\chi}}(C')$ for every loop C', as in Paper, (12.5); by multiplicativity, $\vartheta_t W_\mathcal{C} = W_{\vartheta_t\mathcal{C}}$ and $\prod_{t\in\Lambda} \vartheta_t W_\mathcal{C} = W_\mathcal{D}$. Lemma 2.2 therefore gives

$$|\mathbb{E}_K^{(s)}[W_\mathcal{C}]|^{|\Lambda|} \leq \mathbb{E}_K^{(s)}[W_\mathcal{D}] \,. \tag{3.2}$$

The normalisations. By Paper, Proposition 12.3, $\kappa_K^{(q)}(\varphi_t C_i, \chi) = \kappa_K^{(q)}(C_i,\chi)$ since $\varphi_t \in \mathfrak{G}_q$, and $\kappa_K^{(q)}(C_i,\bar{\chi}_i) = \overline{\kappa_K^{(q)}(C_i,\chi_i)}$. The normalisation of $\mathcal{D}$ is therefore the product over $t \in \Lambda$ of $\prod_i \kappa_K^{(q)}(C_i,\chi_i)$ or of its complex conjugate, and its modulus is $|\prod_i \kappa_K^{(q)}(C_i,\chi_i)|^{|\Lambda|}$, which is not zero because the step factors are not (Paper, Section 1.3). Dividing (3.2) by it gives

$$|W_K^{(s)}(\mathcal{C})|^{|\Lambda|} \leq |W_K^{(s)}(\mathcal{D})| \,. \tag{3.3}$$

The uniform bound for $\mathcal{D}$. The requirements on $\bar{g}$ in Paper, Theorem 8.6, (8.19) and the hypothesis of Paper, Proposition 8.4, involve no representation and are among those which define γ_0, and the requirement (8.40) on γ_q is among those which define $q_\sharp$ (Paper, Section 1.3). That theorem gives, for $K \geq K_0(\mathcal{D})$,

$$|W_K^{(s)}(\mathcal{D})| \leq e^{B_E(\mathcal{D}) + I_*(\mathcal{D})}(1 + \bar{\delta}_{\mathrm{LF}}) \,. \tag{3.4}$$

The number $B_E(\mathcal{D})$ of Paper, (7.32), is a sum over the loops of $\mathcal{D}$ of numbers which depend only on P/r, P_q, and $c_\mathcal{D} = c_\mathcal{C}$, so that $B_E(\mathcal{D}) = |\Lambda| B_E(\mathcal{C})$; and $I_*(\mathcal{D}) \leq I_*^\flat(\mathcal{D}) = |\Lambda| I_*^\flat(\mathcal{C})$, because $P_{K'}(\mathcal{D}) = |\Lambda| P_{K'}(\mathcal{C})$. By Paper, Proposition 8.4, $\bar{\delta}_{\mathrm{LF}} = e^{2\bar{C}_K}(e^{\bar{s}_{\mathrm{LF}}} - 1)$ with $\bar{C}_K \coloneqq C_{\mathrm{fs}} N_q^4 (1 + \frac{1}{2}\log \hat{g}_{K'}^{-2})$ and $\bar{s}_{\mathrm{LF}} \coloneqq 2N_q^4 (MR_{K'})^{-4} \exp(-c_{\mathrm{LF}}\, p_0((1+\beta_0)\bar{g}))$, so that $1 + \bar{\delta}_{\mathrm{LF}} \leq e^{2\bar{C}_K} e^{\bar{s}_{\mathrm{LF}}}$, because $e^{2\bar{C}_K} \geq 1$. Since $N_q = T_s L^{j_q}/u = 2mbL^{j_q}/u$ by Paper, (8.2), we have $N_q^4 = |\Lambda|(bL^{j_q}/u)^4$, and comparison with (1.2) gives $2\bar{C}_K + \bar{s}_{\mathrm{LF}} = |\Lambda|\Gamma_q(b,\bar{g})$. Altogether,

$$|W_K^{(s)}(\mathcal{D})| \leq \exp\Big(|\Lambda|(B_E(\mathcal{C}) + I_*^\flat(\mathcal{C}) + \Gamma_q(b,\bar{g}))\Big) \,. \tag{3.5}$$

Conclusion. By (3.3) and (3.5), $|W_K^{(s)}(\mathcal{C})| \le \exp(B_E(\mathcal{C}) + I_*^\flat(\mathcal{C}) + \Gamma_q(b, \bar g))$ for every $K \ge \max(K_0(\mathcal{C}), K_0(\mathcal{D}))$, and the limit $K \to \infty$, which exists by Paper, Theorem 1.5, gives (3.1). *q.e.d.*

4 Proof of the Theorem

Proof of Theorem 1.2. The set $\mathfrak{E}$ is countable. For each q the loops at level q in $\mathbb{R}^4$ are finite paths in a countable lattice. The compact group G has countably many classes of irreducible unitary representations, by the Peter–Weyl theorem, since $L^2(G)$ is separable and the matrix coefficients of inequivalent irreducible representations are orthogonal, and distinct classes have distinct characters. A countable union of sets of finite families of elements of countable sets is countable.

A bound uniform in s. Let $(\mathcal{C}, q) \in \mathfrak{E}$ and $s \ge s_q(\mathcal{C})$. Put $n := n_q(\mathcal{C})$, so that $-q \le n \le s-1$ and $b := uL^n = b_q(\mathcal{C})$, and let v_μ be the least coordinate x_μ of a site of $\mathcal{C}$ minus uL^{-q}, so that $v \in uL^{-q}\mathbb{Z}^4$. Every site of $\mathcal{C}$ lies in $v + [uL^{-q}, uL^{-q} + E(\mathcal{C})]^4 \subset v + (0,b)^4$, because $uL^{-q} + E(\mathcal{C}) < b$. The ensemble $\pi_s\mathcal{C}$ has admissible loops and $c_{\pi_s\mathcal{C}} = c_\mathcal{C}$, so that $\bar g$ and q are as in Paper, Theorem 1.5, for it, because γ_0 and $q_\sharp$ do not depend on the torus; and B_E and $I_*^\flat$ take the same values on $\pi_s\mathcal{C}$ and on $\mathcal{C}$. Proposition 3.1 gives

$$|W^{(s)}(\mathcal{C})| \le \beta(\mathcal{C}) := \exp(B_E(\mathcal{C}) + I_*^\flat(\mathcal{C}) + \Gamma_q(b_q(\mathcal{C}), \bar g)) \tag{4.1}$$

for every $s \ge s_q(\mathcal{C})$.

The subsequence. Let $e_1, e_2, \dots$ be an enumeration of $\mathfrak{E}$, and let σ_i be the number $s_q(\mathcal{C})$ of $e_i = (\mathcal{C}, q)$. By (4.1) and the Bolzano–Weierstrass theorem we choose infinite sets $\mathbb{N} \supset N_1 \supset N_2 \supset \dots$ with $N_i \subset [\sigma_i, \infty)$ along which $W^{(s)}(e_i)$ converges, and then integers $s_1 < s_2 < \dots$ with $s_k \in N_k$. For every i the sequence $(s_k)_{k \ge i}$ lies in N_i, so that (1.3) exists for e_i, and (i) follows from (4.1).

The symmetries. Let $g(x) = Ox + v$ be as in (ii), and for every s let g_s be the map $x \mapsto Ox + v$ of $\mathbb{T}^{(s)}$, which belongs to the group $\mathfrak{G}_q$ of $\mathbb{T}^{(s)}$ and satisfies $\pi_s \circ g = g_s \circ \pi_s$, so that $\pi_s(g\mathcal{C}) = g_s \pi_s \mathcal{C}$. For $s \ge \max(s_q(\mathcal{C}), s_q(g\mathcal{C}))$, Paper, Corollary 1.6(ii), gives $W^{(s)}(g\mathcal{C}) = W^{(s)}(\mathcal{C})$, and the limit along (s_k) gives $W^\infty(g\mathcal{C}) = W^\infty(\mathcal{C})$. The other three statements of (ii) follow in the same way from Paper, Corollary 1.6(i), (iii), and (iv), the maps h and g_i being replaced by the maps of $\mathbb{T}^{(s)}$ with the same formulae, which are symmetries of the level q of $\mathbb{T}^{(s)}$.

Reflection positivity. Let μ, y, θ, and $\mathcal{D}_1, \dots, \mathcal{D}_m$ be as in (iii). The ensembles $\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b$ are ensembles at level q in $\mathbb{R}^4$, since θ maps $uL^{-q}\mathbb{Z}^4$ onto itself and the loops of $\theta\bar{\mathcal{D}}_b$ lie in $\{x_\mu < y\}$; their loops are admissible, and $c_{\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b} \le c$, since $c_{\bar\pi} = c_\pi$, so that they form pairs of $\mathfrak{E}$ with the level q, because $q_\sharp$ is nondecreasing in its first argument. Let s be so large that s exceeds the numbers s_q of all these ensembles and that $T_s/2 > x_\mu - y$ for every site x of every $\mathcal{D}_a$. Then π_s maps the sites of every $\mathcal{D}_a$ into the open half-torus $\{y < x_\mu < y + T_s/2\}$ of the reflection $\theta_{\mu,y}$ of $\mathbb{T}^{(s)}$, which is a reflection of level q, and $\pi_s \circ \theta = \theta_{\mu,y} \circ \pi_s$, so that $\pi_s(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b) = \pi_s\mathcal{D}_a \cup \theta_{\mu,y}\overline{\pi_s\mathcal{D}_b}$. Paper, Corollary 1.7, applied on $\mathbb{T}^{(s)}$ to $\pi_s\mathcal{D}_1, \dots, \pi_s\mathcal{D}_m$, says that the matrices (1.4) with $W^{(s)}$ in place of W^∞ are Hermitian and positive semidefinite, and that $W^{(s)}(\theta\bar{\mathcal{D}}_a) = \overline{W^{(s)}(\mathcal{D}_a)}$. These are closed conditions on finitely many numbers, and they pass to the limit along (s_k). *q.e.d.*

Proof of Corollary 1.3. (i) Each term of (1.5) satisfies the bound of Theorem 1.2(i) for $P_\sigma\mathcal{C}$, and B_E, $I_*^\flat$, and b_q take the same values on $P_\sigma\mathcal{C}$ as on $\mathcal{C}$, since they depend on the ensemble

only through the ratios P/r, the numbers P_q, the number c, and the extent.

(ii) The first, third, and fourth statements follow term by term from Theorem 1.2(ii), since P_σ commutes with the reordering of the loops, with the conjugation of the characters, and with the reversal of the orientations. For the second, let $\mathcal{C}' = \mathcal{C}+w$. Then $h\mathcal{C}' = O\mathcal{C}+(Ow+v)$ is a translate of $O\mathcal{C}$, and $(O\mathcal{C}, q) \in \mathfrak{E}$, because O, like P_σ, maps $uL^{-q}\mathbb{Z}^4$ onto itself and preserves the distance $|\cdot|_\infty$, the segments, and the corners. By (1.6), $W^{\mathrm{hc}}(h\mathcal{C}') = W^{\mathrm{hc}}(O\mathcal{C})$ and $W^{\mathrm{hc}}(\mathcal{C}') = W^{\mathrm{hc}}(\mathcal{C})$, and it remains to show that $W^{\mathrm{hc}}(O\mathcal{C}) = W^{\mathrm{hc}}(\mathcal{C})$. We write $O = DP_\tau$ with a diagonal matrix D with entries ± 1 and a permutation τ, which is possible in exactly one way. For every σ,

$$P_\sigma O = D_\sigma P_{\sigma\tau} , \qquad D_\sigma \coloneqq P_\sigma D P_\sigma^{-1} , \tag{4.2}$$

where D_σ is diagonal with entries ± 1, since $D_\sigma e_{\sigma(\mu)} = P_\sigma D e_\mu$ is $e_{\sigma(\mu)}$ or $-e_{\sigma(\mu)}$. Hence $P_\sigma(O\mathcal{C}) = D_\sigma(P_{\sigma\tau}\mathcal{C})$, and $W^\infty(P_\sigma O\mathcal{C}) = W^\infty(P_{\sigma\tau}\mathcal{C})$ by Theorem 1.2(ii) for the map $g = D_\sigma$. Summing over σ, and since $\sigma \mapsto \sigma\tau$ is a bijection of the permutations, $W^{\mathrm{hc}}(O\mathcal{C}) = W^{\mathrm{hc}}(\mathcal{C})$.

(iii) Let θ'_σ be the reflection of $\mathbb{R}^4$ which replaces $x_{\sigma(\mu)}$ by $2y - x_{\sigma(\mu)}$. Since $(P_\sigma \theta x)_{\sigma(\mu)} = 2y - x_\mu = 2y - (P_\sigma x)_{\sigma(\mu)}$ and the other coordinates of $P_\sigma\theta x$ and $P_\sigma x$ agree, $P_\sigma\theta = \theta'_\sigma P_\sigma$. Hence the loops of $P_\sigma\mathcal{D}_a$ lie in $\{x_{\sigma(\mu)} > y\}$ and $P_\sigma(\theta\bar{\mathcal{D}}_b) = \theta'_\sigma\overline{P_\sigma\mathcal{D}_b}$, so that

$$W^{\mathrm{hc}}(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b) = \frac{1}{24}\sum_\sigma W^\infty(P_\sigma\mathcal{D}_a \cup \theta'_\sigma\overline{P_\sigma\mathcal{D}_b}) . \tag{4.3}$$

For each σ the matrix of the summands is Hermitian and positive semidefinite by Theorem 1.2(iii), applied to the direction $\sigma(\mu)$, the same y, and the ensembles $P_\sigma\mathcal{D}_1, \dots, P_\sigma\mathcal{D}_m$, which satisfy its hypotheses because P_σ preserves admissibility, the level, and the characters; so the average is Hermitian and positive semidefinite. The same theorem gives $W^\infty(\theta'_\sigma\overline{P_\sigma\mathcal{D}_a}) = \overline{W^\infty(P_\sigma\mathcal{D}_a)}$, and averaging gives $W^{\mathrm{hc}}(\theta\bar{\mathcal{D}}_a) = \overline{W^{\mathrm{hc}}(\mathcal{D}_a)}$. It remains to treat the connected matrix, in which the product of the averages is not the average of the products. We write $w_a^\sigma \coloneqq W^\infty(P_\sigma\mathcal{D}_a)$ and $w_a \coloneqq \frac{1}{24}\sum_\sigma w_a^\sigma = W^{\mathrm{hc}}(\mathcal{D}_a)$; expanding the last sum below and using $\frac{1}{24}\sum_\sigma w_a^\sigma = w_a$ gives

$$W^{\mathrm{hc}}(\mathcal{D}_a \cup \theta\bar{\mathcal{D}}_b) - w_a\overline{w_b} = \frac{1}{24}\sum_\sigma\Big[W^\infty(P_\sigma\mathcal{D}_a \cup \theta'_\sigma\overline{P_\sigma\mathcal{D}_b}) - w_a^\sigma\overline{w_b^\sigma}\Big] + \frac{1}{24}\sum_\sigma (w_a^\sigma - w_a)\overline{(w_b^\sigma - w_b)} . \tag{4.4}$$

The first sum is an average of matrices which are Hermitian and positive semidefinite by Theorem 1.2(iii), and the second is an average of the matrices $(z_a\bar{z}_b)_{a,b}$ with $z_a = w_a^\sigma - w_a$, which are Hermitian and positive semidefinite because $\sum_{a,b}\bar{\xi}_a z_a \bar{z}_b \xi_b = |\sum_a \bar{\xi}_a z_a|^2$; so the connected matrix of W^{hc} is Hermitian and positive semidefinite. *q.e.d.*

Remark 4.1 (What the Theorem Does Not Assert). The theorem gives no lower bound, and the lower bound of the paper does not pass to the limit: its proof, Paper, Proposition 12.5, uses $w_{\ell_0} \geq 1 - \bar{\delta}_{\mathrm{LF}}$ with a threshold $\bar{g}_1$ chosen so that $\bar{\delta}_{\mathrm{LF}} \leq 1/20$, and $\bar{\delta}_{\mathrm{LF}}$ grows with N_q, while a chessboard estimate bounds expectations from above only. The theorem therefore does not exclude that W^∞ vanishes on every nonempty ensemble, and Paper, Theorem 1.8, whose threshold $\bar{g}_0$ depends on T/u, has no counterpart here. The limits may depend on the subsequence.

References

[1] Tadeusz Bałaban, *Renormalization Group Approach to Lattice Gauge Field Theories. I. Generation of Effective Actions in a Small Field Approximation and a Coupling Constant Renormalization in Four Dimensions*, Comm. Math. Phys. **109** (1987), 249–301.

[2] Jürg Fröhlich, Robert Israel, Elliott H. Lieb, and Barry Simon, *Phase Transitions and Reflection Positivity. I. General Theory and Long Range Lattice Models*, Comm. Math. Phys. **62** (1978), 1–34.