

Multi Variable Calculus

Mathias Balling Christiansen

Last updated: December 22, 2023

Contents

1	Fourier	3
2	Laplace	4
2.1	Examples	4
2.1.1	Example 1: Laplace Transform	4
3	Several-Variables	5
4	Double-Integrals	6
4.1	Riemann Sum	6
4.2	Double Integrals over General domains	6
4.3	Iteration of Double Integrals	6
4.4	Double Integrals in Polar Coordinates	6
4.5	Change of Variables in Double Integrals	6
4.6	Examples	7
4.6.1	Example 1: Change of Variables	7
4.6.2	Example 2: Riemann Sum	7
4.6.3	Example 3: By iteration	7
4.6.4	Example 4: Double integral	7
5	Tripple-Integrals	8
5.1	Examples	8
5.1.1	Example 1: Tripple Integral	8
6	Fields-Curve	9
6.1	Examples	9
6.1.1	Example 1: Conservative vector field and potential	9
6.1.2	Example 2: Line integral	9
7	Theorems	10
7.1	Examples	10
7.1.1	Example 1: Div and Curl	10
7.1.2	Example 2: Green's Theorem	10
7.1.3	Example 3: Stokes' Theorem	10
7.1.4	Example 4: Divergence Theorem	10
8	PDE	11
8.1	Classification of PDEs	11
8.2	Characteristics of PDEs	11
8.3	Initial and Boundary Conditions	11
8.4	Wave Equation (1D)	11
8.4.1	D'Alembert's Solution of the Wave Equation	11

8.5	Heat Equation (1D)	12
8.6	Examples	12

1 Fourier

2 Laplace

2.1 Examples

2.1.1 Example 1: Laplace Transform

Using the Laplace transform, find the solution for the following equation:

$$\frac{\partial^2}{\partial t^2}y(t) + 2\frac{\partial}{\partial t}y(t) + 2y(t) = 0$$

with initial conditions $y(0) = 1$ and $Dy(0) = -1$

3 Several-Variables

4 Double-Integrals

4.1 Riemann Sum

$$\sum_{i=1}^n f(x_i^*, y_i^*) \Delta A_i$$
$$I = \iint_D f(x, y) dA \quad \text{where } D \text{ is a region in } \mathbb{R}^2 \text{ and } dA \text{ is } dxdy$$

4.2 Double Integrals over General domains

If $f(x, y)$ is defined and bounded on domain D , then $\hat{f}(x, y)$ is zero outside D .

$$\iint_D f(x, y) dA = \iint_{\mathbb{R}^2} \hat{f}(x, y) dA$$

4.3 Iteration of Double Integrals

4.4 Double Integrals in Polar Coordinates

$$dA = dxdy = r drd\theta$$
$$\begin{aligned} x &= r \cos(\theta) & r^2 &= x^2 + y^2 \\ y &= r \sin(\theta) & \tan(\theta) &= \frac{y}{x} \end{aligned}$$

4.5 Change of Variables in Double Integrals

If x and y are given as a function of u and v :

$$\begin{aligned} x &= x(u, v) \\ y &= y(u, v) \end{aligned}$$

These can be transformed or mapped from points (u, v) in the uv -plane to points (x, y) in the xy -plane.

The inverse transformation is given by:

$$\begin{aligned} u &= u(x, y) \\ v &= v(x, y) \end{aligned}$$

Scaled area element:

$$dA = dxdy = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

where the Jacobian is:

$$\left| \frac{\partial(x, y)}{\partial(u, v)} \right| = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

Let $x(u, v)$ and $y(u, v)$ be a one-to-one transformation from a domain S in the uv -plane onto a domain D in the xy -plane.

Suppose, that function x and y , and first partial derivatives with respect to u and v are continuous in S . If $f(x, y)$ is integrable on D , then $g(u, v) = f(x(u, v), y(u, v))$ is integrable on S and:

$$\iint_D f(x, y) dA = \iint_S g(u, v) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

4.6 Examples

4.6.1 Example 1: Change of Variables

Evaluate the double integral:

$$\iint (x - 3y) dA$$

where R is triangular region with vertices $(0, 0)$, $(2, 1)$, and $(1, 2)$ using the transformation:

$$\begin{aligned}x &= 2u + v \\y &= u + 2v\end{aligned}$$

4.6.2 Example 2: Riemann Sum

4.6.3 Example 3: By iteration

4.6.4 Example 4: Double integral

Find the area within the region:

$$\iint (\sin x + \cos y) dA \quad R : \left\{ x, y \mid \begin{array}{l} 0 \leq x \leq \pi/2 \\ 0 \leq y \leq \pi/2 \end{array} \right.$$

5 Tripple-Integrals

5.1 Examples

5.1.1 Example 1: Tripple Integral

Find $\iiint (x^2 + y^2 + z^2) dV$, where the region is bounded by $z = c\sqrt{(x^2 + y^2)}$ and $x^2 + y^2 + z^2 = a^2$

6 Fields-Curve

6.1 Examples

6.1.1 Example 1: Conservative vector field and potential

Determine whether the given vector field is conservative, and find a potential function if it is:

$$\mathbf{F}(x, y, z) = (2xy - z^2)\mathbf{i} + (2yz + x^2)\mathbf{j} - (2zx - y^2)\mathbf{k}$$

6.1.2 Example 2: Line integral

Evaluate $\oint x^2 y^2 dx + x^3 y dy$ counterclockwise around the square with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$, and $(0, 1)$

7 Theorems

7.1 Examples

7.1.1 Example 1: Div and Curl

Calculate the divergence and curl of the following vector field:

$$\mathbf{F} = \cos x \mathbf{i} - \sin y \mathbf{j} + z \mathbf{k}.$$

Divergence: Curl:

7.1.2 Example 2: Green's Theorem

Using Green's Theorem evaluate $\oint_C (x^2y) dx + (xy^2) dy$, clockwise boundary of the region:

$$0 \leq y \leq \sqrt{9 - x^2}$$

7.1.3 Example 3: Stokes' Theorem

7.1.4 Example 4: Divergence Theorem

Use the Divergence Theorem to calculate the flux of the given vector field out of the sphere s with equation $x^2 + y^2 + z^2 = a^2$, where $a > 0$ and

$$\mathbf{F} = (x^2 + y^2)\mathbf{i} + (y^2 - z^2)\mathbf{j} + z\mathbf{k}$$

8 PDE

Partial Differential Equations are equations with multiple variables and derivatives. They are used to model many physical phenomena, such as heat, sound, and light. The totality of solutions to a PDE is called its general solution, and there can be a lot.

8.1 Classification of PDEs

General representation of a PDE:

$$A \frac{\partial^2 u}{\partial x^2} + B \frac{\partial^2 u}{\partial x \partial y} + C \frac{\partial^2 u}{\partial y^2} + D \frac{\partial u}{\partial x} + E \frac{\partial u}{\partial y} + Fu = G$$
$$Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu = G$$

Conditions:

Linear: A, B, C, D, E, F are constants

Quasi-linear: A, B, C are constants

Fully non-linear: A, B, C, D, E, F are functions of u and its partial derivatives

8.2 Characteristics of PDEs

$B^2 - 4AC > 0$	2 real roots	2 characteristics	Hyperbolic PDE
$B^2 - 4AC = 0$	1 real roots	1 characteristics	Parabolic PDE
$B^2 - 4AC < 0$	0 real roots	0 characteristics	Elliptic PDE

Types of various PDEs:

Wave Equation: Hyperbolic PDE

Heat Equation: Parabolic PDE

Laplace Equation: Elliptic PDE

8.3 Initial and Boundary Conditions

8.4 Wave Equation (1D)

One dimensional wave equation is given by:

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \text{where } c^2 = \left[\frac{T(x,t)}{\mu_x} \right] \quad (1)$$

Steps to solve:

1. Method of Separation of Variables $u(x,t) = X(x)T(t)$
2. Satisfy the Boundary Conditions test
3. Fourier Series Validation

8.4.1 D'Alembert's Solution of the Wave Equation

His solution is given by eq. (1) but extended to two variables:

$$v = (x - ct) \quad w = (x + ct) \quad (2)$$

I.e. $u(v,w)$. Partial derivatives from chain rule:

$$u_x = u_v \cdot v_x + u_w \cdot w_x = u_v + u_w$$

For double derivatives:

$$u_{xx} = (u_v + u_w)_x = (u_v + u_w)_v v_x + (u_v + u_w)_w w_x = u_{vv} + 2u_{vw} + u_{ww}$$

With respect to t :

$$u_{tt} = c^2 u_{xx} = c^2 (u_{vv} + 2u_{vw} + u_{ww})$$

From eq. (1) and eq. (2):

$$u_{vw} = \frac{\partial^2 u}{\partial w \partial v} = 0$$

This can be solved by integrating with respect to v and w :

$$\frac{\partial u}{\partial v} = h(v) \text{ and } u = \int h(v) dv + \psi(w)$$

Here, $h(v)$ and $\psi(w)$ are arbitrary functions of v and w , respectively. The solution in term for x :

$$u = \phi(v) + \psi(w)$$

This is d'Alembert's solution, which is the general solution to the wave equation.

$$u(x, t) = \phi(x + ct) + \psi(x - ct)$$

This solution satisfies the wave equation and the initial conditions:

8.5 Heat Equation (1D)

$$\frac{\partial u}{\partial t} = c \frac{\partial^2 u}{\partial x^2} \quad (3)$$

Conditions:

- PDE is linear and homogeneous.
- Boundary conditions are linear and homogeneous. The two for $u(x, t)$ is $u(0, t) = 0$ and $u(L, t) = 0$ for all $t > 0$.
- One initial condition at time ($t = 0$): $u(x, 0) = f(x)$.

Solve the

Solution:

$$u(x, t) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) e^{-\lambda_n^2 t}$$

where

$$\lambda_n = \frac{cn\pi}{L}$$

and

$$B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx \quad \text{for } n = 1, 2, 3, \dots$$

8.6 Examples