

Multi Variable Calculus

Mathias Balling Christiansen

Last updated: December 30, 2023

Contents

1	Fourier	3
1.1	Fourier Series	3
1.2	Fourier Transform	3
1.3	Examples	3
1.3.1	Example 1: Fourier Series	3
1.3.2	Example 2: Fourier Transform	3
1.3.3	Example 3: Inverse Fourier Transform	4
2	Laplace	5
2.1	Examples	5
2.1.1	Example 1: Laplace Transform	5
2.1.2	Example 2: Trabsfer Function	5
3	Several-Variables	6
3.1	Examples	6
3.1.1	Example 1: Gradient	6
3.1.2	Example 2: Jacobian	6
3.1.3	Example 3: Chain Rule	6
3.1.4	Example 4: Substitution	6
3.1.5	Example 5: Partial Differentiation	6
4	Double-Integrals	7
4.1	Riemann Sum	7
4.2	Double Integrals over General domains	7
4.3	Iteration of Double Integrals	7
4.4	Double Integrals in Polar Coordinates	7
4.4.1	Limits for Polar Coordinates	7
4.5	Change of Variables in Double Integrals	7
4.6	Examples	8
4.6.1	Example 1: Change of Variables	8
4.6.2	Example 2: Double integral	9
4.6.3	Example 3: By iteration	9
4.6.4	Example 4: Polar coordinates	10
5	Tripple-Integrals	11
5.1	Riemann Sum	11
5.2	Tripple Integrals over General domains	11
5.3	Change of Variables	11
5.4	Cylyndrical Coordinates	11
5.4.1	Limits for Cylyndrical Coordinates	11

5.5	Spherical Coordinates	11
5.5.1	Limits for Spherical Coordinates	11
5.6	Examples	12
5.6.1	Example 1: Tripple Integral	12
5.6.2	Example 2: Sphereical Coordinates	12
5.6.3	Example 3: Transformations	12
6	Fields-Curve	13
6.1	Curve & Parameterization	13
6.2	Vector Fields	13
6.2.1	Scalar field	13
6.2.2	Field lines	13
6.2.3	Conervation field	13
6.2.4	Vector field in Polar Coordinates	13
6.3	Line Integral	14
6.3.1	Line integral of a vector field	14
6.4	Examples	14
6.4.1	Example 1: Conservative vector field and potential	14
6.4.2	Example 2: Line integral	14
6.4.3	Example 3: Line integral	14
6.4.4	Example 4: Line integral vector field	14
6.4.5	Example 5: Gradient	14
6.4.6	Example 6: Parametrize a curve	14
7	Surface-Integrals	15
7.1	Parametric Surface	15
7.2	Surface Area	15
7.3	Oriented Surface	15
7.4	Flux	16
7.5	Examples	16
8	Theorems	17
8.1	Differential Operators	17
8.1.1	Gradient	17
8.1.2	Divergence	17
8.1.3	Curl	17
8.2	Green's Theorem	17
8.3	Stokes' Theorem	17
8.4	Divergence Theorem	17
8.5	Examples	18
8.5.1	Example 1: Div and Curl	18
8.5.2	Example 2: Green's Theorem	18
8.5.3	Example 3: Stokes' Theorem	18
8.5.4	Example 4: Divergence Theorem	18
9	PDE	19
9.1	Classification of PDEs	19
9.2	Characteristics of PDEs	19
9.3	Important Second-Order PDEs	20
9.4	Initial and Boundary Conditions	20
9.5	Wave Equation (1D)	20
9.5.1	D'Alembert's Solution of the Wave Equation	20
9.6	Heat Equation (1D)	21
9.7	Examples	21
9.7.1	21

1 Fourier

1.1 Fourier Series

A periodic function with period $2L$ and let $f(x)$ and $f'(x)$ be piecewise continuous on the interval $-L < x < L$

$$S_N(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos\left(\frac{n\pi x}{L}\right) + b_n \sin\left(\frac{n\pi x}{L}\right) \right) = \sum_{n=-\infty}^{\infty} c_n e^{jn\pi x/L}$$

The coefficients are given by:

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx \quad n \geq 0$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx \quad n > 0$$

$$c_n = \frac{1}{2}(a_n - jb_n) \quad n > 0$$

1.2 Fourier Transform

If $h(t)$ is a periodic function then the Fourier transform is given by:

$$H(\omega) = \int_{-\infty}^{\infty} h(t) e^{-j\omega t} dt$$

Inverse Fourier transformation of $H(\omega)$:

$$h(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} H(\omega) e^{j\omega t} d\omega$$

1.3 Examples

1.3.1 Example 1: Fourier Series

Find the Fourier coefficients and Fourier Series for the square wave shown below:

$$f(t) = \begin{cases} 1 & \text{for } t - a > 0 \\ 0 & \text{for } t - a < 0 \end{cases}$$

and

$$f(x+2) = f(x)$$

Ans:

$$f(x) = \frac{1}{2} + \sum_{n=1,3,5,\dots} \frac{2}{n\pi} \sin(n\pi x)$$

1.3.2 Example 2: Fourier Transform

The unit step function is defined as:

$$u(t-a) = \begin{cases} 1 & \text{for } t-a > 0 \\ 0 & \text{for } t-a < 0 \end{cases}$$

is used to define the rectangular pulse function:

$$x(t) = u(t-a) - u(t-b) \quad \text{where } a < b$$

Ans:

$$X(\omega) = \frac{e^{-j\omega a} - e^{-j\omega b}}{j\omega}$$

1.3.3 Example 3: Inverse Fourier Transform

Consider the signal: $x(t) = \sin(\omega_0 t)$ where ω_0 is a constant. Find the Fourier Transform of this signal to find $X(\omega)$.

Ans:

$$X(\omega) = -j\pi(\delta(\omega - \omega_0) - \delta(\omega + \omega_0))$$

2 Laplace

Is a generalisation of the Fourier transform and defined as:

$$H(s) = \mathcal{L}\{h(t)\} = \int_{0^-}^{\infty} h(t)e^{-st} dt \quad s \in \mathbb{C}$$

s is a complex number $s = \sigma + j\omega$ and is identical with Fourier transform, if s is set to $j\omega$.
Inverse Laplace transformation:

$$f(t) = \frac{1}{2\pi j} \int_{\sigma_c - j\infty}^{\sigma_c + j\infty} F(s)e^{st} ds$$

2.1 Examples

2.1.1 Example 1: Laplace Transform

Using the Laplace transform, find the solution for the following equation:

$$\frac{\partial^2}{\partial t^2}y(t) + 2\frac{\partial}{\partial t}y(t) + 2y(t) = 0$$

with initial conditions $y(0) = 1$ and $Dy(0) = -1$

2.1.2 Example 2: Transfer Function

Consider a mass-spring-damper system with the following differential equation:

$$m\ddot{x} = -kx - b\dot{x} + f$$

Find the transfer function for the system with input f and output x .

Ans:

$$G(s) = \frac{X(s)}{F(s)} = \frac{1}{ms^2 + bs + k}$$

3 Several-Variables

3.1 Examples

3.1.1 Example 1: Gradient

Find the rate of change of $f(x, y) = y^4 + 2xy^3 + x^2y^2$ at $(0, 1)$ in each of the following directions:

1. $\mathbf{i} + 2\mathbf{j}$
2. $\mathbf{j} - 2\mathbf{i}$
3. $3\mathbf{i}$
4. $\mathbf{i} + \mathbf{j}$

3.1.2 Example 2: Jacobian

Find the Jacobian $Df(1, 0)$ for the transformation from \mathbb{R}^2 to \mathbb{R}^3 given by:

$$\mathbf{f}(x, y) = (xe^y + \cos(\pi y), x^2, x - e^y)$$

3.1.3 Example 3: Chain Rule

If $z = \sin(x^2y)$ where $x = st^2$ and $y = s^2 + \frac{1}{t}$, $y = e^{-t}$, find $\frac{\partial z}{\partial s}$ and $\frac{\partial z}{\partial t}$ using chain rule.

3.1.4 Example 4: Substitution

If $z = \sin(x^2y)$ where $x = st^2$ and $y = s^2 + \frac{1}{t}$, $y = e^{-t}$, find $\frac{\partial z}{\partial s}$ and $\frac{\partial z}{\partial t}$ using substitution.

3.1.5 Example 5: Partial Differentiation

Calculate $f_{223}(x, y, z)$, $f_{232}(x, y, z)$, and $f_{322}(x, y, z)$ of the following function:

$$f(x, y, z) = e^{x-2y+3z}$$

4 Double-Integrals

4.1 Riemann Sum

$$\sum_{i=1}^n f(x_i^*, y_i^*) \Delta A_i$$

$$I = \iint_D f(x, y) dA \quad \text{where } D \text{ is a region in } \mathbb{R}^2 \text{ and } dA \text{ is } dx dy$$

4.2 Double Integrals over General domains

If $f(x, y)$ is defined and bounded on domain D , then $\hat{f}(x, y)$ is zero outside D .

$$\iint_D f(x, y) dA = \iint_R \hat{f}(x, y) dA$$

4.3 Iteration of Double Integrals

If $f(x, y)$ is continuous on the bounded y-simple domain D given by $a \leq x \leq b$ and $c(x) \leq y \leq d(x)$, then:

$$\iint f(x, y) dA = \int_a^b dx \int_{c(x)}^{d(x)} f(x, y) dy$$

If $f(x, y)$ is continuous on the bounded x-simple domain D given by $c \leq x \leq d$ and $a(x) \leq y \leq b(x)$, then:

$$\iint f(x, y) dA = \int_c^d dx \int_{a(x)}^{b(x)} f(x, y) dy$$

4.4 Double Integrals in Polar Coordinates

$$dA = dx dy = r dr d\theta$$

$$x = r \cos(\theta) \quad r^2 = x^2 + y^2$$

$$y = r \sin(\theta) \quad \tan(\theta) = \frac{y}{x}$$

4.4.1 Limits for Polar Coordinates

r is the radius from origin to the point

$$r \geq 0$$

θ is the angle in the positive direction of the xy-plane

$$0 \leq \theta \leq 2\pi$$

4.5 Change of Variables in Double Integrals

If x and y are given as a function of u and v :

$$x = x(u, v)$$

$$y = y(u, v)$$

These can be transformed or mapped from points (u, v) in the uv -plane to points (x, y) in the xy -plane.

The inverse transformation is given by:

$$u = u(x, y)$$

$$v = v(x, y)$$

Scaled area element:

$$dA = dx dy = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

where the Jacobian is:

$$\left| \frac{\partial(x, y)}{\partial(u, v)} \right| = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

Let $x(u, v)$ and $y(u, v)$ be a one-to-one transformation from a domain S in the uv -plane onto a domain D in the xy -plane.

Suppose, that function x and y , and first partial derivatives with respect to u and v are continuous in S . If $f(x, y)$ is integrable on D , then $g(u, v) = f(x(u, v), y(u, v))$ is integrable on S and:

$$\iint_D f(x, y) dA = \iint_S g(u, v) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

4.6 Examples

4.6.1 Example 1: Change of Variables

Evaluate the double integral:

$$\iint (x - 3y) dA$$

where R is triangular region with vertices $(0, 0)$, $(2, 1)$, and $(1, 2)$ using the transformation:

$$x = 2u + v$$

$$y = u + 2v$$

As x and y are dependent on u and v the transformation of dA is given by

$$dA = dx dy = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

Using:

$$\left| \frac{\partial(x, y)}{\partial(u, v)} \right| = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

the Jacobian can be calculated:

$$\frac{\partial x}{\partial u} = 2 \quad \frac{\partial y}{\partial v} = 2$$

$$\frac{\partial x}{\partial v} = 1 \quad \frac{\partial y}{\partial u} = 1$$

$$2 * 2 - 1 * 1 = 3$$

Find the boundaries for the double integral

$$y_1 : (0, 0) \rightarrow (2, 1) \text{ is } y = \frac{1}{2}x$$

$$y_2 : (0, 0) \rightarrow (1, 2) \text{ is } y = 2x$$

$$y_3 : (1, 2) \rightarrow (2, 1) \text{ is } y = 3 - x$$

Replace x and y with their transformation:

$$u + 2v = \frac{2u+v}{2} \quad v = 0$$

$$u + 2v = 4u + 2v \quad u = 0$$

$$u + 2v = 3 - 2u - v \quad u = 1 - v$$

Therefore $0 \leq u \leq 1 - v$ and $0 \leq v \leq 1$

Now transform the original function $x - 3y$:

$$\begin{aligned} x - 3y &= 2u + v - 3(u + 2v) \\ &= 2u - 3u + v - 6v \\ &= -u - 5v \end{aligned}$$

$$\begin{aligned} \int_0^1 \int_0^{1-v} (-u - 5v) \cdot 3 \, du dv &= -3 \int_0^1 \left[\frac{u^2}{2} + 5uv \right]_0^{1-v} dv \\ &= -\frac{3}{2} \int_0^1 \left(\frac{27v^2}{2} - 12v \right) dv \\ &= \left[\frac{9v^3}{2} - 6v^2 - \frac{3v}{2} \right]_0^1 \\ &= \boxed{-3} \end{aligned}$$

4.6.2 Example 2: Double integral

Evaluate the double integral by iteration

$$\iint_R (x^2 + y^2) \, dA$$

where R is the rectangle $0 \leq x \leq a$, $0 \leq y \leq b$

Insert the limits and solve the integral:

$$\begin{aligned} \int_0^b \int_0^a (x^2 + y^2) dx dy &= \int_0^b \left[\frac{x^3}{3} + xy^2 \right]_0^a dy \\ &= \int_0^b \left(\frac{a^3}{3} + ay^2 \right) dy \\ &= \boxed{\frac{a^3b}{3} + \frac{ab^3}{3}} \end{aligned}$$

4.6.3 Example 3: By iteration

Evaluate the double integral by iteration:

$$\iint_D x \cos y \, dA$$

where D is the finite region in the first quadrant bounded by the coordinate axes and the curve $y = 1 - x^2$.

Given the region the minimum for x and y must be 0 and for x the maximum is 1:

$$\begin{aligned}
\int_0^1 \int_0^{1-x^2} (x \cos(y)) dy dx &= \int_0^1 x \sin(1-x^2) dx \\
&= \left[\frac{1}{2} \sin(1) \sin(x^2) + \frac{1}{2} \cos(1) \cos(x^2) \right]_0^1 \\
&= \boxed{\sin^2\left(\frac{1}{2}\right)}
\end{aligned}$$

4.6.4 Example 4: Polar coordinates

5 Tripple-Integrals

5.1 Riemann Sum

5.2 Tripple Integrals over General domains

5.3 Change of Variables

$$dV = dx \, dy \, dz = \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du \, dv \, dw$$
$$\iiint f(x, y, z) \, dx \, dy \, dz = \iiint g(u, v, w) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du \, dv \, dw$$

5.4 Cylyndrical Coordinates

$$x = r \cos(\theta) \quad r^2 = x^2 + y^2$$
$$y = r \sin(\theta) \quad \tan \theta = \left(\frac{y}{x}\right)$$
$$z = z$$

From the Jacobian using change of variables:

$$\left| \frac{\partial(x, y, z)}{\partial(r, \theta, z)} \right| = r$$
$$dV = dx \, dy \, dz = r \, dr \, d\theta \, dz$$

5.4.1 Limits for Cylyndrical Coordinates

r is the radius from origin to the point

$$r \geq 0$$

θ is the angle in the positive direction of the x-axis

$$0 \leq \theta \leq 2\pi$$

5.5 Spherical Coordinates

$$x = \rho \sin(\phi) \cos(\theta) \quad \rho^2 = x^2 + y^2 + z^2$$
$$y = \rho \sin(\phi) \sin(\theta) \quad \cos(\phi) = \frac{z}{\rho}$$
$$z = \rho \cos(\phi) \quad \tan \theta = \frac{y}{x}$$
$$dx dy dz = dV = \rho^2 \sin(\phi) d\rho d\phi d\theta$$

5.5.1 Limits for Spherical Coordinates

ρ is the distance from the origin O to the point P

$$\rho \geq 0$$

ϕ is the angle by the radial line OP to the positive direction of the z-axis

$$0 \leq \phi \leq \pi$$

θ is the angle in the positive direction of the x-axis to the point P in the xy-plane

$$0 \leq \theta \leq 2\pi$$

5.6 Examples

5.6.1 Example 1: Tripple Integral

Find $\iiint (x^2 + y^2 + z^2) dV$, where the region is bounded by $z = c\sqrt{(x^2 + y^2)}$ and $x^2 + y^2 + z^2 = a^2$

5.6.2 Example 2: Spherical Coordinates

Find the volume of:

$$\iiint \sqrt{x^2 + y^2 + z^2} dx dy dz$$

where the region is bounded by $x^2 + y^2 + z^2 \leq 1$ in spherical domain.

Transform the region:

$$\rho^2 = x^2 + y^2 + z^2 \leq 1 \quad \text{Therefore: } 0 \leq \rho \leq 1$$

For θ and ϕ we have:

$$0 \leq \theta \leq 2\pi \quad 0 \leq \phi \leq \pi$$

and for dV :

$$dV = dx dy dz = \rho^2 \sin \phi d\rho d\phi d\theta$$

Transform the function:

$$\sqrt{x^2 + y^2 + z^2} = \sqrt{\rho^2} = \rho$$

Solve the integral:

$$\begin{aligned} \int_0^{2\pi} \int_0^\pi \int_0^1 \rho \cdot \rho^2 \sin(\phi) d\rho d\phi d\theta &= \int_0^{2\pi} \int_0^\pi \left[\frac{1}{4} \rho^4 \sin(\phi) \right]_0^1 d\phi d\theta \\ &= \int_0^{2\pi} \int_0^\pi \frac{\sin(\phi)}{4} d\phi d\theta \\ &= \int_0^{2\pi} \left[-\frac{\cos(\phi)}{4} \right]_0^\pi d\theta \\ &= \frac{1}{2} \int_0^{2\pi} d\theta \\ &= \frac{1}{2} \cdot 2\pi \\ &= \boxed{\pi} \end{aligned}$$

5.6.3 Example 3: Transformations

6 Fields-Curve

6.1 Curve & Parameterization

Representation of a curve in 3 space by using its position vector is given as:

$$r = r(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k} \quad \text{where } a \leq t \leq b$$

6.2 Vector Fields

$$\mathbf{F}(x, y, z) = \underbrace{f_x(x, y, z)}_{\text{Scaler function}} \mathbf{i} + \underbrace{f_y(x, y, z)}_{\text{Scaler function}} \mathbf{j} + \underbrace{f_z(x, y, z)}_{\text{Scaler function}} \mathbf{k}$$

$$\frac{\partial f}{\partial x} = f_1 = f_x \quad \frac{\partial f}{\partial y} = f_2 = f_y \quad \frac{\partial f}{\partial z} = f_3 = f_z$$

Position vector:

$$r = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$$

Unit vector with magnitude 1:

$$r = \mathbf{i} + \mathbf{j} + \mathbf{k}$$

6.2.1 Scalar field

$$f(x, y, z) = f_1(x, y, z) = f_2(x, y, z) = f_3(x, y, z)$$

The gradient of a scalar field is a vector field:

$$\nabla f = \text{grad } f(x, y, z) = f_x(x, y, z)\mathbf{i} + f_y(x, y, z)\mathbf{j} + f_z(x, y, z)\mathbf{k}$$

6.2.2 Field lines

6.2.3 Conervation field

If $\mathbf{F}(x, y, z) = \nabla\phi(x, y, z)$ in a domain D , then \mathbf{F} is a conservative vector field in D and function ϕ is the potential function.

$$\mathbf{F}(x, y, z) = \nabla\phi(x, y, z) = \phi_x(x, y, z)\mathbf{i} + \phi_y(x, y, z)\mathbf{j} + \phi_z(x, y, z)\mathbf{k}$$

If the vector field is conservative, then all the following equations are true:

$$\begin{aligned} \frac{\partial f_x}{\partial y} &= \frac{\partial f_y}{\partial x} \\ \frac{\partial f_x}{\partial z} &= \frac{\partial f_z}{\partial x} \\ \frac{\partial f_y}{\partial z} &= \frac{\partial f_z}{\partial y} \end{aligned}$$

6.2.4 Vector field in Polar Coordinates

$$\mathbf{F} = f(r, \theta) = f_r(r, \theta)\hat{\mathbf{r}} + f_\theta(r, \theta)\hat{\boldsymbol{\theta}}$$

where:

$$\begin{aligned} \hat{\mathbf{r}} &= \cos(\theta)\mathbf{i} + \sin(\theta)\mathbf{j} \\ \hat{\boldsymbol{\theta}} &= -\sin(\theta)\mathbf{i} + \cos(\theta)\mathbf{j} \end{aligned}$$

6.3 Line Integral

$$f(x, y)ds = \text{Area (tiny point)}$$

$$\text{Length of } \mathcal{C} = \int_{\mathcal{C}} f(x, y, z) ds = \int_a^b f(r(t)) \left| \frac{dr}{dt} \right| dt$$

6.3.1 Line integral of a vector field

$$W = \int_{\mathcal{C}} \mathbf{F} \cdot \hat{T} ds = \int F dr = \int_{\mathcal{C}} f_1(x, y, z)dx + f_2(x, y, z)dy + f_3(x, y, z)dz$$

6.4 Examples

6.4.1 Example 1: Conservative vector field and potential

Determine whether the given vector field is conservative, and find a potential function if it is:

$$\mathbf{F}(x, y, z) = (2xy - z^2)\mathbf{i} + (2yz + x^2)\mathbf{j} - (2zx - y^2)\mathbf{k}$$

6.4.2 Example 2: Line integral

Evaluate $\oint x^2y^2 dx + x^3y dy$ counterclockwise around the square with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$, and $(0, 1)$

6.4.3 Example 3: Line integral

Evaluate the line integral for $f(x, y) = x^2y^2$ along a straight line from origin to the point $(2, 1)$

Ans: $(5\sqrt{5})/3$

6.4.4 Example 4: Line integral vector field

Evaluate the line integral for $\mathbf{F}(x, y) = y^2\mathbf{i} + 2xy\mathbf{j}$ from $(0, 0)$ to $(1, 1)$

Along the line $y = x$

Along the line $y = x^2$

6.4.5 Example 5: Gradient

6.4.6 Example 6: Parametrize a curve

Use $t = y$ to parametrize the part of the line of intersection of the two planes:

Plane 1: $y = 2x - 4$

Plane 2: $z = 3x + 1$ from $(2, 0, 7)$ to $(3, 2, 10)$

Ans: $r(t) = \left(\frac{1}{2}(t+4)\right)\mathbf{i} + (t)\mathbf{j} + \left(\frac{3}{2}t+7\right)\mathbf{k}$

7 Surface-Integrals

7.1 Parametric Surface

For curve parametrization:

$$r = r(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k} \quad \text{where } a \leq t \leq b$$

For surface parametrization:

$$r = r(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k} \quad \text{where } a \leq u \leq b, \quad c \leq v \leq d$$

7.2 Surface Area

For a surface the area is given by:

$$\iint_S f(x, y, z) dS$$
$$dS = \left| \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} \right| du dv = \sqrt{\left(\frac{\partial(y, z)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(z, x)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(x, y)}{\partial(u, v)} \right)^2} du dv$$

For a parametrized surface S given by $r = r(u, v)$, where (u, v) is in the domain D in the uv -plane, the surface area is given by:

$$\iint_S f dS = \iint_D f(r(u, v)) \left| \frac{\partial r}{\partial u} \times \frac{\partial r}{\partial v} \right| du dv$$
$$= \iint_D f(x(u, v), y(u, v), z(u, v)) \sqrt{\left(\frac{\partial(y, z)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(z, x)}{\partial(u, v)} \right)^2 + \left(\frac{\partial(x, y)}{\partial(u, v)} \right)^2} du dv$$

For a surface S given by $z = g(x, y)$, where (x, y) is in the domain D in the xy -plane, the surface area is given by:

$$\iint_S f(x, y, z) dS = \iint_D f(x, y, z(x, y)) \sqrt{1 + \left(\frac{\partial g(x, y)}{\partial x} \right)^2 + \left(\frac{\partial g(x, y)}{\partial y} \right)^2} dx dy$$

The projection of normal vector onto the xy -plane is given by:

$$\cos(\gamma) = \frac{1}{\sqrt{1 + \left(\frac{\partial g(x, y)}{\partial x} \right)^2 + \left(\frac{\partial g(x, y)}{\partial y} \right)^2}} \quad \text{hence } dS = \frac{1}{\cos(\gamma)} dx dy$$

7.3 Oriented Surface

- A smooth surface S in 3-space is said to be orientable if there exists a unit vector field $\hat{N}(P)$.
- $\hat{N}(P)$ defined on S that varies continuously as P ranges over S and that is everywhere normal to S .
- Any such vector field $\hat{N}(P)$ determines an orientation of S .
- The oriented surface must have two sides.
- $\hat{N}(P)$ can have only one value at each point P with two sides.

7.4 Flux

$$\mathbf{F} = f_1(x, y, z)\mathbf{i} + f_2(x, y, z)\mathbf{j} + f_3(x, y, z)\mathbf{k}$$

Given any continuous vector field \mathbf{F} , flux of \mathbf{F} across the orientable surface S is integral of the normal component of \mathbf{F} over S

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S (\mathbf{F} \cdot \hat{\mathbf{N}}) dS$$

If the surface is closed, then the flux is given by:

$$\oiint_S \mathbf{F} \cdot d\mathbf{S} = \oiint_S (\mathbf{F} \cdot \hat{\mathbf{N}}) dS$$

If S is a parametrized surface given by $\mathbf{r} = \mathbf{r}(u, v)$, where (u, v) is in the domain D in the uv -plane, then the flux is given by:

$$\begin{aligned} \iint_S \mathbf{F} \cdot d\mathbf{S} &= \iint_D \mathbf{F} \cdot \left(\frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} \right) du dv \\ &= \iint_D \left(f_1 \frac{\partial(y, z)}{\partial(u, v)} + f_2 \frac{\partial(z, x)}{\partial(u, v)} + f_3 \frac{\partial(x, y)}{\partial(u, v)} \right) du dv \end{aligned}$$

For a surface S given by $z = g(x, y)$, where (x, y) is in the domain D in the xy -plane, the flux is given by:

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D \left(-f_1 \frac{\partial z}{\partial x} \mathbf{i} - f_2 \frac{\partial z}{\partial y} \mathbf{j} + f_3 \mathbf{k} \right) dx dy$$

7.5 Examples

8 Theorems

8.1 Differential Operators

8.1.1 Gradient

The gradient of a scalar field is a vector field that points in the direction of the steepest increase of the scalar field.

$$\begin{aligned}\text{grad } f(x, y, z) &= \nabla f(x, y, z) = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} + \frac{\partial f}{\partial z} \mathbf{k} \\ \mathbf{F}(x, y, z) &= f_x(x, y, z) \mathbf{i} + f_y(x, y, z) \mathbf{j} + f_z(x, y, z) \mathbf{k}\end{aligned}$$

8.1.2 Divergence

The divergence of a velocity field represents the net flow of fluid out of a small volume in a scalar field.

$$\text{div } \mathbf{F}(x, y, z) = \nabla \cdot \mathbf{F}(x, y, z) = \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} + \frac{\partial f_z}{\partial z}$$

8.1.3 Curl

The curl or field circulation of the electric field gives the rate of change of the magnetic field.

$$\begin{aligned}\text{curl } \mathbf{F}(x, y, z) &= \nabla \times \mathbf{F}(x, y, z) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ f_x & f_y & f_z \end{vmatrix} \\ &= \left(\frac{\partial f_z}{\partial y} - \frac{\partial f_y}{\partial z} \right) \mathbf{i} + \left(\frac{\partial f_x}{\partial z} - \frac{\partial f_z}{\partial x} \right) \mathbf{j} + \left(\frac{\partial f_y}{\partial x} - \frac{\partial f_x}{\partial y} \right) \mathbf{k}\end{aligned}$$

8.2 Green's Theorem

Let R be a regular, closed region in the xy -plane whose boundary, C , consists of one or more piecewise smooth, simple closed curves that are positively oriented with respect to R .

$$\oint_C f_1(x, y) dx + f_2(x, y) dy = \iint_R \left(\frac{\partial f_2}{\partial x} - \frac{\partial f_1}{\partial y} \right) dA$$

8.3 Stokes' Theorem

Let S be a piecewise smooth, oriented surface in 3-space, having unit normal field \hat{N} and boundary C consisting of one or more piecewise smooth, closed curves with orientation inherited from S .

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \text{curl } \mathbf{F} \cdot \hat{N} dS$$

8.4 Divergence Theorem

Let S , be a closed piecewise smooth surface, which is the boundary of V with normal \hat{N} pointing outwards.

$$\oint_s (\mathbf{F} \cdot \hat{N}) dS = \iiint_V \text{div } \mathbf{F} dV$$

More variants:

$$\begin{aligned}\iiint_D \text{curl } \mathbf{F} dV &= - \oint_s (\mathbf{F} \times \hat{N}) dS \\ \iiint_D \text{grad } \phi dV &= \oint_s \phi dS\end{aligned}$$

8.5 Examples

8.5.1 Example 1: Div and Curl

Calculate the divergence and curl of the following vector field:

$$\mathbf{F} = \cos x \mathbf{i} - \sin y \mathbf{j} + z \mathbf{k}.$$

Divergence: Curl:

8.5.2 Example 2: Green's Theorem

Using Green's Theorem evaluate $\oint_C (x^2y) dx + (xy^2) dy$, clockwise boundary of the region:

$$0 \leq y \leq \sqrt{9 - x^2}$$

8.5.3 Example 3: Stokes' Theorem

8.5.4 Example 4: Divergence Theorem

Use the Divergence Theorem to calculate the flux of the given vector field out of the sphere s with equation $x^2 + y^2 + z^2 = a^2$, where $a > 0$ and

$$\mathbf{F} = (x^2 + y^2)\mathbf{i} + (y^2 - z^2)\mathbf{j} + z\mathbf{k}$$

9 PDE

Partial Differential Equations are equations with multiple variables and derivatives. They are used to model many physical phenomena, such as heat, sound, and light. The totality of solutions to a PDE is called its general solution, and there can be a lot.

9.1 Classification of PDEs

General representation of a PDE:

$$A \frac{\partial^2 u}{\partial x^2} + B \frac{\partial^2 u}{\partial x \partial y} + C \frac{\partial^2 u}{\partial y^2} + D \frac{\partial u}{\partial x} + E \frac{\partial u}{\partial y} + Fu = G$$

$$Au_{xx} + Bu_{xy} + Cu_{yy} + Du_x + Eu_y + Fu = G$$

Conditions:

Linear: A, B, C, D, E, F are only function of x, y variables, not u .

Quasi-linear: A, B, C, D, E, F may be function of (x, y, u, u_x, u_y)

Fully non-linear: A, B, C, D, E, F may be function of $(x, y, u, u_x, u_y, u_{xx}, u_{yy}, u_{xy})$

9.2 Characteristics of PDEs

$B^2 - 4AC > 0$ 2 real roots 2 characteristics **Hyperbolic PDE**

$B^2 - 4AC = 0$ 1 real roots 1 characteristics **Parabolic PDE**

$B^2 - 4AC < 0$ 0 real roots 0 characteristics **Elliptic PDE**

Types of various PDEs:

Wave Equation: Hyperbolic PDE

Heat Equation: Parabolic PDE

Laplace Equation: Elliptic PDE

$B^2 - 4AC > 0$ 2 real roots 2 characteristics **Hyperbolic PDE**

$B^2 - 4AC = 0$ 1 real roots 1 characteristics **Parabolic PDE**

$B^2 - 4AC < 0$ 0 real roots 0 characteristics **Elliptic PDE**

9.3 Important Second-Order PDEs

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \text{One-dimensional wave equation}$$

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \text{One-dimensional heat equation}$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad \text{Two-dimensional Laplace equation}$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f(x, y) \quad \text{Two-dimensional Poisson equation}$$

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) \quad \text{Two-dimensional wave equation}$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = 0 \quad \text{Three-dimensional Laplace equation}$$

9.4 Initial and Boundary Conditions

9.5 Wave Equation (1D)

One dimensional wave equation is given by:

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \quad \text{where } c^2 = \left[\frac{T(x, t)}{\mu_x} \right] \quad (1)$$

With two boundary conditions $x = 0$ and $x = L$:

$$u(0, t) = 0 \quad u(L, t) = 0 \quad \text{For all } t \geq 0$$

And two initial conditions, initial displacement and initial velocity at time $t = 0$:

$$u(x, 0) = f(x) \quad \frac{\partial u}{\partial t}(x, 0) = g(x) \quad \text{For all } 0 \leq x \leq L$$

Steps to solve:

1. Method of Separation of Variables $u(x, t) = X(x)T(t)$
2. Satisfy the Boundary Conditions test
3. Fourier Series Validation

9.5.1 D'Alembert's Solution of the Wave Equation

His solution is given by eq. (1) but extended to two variables:

$$v = (x - ct) \quad w = (x + ct) \quad (2)$$

I.e. $u(v, w)$. Partial derivatives from chain rule:

$$u_x = u_v \cdot v_x + u_w \cdot w_x = u_v + u_w$$

For double derivatives:

$$u_{xx} = (u_v + u_w)_x = (u_v + u_w)_v v_x + (u_v + u_w)_w w_x = u_{vv} + 2u_{vw} + u_{ww}$$

With respect to t :

$$u_{tt} = c^2 u_{xx} = c^2(u_{vv} + 2u_{vw} + u_{ww})$$

From eq. (1) and eq. (2):

$$u_{vw} = \frac{\partial^2 u}{\partial w \partial v} = 0$$

This can be solved by integrating with respect to v and w :

$$\frac{\partial u}{\partial v} = h(v) \text{ and } u = \int h(v) dv + \psi(w)$$

Here, $h(v)$ and $\psi(w)$ are arbitrary functions of v and w , respectively. The solution in term for x :

$$u = \phi(v) + \psi(w)$$

This is d'Alembert's solution, which is the general solution to the wave equation.

$$u(x, t) = \phi(x + ct) + \psi(x - ct)$$

This solution satisfies the wave equation and the initial conditions:

9.6 Heat Equation (1D)

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \quad (3)$$

Conditions:

- PDE is linear and homogeneous.
- Boundary conditions are linear and homogeneous. The two for $u(x, t)$ is $u(0, t) = 0$ and $u(L, t) = 0$ for all $t > 0$.
- One initial condition at time ($t = 0$): $u(x, 0) = f(x)$.

Solve the

Solution:

$$u(x, t) = \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right) e^{-\lambda_n^2 t}$$

where

$$\lambda_n = \frac{cn\pi}{L}$$

and

$$B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx \quad \text{for } n = 1, 2, 3, \dots$$

9.7 Examples

9.7.1