

EXPONENTIAL SUM ESTIMATES USING VINOGRADOV MEAN VALUE THEOREM

Mathias Garnier

Supervised by Julia Brandes

March 3 to May 30, 2025

Contents

1. **Motivations**

1.1 Exponential sums, plots

1.2 Some applications

2. **The classical theory in a nutshell**

3. **Improvements à la Heath-Brown**

3.1 The Vinogradov Mean Value Theorem

3.2 Improved k -th derivative estimate

3.3 New improvements, $k = 4$

3.4 Growth rate of the Riemann zeta function on vertical lines

4. **What's next?**

1. – Motivations

1.1 – Exponential sums

Object of interest:

$$\sum_{n \in I} e(f(n)) := \sum_{n \in I} \exp(2i\pi f(n))$$

for $I \subset \mathbf{R}$ a compact interval and $f : I \rightarrow \mathbf{R}$ a *nice* function.

1.1 – Exponential sums

Object of interest:

$$\sum_{n \in I} e(f(n)) := \sum_{n \in I} \exp(2i\pi f(n))$$

for $I \subset \mathbf{R}$ a compact interval and $f : I \rightarrow \mathbf{R}$ a *nice* function.

Objective: bound it non-trivially, i.e. get better estimates than

$$\sum_{n \in I} e(f(n)) \ll \#I.$$

Vinogradov notation: $f(x) \ll g(x)$ if and only if $f(x) = \mathcal{O}(g(x))$.

1.1 – Exponential sums

Object of interest:

$$\sum_{n \in I} e(f(n)) := \sum_{n \in I} \exp(2i\pi f(n))$$

for $I \subset \mathbf{R}$ a compact interval and $f : I \rightarrow \mathbf{R}$ a *nice* function.

Objective: bound it non-trivially, i.e. get better estimates than

$$\sum_{n \in I} e(f(n)) \ll \#I.$$

Vinogradov notation: $f(x) \ll g(x)$ if and only if $f(x) = \mathcal{O}(g(x))$.

Is it a regular object ? symmetric ? an erratic one ? smooth ?
diffuse or concentrated ? ...

1.1 – Exponential sums

Let $\Sigma_N = \sum_{n=1}^N e(f(n))$.

1.1 – Exponential sums

Let $\Sigma_N = \sum_{n=1}^N e(f(n))$.





Figure: Scatter plot $\{(\Re(\Sigma_i), \Im(\Sigma_i))\}_{i=1, \dots, 100'000}$ for different f .

1.1 – Exponential sums

Other representations.

1.1 – Exponential sums

Other representations.




Figure: Magnitude of $\sum_{n=1}^N e(an^2 + bn)$ with $-2 \leq a, b \leq 2$,
 $N = 5, 10, 25$.

1.1 – Exponential sums

Obviously not restricted to polynomials.

1.1 – Exponential sums

Obviously not restricted to polynomials.




Figure: Plots of exponential sums with $f(x) = \log(x)^i$ for $i = 1, \dots, 6$.

1.2 – Some applications

1.2 – Some applications

- Bound on the Riemann zeta function

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s} \ll_{\varepsilon} \log(2 + |t|) \sup_{1 \leq M \leq N \ll |t|} N^{1-\sigma} \left| \frac{1}{N} \sum_{N \leq n < N+M} e\left(-\frac{t}{2\pi} \log n\right) \right|$$

for $s = \sigma + it$ with $0 < \varepsilon \leq \sigma \ll 1$ and $1 \ll |t|$.

1.2 – Some applications

- Bound on the Riemann zeta function

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s} \ll_{\varepsilon} \log(2 + |t|) \sup_{1 \leq M \leq N \ll |t|} N^{1-\sigma} \left| \frac{1}{N} \sum_{N \leq n < N+M} e\left(-\frac{t}{2\pi} \log n\right) \right|$$

for $s = \sigma + it$ with $0 < \varepsilon \leq \sigma \ll 1$ and $1 \ll |t|$.

- Lindelöf hypothesis: let

$$\mu(\sigma) = \inf\{\xi \in \mathbf{R}_+ : \zeta(\sigma + it) = \mathcal{O}(|t|^{\xi})\}$$

it is conjectured that $\mu(1/2) = 0$ (today's best: Bourgain (2017), $\mu(1/2) \leq 13/84$).

1.2 – Some applications

- Bound on the Riemann zeta function

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s} \ll_{\varepsilon} \log(2 + |t|) \sup_{1 \leq M \leq N \ll |t|} N^{1-\sigma} \left| \frac{1}{N} \sum_{N \leq n < N+M} e\left(-\frac{t}{2\pi} \log n\right) \right|$$

for $s = \sigma + it$ with $0 < \varepsilon \leq \sigma \ll 1$ and $1 \ll |t|$.

- Lindelöf hypothesis: let

$$\mu(\sigma) = \inf\{\xi \in \mathbf{R}_+ : \zeta(\sigma + it) = \mathcal{O}(|t|^{\xi})\}$$

it is conjectured that $\mu(1/2) = 0$ (today's best: Bourgain (2017), $\mu(1/2) \leq 13/84$).

Using the theory of exponential sums, one can show that

$$\forall \varepsilon > 0, \zeta(11/15 + it) \ll_{\varepsilon} |t|^{1/15+\varepsilon}.$$

As a consequence, $\mu(11/15) \leq 1/15$.

1.2 – Some applications

- Bound on the Riemann zeta function

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s} \ll_{\varepsilon} \log(2 + |t|) \sup_{1 \leq M \leq N \ll |t|} N^{1-\sigma} \left| \frac{1}{N} \sum_{N \leq n < N+M} e\left(-\frac{t}{2\pi} \log n\right) \right|$$

for $s = \sigma + it$ with $0 < \varepsilon \leq \sigma \ll 1$ and $1 \ll |t|$.

- Lindelöf hypothesis: let

$$\mu(\sigma) = \inf\{\xi \in \mathbf{R}_+ : \zeta(\sigma + it) = \mathcal{O}(|t|^{\xi})\}$$

it is conjectured that $\mu(1/2) = 0$ (today's best: Bourgain (2017), $\mu(1/2) \leq 13/84$).

Using the theory of exponential sums, one can show that

$$\forall \varepsilon > 0, \zeta(11/15 + it) \ll_{\varepsilon} |t|^{1/15+\varepsilon}.$$

As a consequence, $\mu(11/15) \leq 1/15$. **Improvements?**

2. – The classical theory in a nutshell

2. – The classical theory in a nutshell (tricks on sums)

Objective: estimate exponential sums.

2. – The classical theory in a nutshell (tricks on sums)

Objective: estimate exponential sums.

Theorem (van der Corput, $k = 2$)

Suppose f is a real valued function and has two continuous derivatives on an interval I . Suppose further that there exists λ_2 and $A \geq 1$ such that $0 < \lambda_2 \leq |f^{(2)}(x)| \leq A\lambda_2$ on I . Then

$$\sum_{n \in I} e(f(n)) \ll A|I|\lambda_2^{1/2} + \lambda_2^{-1/2}.$$

2. – The classical theory in a nutshell (tricks on sums)

Objective: estimate exponential sums.

Theorem (van der Corput, $k = 2$)

Suppose f is a real valued function and has two continuous derivatives on an interval I . Suppose further that there exists λ_2 and $A \geq 1$ such that $0 < \lambda_2 \leq |f^{(2)}(x)| \leq A\lambda_2$ on I . Then

$$\sum_{n \in I} e(f(n)) \ll A|I|\lambda_2^{1/2} + \lambda_2^{-1/2}.$$

Theorem (Classical k -th derivative estimate)

Let $N \in \mathbf{N}$, $k \geq 2$ an integer and suppose that $f(x) : [0, N] \rightarrow \mathbf{R}$ has continuous derivatives of order up to k on $(0, N)$. Suppose further that there exists λ_k and $A \geq 1$ such that $0 < \lambda_k \leq |f^{(k)}(x)| \leq A\lambda_k$ on $(0, N)$. Then

$$\sum_{n \leq N} e(f(n)) \ll A^{2^{2-k}} N \lambda_k^{1/(2^k-2)} + N^{1-2^{2-k}} \lambda_k^{-1/(2^k-2)}.$$

Can we do better ?

3. – Improvements à la Heath-Brown

3. – Improvements à la Heath-Brown

Plan.

- Use Vinogradov Mean Value Theorem (VMVT), count number of solutions of systems of polynomial equations.
- Improve the k -th derivative estimate.
- Improved improvements for $k = 4$.
- Application: study the growth rate of the Riemann zeta function ζ on vertical lines.

3.1 – The Vinogradov Mean Value Theorem

In the 40s and 50s, Vinogradov developed a method.

3.1 – The Vinogradov Mean Value Theorem

In the 40s and 50s, Vinogradov developed a method. To bound exponential sums, he introduced

$$J_{s,k}(X) = \int_{[0,1]^k} \left| \sum_{n \leq X} e(\alpha_1 n + \cdots + \alpha_k n^k) \right|^{2s} d\alpha.$$

3.1 – The Vinogradov Mean Value Theorem

In the 40s and 50s, Vinogradov developed a method. To bound exponential sums, he introduced

$$J_{s,k}(X) = \int_{[0,1]^k} \left| \sum_{n \leq X} e(\alpha_1 n + \cdots + \alpha_k n^k) \right|^{2s} d\alpha.$$

In the 70s, the problem was reduced to count solutions of system of polynomial equations modulo prime powers. Pave the way for p -adic methods (Linnik, Karatsuba, Stetchkin, Wooley).

3.1 – The Vinogradov Mean Value Theorem

In the 40s and 50s, Vinogradov developed a method. To bound exponential sums, he introduced

$$J_{s,k}(X) = \int_{[0,1]^k} \left| \sum_{n \leq X} e(\alpha_1 n + \cdots + \alpha_k n^k) \right|^{2s} d\alpha.$$

In the 70s, the problem was reduced to count solutions of system of polynomial equations modulo prime powers. Pave the way for p -adic methods (Linnik, Karatsuba, Stetchkin, Wooley). \implies **VMVT**.

3.1 – The Vinogradov Mean Value Theorem

Theorem (VMVT)

For all integers $s, k \geq 1$,

$$J_{s,k}(X) \ll_{s,k,\varepsilon} X^\varepsilon \left(X^s + X^{2s-k(k+1)/2} \right)$$

for all $X \geq 1$ and every $\varepsilon > 0$.

3.1 – The Vinogradov Mean Value Theorem

Theorem (VMVT)

For all integers $s, k \geq 1$,

$$J_{s,k}(X) \ll_{s,k,\varepsilon} X^\varepsilon \left(X^s + X^{2s-k(k+1)/2} \right)$$

for all $X \geq 1$ and every $\varepsilon > 0$.

In the 2010s, Wooley and Bourgain, Demeter, Guth independently proved the Vinogradov Mean Value Theorem.

3.1 – The Vinogradov Mean Value Theorem

Theorem (VMVT)

For all integers $s, k \geq 1$,

$$J_{s,k}(X) \ll_{s,k,\varepsilon} X^\varepsilon \left(X^s + X^{2s-k(k+1)/2} \right)$$

for all $X \geq 1$ and every $\varepsilon > 0$.

In the 2010s, Wooley and Bourgain, Demeter, Guth independently proved the Vinogradov Mean Value Theorem.

Proof.

Nested efficient congruencing *and* / *or* decoupling theory. □

3.2 – Improved k -th derivative estimate

Recall that, under appropriate hypotheses, the classical k -th derivative estimate gives

$$\sum_{n \leq N} e(f(n)) \ll_{A,k} N \lambda_k^{1/(2^k-2)} + N^{1-2^{2-k}} \lambda_k^{-1/(2^k-2)}.$$

3.2 – Improved k -th derivative estimate

Recall that, under appropriate hypotheses, the classical k -th derivative estimate gives

$$\sum_{n \leq N} e(f(n)) \ll_{A,k} N \lambda_k^{1/(2^k-2)} + N^{1-2^{2-k}} \lambda_k^{-1/(2^k-2)}.$$

Using VMVT, Heath-Brown gets

$$\sum_{n \leq N} e(f(n)) \ll_{A,k,\varepsilon} N^{1+\varepsilon} \left(\lambda_k^{1/(k(k-1))} + N^{-1/(k(k-1))} + N^{-2/(k(k-1))} \lambda_k^{-2/(k^2(k-1))} \right).$$

3.2 – Improved k -th derivative estimate

Recall that, under appropriate hypotheses, the classical k -th derivative estimate gives

$$\sum_{n \leq N} e(f(n)) \ll_{A,k} N \lambda_k^{1/(2^k-2)} + N^{1-2^{2-k}} \lambda_k^{-1/(2^k-2)}.$$

Using VMVT, Heath-Brown gets

$$\sum_{n \leq N} e(f(n)) \ll_{A,k,\varepsilon} N^{1+\varepsilon} \left(\lambda_k^{1/(k(k-1))} + N^{-1/(k(k-1))} + N^{-2/(k(k-1))} \lambda_k^{-2/(k^2(k-1))} \right).$$

3.2 – Improved k -th derivative estimate

Recall that, under appropriate hypotheses, the classical k -th derivative estimate gives

$$\sum_{n \leq N} e(f(n)) \ll_{A,k} N \lambda_k^{1/(2^k-2)} + N^{1-2^{2-k}} \lambda_k^{-1/(2^k-2)}.$$

Using VMVT, Heath-Brown gets

$$\sum_{n \leq N} e(f(n)) \ll_{A,k,\varepsilon} N^{1+\varepsilon} \left(\lambda_k^{1/(k(k-1))} + N^{-1/(k(k-1))} + N^{-2/(k(k-1))} \lambda_k^{-2/(k^2(k-1))} \right).$$

3.2 – Improved k -th derivative estimate

Recall that, under appropriate hypotheses, the classical k -th derivative estimate gives

$$\sum_{n \leq N} e(f(n)) \ll_{A,k} N \lambda_k^{1/(2^k-2)} + N^{1-2^{2-k}} \lambda_k^{-1/(2^k-2)}.$$

Using VMVT, Heath-Brown gets

$$\sum_{n \leq N} e(f(n)) \ll_{A,k,\varepsilon} N^{1+\varepsilon} \left(\lambda_k^{1/(k(k-1))} + N^{-1/(k(k-1))} + N^{-2/(k(k-1))} \lambda_k^{-2/(k^2(k-1))} \right).$$

3.2 – Improved k -th derivative estimate

Recall that, under appropriate hypotheses, the classical k -th derivative estimate gives

$$\sum_{n \leq N} e(f(n)) \ll_{A,k} N \lambda_k^{1/(2^k-2)} + N^{1-2^{2-k}} \lambda_k^{-1/(2^k-2)}.$$

Using VMVT, Heath-Brown gets

$$\sum_{n \leq N} e(f(n)) \ll_{A,k,\varepsilon} N^{1+\varepsilon} \left(\lambda_k^{1/(k(k-1))} + N^{-1/(k(k-1))} + N^{-2/(k(k-1))} \lambda_k^{-2/(k^2(k-1))} \right)$$

if $f \in \mathcal{C}^k((0, N), \mathbf{R})$ and λ_k, A s.t. $0 < \lambda_k \leq |f^{(k)}(x)| \leq A \lambda_k$ on $(0, N)$.

3.3 – New improvements, $k = 4$

Fix $k = 4$, let $\lambda_4 = N^{-\varpi}$. Consider the regime $N^{-2} \leq \lambda_4 \leq N^{-1}$.

3.3 – New improvements, $k = 4$

Fix $k = 4$, let $\lambda_4 = N^{-\varpi}$. Consider the regime $N^{-2} \leq \lambda_4 \leq N^{-1}$.

Using VMVT type theorems (Demeter, Guth, Wang), Heath-Brown proved that

$$\sum_{n \leq N} e(f(n)) \ll_{A,\varepsilon} N^{1-\varpi/(4\varpi+8)+\varepsilon} + N^{8/9+\varepsilon}.$$

3.3 – New improvements, $k = 4$

Fix $k = 4$, let $\lambda_4 = N^{-\varpi}$. Consider the regime $N^{-2} \leq \lambda_4 \leq N^{-1}$.

Using VMVT type theorems (Demeter, Guth, Wang), Heath-Brown proved that

$$\sum_{n \leq N} e(f(n)) \ll_{A,\varepsilon} N^{1-\varpi/(4\varpi+8)+\varepsilon} + N^{8/9+\varepsilon}.$$

Using VMVT type theorems (Maldague, Oh), one gets better bounds on $J_{s,4-1}(N)$ and then better exponential sum estimates

$$\sum_{n \leq N} e(f(n)) \ll_{A,\varepsilon} N^{1-\varpi/(4\varpi+8)+\varepsilon} + N^{6/7+\varepsilon}.$$

3.3 – New improvements, $k = 4$

Fix $k = 4$, let $\lambda_4 = N^{-\varpi}$. Consider the regime $N^{-2} \leq \lambda_4 \leq N^{-1}$.

Using VMVT type theorems (Demeter, Guth, Wang), Heath-Brown proved that

$$\sum_{n \leq N} e(f(n)) \ll_{A,\varepsilon} N^{1-\varpi/(4\varpi+8)+\varepsilon} + N^{8/9+\varepsilon}.$$

Using VMVT type theorems (Maldague, Oh), one gets better bounds on $J_{s,4-1}(N)$ and then better exponential sum estimates

$$\sum_{n \leq N} e(f(n)) \ll_{A,\varepsilon} N^{1-\varpi/(4\varpi+8)+\varepsilon} + N^{6/7+\varepsilon}.$$

Application?

3.4 – Growth rate of the Riemann zeta function on vertical lines

Recall $\mu(\sigma) = \inf\{\xi \in \mathbf{R}_+ : \zeta(\sigma + it) = \mathcal{O}(|t|^\xi)\}$.




Figure: Critical strip, $\sigma = 1/2$. Source: hdd23.com

3.4 – Growth rate of the Riemann zeta function on vertical lines

Heath-Brown interest: $\sigma = 11/15$. He proved that

$$\forall \varepsilon > 0, \zeta(11/15 + it) \ll_{\varepsilon} |t|^{1/15 + \varepsilon}.$$

3.4 – Growth rate of the Riemann zeta function on vertical lines

Heath-Brown interest: $\sigma = 11/15$. He proved that

$$\forall \varepsilon > 0, \zeta(11/15 + it) \ll_{\varepsilon} |t|^{1/15+\varepsilon}.$$

Proof idea: sufficient to prove that

$$\sum_{N < n \leq 2N} n^{it} \ll_{\varepsilon} N^{11/15} t^{1/15+\varepsilon}$$

for every $1 = t^0 \leq N \leq t^{1/2}$. Then, use five different estimates on five different subintervals for N in

$$[t^0, t^{1/4}], [t^{1/4}, t^{1/3}], [t^{1/3}, t^{5/12}], [t^{5/12}, t^{3/7}], [t^{3/7}, t^{1/2}].$$

3.4 – Growth rate of the Riemann zeta function on vertical lines

Generalize Heath-Brown method.

3.4 – Growth rate of the Riemann zeta function on vertical lines

Generalize Heath-Brown method. Under technical assumptions on σ_0 and α , we have

$$\forall \varepsilon > 0, \zeta(\sigma_0 + it) \ll_{\varepsilon} |t|^{\alpha + \varepsilon}.$$

3.4 – Growth rate of the Riemann zeta function on vertical lines

Generalize Heath-Brown method. Under technical assumptions on σ_0 and α , we have

$$\forall \varepsilon > 0, \zeta(\sigma_0 + it) \ll_{\varepsilon} |t|^{\alpha+\varepsilon}.$$

In particular

$$\forall \varepsilon > 0, \zeta(11/15 + it) \ll_{\varepsilon} |t|^{11/180+\varepsilon}.$$

Hence $\mu(11/15) \leq 11/180 \approx 0.0611 < 1/15 \approx 0.0666$.

3.4 – Growth rate of the Riemann zeta function on vertical lines

Generalize Heath-Brown method. Under technical assumptions on σ_0 and α , we have

$$\forall \varepsilon > 0, \zeta(\sigma_0 + it) \ll_{\varepsilon} |t|^{\alpha+\varepsilon}.$$

In particular

$$\forall \varepsilon > 0, \zeta(11/15 + it) \ll_{\varepsilon} |t|^{11/180+\varepsilon}.$$

Hence $\mu(11/15) \leq 11/180 \approx 0.0611 < 1/15 \approx 0.0666$.

Proof idea: Optimize Heath-Brown procedure and use previous improvements. Sufficient to prove that

$$\sum_{N < n \leq 2N} n^{it} \ll_{\varepsilon} N^{\sigma_0} t^{\alpha+\varepsilon}.$$

3.4 – Growth rate of the Riemann zeta function on vertical lines

Proof idea (continued). Sufficient to prove that

$$\sum_{N < n \leq 2N} n^{it} \ll_{\varepsilon} N^{\sigma_0} t^{\alpha + \varepsilon}.$$

- it holds trivially for $t^0 \leq N \leq t^{11/48}$,
- by fourth derivative estimates, it holds for $t^{11/48} \leq N \leq t^{1/3}$,
- by the 6/7 improvement, it holds for $t^{1/3} \leq N \leq t^{1/2}$.

This "concludes" the "proof".

4. – **What's next?**

4. – What's next?

- Density Functional Theory and exponential sums.

See Next-order correction to the Dirac exchange energy of the free electron gas in the thermodynamic limit and generalized gradient approximations by T. C.

Corso and G. Friesecke.

4. – What's next?

- Density Functional Theory and exponential sums.

See Next-order correction to the Dirac exchange energy of the free electron gas in the thermodynamic limit and generalized gradient approximations by T. C.

Corso and G. Friesecke.

- Study ℓ^2 -decoupling and more generally harmonic analysis (especially the link with PDEs).

4. – What's next?

- Density Functional Theory and exponential sums.

See Next-order correction to the Dirac exchange energy of the free electron gaz in the thermodynamic limit and generalized gradient approximations by T. C.

Corso and G. Friesecke.

- Study ℓ^2 -decoupling and more generally harmonic analysis (especially the link with PDEs).
- Continue to discover analytic number theory with Julia Brandes, Régis de la Bretèche and, why not, be introduced to probabilistic number theory...

Thanks!

Any questions?