

# The Goldbach–Chen Frontier: How the Semiprime Channel Inherits the Singular Series

Mathias Rodríguez Castro

Working draft — June 17, 2026

## Abstract

The binary Goldbach conjecture asks whether every even  $N > 2$  is a sum of two primes; Chen’s theorem replaces the second prime by a prime or semiprime. We study how the Goldbach count  $R_1(N)$  (prime  $q$ ) and the Chen-surplus count  $R_2(N)$  (semiprime  $q$ ) respond to the Hardy–Littlewood singular series  $\mathfrak{S}(N)$ . Decomposing the semiprime channel  $q = rs$  by its small prime factor  $r$ , each channel carries its own two-form singular series, and in the ratio  $R_2/R_1$  the singular series *cancels* (an unconditional Euler-product identity), leaving a Mertens channel sum  $\widetilde{W}(N) \sim \log \log N$ . Thus  $\mathfrak{S}(N)$  enters  $R_2$  with exponent 1, exactly as in  $R_1$ :  $R_2$  inherits the full singular series, and the apparent “half power” reported by finite-range log–log fits is not an exponent but a regression *slope* of a non-power-law object, obeying  $\beta_2(N) = 1 - c(N)/\langle R_2/R_1 \rangle \rightarrow 1$  with deficit  $\sim 1/\log \log N$ . A circle-method second-moment bound—only Vaughan’s classical *prime* minor-arc estimate and the trivial  $L^2$  size of the semiprime sum, no sieve—gives the *linear* almost-all asymptotic  $R_2(N) = \text{main}(N)(1 + o(1))$ ; the passage to the log-regression statement  $\beta_2 \rightarrow 1$  is isolated as one explicitly stated lower-tail hypothesis rather than overclaimed. The goal is not a proof of Goldbach but a clean new identity, reproducible evidence, and a bounded reduction; a gallery of cross-disciplinary reframings, all governed by the same  $\mathfrak{S}(N)$ , is collected in an appendix.

## Contents

|          |                                                                     |          |
|----------|---------------------------------------------------------------------|----------|
| <b>1</b> | <b>Introduction</b>                                                 | <b>2</b> |
| <b>2</b> | <b>Basic notation</b>                                               | <b>4</b> |
| <b>3</b> | <b>The Goldbach–Chen frontier</b>                                   | <b>4</b> |
| 3.1      | Chen surplus and Goldbach fragility . . . . .                       | 4        |
| 3.2      | Prime versus semiprime share . . . . .                              | 5        |
| <b>4</b> | <b>Distribution of the second summand <math>q</math></b>            | <b>5</b> |
| 4.1      | Location of $q$ inside $[0, N]$ . . . . .                           | 6        |
| 4.2      | Type-labeled empirical measures . . . . .                           | 6        |
| 4.3      | Factor geometry of semiprime $q$ . . . . .                          | 6        |
| <b>5</b> | <b>Heuristic asymptotics (classical recovery)</b>                   | <b>7</b> |
| <b>6</b> | <b>Selection formulations</b>                                       | <b>7</b> |
| <b>7</b> | <b>Empirical results and a new regularity</b>                       | <b>8</b> |
| 7.1      | Computational setup and validation . . . . .                        | 8        |
| 7.2      | The semiprime channel and the half-power mirage . . . . .           | 8        |
| 7.3      | The singular exponent across arithmetic-complexity layers . . . . . | 12       |
| 7.4      | Diagnostics . . . . .                                               | 13       |

|          |                                                                                |           |
|----------|--------------------------------------------------------------------------------|-----------|
| 7.5      | Optimization experiments . . . . .                                             | 13        |
| 7.6      | Weak continuity of the location measures . . . . .                             | 14        |
| 7.7      | The residual Goldbach–Chen balance exponent . . . . .                          | 15        |
| 7.8      | Three further probes: transport, topology, dynamics . . . . .                  | 16        |
| 7.9      | An empirical circle method: the spectrum of the comet . . . . .                | 18        |
| 7.10     | Primorial bases and carry geometry . . . . .                                   | 19        |
| <b>8</b> | <b>Conclusion</b>                                                              | <b>20</b> |
| <b>A</b> | <b>Towards a proof that <math>\beta_2 \rightarrow 1</math></b>                 | <b>20</b> |
| A.1      | Notation and the statistic . . . . .                                           | 21        |
| A.2      | Unconditional lemmas . . . . .                                                 | 21        |
| A.3      | The conditional theorem . . . . .                                              | 22        |
| A.4      | Confirmation and the unconditional model . . . . .                             | 22        |
| A.5      | A first-moment reduction: replacing pointwise HL by equidistribution . . . . . | 23        |
| A.6      | The variance bound and the exceptional set . . . . .                           | 25        |
| A.7      | Truncated logarithms and the blocked-channel model . . . . .                   | 27        |
| A.8      | The $\Omega$ -generating function and singular-series response . . . . .       | 29        |
| A.9      | Weak uniformity of the location measures . . . . .                             | 30        |
| A.10     | The least-unblocked-prime law and residual collapse . . . . .                  | 31        |
| A.11     | Local carry excess from prime residue restrictions . . . . .                   | 32        |
| A.12     | The major arcs: semiprimes in progressions . . . . .                           | 33        |
| <b>B</b> | <b>The infinite formulation and column generation</b>                          | <b>34</b> |
| <b>C</b> | <b>Supplementary reframings</b>                                                | <b>35</b> |
| C.1      | Goldbach as arithmetic ground-state energy . . . . .                           | 35        |
| C.2      | Adversarial robustness of the Goldbach graph . . . . .                         | 36        |
| C.3      | Goldbach as a market: liquidity and the fair-weather substitute . . . . .      | 36        |
| C.4      | Persistent topology of additive decompositions (a negative note) . . . . .     | 38        |
| C.5      | Goldbach as control: the value of arithmetic information . . . . .             | 38        |
| C.6      | The information content of the comet . . . . .                                 | 38        |
| C.7      | Discrete geometry of the Chen surface . . . . .                                | 39        |

## 1 Introduction

The binary Goldbach conjecture states that every even integer  $N > 2$  can be written as the sum of two primes. The conjecture remains open. However, several nearby facts are known. Helfgott proved the ternary, or weak, Goldbach conjecture: every odd integer greater than 5 is the sum of three primes [4]. Oliveira e Silva, Herzog and Pardi verified the binary Goldbach conjecture computationally up to  $4 \cdot 10^{18}$  [9]. Chen’s theorem proves that every sufficiently large even integer can be written as

$$N = p + q,$$

where  $p$  is prime and  $q$  is either prime or the product of at most two primes [1]; the semiprime channel and the “ $E_2$ ” numbers it counts are classical sieve objects [2], and for the broader additive-bases context we refer to Nathanson [8]. Recent sieve-theoretic work also studies intermediate statements between Chen’s theorem and the binary Goldbach conjecture [6].

This project focuses on the transition between the Goldbach representation

$$N = p + \text{prime}$$

and the Chen representation

$$N = p + \text{semiprime}.$$

The guiding question is not merely whether a representation exists, but how the available representations are distributed, how often the second term is genuinely prime, how often it is semiprime, and whether a representation can be selected in a smooth way as  $N$  varies.

**Main idea.** For each even  $N$ , define the number of representations with  $q$  prime and with  $q$  semiprime. Then study the empirical object

$$N \mapsto \{(p, q) : p + q = N, p \in \mathbb{P}, q \in \mathbb{P} \cup \mathcal{S}_2\}.$$

This is a discrete object, so it is not continuous in the ordinary analytic sense. Nevertheless, one can ask whether the induced empirical measures converge weakly, whether normalized counts have stable asymptotics, and whether a mixed-integer model can select one representation per even integer with small variation. This gives a precise mathematical language for the intuition that the representation may “jump” between prime and semiprime behavior.

**Summary of findings.** Sections 7–7.7 report the results of implementing this program; all numbers are reproducible from the accompanying code. The main outcomes are:

1. **The singular series of  $R_2$ , and the half-power mirage (§7.2).** Counts are computed for all  $N \leq 2 \cdot 10^6$  by FFT and locally in windows up to  $10^9$  by a block-convolution sieve. Both  $R_1$  and  $R_2$  carry  $\mathfrak{S}(N)$  with exponent 1: in the ratio  $R_2/R_1$  the singular series *cancels* (Lemma 1), leaving the Mertens channel sum  $\widetilde{W}(N) \sim \log \log N$ . The number  $\beta_2 \approx \frac{1}{2}$  that a finite-range log–log fit reports is therefore not an exponent but a regression slope of a non-power-law object; a derivation and the data agree on  $\beta_2(N) = 1 - c/\langle R_2/R_1 \rangle_N$  with  $\langle R_2/R_1 \rangle \sim \log \log N$ , so  $\beta_2 \rightarrow 1$  with an  $\mathcal{O}(1/\log \log N)$  deficit (validation  $(1 - \beta_2)\langle R_2/R_1 \rangle \approx 1.09$  to 1%). The slope is multi-channel: deleting the heaviest channel ( $3 \mid N$ ) removes only  $\sim 10\%$  of it (§7.2), refuting a single-atom reading.
2. **Layers and sign reversal (§7.3).** For  $\Omega(q) = k$ , the slope  $\beta_k$  falls through  $1, \frac{1}{2}, \dots$  and becomes negative for  $k \geq 3$ :  $\mathfrak{S}(N)$  enhances low-complexity and suppresses high-complexity channels.
3. **Diagnostics, continuity, optimization (§7.4–7.7).** Factor geometry favours unbalanced semiprimes; the  $q/N$  measures stabilize weakly to a near-uniform law; minimum-variation selection is a Pareto trade-off between geometric and type smoothness; the Chen-rescue threshold under a balance constraint is unimodal; and the residual balance exponent  $a_{\text{Res}}(N)$  correlates positively with  $\mathfrak{S}(N)$ .
4. **An almost-all theorem, honestly bounded (Appendix A).** The cancellation identity (Lemma 1) is unconditional; a circle-method second-moment bound—Vaughan’s classical *prime* minor-arc estimate plus the trivial  $L^2$  size of the semiprime sum, no sieve—gives the *linear* asymptotic  $R_2(N) = \text{main}(N)(1 + o(1))$  for all but  $o(X)$  even  $N$ , unconditional in outline. Upgrading this to the log-regression statement  $\beta_2(N) \rightarrow 1$  needs one further input (lower-tail control of  $R_2/R_1$ , beyond  $L^2$ ); we state that gap rather than claim past it.
5. **Many mirrors, one fact (§7.9–7.10; Appendix C).** The frontier can be recast as a thermodynamic ground state, an adversarial network, a liquidity market, a control problem, an empirical circle method, an information channel, and a discrete surface. We are explicit that this is *one* finding— $\mathfrak{S}(N)$  sets the local densities—observed many times, not many findings; most reframings are deterministic functions of the same  $\mathfrak{S}/R_2$  data and are collected in Appendix C (with the honest negatives: an empty additive-complexity topology and a Chen substitute with no resilience premium). Two earn a place in the body by yielding

a *new computation*: the empirical circle method (§7.9), whose comet power spectrum is literally the major arcs, and the carry geometry (§7.10), which surfaces a digit/Kummer signature of primality.

## 2 Basic notation

Let  $\mathbb{P}$  denote the set of primes. For an integer  $n \geq 2$ , let  $\Omega(n)$  be the number of prime factors of  $n$  counted with multiplicity. Thus  $\Omega(12) = 3$  because  $12 = 2^2 \cdot 3$ .

**Definition 1** (Semiprimes and Chen numbers). *Define*

$$\mathcal{S}_2 := \{n \geq 2 : \Omega(n) = 2\}$$

*to be the set of semiprimes, allowing squares of primes. Define also*

$$\mathcal{P}_2 := \{n \geq 2 : \Omega(n) \leq 2\} = \mathbb{P} \cup \mathcal{S}_2.$$

*Thus  $\mathcal{P}_2$  is the set of integers with at most two prime factors counted with multiplicity.*

For an even integer  $N$ , define the prime–prime representation count

$$R_1(N) := \sum_{p < N} \mathbf{1}_{\mathbb{P}}(p) \mathbf{1}_{\mathbb{P}}(N - p). \quad (1)$$

This counts ordered representations by choosing the first summand  $p$ . If unordered counts are desired, one may restrict to  $p \leq N/2$ .

Define the prime–semiprime representation count

$$R_2(N) := \sum_{p < N} \mathbf{1}_{\mathbb{P}}(p) \mathbf{1}_{\mathcal{S}_2}(N - p). \quad (2)$$

Finally, define the Chen count

$$R_{\leq 2}(N) := R_1(N) + R_2(N) = \sum_{p < N} \mathbf{1}_{\mathbb{P}}(p) \mathbf{1}_{\mathcal{P}_2}(N - p). \quad (3)$$

**Remark 1.** *The binary Goldbach conjecture is equivalent to  $R_1(N) > 0$  for every even  $N > 2$ . Chen’s theorem implies  $R_{\leq 2}(N) > 0$  for all sufficiently large even  $N$ .*

## 3 The Goldbach–Chen frontier

The central empirical object is the decomposition of Chen representations into genuinely Goldbach and genuinely semiprime representations:

$$R_{\leq 2}(N) = R_1(N) + R_2(N).$$

This suggests several diagnostics.

### 3.1 Chen surplus and Goldbach fragility

**Definition 2** (Chen surplus). *For even  $N$ , define the Chen surplus ratio*

$$C(N) := \frac{R_2(N)}{R_1(N) + 1}. \quad (4)$$

*Large values of  $C(N)$  identify integers for which semiprime representations are much more abundant than prime–prime representations.*

**Definition 3** (Goldbach fragility). *For a threshold  $k \in \mathbb{N}$ , define the set of  $k$ -fragile even integers up to  $X$  by*

$$\mathcal{F}_k(X) := \{N \leq X : N \equiv 0 \pmod{2}, R_1(N) \leq k, R_2(N) > 0\}. \quad (5)$$

*These are integers for which Goldbach representations exist only sparsely, while Chen-type representations remain available.*

**Research question.** How large is  $\mathcal{F}_k(X)$ ? Are fragile integers concentrated in specific congruence classes? Does fragility correlate with the singular series correction in the Hardy–Littlewood heuristic?

### 3.2 Prime versus semiprime share

Define

$$\theta(N) := \frac{R_1(N)}{R_1(N) + R_2(N)}, \quad (6)$$

with the convention  $\theta(N) = 0$  if the denominator is zero.

Here  $\theta(N)$  is the empirical share of Chen representations in which  $q$  is prime rather than semiprime. A small value of  $\theta(N)$  means that most Chen representations use a genuinely semiprime second summand.

A naive density heuristic suggests

$$R_1(N) \asymp \frac{N}{\log^2 N}, \quad R_2(N) \asymp \frac{N \log \log N}{\log^2 N},$$

because primes have local density roughly  $1/\log N$ , while semiprimes have density roughly  $\log \log N / \log N$ . Therefore, at a first heuristic level, semiprime representations should outnumber prime–prime representations by a factor of order  $\log \log N$ . This does not prove Goldbach or Chen, but it motivates the following empirical conjecture.

**Conjecture 1** (Average semiprime dominance). *There exists a slowly varying normalization  $L(X) \sim \log \log X$  such that*

$$\frac{1}{X/2} \sum_{\substack{N \leq X \\ N \equiv 0 \pmod{2}}} \frac{R_2(N)}{R_1(N) + 1} \asymp L(X).$$

*Equivalently, prime–semiprime representations should become increasingly abundant relative to prime–prime representations on average.*

## 4 Distribution of the second summand $q$

For each even  $N$ , define the representation sets

$$\mathcal{R}_1(N) := \{(p, q) : p + q = N, p \in \mathbb{P}, q \in \mathbb{P}\}, \quad (7)$$

$$\mathcal{R}_2(N) := \{(p, q) : p + q = N, p \in \mathbb{P}, q \in \mathcal{S}_2\}, \quad (8)$$

$$\mathcal{R}_{\leq 2}(N) := \mathcal{R}_1(N) \cup \mathcal{R}_2(N). \quad (9)$$

The object  $q = N - p$  can be studied in several ways.

#### 4.1 Location of $q$ inside $[0, N]$

Define empirical measures on  $[0, 1]$ :

$$\mu_N^{(1)} := \frac{1}{R_1(N)} \sum_{(p,q) \in \mathcal{R}_1(N)} \delta_{q/N}, \quad (10)$$

$$\mu_N^{(2)} := \frac{1}{R_2(N)} \sum_{(p,q) \in \mathcal{R}_2(N)} \delta_{q/N}, \quad (11)$$

$$\mu_N^{(\leq 2)} := \frac{1}{R_{\leq 2}(N)} \sum_{(p,q) \in \mathcal{R}_{\leq 2}(N)} \delta_{q/N}, \quad (12)$$

whenever the denominators are nonzero.

**Question 1** (Weak continuity). *Do the measures  $\mu_N^{(1)}$ ,  $\mu_N^{(2)}$ , or  $\mu_N^{(\leq 2)}$  converge weakly along typical even  $N$ ? If so, is the limit close to the uniform distribution on  $(0, 1)$  after removing small endpoint effects?*

This is a precise version of the intuition that the representation function might become “continuous” after smoothing. The pointwise object is discontinuous, because primality is discrete; but the normalized cloud of representations may stabilize in distribution.

#### 4.2 Type-labeled empirical measures

Let the type of  $q$  be

$$\tau(q) := \begin{cases} 1, & q \in \mathbb{P}, \\ 2, & q \in \mathcal{S}_2. \end{cases}$$

Define a type-labeled empirical measure on  $[0, 1] \times \{1, 2\}$  by

$$\nu_N := \frac{1}{R_{\leq 2}(N)} \sum_{(p,q) \in \mathcal{R}_{\leq 2}(N)} \delta_{(q/N, \tau(q))}. \quad (13)$$

**Conjecture 2** (Weak type stabilization). *For typical large even  $N$ , the measures  $\nu_N$  are close to a product-like law*

$$\nu_N \approx \int f_N(x, t) dx d\tau,$$

where the spatial component in  $x = q/N$  varies slowly, while the type marginal increasingly favors  $t = 2$  because semiprimes have larger density than primes.

This is not ordinary continuity. It is closer to convergence of empirical distributions, or continuity after smoothing over windows  $N \in [X, X + H]$ .

#### 4.3 Factor geometry of semiprime $q$

If  $q \in \mathcal{S}_2$ , write

$$q = ab, \quad a, b \in \mathbb{P}, \quad a \leq b.$$

Then define the balance statistic

$$B(q) := \frac{\log a}{\log q} \in \left(0, \frac{1}{2}\right]. \quad (14)$$

A value near  $1/2$  means that  $q$  is a balanced semiprime; a value near  $0$  means that  $q$  has a very small prime factor.

**Question 2** (Semiprime factor geometry). *Among representations  $N = p + q$  with  $q \in \mathcal{S}_2$ , how is  $B(q)$  distributed? Are Chen representations dominated by unbalanced semiprimes with a small factor, or by balanced semiprimes?*

This question is interesting because Chen’s theorem is obtained by sieve methods. Empirically, one can ask whether the extra freedom in  $q$  comes mostly from allowing small prime divisors, or whether semiprime representations are broadly distributed across factor shapes.

## 5 Heuristic asymptotics (classical recovery)

The asymptotics in this section are classical; we recall them only to fix normalisation and as a control for the computations of §7, not as new derivations. The Hardy–Littlewood circle-method heuristic for Goldbach [3] predicts a main term of the form

$$R_1(N) \approx 2C_2 \mathfrak{S}(N) \frac{N}{\log^2 N}, \quad (15)$$

where  $C_2$  is the twin-prime constant and  $\mathfrak{S}(N) = \prod_{p|N, p>2} \frac{p-1}{p-2}$  is the singular-series correction depending on the odd prime divisors of  $N$  (the factor  $2C_2$  already absorbs the  $p = 2$  local density).

For semiprime  $q$ , Landau’s density of numbers with two prime factors [5],  $\#\{n \leq x : \Omega(n) = 2\} \sim x \log \log x / \log x$ , gives the rough estimate

$$\#\{n \leq x : \Omega(n) = 2\} \sim \frac{x \log \log x}{\log x}. \quad (16)$$

Therefore, a first-order independence heuristic gives

$$R_2(N) \approx K_2(N) \frac{N \log \log N}{\log^2 N}, \quad (17)$$

where  $K_2(N)$  is a correction factor capturing congruence constraints and local arithmetic structure.

**Normalization.** The empirical project should therefore compare

$$A_1(N) := \frac{R_1(N)}{N / \log^2 N}, \quad (18)$$

$$A_2(N) := \frac{R_2(N)}{N \log \log N / \log^2 N}. \quad (19)$$

If the heuristic is meaningful,  $A_1(N)$  and  $A_2(N)$  should be much more stable than the raw counts.

## 6 Selection formulations

*Scope note.* This section records the optimization *language* used by two diagnostics measured in §7.5 (a minimum-variation selection and a Chen-rescue threshold). It is deliberately short: the feasibility version is trivial and no MIP solver will prove Goldbach (Remark 2). The infinite-dimensional and column-generation readings are recorded in Appendix B; a reader after the empirical results may skip to §7.

Fix an even  $N \leq X$  and let  $\mathcal{A}_N^{(1)} = \{p \in \mathbb{P} : N - p \in \mathbb{P}\}$  and  $\mathcal{A}_N^{(2)} = \{p \in \mathbb{P} : N - p \in \mathcal{S}_2\}$ . Binary variables  $z_{N,p,t}$  select one representation  $N = p + q$  of type  $t$  (prime  $t = 1$ , semiprime

$t = 2$ ); the cover constraint  $\sum_{p \in \mathcal{A}_N^{(1)}} z_{N,p,1} + \sum_{p \in \mathcal{A}_N^{(2)}} z_{N,p,2} \geq 1$  for every even  $N$  in a window  $\mathcal{E}_X$  is Chen feasibility, and dropping the type-2 terms is Goldbach feasibility. With the sets  $\mathcal{A}_N^{(t)}$  precomputed by a sieve this is trivial as a feasibility problem; it becomes informative only under added structure.

**Remark 2** (Why this is not a proof strategy). *These mixed-integer programs do not make Goldbach easier by themselves. They translate the conjecture into a covering problem over a hypergraph whose vertices are even integers and whose hyperedges are prime-pair certificates; the value is conceptual, letting one import selection, smoothness, total variation, and column generation into the empirical study of Goldbach-type representations (Appendix B).*

**Minimum-variation selection.** Selecting one representation per even  $N$  in a window, with type  $u_N = \sum_{p \in \mathcal{A}_N^{(2)}} z_{N,p,2} \in \{0, 1\}$  and normalized location  $s_N = q_N/N = \sum_{p,t} \frac{N-p}{N} z_{N,p,t}$  (linear in the selection variables), the minimum-variation MIP minimizes  $\alpha \sum_N v_N + \beta \sum_N r_N + \gamma \sum_N |s_N - \frac{1}{2}|$  subject to one selection per  $N$  and  $v_N \geq |u_N - u_{N-2}|$ ,  $r_N \geq |s_N - s_{N-2}|$ . The term  $\sum v_N$  measures type oscillation,  $\sum r_N$  geometric oscillation, and  $|s_N - \frac{1}{2}|$  rewards balance. The guiding question is whether the minimum type-switch rate  $|\mathcal{E}_X|^{-1} \min \sum_N v_N$  decreases or stays bounded away from 0 as  $X \rightarrow \infty$ ; §7.5 reports the answer (a smooth-location branch that nonetheless keeps switching type).

**Chen-rescue threshold.** Restricting representations to a balanced band  $p \in [N/2 - k, N/2]$  and counting how many  $N$  require a semiprime (no prime pair in the band) defines the rescue fraction, whose unimodal profile in  $k$  is reported in §7.5; if the minimum number of semiprime selections is zero over a finite range, Goldbach holds there.

## 7 Empirical results and a new regularity

This section reports the outcome of implementing the program above. All numbers are reproducible from the accompanying code (see the Data and code availability statement; the drivers `run_counts.py`, `run_scaling.py`, `run_betalaw.py`, `run_estimator_audit.py`); unless stated otherwise they use the range  $X = 2 \cdot 10^6$ .

### 7.1 Computational setup and validation

The counts  $R_1$  and  $R_2$  are obtained for *all*  $N \leq X$  at once as additive convolutions,

$$R_1 = \mathbf{1}_{\mathbb{P}} * \mathbf{1}_{\mathbb{P}}, \quad R_2 = \mathbf{1}_{\mathbb{P}} * \mathbf{1}_{\mathcal{S}_2},$$

evaluated by FFT in  $O(X \log X)$ . For  $X = 2 \cdot 10^6$  this takes about 15 seconds including sieves, diagnostics and figures. The FFT counts agree exactly with a brute-force enumeration on random samples (the rounding error of the transform is far below  $1/2$ ), and Goldbach's condition  $R_1(N) > 0$  is verified throughout the range.

### 7.2 The semiprime channel and the half-power mirage

The Hardy–Littlewood prediction (15) attaches to  $R_1$  the full singular series  $\mathfrak{S}(N) = \prod_{p|N, p>2} (p-1)/(p-2)$ . Regressing  $\log(R_1/(N/\log^2 N))$  against  $\log \mathfrak{S}(N)$  recovers, empirically, a slope 1.000, and dividing by  $\mathfrak{S}(N)$  collapses the coefficient of variation of the normalized count from 0.39 to 0.015 (a 25× reduction); this is a control that reproduces (15) and the band structure of the Goldbach comet.



The new observation concerns  $R_2$ . We measure how strongly the relative semiprime share tracks  $\mathfrak{S}(N)$  by the *regression slope*

$$\beta_2 - 1 := \text{slope of } \log \frac{R_2(N)}{R_1(N)} \text{ against } \log \mathfrak{S}(N),$$

which uses  $R_1$  itself as a meter for  $\mathfrak{S}(N)$  and is therefore independent of any chosen normalization. We call  $\beta_2$  the *empirical singular exponent*, but stress at the outset that  $\beta_2$  **is a regression slope, not an exponent of a power law**: as Lemma 1 shows below,  $\mathfrak{S}(N)$  enters  $R_2$  with exponent 1 (exactly as in  $R_1$ ) and *cancels* in the ratio  $R_2/R_1$ , so  $\beta_2$  is the slope of a non-power-law object and the value  $\approx \frac{1}{2}$  reported in finite ranges is a finite-size correction, not a structural exponent (developed after Table 1). We adopt one estimator throughout: ordinary least squares of  $\log(R_2/R_1)$  on  $\log \mathfrak{S}$ , taken *cumulatively* over all even  $5000 \leq N \leq X$ . Table 1 reports it across several ranges; standard errors are  $\leq 10^{-4}$  (so the digits shown are significant), and the appendix (§A.6) instead uses a *local* window centred at  $X$ , which runs  $\approx 0.02$ – $0.03$  higher at the same  $X$ —the distinction is between an average over  $[5000, X]$  and a measurement at the leading edge.

| $X$            | $\beta_2$ | $R^2$ of the regression |
|----------------|-----------|-------------------------|
| $5 \cdot 10^5$ | 0.452     | 0.91                    |
| $1 \cdot 10^6$ | 0.475     | 0.91                    |
| $2 \cdot 10^6$ | 0.497     | 0.92                    |
| $4 \cdot 10^6$ | 0.513     | 0.93                    |
| $8 \cdot 10^6$ | 0.529     | 0.93                    |

Table 1: Cumulative regression slope  $\beta_2$  of  $\log(R_2/R_1)$  on  $\log \mathfrak{S}$  over  $[5000, X]$  (standard error  $\leq 10^{-4}$ ). The value sits near  $\frac{1}{2}$  in this range and *rises* with  $X$ —already a warning that it is not a fixed exponent (contrast  $\beta_1 = 1.000$ , which is genuinely scale-invariant). The singular series explains about 92% of the variance of  $\log(R_2/R_1)$ .




Figure 1: Left:  $\log[R_1/(N/\log^2 N)]$  against  $\log \mathfrak{S}(N)$  has slope 1.000 (Hardy–Littlewood). Right: the same for  $R_2$  has slope 0.497 at  $X = 2 \cdot 10^6$ , near the reference line  $\beta = 1/2$ —but this is the *finite-range* slope of a non-power-law object (it drifts toward 1; Fig. 2), not a fixed exponent. The vertical striations are the discrete values of  $\mathfrak{S}(N)$ .

Over this range, the best finite-range log–log normalization is consistent with the effective fit

(near  $X \sim 10^6$ , the exponent  $\frac{1}{2}$  drifting to 1; §7.2 below)

$$R_2(N) \propto \mathfrak{S}(N)^{1/2} \frac{N}{\log^2 N} W(N), \quad W(N) = \sum_{\substack{3 \leq r \leq \sqrt{N} \\ r \text{ prime}}} \frac{r-1}{r-2} \frac{1}{r} \frac{\log N}{\log(N/r)}, \quad (20)$$

where  $W(N) \sim \log \log N$  by Mertens' theorem [12]. The weight  $W$  makes precise the  $\log \log N$  factor of (17): it is a Mertens sum over the *small prime factor*  $r$  of the semiprime  $q = rs$ , weighted by the twin-prime-type local factor  $(r-1)/(r-2)$ . Table 2 compares the dispersion of  $R_2$  under three normalizations.

| normalization of $R_2$                                            | CV ( $X = 2 \cdot 10^6$ ) |
|-------------------------------------------------------------------|---------------------------|
| $N \log \log N / \log^2 N$ (heuristic of §5)                      | 0.184                     |
| $2C_2 \mathfrak{S}(N)^1 N W / \log^2 N$ (full inheritance)        | 0.174                     |
| $2C_2 \mathfrak{S}(N)^{1/2} N W / \log^2 N$ ( <b>half power</b> ) | <b>0.018</b>              |

Table 2: At  $X = 2 \cdot 10^6$  the half-power normalization fits about 10× tighter than the  $\log \log N$  heuristic, and tighter than the full singular series ( $\beta = 1$ )—but this is a statement about the *finite-range* value of the slope, not about the limiting exponent. As  $X$  grows the best-fitting  $\beta$  rises toward 1 (Fig. 2); the local optimality of  $\beta = \frac{1}{2}$  here reflects the  $1/\log \log N$  deficit at  $X \sim 10^6$ , not a failure of  $R_2$  to inherit  $\mathfrak{S}(N)$ .

**Interpretation: a quantitative Chen rescue effect.** Because the finite-range slope has  $\beta_2 < 1$ , over this range the relative semiprime share falls with the singular series,  $R_2/R_1 \propto \mathfrak{S}(N)^{-(1-\beta_2)}$  (locally  $\approx \mathfrak{S}(N)^{-1/2}$ ): where Goldbach is locally rich (large  $\mathfrak{S}$ ), the prime channel dominates, whereas where Goldbach is *fragile* (small  $\mathfrak{S}$ ) the semiprime channel is proportionally most abundant. This makes the Chen-rescue intuition—the semiprime channel matters most exactly where primes are scarce—quantitative as an *effective finite-window scaling*, with the deficit  $1 - \beta_2 \asymp 1/\log \log N$  vanishing asymptotically (so it is a slope, not a fixed power; §7.2 opening). Heuristically the slope sits below 1 because writing  $q = rs$  couples the divisors of  $N$  to the factor  $r$ : when a small prime  $\ell \mid N$ , the channel  $r = \ell$  is *blocked* (then  $p = N - \ell s \equiv 0 \pmod{\ell}$  forces  $p = \ell$ ), removing its term from the sum over channels precisely when  $\mathfrak{S}(N)$  is large (Lemma 1). This explains the direction  $\beta_2 < 1$ ; the precise value is not  $1/2$ , as the scaling analysis below shows.

**Scaling to  $10^9$ : the slope grows like  $\log \log X$ .** Measuring  $\beta_2$  *locally* in a window of  $2 \cdot 10^6$  even integers centred at  $X$  (computed by a block-convolution sieve, so that  $X$  can reach  $10^9$  while only the window is held in memory), the slope does *not* settle at  $1/2$ : it rises monotonically (Fig. 2). Eight windows from  $6 \cdot 10^6$  to  $10^9$  are fit to within 0.0013 by

$$\beta_2(X) \approx -0.094(7) + 0.234(3) \log \log X \quad (R^2 = 0.9987, \text{ 8 windows}), \quad (21)$$

the parenthesised figures being the last-digit standard errors of the two-parameter fit. Thus the half-power  $\beta_2 \approx \frac{1}{2}$  is only the *local* value near  $X \sim 10^6$ , and the slope is not a constant. The fit reaches  $\beta_2 \approx 0.61$  at  $10^9$  (measured). We deliberately *do not* read a numerical limit off (21): extrapolating a  $\log \log X$  trend fitted over three decades to, say,  $X \sim 10^{46}$  (where it would touch  $\beta_2 = 1$ ) is well outside the data and is not evidence—the robust content is the *functional form*  $1 - \beta_2 \asymp 1/\log \log X$  and its derivation below, not a forecast of where the slope crosses a particular value. The regression quality stays high throughout ( $R^2 \approx 0.997$ , CV of the half-power normalization  $\lesssim 0.04$ ).




Figure 2: Local regression slope  $\beta_2(X)$  for windows up to  $10^9$ , with the  $\log \log X$  fit (21) (the dashed extrapolation beyond  $10^9$  is shown only to display the functional form, not as a prediction). The slope drifts steadily above  $1/2$  without sign of leveling off, consistent with  $\beta_2 \rightarrow 1$ .

**A derivation:  $\beta_2 \rightarrow 1$  with an  $1/\log \log N$  deficit.** The drift is not arbitrary. Writing  $q = N - p = rs$  and summing over the small factor  $r$ , each channel is a Goldbach count for the linear form  $(s, N - rs)$ , whose two-form singular series factorizes as

$$\mathfrak{S}_2(r, N) = \begin{cases} 2C_2 \mathfrak{S}(N) \frac{r-1}{r-2}, & r \nmid N, \\ 0, & r \mid N \end{cases} \quad (22)$$

(Lemma 1; the  $r \mid N$  channel is identically blocked). Since  $R_1 = 2C_2 \mathfrak{S}(N) I_1$  and  $R_2 = 2C_2 \mathfrak{S}(N) \sum_{r \nmid N} \frac{r-1}{r-2} I_r$ , the ratio *cancels the singular series exactly*,

$$\frac{R_2(N)}{R_1(N)} = \widetilde{W}(N) := \sum_{\substack{3 \leq r \leq \sqrt{N} \\ r \nmid N}} \frac{r-1}{r-2} w_r(N), \quad w_r = \frac{1}{r} \frac{\log N}{\log(N/r)}. \quad (23)$$

Hence the measured exponent is a regression slope,  $\beta_2 - 1 = \text{Cov}(\log \widetilde{W}, \log \mathfrak{S}) / \text{Var}(\log \mathfrak{S})$ , and the only coupling to  $\mathfrak{S}(N)$  comes from the blocked channels  $r \mid N$ , each *removed* from  $\widetilde{W}$ . Modelling  $\mathbf{1}[\ell \mid N]$  as independent with probability  $1/\ell$ ,  $\text{Var}(\log \mathfrak{S}) =: B$  is a convergent constant, while the removed terms contribute  $\text{Cov}(\log \widetilde{W}, \log \mathfrak{S}) = -A/\widetilde{W}_0$  with  $A > 0$  constant and  $\widetilde{W}_0 \approx \langle R_2/R_1 \rangle \sim \log \log N$  the baseline sum (Theorem 1). Therefore

$$\beta_2(N) = 1 - \frac{c(N)}{\langle R_2/R_1 \rangle_N} + o\left(\frac{1}{\log \log N}\right), \quad c(N) = \frac{A}{B} = O(1) (\approx 1.1 \text{ in range; Fig. 15}), \quad (24)$$

so  $\beta_2 \rightarrow 1$ :  $R_2$  *asymptotically inherits the full Goldbach singular series*, the approach being governed by  $1/\log \log N$ . This explains why finite ranges show  $\beta_2 \approx \frac{1}{2}$  and a slow apparent drift.

**Proposition 1** (Validation, non-circular). *The law (24) predicts that  $(1 - \beta_2) \langle R_2/R_1 \rangle$  is constant. Measured over windows from  $6 \cdot 10^6$  to  $2 \cdot 10^8$  it equals 1.11, 1.11, 1.10, 1.09, 1.09, 1.08, a coefficient of variation of 1% (versus 1.7% for  $(1 - \beta_2) \log \log N$ ): the correct “clock” is  $\langle R_2/R_1 \rangle$ , exactly as (24) requires (Fig. 3).*




Figure 3: The derived law:  $\beta_2$  against  $1/\langle R_2/R_1 \rangle \sim 1/\log \log N$  is a straight line through the limit point  $(0, 1)$ , i.e.  $\beta_2 \rightarrow 1$ .

**Not a single-atom effect.** A natural worry is that the whole slope is the arithmetic of one prime:  $3 \mid N$  simultaneously doubles  $\mathfrak{S}(N)$  (the factor  $\frac{3-1}{3-2} = 2$ ) and deletes the heaviest semiprime channel  $r = 3$ , so one might suspect  $\beta_2 \approx \frac{1}{2}$  is just “what divisibility by 3 does” near  $X \sim 10^6$ . The data refute this. Recomputing the slope on the subpopulation  $3 \nmid N$  (the  $r = 3$  atom removed) gives, at  $X = 2 \cdot 10^6$ ,  $\beta_2 = 0.547 \pm 0.0004$  against 0.495 on all  $N$ : the deficit  $1 - \beta_2$  falls by only about 10% (`run_estimator_audit.py`). The mechanism is real—conditional means confirm  $\langle \log \mathfrak{S} \mid 3 \mid N \rangle - \langle \log \mathfrak{S} \mid 3 \nmid N \rangle = 0.69 \approx \log 2$  and a depressed  $R_2/R_1$  on  $3 \mid N$ —but it accounts for a tenth of the slope, not the bulk. The negativity of  $\beta_2 - 1$  is genuinely a *sum over channels*  $r = 3, 5, 7, \dots$ , each blocked on the sparse set  $r \mid N$ , exactly as  $A(X) = \sum_r(\dots)$  in (24); it is not the signature of a single residue class.

### 7.3 The singular exponent across arithmetic-complexity layers

The decomposition  $R_1, R_2$  extends to layers  $R_k(N) = \#\{p < N : \Omega(N - p) = k\}$ , counting representations whose second summand has exactly  $k$  prime factors (Goldbach is  $k = 1$ , Chen’s semiprime channel is  $k = 2$ ). Measuring the singular exponent  $\beta_k$  of each layer – the slope of  $\log(R_k/R_1)$  against  $\log \mathfrak{S}(N)$ , plus one – reveals a clean pattern (Table 3, Fig. 4):

| $k$             | 1     | 2     | 3     | 4     | 5     | 6     |
|-----------------|-------|-------|-------|-------|-------|-------|
| share $\rho(k)$ | 15.5% | 33.5% | 28.7% | 14.6% | 5.5%  | 1.8%  |
| $\beta_k$       | 1.00  | 0.50  | −0.30 | −1.38 | −2.73 | −4.04 |

Table 3: Layer profile and singular exponent at  $X = 2 \cdot 10^6$  (each regression has  $R^2 \approx 0.92$ ). The exponent decreases monotonically and crosses zero between  $k = 2$  and  $k = 3$ ; it is *not*  $1/k$ .

**Sign reversal of the singular response.** The Goldbach singular series *enhances* the low-complexity channels ( $\beta_1 = 1$ ,  $\beta_2 \approx \frac{1}{2} > 0$ ) but *suppresses* the high-complexity ones ( $\beta_k < 0$  for  $k \geq 3$ ): where  $\mathfrak{S}(N)$  is large, representations with a many-factored  $q$  become rarer. The mechanism is the same divisor effect seen for  $R_2$ , now run in reverse. When  $\mathfrak{S}(N)$  is large,  $N$  is




Figure 4: Left: the arithmetic-complexity profile  $\rho(k)$  peaks at  $k = 2$ . Right: the singular exponent  $\beta_k$  decreases through  $1, \frac{1}{2}, \dots$  and becomes negative for  $k \geq 3$ , diverging from the naive  $1/k$  reference.

divisible by many small primes; the admissible  $p$  avoid those residues, which forces  $q = N - p$  to also avoid small prime factors, so  $q$  tends to have *fewer* prime factors. Thus a large singular series biases  $q$  toward low arithmetic complexity. All  $\beta_k$  drift upward with  $X$  at the same slow rate as  $\beta_2$  (e.g.  $\beta_3$  moves from  $-0.30$  at  $2 \cdot 10^6$  to  $-0.20$  at  $8 \cdot 10^6$ ), so the crossing point of  $\beta_k = 0$  moves slowly to the right. We also find that the strongest “rescue” channel,  $q = \text{composite} \times \text{composite}$  (both factors composite), is available for *every* even  $N$  in the range with on average  $\sim 1.8 \cdot 10^4$  representations: the additive hierarchy is extremely redundant below the prime layer. The whole sign reversal is the coefficient-level shadow of a single generating-function response in the blocked-channel model, made precise in Appendix A.8.

## 7.4 Diagnostics

The prime share has mean  $\bar{\theta} = 0.311$  and decreases slowly with  $N$  (the semiprime channel gains ground, as the  $\log \log N$  factor predicts); the Chen surplus has mean  $\bar{C} = 2.27$ . Goldbach fragility is extremely rare:  $|\mathcal{F}_1| = 1$ ,  $|\mathcal{F}_2| = 3$ ,  $|\mathcal{F}_3| = 5$ ,  $|\mathcal{F}_5| = 14$ ,  $|\mathcal{F}_{10}| = 49$  up to  $X$ ; and *every* fragile integer observed has  $R_2 > 0$ , so Chen always rescues. The semiprime factor balance  $B(q) = \log a / \log q$  has weighted mean 0.25, with 16% of the representation mass at  $B < 0.1$  (a tiny prime factor) and 18% at  $B > 0.4$  (near balanced): Chen representations are dominated by *unbalanced* semiprimes with a small prime factor (Fig. 5), answering the semiprime factor-geometry question of §4.

## 7.5 Optimization experiments

**Smoothness is a Pareto trade-off.** Minimizing the total spatial variation  $\sum |s_N - s_{N-2}|$  of the selected location  $s_N = q_N/N$  yields a geometrically very smooth branch (total variation 0.0040 over 201 consecutive even integers) that nonetheless *switches type in 28% of its steps* and uses semiprimes 72% of the time (Fig. 6). Forbidding semiprimes (the pure Goldbach branch, with zero type switches) raises the variation to 0.0246, a factor of 6.2 rougher. One cannot simultaneously maximize geometric and type smoothness: continuity in location and chaos in type coexist.

**A unimodal Chen-rescue threshold.** Restricting representations to a balanced band  $p \in [N/2 - k, N/2]$  and measuring the fraction of  $N$  that *require* a semiprime (no prime pair in the band) gives a unimodal curve in  $k$  (Fig. 7). For  $k \gtrsim 0.005 N$  the prime channel




Figure 5: Representation-weighted distribution of the factor balance  $B(q)$  of the semiprime second summand. Mass concentrates toward small  $B$  (small prime factor), not toward the balanced value  $1/2$ .

essentially never fails; the rescue fraction peaks near 38–40% at an intermediate half-width  $k \sim 20$ –40; and for  $k = 1, 2$  it falls again because often no representation exists at all. The peak shifts right with  $N$  (roughly as  $\log^2 N$ ), marking where the expected number of balanced prime pairs,  $\sim 2C_2 \mathfrak{S}(N) k / \log^2(N/2)$ , drops to  $O(1)$  – a Poisson-type transition. This is the regime captured by the residual exponent  $a_{\text{Res}}(N)$  of §7.7: Chen’s semiprime flexibility becomes *necessary* precisely when near-perfect balance is imposed.

## 7.6 Weak continuity of the location measures

Finally we test the weak-continuity intuition of §4 (Question 1 and the weak type-stabilization hypothesis there) directly. The pointwise measure  $\mu_N$  of  $q/N$  is noisy, but averaging over a window  $N \in [X, X + H]$  produces a stable density (Fig. 8, left). Three findings, at  $X = 10^6$ :

- **The smoothed measure converges.** Splitting a window into two same-scale subsamples, the total-variation distance between their densities falls monotonically from  $2.1 \cdot 10^{-3}$  to  $3 \cdot 10^{-4}$  as  $H$  grows from 125 to 4000 (Fig. 8, right), i.e. the per- $N$  noise averages out – a concrete form of weak convergence.
- **The limit is essentially uniform.** The prime- $q$  density is within total variation 0.028 of the uniform law on  $(0, 1)$ , with only a mild upturn near the endpoints (the  $1/(\log(xN) \log((1 - x)N))$  effect). This answers Question 1 affirmatively: after removing endpoint effects,  $\mu_N^{(1)}$  is close to uniform.
- **Prime and semiprime locations nearly coincide.** The two densities are within total variation 0.019 of each other (Fig. 8, center), confirming the weak type-stabilization picture. The semiprime location is slightly skewed toward larger  $q$  (mean  $q/N = 0.513$  versus the exactly symmetric 0.500 for primes), the only visible signature that  $p$  and  $q$  play asymmetric roles in  $R_2$ .

Appendix A.9 gives the weighted Hardy–Littlewood explanation: after normalization, both the inherited singular series and the Mertens channel sum cancel, leaving the uniform law for  $q/N$  in either channel (conditional on the weighted channel asymptotics).




Figure 6: Minimum-variation branch (Gurobi). The optimum hugs  $s_N \approx 1$  (the “least partner” branch,  $p$  a small prime) and is smooth in  $s_N$ , yet its type (blue prime / red semiprime) alternates.

## 7.7 The residual Goldbach–Chen balance exponent

Inspired by the  $(1+a)$  interpolation of Li and Liu (every large even  $N$  is  $p + rq$  with  $r \leq q^{a-1}$ , so that  $(1+1)$  is Goldbach and  $(1+2)$  is Chen), one can ask not whether a *fixed* exponent works, but the smallest exponent *needed* for each  $N$ . For the nondegenerate Chen layer  $N = p + rs$  ( $p, r, s$  prime,  $r \leq s$ ), define the residual balance exponent and minimal multiplier

$$a_{\text{Res}}(N) = 1 + \min_{N=p+rs} \frac{\log r}{\log s}, \quad r_*(N) = \min\{r \in \mathbb{P} : \exists p, s \in \mathbb{P}, N = p + rs\}.$$

We compute both exactly for all even  $N \leq 2 \cdot 10^6$  ( $r_*$  by FFT channel coverage;  $a_{\text{Res}}$  by a small- $p$  scan, validated against full enumeration to 0 error). Findings (Fig. 9,  $N \geq 1000$ ):

- **Typical residual collapse.** The median  $a_{\text{Res}}$  *decreases* with  $X$  ( $1.115 \rightarrow 1.099 \rightarrow 1.088$  at  $X = 10^5, 5 \cdot 10^5, 2 \cdot 10^6$ ), with  $p_{99} = 1.182$  and a finite-range maximum of 1.385 – far below the Li–Liu threshold 1.9. The worst case is pinned by a few small “hard”  $N$ .
- **The minimal multiplier is tiny.**  $r_* = 3$  for 68% of  $N$  and the dictionary  $\{2, 3, 5, 7\}$  already covers 99.7%;  $\{2, \dots, 11\}$  covers 99.99%. This settles the multiplier-dictionary question empirically. Appendix A.10 explains the dictionary through a least-unblocked-prime law: conditionally on fixed-channel almost-all asymptotics,  $r_*(N)$  equals the least odd prime not dividing  $N$  for almost all even  $N$  (so  $\Pr(r_* \in \{3, 5, 7\}) \approx 0.99$  in the divisibility model).
- **Local obstruction.**  $r_*$  jumps when a small prime divides  $N$ : if  $3 \mid N$  the channel  $r = 3$  is blocked (then  $p = N - 3s \equiv 0 \pmod{3}$ ), and indeed the mean  $r_*$  is 4.59 for  $3 \mid N$  versus 2.94 otherwise.
- **Link to the singular series.**  $a_{\text{Res}}$  correlates *positively* with  $\log \mathfrak{S}(N)$  (correlation +0.59). The very divisibility of  $N$  by small primes that enlarges  $\mathfrak{S}(N)$  (boosting Goldbach) also blocks the small- $r$  Chen channels, so the residual balance is *worst exactly where Goldbach is locally richest*. This ties the balance exponent to the singular-series theme of §7.3.






Figure 7: Fraction of  $N$  that require a semiprime rescue as a function of the band half-width  $k$ , for three scales of  $N$ . The curve is unimodal and its peak scales like  $\log^2 N$ .




Figure 8: Weak continuity of  $q/N$ . Left: noisy pointwise  $\mu_N$  (gray) versus the window-smoothed density (blue). Center: prime versus semiprime location densities, both close to uniform. Right: same-scale sampling distance decreases with the window size  $H$  (log–log).

The full geometry of the Chen surface  $p + rs = N$  on which the exponent  $a_{\text{Res}}$  lives—a 2D cloud of solutions organised by the small prime factor  $r$ , with mass against the Goldbach edge—is developed in Appendix C.

## 7.8 Three further probes: transport, topology, dynamics

We close with three geometric and statistical probes of the frontier; each, perhaps surprisingly, returns to the singular series.

**Optimal transport (reflection defect).** Under the reflection  $T(x) = 1 - x$  (which sends  $p/N$  to  $q/N$ ), compare the prime location law  $\alpha$  (density of  $p/N$ ) with the reflected layer laws  $T_{\#}\beta_k$ . The reflection defects in  $W_1$  are  $\Delta_1 = W_1(\alpha, T_{\#}\beta_1) = 0.040$  and  $\Delta_2 = W_1(\alpha, T_{\#}\beta_2) = 0.032$ , and the optimal Chen mixture attains  $\Delta_{\leq 2} = 0.032$  at  $\lambda^* = 0$ : the reflected *semiprime* cloud matches the primes *better* than the reflected primes do, a 22% “transport collapse” (stable in  $X$ ), driven by unbalanced semiprimes with a small factor (Fig. 10).






Figure 9: Left: percentile envelope of  $\lambda_{\text{Res}} = a_{\text{Res}} - 1$  versus  $N$ , far below the Li-Liu line 0.9. Centre: the minimal multiplier  $r_*$  ( $r = 3$  dominates). Right:  $\lambda_{\text{Res}}$  rises with the Goldbach singular series  $\mathfrak{S}(N)$  (positive correlation).




Figure 10: Reflection transport. Left: prime law  $\alpha$  and the reflected prime and semiprime laws. Right: admitting semiprimes lowers the reflection defect by 22%.

**Topology of valleys.** Normalising each count by its own singular factor ( $R_1/\mathfrak{S}$ ,  $R_{\leq 2}/(\mathfrak{S} + \mathfrak{S}^{1/2}W)$ ) and computing the 0-dimensional sublevel persistence of the detrended sequences, the valleys of the Goldbach and Chen comets largely *coincide*: the anomalies correlate at +0.64, and at the 200 most fragile  $N$  the Chen anomaly (0.29) is as deep as the Goldbach one (0.30). Chen does *not* topologically fill the Goldbach valleys (only the deepest 30% are softened, Fig. 11); the rescue is one of magnitude, not of valley structure, because both channels share the constraint that  $p$  be prime.

**Statistical dynamics of the frontier.** The prime share is almost a deterministic function of the singular series:  $\theta(N)$  is 94% explained by  $\mathfrak{S}(N)$  alone (adding  $N \bmod 30$  contributes nothing further), and the increment  $\Delta\theta_N$  is 98% explained by the jump  $\mathfrak{S}(N) \rightarrow \mathfrak{S}(N+2)$ . The residual is small (25% of the spread of  $\theta$ ) and is *not* white noise (short-range lag-1 autocorrelation  $-0.21$  after detrending): the Goldbach-Chen frontier carries little irreducible randomness, being governed by the local arithmetic of  $N$  (Fig. 12).

## Two reframings; the rest in an appendix

The Goldbach-Chen frontier can be recast in many external languages. We keep here only the two that yield a genuinely *new* computation. The remaining reformulations—an arithmetic ground-state energy, the adversarial robustness of the Goldbach graph, a market/liquidity reading,




Figure 11: Valley persistence. Left: Chen has nearly as many valleys as Goldbach (slightly fewer deep ones). Right: the normalised anomalies coincide (+0.64).




Figure 12: Dynamics. Left:  $\theta(N)$  tracks its singular-series prediction. Right: the residual autocorrelation, small and non-white.

a simplicial-topology probe, a control policy, an information budget, and the discrete geometry of the Chen surface—are collected in Appendix C, since they mostly re-express singular-series governance rather than constitute independent findings (we record there the honest negatives too: the topology probe is empty and the Chen substitute carries no resilience premium).

## 7.9 An empirical circle method: the spectrum of the comet

The circle method decomposes additive problems by Fourier analysis into major arcs (rationals  $a/q$  with small  $q$ , carrying the singular series) and minor arcs (the pseudorandom remainder). One can see this decomposition *empirically*. Detrending  $R_1$  over a window of consecutive even integers and taking its power spectrum (Fig. 13, left) reveals a forest of sharp lines exactly at the arithmetic frequencies: the dominant peak is at  $\frac{1}{3}$  (period  $T_N = 6$ , the modulus 3), followed by  $\frac{1}{5}, \frac{2}{5}$  (modulus 5), with heights that decay across primes in step with the singular-series factors  $(\ell - 1)/(\ell - 2)$ . These are the major arcs made visible.

Dually, one can ask how much of the comet’s oscillation is captured by purely local information. Predicting the detrended  $\log R_1(N)$  from the residue  $N \bmod \prod_{\ell \leq B} \ell$  explains 84.6% of the variance at  $B = 3$  (i.e.  $N \bmod 6$  alone), 95.0% at  $B = 5$ , and saturates toward the singular-series ceiling  $R^2 = 0.999$  (Fig. 13, right). The comet is, to first order, a *modulo-6 phenomenon*: the

prime 3 supplies 85% of all explainable structure, and successive primes contribute the rapidly shrinking remainder predicted by  $\mathfrak{S}(N)$ .




Figure 13: Empirical circle method. Left: the power spectrum of the detrended Goldbach comet, with peaks at the arithmetic frequencies  $a/q$  (the major arcs), dominated by  $\frac{1}{3}$  (modulus 3). Right: the fraction of the comet’s oscillation explained by  $N$  modulo the small primes, saturating at the singular-series ceiling;  $N \bmod 6$  already gives 85%.

## 7.10 Primorial bases and carry geometry

Two further reframings come from writing integers in a base. In a primorial base  $B_y = \prod_{p \leq y} p$ , every large prime is a unit modulo  $B_y$ , so the residues admissible for  $N$  are  $A_y(N) = \{a \in (\mathbb{Z}/B_y)^\times : N - a \in (\mathbb{Z}/B_y)^\times\}$ , with local coverage  $\lambda_y(N) = |A_y(N)|/\phi(B_y)$ . By the Chinese remainder theorem this factorises, and the *relative* coverage is precisely the truncated singular series,

$$\frac{\lambda_y(N)}{\lambda_y(\text{generic})} = \prod_{\substack{p|N \\ 3 \leq p \leq y}} \frac{p-1}{p-2} = \mathfrak{S}_y(N).$$

So primorial coverage *is* the singular series, viewed as accumulated local obstructions; predicting  $\log R_1$  from  $\mathfrak{S}_y$  explains 84.6% of the comet’s oscillation at  $y = 3$  ( $B_3 = 6$ ), rising to 99.8% by  $y = 30$ —the same saturation as §7.9, now read as “the comet is the shadow of primorial local coverage”.

The genuinely new object is the *carry geometry*. Adding  $p + q = N$  in base  $b$  produces carries  $c_i \in \{0, 1\}$ , and their number is fixed by the digit sums,  $\#\text{carries}(p, N - p) = (S_b(p) + S_b(N - p) - S_b(N))/(b - 1)$ ; in base 2 this is  $v_2\binom{N}{p}$  (Kummer). Comparing Goldbach representations to random pairs  $a + (N - a) = N$  reveals a systematic *carry excess*: Goldbach pairs carry more (Fig. 14). In base 2 the excess is +0.97 carries (Goldbach 9.9 versus 8.9), and a positive excess persists in every base tested ( $b = 2, \dots, 16$ ), largest in the small bases. The origin is arithmetic: a prime’s last digit is coprime to  $b$ , which biases the digit sums  $S_b(p) \equiv p \pmod{b-1}$  away from the values that would avoid carries. Thus the same local constraints that produce the singular series leave a *digital signature*: Goldbach decompositions are, bit for bit, slightly carry-heavier than chance—a statistical-mechanical fingerprint of primality in the geometry of addition. A local model theorem in Appendix A.11 shows that restricting both terminal digits to reduced residue classes already increases the least-digit carry probability relative to unrestricted random summands.




Figure 14: Carry geometry. Left: the distribution of binary carries  $v_2(N/p)$  for Goldbach representations (red) versus random pairs (grey)—Goldbach is shifted to more carries (+0.97). Right: the carry excess is positive in every base.

## 8 Conclusion

The one durable result of this study is a single Euler-product identity and its consequence: decomposing the semiprime channel by its small prime factor shows that the Goldbach singular series  $\mathfrak{S}(N)$  enters  $R_2$  with exponent 1, exactly as it enters  $R_1$ , and *cancels* in the ratio  $R_2/R_1 = \widehat{W}(N)$  (Lemma 1). The widely tabulated “ $\beta_2 \approx \frac{1}{2}$ ” is therefore not an exponent but a regression slope of a non-power-law object, drifting to 1 like  $1 - c/\log \log N$ ; we verify this non-circularly, show it is a multi-channel effect (not the  $3 \mid N$  atom), and reduce the limit to classical inputs—unconditionally for the *linear* almost-all asymptotic  $R_2 = \text{main}(1 + o(1))$ , and, for the log-regression statement  $\beta_2 \rightarrow 1$ , modulo one explicitly stated lower-tail hypothesis (Corollary 1). A truncated-log proposition and a blocked-channel model (§A.7) sharpen this gap: the second moment already controls a stable logarithmic statistic and the model proves the  $1/\log \log X$  deficit, while only the untruncated log-regression still requires lower-tail input.

We have deliberately resisted the temptation to read this one fact as many. The optimization formulations give a language, not a proof strategy (Remark 2); the dozen physical, economic, and information-theoretic reframings are, with two exceptions (the empirical circle method and the carry geometry), the same singular-series governance re-photographed in different filters, and we have said so rather than count them as separate findings. The negatives—an empty additive-complexity topology, a Chen substitute with no resilience premium—are reported as negatives.

This is not a proof of Goldbach. It is reproducible evidence, one clean new identity, and a tightly bounded reduction, at the boundary between additive number theory, computational experimentation, and discrete optimization.

## A Towards a proof that $\beta_2 \rightarrow 1$

We make the law (24) precise. We cannot prove it unconditionally: the asymptotics of  $R_1, R_2$  are themselves the Hardy–Littlewood conjecture, and even  $R_1(N) > 0$  is open. We therefore isolate what is unconditional (Lemmas 1–2), prove the law *conditionally* on a quantitative Hardy–Littlewood hypothesis (Theorem 1), and note that it holds unconditionally in the Cramér random model.

## A.1 Notation and the statistic

Fix a window  $\mathcal{N}_X = \{N \equiv 0 \pmod{2} : X \leq N < X + H\}$  with  $H = H(X) \rightarrow \infty$ ,  $H = o(X)$ . Write  $\mathbb{E}_X, \text{Var}_X, \text{Cov}_X$  for empirical moments over  $\mathcal{N}_X$ ,  $g_\ell = \log \frac{\ell-1}{\ell-2}$ , and define the local singular exponent through the (least-squares) regression slope

$$\beta_2(X) - 1 := \frac{\text{Cov}_X(\log(R_2/R_1), \log \mathfrak{S})}{\text{Var}_X(\log \mathfrak{S})}. \quad (25)$$

## A.2 Unconditional lemmas

**Lemma 1** (Singular series of a channel). *Let  $r$  be an odd prime and  $N$  even. The singular series of the pair of linear forms  $s \mapsto (s, N - rs)$  is*

$$\mathfrak{S}_2(r, N) = \prod_{\ell} \frac{1 - \nu_r(\ell)/\ell}{(1 - 1/\ell)^2}, \quad \nu_r(\ell) = \#\{s \bmod \ell : s(N - rs) \equiv 0\},$$

and equals

$$\mathfrak{S}_2(r, N) = \begin{cases} 2C_2 \mathfrak{S}(N) \frac{r-1}{r-2}, & r \nmid N, \\ 0, & r \mid N, \end{cases} \quad C_2 = \prod_{p>2} \left(1 - \frac{1}{(p-1)^2}\right).$$

*Proof.* Compute  $\nu_r(\ell)$ . For  $\ell = 2$ : as  $s$  and  $N - rs$  are both odd exactly when  $s$  is odd,  $\nu_r(2) = 1$  and the factor is  $(1 - \frac{1}{2})/(1 - \frac{1}{2})^2 = 2$ . For odd  $\ell \nmid rN$ :  $s \equiv 0$  and  $s \equiv Nr^{-1} \not\equiv 0$  are two distinct roots,  $\nu_r = 2$ , factor  $(1 - 2/\ell)/(1 - 1/\ell)^2$ . For odd  $\ell \mid N$ ,  $\ell \nmid r$ :  $N - rs \equiv -rs$  vanishes iff  $s \equiv 0$ , the same root as  $s \equiv 0$ , so  $\nu_r = 1$ , factor  $(1 - 1/\ell)/(1 - 1/\ell)^2$ . For  $\ell = r \nmid N$ :  $N - rs \equiv N \not\equiv 0$ , so only  $s \equiv 0$ ,  $\nu_r = 1$ , same factor. Taking  $\prod_{\ell>2} (1 - 2/\ell)/(1 - 1/\ell)^2 = C_2$  as the base and converting each special factor relative to it multiplies by  $(\ell - 1)/(\ell - 2)$ ; the special primes are exactly those dividing  $N$  (giving  $\prod_{\ell \mid N} \mathfrak{S}(N)$  when  $r \nmid N$ ) together with  $\ell = r$  (giving  $(r - 1)/(r - 2)$ ). This proves the first case. If  $r \mid N$  then  $N - rs \equiv 0 \pmod{r}$  for all  $s$ , so  $\nu_r(r) = r$ , the factor  $1 - r/r = 0$  vanishes, and  $\mathfrak{S}_2(r, N) = 0$ : the channel is identically blocked because  $p = N - rs \equiv 0 \pmod{r}$  forces  $p = r$ .  $\square$

**Lemma 2** (Moments of  $\log \mathfrak{S}$ ). *For distinct odd primes  $\ell_1, \dots, \ell_k$ ,  $\#\{N \in \mathcal{N}_X : \ell_i \mid N \ \forall i\} = |\mathcal{N}_X| \prod_i \ell_i^{-1} + O(2^k)$ . Consequently the indicators  $e_\ell = \mathbf{1}[\ell \mid N]$  are, as  $X \rightarrow \infty$ , independent Bernoulli( $1/\ell$ ) in all joint moments, and for any weights  $a_\ell$  with  $\sum a_\ell^2/\ell < \infty$ ,*

$$\mathbb{E}_X \left[ \sum_{\ell} a_\ell e_\ell \right] \rightarrow \sum_{\ell} \frac{a_\ell}{\ell}, \quad \text{Var}_X \left[ \sum_{\ell} a_\ell e_\ell \right] \rightarrow \sum_{\ell} a_\ell^2 \frac{1}{\ell} \left(1 - \frac{1}{\ell}\right).$$

In particular  $\text{Var}_X(\log \mathfrak{S}) \rightarrow B := \sum_{\ell>2} g_\ell^2 \ell^{-1} (1 - \ell^{-1})$ .

*Proof.* The count is inclusion of the arithmetic progression  $0 \pmod{\prod \ell_i}$  in an interval of  $\frac{1}{2}|\mathcal{N}_X|$  even integers, giving the stated main term and  $O(2^k)$  boundary error. Hence joint factorial moments of the  $e_\ell$  converge to those of independent Bernoulli( $1/\ell$ ), and the moment formulas follow by the dominated convergence afforded by  $\sum a_\ell^2/\ell < \infty$  (here  $a_\ell = g_\ell \asymp 1/\ell$ ).  $\square$

**Remark 3.** *Lemma 2 is sharp numerically: at  $X = 10^8$ ,  $H = 2 \cdot 10^6$  one finds  $\text{Var}_X(\log \mathfrak{S}) = 0.12620$  against  $B = 0.12620$ , and  $\mathbb{E}_X[\log \mathfrak{S}] = 0.34842 = \sum_{\ell} g_\ell/\ell$  to five places.*

### A.3 The conditional theorem

**Assumption 1** (Quantitative Hardy–Littlewood). *There is  $\delta > 0$  such that, uniformly for odd primes  $r \leq \sqrt{N}$  with  $r \nmid N$ ,*

$$C'_r(N) := \#\{p, s \in \mathbb{P} : p + rs = N, s \geq r\} = \mathfrak{S}_2(r, N) I_r(N) (1 + O(\log^{-\delta} N)),$$

with  $I_r(N) = \int_2^{N/r} \frac{dt}{\log t \log(N-rt)}$ , and the same for  $R_1(N) = 2C_2 \mathfrak{S}(N) I_1(N) (1 + O(\log^{-\delta} N))$ ,  $I_1(N) = \int_2^N \frac{dt}{\log t \log(N-t)}$ .

This is the standard prime  $k$ -tuple heuristic made quantitative and *uniform in the coefficient  $r$* ; the uniformity is the genuine analytic difficulty (it is the arena of the sieve work surrounding Proposition (1 + a)).

**Theorem 1** ( $\beta_2 \rightarrow 1$ , conditional on Assumption 1). *Under Assumption 1, for  $N \in \mathcal{N}_X$ ,*

$$\frac{R_2(N)}{R_1(N)} = \widetilde{W}(N)(1+o(1)), \quad \widetilde{W}(N) = \sum_{\substack{3 \leq r \leq \sqrt{N} \\ r \nmid N}} \frac{r-1}{r-2} w_r(N), \quad w_r = \frac{I_r}{I_1} = \frac{1}{r} \frac{\log N}{\log(N/r)} (1+o(1)),$$

the Goldbach singular series  $\mathfrak{S}(N)$  cancelling by Lemma 1. Writing  $\widetilde{W}_0(X) = \mathbb{E}_X[\widetilde{W}] \sim \log \log X$  and linearising  $\log \widetilde{W}$  about  $\widetilde{W}_0$ ,

$$\beta_2(X) - 1 = -\frac{c(X)}{\widetilde{W}_0(X)} + O(\widetilde{W}_0^{-2}), \quad c(X) = \frac{A(X)}{B}, \quad A(X) = \sum_{r>2} \frac{r-1}{r-2} w_r(X) g_r \frac{1}{r} \left(1 - \frac{1}{r}\right),$$

both  $A(X)$  and  $B$  being absolutely convergent (and  $A$  formed with the exact channel weights  $I_r/I_1$ , not their leading approximation). Thus  $c(X)$  is an explicit  $O(1)$  constant: the leading-order weights give  $c \approx 1.0$ , the exact  $I_r/I_1$  raise it to the measured  $c \approx 1.1$ , and  $c(X)$  decreases slowly with  $X$ . The precise limit  $c_0 = \lim_X c(X)$  is delicate (it depends on the exact weights and averaging convention), but is immaterial to the conclusion, since the order is robust:

$$\boxed{\beta_2(X) = 1 - \frac{c(X)}{\langle R_2/R_1 \rangle_X} \rightarrow 1, \quad c(X) \asymp 1, \quad 1 - \beta_2(X) \asymp \frac{1}{\log \log X}.$$

*Proof sketch.* By Lemma 1,  $R_2 = \sum_{r \nmid N} \mathfrak{S}_2(r, N) I_r = 2C_2 \mathfrak{S}(N) \sum_{r \nmid N} \frac{r-1}{r-2} I_r$  and  $R_1 = 2C_2 \mathfrak{S}(N) I_1$ , so under Assumption 1 the ratio is  $\widetilde{W}(N)(1 + O(\log^{-\delta} N))$  and  $\mathfrak{S}(N)$  cancels. Only the channels  $r \mid N$  are special, each *removed* from  $\widetilde{W}$  (not merely reweighted); thus  $\widetilde{W}(N) = \widetilde{W}_0 - \sum_r \frac{r-1}{r-2} w_r e_r$  up to the slow variation of  $w_r$ . Linearising,  $\log \widetilde{W} = \log \widetilde{W}_0 - \widetilde{W}_0^{-1} \sum_r \frac{r-1}{r-2} w_r e_r + O(\widetilde{W}_0^{-2})$ . Insert into (25): with  $\log \mathfrak{S} = \sum_\ell g_\ell e_\ell$  and the asymptotic independence of Lemma 2, only the diagonal  $r = \ell$  terms survive, giving  $\text{Cov}_X = -\widetilde{W}_0^{-1} A(X)$  and  $\text{Var}_X = B$ , whence the displayed slope. Finally  $\widetilde{W}_0(X) = \mathbb{E}_X[\widetilde{W}] = \sum_{3 \leq r \leq \sqrt{X}} \frac{r-1}{r-2} \frac{1}{r} \frac{\log X}{\log(X/r)} (1 - 1/r) = \log \log X + O(1)$  by Mertens, and equals  $\langle R_2/R_1 \rangle_X$  up to  $o(1)$ .  $\square$

### A.4 Confirmation and the unconditional model

Numerically (Fig. 3; `run_betalaw.py`): the prediction  $\widetilde{W}(N)$  regressed against  $\log \mathfrak{S}$  reproduces the measured  $\beta_2$  with the correct drift (0.554, 0.583, 0.602 at  $X = 2 \cdot 10^6, 2 \cdot 10^7, 10^8$  versus measured 0.531, 0.567, 0.589), and the explicit constant  $c(X) = A/B$  matches the measured  $(1 - \beta_2) \langle R_2/R_1 \rangle \approx 1.1$  in order and sign; with the leading weights  $w_r$  one gets  $c \approx 1.0$ , the residual gap being the crude approximation  $w_r \approx I_r/I_1$  (exact agreement to 0.4% once  $I_r$  is used, Remark in §A.12) together with the linearisation remainder. In the Cramér model, where  $\mathbf{1}_\mathbb{P}, \mathbf{1}_{\mathcal{S}_2}$  are replaced by independent variables with the correct local densities and divisibility, Assumption 1 holds by construction and Theorem 1 is unconditional.

**The rate constant is not cleanly determined.** The natural “constant” admits two forms,  $\kappa(X) = (1 - \beta_2) \log \log X$  and  $c(X) = (1 - \beta_2) \langle R_2/R_1 \rangle$ , which agree in order but not in value because  $\langle R_2/R_1 \rangle = \log \log X + O(1)$ . Both *decrease* over the computed range (Fig. 15):  $\kappa = 1.25, 1.22, 1.20, 1.18$  and  $c = 1.12, 1.10, 1.09, 1.08$  at  $X = 2 \cdot 10^6, 2 \cdot 10^7, 10^8, 5 \cdot 10^8$ . A linear fit in  $1/\log \log X$  extrapolates to  $\kappa_\infty \approx 0.54$  and  $c_\infty \approx 0.73$ —with a correction term of slope  $\approx 1.9$ , larger than the limit itself. Two conclusions follow. First, the limiting constant is *normalisation-dependent* (0.54 versus 0.73), so no single number is canonical. Second, the approach is correction-dominated: the finite-range value  $\approx 1.1$  is mostly the  $1/\log \log X$  term, and the true limit ( $\approx 0.5$ – $0.7$ , itself uncertain) lies far below it and is *not reliably determinable* from data up to  $10^9$ . (Our two back-of-envelope analytic estimates, 1.0 and 1.8, bracket the difficulty rather than resolve it.) Only the order  $1 - \beta_2 \asymp 1/\log \log X$ , and hence  $\beta_2 \rightarrow 1$ , is robust—which is all Theorem 1 asserts.




Figure 15: The rate constant versus  $1/\log \log X$ . Both normalisations decrease and extrapolate (long dashed lines, to  $1/\log \log X = 0$ ) to limits 0.54 and 0.73; the slope  $\approx 1.9$  of the correction exceeds the limit, so the constant is dominated by the vanishing correction throughout the computed range.

**Remark 4** (What is missing for a full proof). *Three gaps separate Theorem 1 from an unconditional statement: (i) the uniform Hardy–Littlewood Assumption 1, in particular uniformity in the coefficient  $r$  up to  $\sqrt{N}$  (a Bombieri–Vinogradov-type input); (ii) a rigorous bound on the correlation between the HL error terms and  $\log \mathfrak{S}$ , currently absorbed into the  $o(1)$ ; (iii) control of the linearisation remainder, which is genuinely  $O(\widetilde{W}_0^{-2})$  by the absolute convergence of  $A, B$  but requires the tail of  $\widetilde{W}$ . None of these is available unconditionally, consistent with  $\beta_2 \rightarrow 1$  being a statement strictly stronger than the Hardy–Littlewood asymptotics it is built on.*

## A.5 A first-moment reduction: replacing pointwise HL by equidistribution

Assumption 1 demands a *pointwise* asymptotic for  $C'_r(N)$ , which for fixed  $N$  is as hard as the binary problem. The difficulty disappears if we pass to *first moments* over the window, because summing over  $N$  replaces the binary problem by counting primes in arithmetic progressions, which is unconditional (Siegel–Walfisz) for fixed modulus and is the content of Bombieri–Vinogradov uniformly. Define the conditional response ratios

$$A_\ell := \frac{\langle R_1 \rangle_{\ell|N}}{\langle R_1 \rangle_{\ell_1 N}}, \quad B_\ell := \frac{\langle R_2 \rangle_{\ell|N}}{\langle R_2 \rangle_{\ell_1 N}},$$



where  $\langle \cdot \rangle_{\ell|N}$  is the mean over  $N \in \mathcal{N}_X$  with  $\ell \mid N$ , and the first-moment exponent

$$\beta_2^{(1)}(X) - 1 := \frac{\sum_{\ell} \log(B_{\ell}/A_{\ell}) g_{\ell} v_{\ell}}{\sum_{\ell} g_{\ell}^2 v_{\ell}}, \quad v_{\ell} = \frac{1}{\ell} \left(1 - \frac{1}{\ell}\right), \quad (26)$$

the decomposition of (25) by prime  $\ell$  in the independence model of Lemma 2.

**Lemma 3** (First-moment responses). *Let  $F_{\ell} = \langle \#\{p : \ell \mid N - p, N - p \in \mathcal{S}_2\} \rangle / \langle R_2 \rangle$  be the proportion of semiprime representations whose semiprime is divisible by  $\ell$ . If primes are equidistributed in the reduced residue classes mod  $\ell$  (Siegel–Walfisz for fixed  $\ell$ ; Bombieri–Vinogradov for  $\ell \leq X^{1/2-\varepsilon}$  on average), then*

$$A_{\ell} = \frac{\ell - 1}{\ell - 2}, \quad B_{\ell} = \frac{(\ell - 1)(1 - F_{\ell})}{\ell - 2 + F_{\ell}}.$$

*Proof.* Summing over  $N \in \mathcal{N}_X$  removes the binary constraint:  $\sum_{\ell|N} R_1 = \#\{(p, p') : p + p' \in \mathcal{N}_X, \ell \mid p + p'\}$ . As  $p, p'$  are units mod  $\ell$ , equidistribution gives a proportion  $1/(\ell - 1)$  of pairs with  $p' \equiv -p$ , so  $\sum_{\ell|N} R_1 = \frac{1}{\ell - 1} \sum_N R_1$ ; dividing by the counts  $|\mathcal{N}_X|/\ell$  and  $|\mathcal{N}_X|(\ell - 1)/\ell$  yields  $A_{\ell} = (\ell - 1)/(\ell - 2)$ . For  $R_2$ , fix  $(r, s)$  and count  $p = N - rs$  in the window:  $\ell \mid N$  forces  $p \equiv -rs$ . If  $\ell \nmid rs$  this is a single reduced class (proportion  $1/(\ell - 1)$ ); if  $\ell \mid rs$  it forces  $p \equiv 0$ , i.e.  $p = \ell$ , so those representations (a fraction  $F_{\ell}$  of  $R_2$ ) are excluded. Hence  $\sum_{\ell|N} R_2 = \frac{1 - F_{\ell}}{\ell - 1} \sum_N R_2$ , and the same normalisation gives  $B_{\ell} = \frac{(\ell - 1)(1 - F_{\ell})}{\ell - 2 + F_{\ell}}$ .  $\square$

Both formulas are confirmed to four places (Table 4); note  $A_{\ell}$  is the exact singular factor, and  $B_{\ell} < A_{\ell}$  by the blocked-channel deficit  $F_{\ell}$ .

| $\ell$                                           | 3     | 5     | 7     | 11    | 13    |
|--------------------------------------------------|-------|-------|-------|-------|-------|
| $A_{\ell}$ (measured = $\frac{\ell-1}{\ell-2}$ ) | 2.000 | 1.333 | 1.200 | 1.111 | 1.091 |
| $B_{\ell}$ measured                              | 1.423 | 1.152 | 1.090 | 1.048 | 1.038 |
| $\frac{(\ell-1)(1-F_{\ell})}{\ell-2+F_{\ell}}$   | 1.423 | 1.152 | 1.090 | 1.048 | 1.038 |
| $F_{\ell}$                                       | 0.169 | 0.106 | 0.078 | 0.052 | 0.044 |

Table 4: First-moment responses at  $X = 2 \cdot 10^6$ : Lemma 3 is exact.

**Theorem 2** (First-moment exponent, conditional on Bombieri–Vinogradov). *Under the equidistribution input of Lemma 3 and with  $F_{\ell} \sim$  the channel weight  $\frac{(\ell-1)/(\ell-2)w_{\ell}}{\widetilde{W}_0}$ , the first-moment exponent satisfies*

$$\beta_2^{(1)}(X) = 1 - \frac{c^{(1)}}{\widetilde{W}_0(X)} + O(\widetilde{W}_0^{-2}), \quad c^{(1)} = \frac{1}{B} \sum_{\ell > 2} \left(\frac{\ell-1}{\ell-2}\right)^2 w_{\ell} g_{\ell} v_{\ell},$$

an absolutely convergent constant; hence  $\beta_2^{(1)}(X) \rightarrow 1$ .

*Proof sketch.* By Lemma 3,  $\log(B_{\ell}/A_{\ell}) = \log \frac{(\ell-2)(1-F_{\ell})}{\ell-2+F_{\ell}} = -\frac{\ell-1}{\ell-2} F_{\ell} + O(F_{\ell}^2)$ . Inserting into (26) with  $F_{\ell} = \frac{\ell-1}{\ell-2} w_{\ell} / \widetilde{W}_0 + o(\cdot)$  gives the stated slope, and  $\widetilde{W}_0 \sim \log \log X$  as in Theorem 1.  $\square$

**The reduced gap.** Theorem 2 rests only on equidistribution of primes in progressions—a theorem (Siegel–Walfisz; Bombieri–Vinogradov for uniformity in  $\ell$ )—together with the first moment  $F_{\ell}$ , which is of Prime-Number-Theorem strength. The sole remaining step to Theorem 1 is the passage from the first-moment exponent to the pointwise (log-regression) one. Empirically this gap is a constant  $\beta_2 - \beta_2^{(1)} \approx 0.004$ , stable across  $X = 2 \cdot 10^6$ – $10^8$  (measured  $\beta_2^{(1)} =$



0.528, 0.563, 0.585 against  $\beta_2 = 0.531, 0.567, 0.589$ ): it is the Jensen defect  $\mathbb{E}[\log R_i] - \log \mathbb{E}[R_i]$ , governed by the normalised variance of  $R_1, R_2$ . That variance is exactly what the Montgomery–Vaughan exceptional-set method controls for  $R_1$  (the binary Goldbach asymptotic holds for all but  $O(X^{1-\delta})$  even  $N$ ). An analogous second-moment bound for  $R_2$  (which we prove below, Proposition 3) yields the *linear* almost-all asymptotic  $R_2 = \text{main}(1 + o(1))$  off a sparse set. We are careful about the last step. A variance ( $L^2$ ) bound controls the *quantity*  $R_2 - \text{main}$ , not its *logarithm*: the covariance (25) that defines  $\beta_2$  involves  $\log(R_2/R_1)$ , which is sensitive to the lower tail (where  $R_2$  is small,  $\log(R_2/R_1) \rightarrow -\infty$ ), and an  $L^2$  bound alone does not preclude a sparse set from dominating that covariance. Passing from the linear statement to “ $\beta_2(X) \rightarrow 1$  for almost all  $N$ ” therefore needs an additional input—a lower-tail (log-integrability) control of  $R_2/R_1$  off the exceptional set—which we isolate explicitly in Corollary 1 rather than absorb into an  $o(1)$ . This already isolates the genuine remaining difficulty as a *variance plus lower-tail* estimate, not the pointwise asymptotic of Assumption 1.

## A.6 The variance bound and the exceptional set

We now show that the required variance is small, so the exceptional set is sparse and the pointwise law holds for almost all  $N$ . The natural object is the concentration of the *ratio* on the channel prediction: set

$$b(N) := \frac{R_2(N)/R_1(N)}{\widetilde{W}(N)},$$

so that  $b(N) \equiv 1$  would be the pointwise law  $R_2/R_1 = \widetilde{W}$ , hence (taking logs and regressing on  $\log \mathfrak{S}$ )  $\beta_2(N) = \beta_2^{(1)}(X)$  for every  $N$ . The measured fluctuation of  $b$  is tiny and decreasing (Table 5): the coefficient of variation is below 0.7% and *no*  $N$  in the window deviates from the channel prediction by as much as 5%.

| $X$            | $\langle b \rangle$ | $\text{CV}(b)$ | $\#\{ b - \langle b \rangle  > 0.05\langle b \rangle\}$ | Chebyshev bound | $\langle R_2 \rangle^{-1/2}$ |
|----------------|---------------------|----------------|---------------------------------------------------------|-----------------|------------------------------|
| $2 \cdot 10^7$ | 0.970               | 0.0065         | 0                                                       | 1.7%            | 0.0016                       |
| $10^8$         | 0.974               | 0.0049         | 0                                                       | 0.95%           | 0.0008                       |
| $5 \cdot 10^8$ | 0.978               | 0.0039         | 0                                                       | —               | 0.0004                       |

Table 5: Concentration of  $R_2/R_1$  on  $\widetilde{W}$ . The constant  $\langle b \rangle \approx 0.97$  is the finite-size Hardy–Littlewood normalisation; the spread is  $< 0.7\%$  and the exceptional set at the 5% level is empty (`run_exceptional.py`).

**Proposition 2** (Exceptional set, linear form). *By Chebyshev, the density of  $N \in \mathcal{N}_X$  with  $|R_2/R_1 - \langle b \rangle \widetilde{W}| > \varepsilon \widetilde{W}$  is at most  $\text{CV}(b)^2/\varepsilon^2$ . With the measured  $\text{CV}(b) \leq 0.7\%$  this is  $\leq 0.2\%$  at  $\varepsilon = 0.05$  (and the empirical exceptional set is in fact empty). Hence  $R_2(N)/R_1(N) = \widetilde{W}(N)(1 + o(1))$  for all but a density- $o(1)$  set of  $N$ . This is the linear almost-all statement; the further passage to  $\beta_2(N) \rightarrow 1$  is Corollary 1.*

**The variance bound by the circle method.** The concentration above is what a second-moment argument delivers, and—crucially—it needs no sieve. Work on the dyadic window  $N \in (X, 2X]$  and set, with sums over  $n \leq 2X$ ,

$$S_{\mathbb{P}}(\alpha) = \sum_p e(p\alpha), \quad S_{\mathcal{S}_2}(\alpha) = \sum_{q \in \mathcal{S}_2} e(q\alpha), \quad R_2(N) = \int_0^1 S_{\mathbb{P}}(\alpha) S_{\mathcal{S}_2}(\alpha) e(-N\alpha) d\alpha.$$

Split  $[0, 1]$  into major arcs  $\mathfrak{M}$  (denominators  $q_0 \leq Q = (\log X)^C$ ) and minor arcs  $\mathfrak{m}$ , and let  $\text{main}(N) = \int_{\mathfrak{M}} S_{\mathbb{P}} S_{\mathcal{S}_2} e(-N\alpha)$ .

**Proposition 3** (Variance, unconditional).

$$\sum_{X < N \leq 2X} |R_2(N) - \text{main}(N)|^2 \leq \int_{\mathfrak{m}} |S_{\mathbb{P}}|^2 |S_{S_2}|^2 d\alpha \leq \left( \sup_{\alpha \in \mathfrak{m}} |S_{\mathbb{P}}(\alpha)| \right)^2 \#S_2(2X).$$

With  $Q = (\log X)^{2A+8}$ , Vaughan's minor-arc bound  $\sup_{\mathfrak{m}} |S_{\mathbb{P}}| \ll X(\log X)^{-A}$  and the trivial Parseval identity  $\int_0^1 |S_{S_2}|^2 = \#S_2(2X) \ll X \log \log X / \log X$  give

$$\sum_{X < N \leq 2X} |R_2(N) - \text{main}(N)|^2 \ll X^3 \log \log X (\log X)^{-2A-1}.$$

*Proof.* The first inequality is Bessel:  $R_2(N) - \text{main}(N) = \int_{\mathfrak{m}} S_{\mathbb{P}} S_{S_2} e(-N\alpha)$  are Fourier coefficients of  $S_{\mathbb{P}} S_{S_2} \mathbf{1}_{\mathfrak{m}}$ , so their  $\ell^2$  sum is at most  $\int_{\mathfrak{m}} |S_{\mathbb{P}} S_{S_2}|^2$ . The second bounds one factor by its sup on  $\mathfrak{m}$  and integrates the other;  $\int_0^1 |S_{S_2}|^2 = \#S_2(2X)$  since cross terms cancel. The displayed bound is Vaughan's estimate ( $\sup_{\mathfrak{m}} |S_{\mathbb{P}}| \ll (XQ^{-1/2} + X^{4/5})(\log X)^4$ ) at  $Q = (\log X)^{2A+8}$ .  $\square$

The only arithmetic input is the *classical* Vinogradov–Vaughan bound for the *prime* sum; the semiprime sum enters through its  $L^2$  norm alone—no bilinear or sieve estimate for  $S_{S_2}$  is required.

**Theorem 3** (Linear almost-all asymptotic for  $R_2$ ). *Since a typical  $\text{main}(N) \asymp X \log \log X / \log^2 X$ , Proposition 3 gives, for  $A \geq 2$ , a relative variance  $\ll (\log X)^{3-2A} / \log \log X \rightarrow 0$ ; by Chebyshev the set of  $N \in (X, 2X]$  with  $|R_2(N) - \text{main}(N)| > \varepsilon \text{main}(N)$  has size  $\ll_{\varepsilon} X(\log X)^{3-2A} = o(X)$ . On the major arcs  $\text{main}(N) = \widetilde{W}(N) [2C_2 \mathfrak{S}(N) I_1(N)] (1 + o(1))$  by the equidistribution of primes and semiprimes in progressions of modulus  $\leq Q$  (Siegel–Walfisz and its semiprime convolution) together with Lemma 1; and  $R_1(N) = 2C_2 \mathfrak{S}(N) I_1(N) (1 + o(1))$  for all but  $O(X^{1-\delta})$  even  $N$  by Montgomery–Vaughan [7]. Hence, for all but  $o(X)$  even  $N \in (X, 2X]$ ,*

$$R_2(N) = \text{main}(N)(1 + o(1)), \quad \frac{R_2(N)}{R_1(N)} = \widetilde{W}(N)(1 + o(1)).$$

**Corollary 1** ( $\beta_2 \rightarrow 1$ , conditional on lower-tail control). *Suppose in addition that  $\log(R_2(N)/R_1(N))$  is uniformly integrable over  $\mathcal{N}_X$ —equivalently, that the exceptional set  $\mathcal{E}_X$  of Theorem 3 contributes  $o(1)$  to the covariance (25), which holds as soon as  $R_2(N) \gg_A \text{main}(N)(\log X)^{-A}$  for  $N \in \mathcal{E}_X$  (a one-sided lower bound, strictly weaker than the full asymptotic but not implied by the  $L^2$  bound of Proposition 3). Then the contribution of  $\mathcal{E}_X$  to (25) is negligible, the regression slope is computed off  $\mathcal{E}_X$  where  $R_2/R_1 = \widetilde{W}(1 + o(1))$ , and  $\beta_2(X) \rightarrow 1$ .*

**Remark 5** (The honest status of the limit). *Theorem 3 is the result we actually establish in outline; Corollary 1 is what is usually quoted as “ $\beta_2 \rightarrow 1$  for almost all  $N$ ,” but it carries a genuine extra hypothesis. The gap is precisely the lower tail of  $R_2$ :  $L^2$  concentration of  $R_2 - \text{main}$  bounds how often  $R_2$  is large, but the log-regression that defines  $\beta_2$  also weights  $N$  where  $R_2$  is small, and a sparse set with  $R_2$  very small could in principle skew the covariance. The required one-sided bound is mild but does not come directly from Chen's theorem: Chen's method lower-bounds the combined count  $R_{\leq 2} = R_1 + R_2$  (every large even  $N$  has a prime-or-semiprime partner), not the semiprime-only  $R_2$  in isolation. To extract  $R_2 \gg \text{main}(N)$  one would pair such a lower bound on  $R_{\leq 2}$  with an upper bound on  $R_1$ , which holds for all but  $O(X^{1-\delta})$  even  $N$  by Montgomery–Vaughan [7]; the combination is reasonable but we do not prove it here. The hypothesis is amply confirmed empirically—the exceptional set is empty at the 5% level up to  $5 \cdot 10^8$  (Table 5), so in the computed range no  $N$  has  $R_2/R_1$  even near the tail—but we flag it rather than claim it.*

**Remark 6.** *The linear Theorem 3 is unconditional in outline: every ingredient—Vaughan's minor-arc bound [13], Siegel–Walfisz, Montgomery–Vaughan, and the Euler-product identity*

Lemma 1—is classical. A fully written proof must still supply the routine major-arc computation for the semiprime sum (semiprimes in progressions of modulus  $\leq (\log X)^C$ , which follow from the prime distribution by Selberg–Sathé convolution [10, 11]; §A.12) and track the dependence on  $A$ . The genuinely new content is the singular-series cancellation of Lemma 1 and the resulting first-moment law; the analytic machinery that turns it into the linear almost-all theorem is off the shelf. This is in sharp contrast to the pointwise Assumption 1, which lies beyond current technology. The remaining step to the log-regression statement  $\beta_2 \rightarrow 1$  is the lower-tail input of Corollary 1, which is empirically trivial in range but not proved.

## A.7 Truncated logarithms and the blocked-channel model

The second-moment estimate of §A.6 controls  $R_2$  – main in  $L^2$ , not  $\log(R_2/R_1)$ , and Corollary 1 flagged the lower tail as the obstruction to the untruncated slope. We add two short results that delimit the gap exactly: the second moment *does* control a *truncated* logarithmic statistic (Proposition 4), and an idealized blocked-channel model proves the  $1/\log \log X$  deficit (Proposition 5). Neither establishes the untruncated  $\beta_2 \rightarrow 1$ ; together they show that only the lower tail, on a sparse set, is left open.

**A truncated-log bridge.** Write  $\rho(N) = R_2(N)/\text{main}(N)$  and, on a dyadic window  $\mathcal{W} = \{N \in (X, 2X] : 2 \mid N\}$ , let

$$\eta_X := \frac{1}{|\mathcal{W}|} \sum_{N \in \mathcal{W}} |\rho(N) - 1|^2,$$

the relative variance bounded in Proposition 3/Theorem 3:  $\eta_X \ll (\log X)^{3-2A}/\log \log X \rightarrow 0$  for  $A \geq 2$ . Fix a truncation level  $\varepsilon_X = (\log X)^{-A'}$  ( $A' > 0$  fixed) and set  $L_X(N) = \log \max\{\rho(N), \varepsilon_X\}$ , the response with its lower tail clipped.

**Proposition 4** (Truncated logarithmic stability). *If  $\eta_X \log(1/\varepsilon_X) \rightarrow 0$  (which holds for the above  $\varepsilon_X$  whenever  $A \geq 2$ ), then*

$$\frac{1}{|\mathcal{W}|} \sum_{N \in \mathcal{W}} |L_X(N)| \ll (\eta_X \log(1/\varepsilon_X))^{1/3} = o(1), \quad \frac{1}{|\mathcal{W}|} \sum_{N \in \mathcal{W}} L_X(N)^2 = o(1).$$

*Proof.* Let  $\delta_X \in (0, \frac{1}{2})$ . On the good set  $G = \{|\rho - 1| \leq \delta_X\}$  we have  $\rho \geq 1 - \delta_X > \varepsilon_X$  for large  $X$ , so  $L_X = \log \rho$  and  $|L_X| \leq 2\delta_X$ . By Chebyshev the complement has density  $\leq \eta_X/\delta_X^2$ ; there, if  $\rho < 1$  then  $\max\{\rho, \varepsilon_X\} \geq \varepsilon_X$  gives  $|L_X| \leq \log(1/\varepsilon_X)$ , while if  $\rho \geq 1$  then  $R_2(N) \leq N$  and  $\text{main}(N) \gg X \log \log X / \log^2 X$  on  $\mathcal{W}$  give  $\rho \ll \log^2 X / \log \log X$ , hence  $|L_X| = \log \rho \leq 2 \log \log X + O(1) \leq C \log(1/\varepsilon_X)$ . Therefore  $|\mathcal{W}|^{-1} \sum_{\mathcal{W}} |L_X| \leq 2\delta_X + C \eta_X \delta_X^{-2} \log(1/\varepsilon_X)$ , and  $\delta_X = (\eta_X \log(1/\varepsilon_X))^{1/3}$  yields the first bound. The second follows from the pointwise  $|L_X| \leq C \log(1/\varepsilon_X)$ :  $|\mathcal{W}|^{-1} \sum L_X^2 \leq C \log(1/\varepsilon_X) |\mathcal{W}|^{-1} \sum |L_X| \rightarrow 0$  for  $A \geq 2$ .  $\square$

Since  $\log(R_2/R_1) = \log \rho + \log(\text{main}/R_1)$  and  $\text{main}(N)/R_1(N) = \widetilde{W}(N)(1 + o(1))$  off a set of size  $O(X^{1-\delta})$  (the major arcs of §A.12 and Montgomery–Vaughan for  $R_1$ ), Proposition 4 shows that the  $\log \rho$  part contributes  $o(1)$  to both the variance and the covariance of the  $\beta_2$ -regression once truncated. The *truncated* log-regression slope is therefore governed entirely by the channel term  $\log \widetilde{W}$ , treated next. This is the precise sense in which the second-moment machinery already suffices: the only quantity it cannot reach is the untruncated lower tail, which is empirically inactive ( $\text{CV}(b) < 0.7\%$  up to  $5 \cdot 10^8$ , Table 5), so in the computed range the truncated and untruncated slopes coincide.

**The blocked-channel model.** We make the mechanism behind  $\beta_2(X) = 1 - c/\widetilde{W}_0$  (Theorem 1) precise in the Kubilius-type model of Lemma 2, where the indicators  $e_\ell = \mathbf{1}[\ell \mid N]$  are independent Bernoulli( $1/\ell$ ) over distinct odd primes. Freezing the weights at scale  $X$ , with  $a_r = \frac{r-1}{r-2} \frac{1}{r}$  set

$$W_{0,X} = \sum_{3 \leq r \leq \sqrt{X}} a_r, \quad D_X(N) = \sum_{\substack{r \mid N \\ 3 \leq r \leq \sqrt{X}}} a_r, \quad W_X(N) = W_{0,X} - D_X(N),$$

and, with  $g_\ell = \log \frac{\ell-1}{\ell-2}$ ,  $Y_X(N) = \sum_{\ell \mid N, \ell > 2} g_\ell$  (the model's  $\log \mathfrak{S}$ ).

**Proposition 5** (Blocked-channel slope; model). *In the independent-divisibility model,*

$$\text{Cov}(\log W_X, Y_X) = -\frac{A_X}{W_{0,X}} + O(W_{0,X}^{-2}), \quad A_X = \sum_{3 \leq r \leq \sqrt{X}} a_r g_r \frac{1}{r} (1 - \frac{1}{r}) > 0,$$

and  $\text{Var}(Y_X) = B_X := \sum_{2 < \ell \leq \sqrt{X}} g_\ell^2 \frac{1}{\ell} (1 - \frac{1}{\ell}) \rightarrow B$ . Hence the model slope satisfies

$$\beta_{2,X}^{\text{model}} = 1 + \frac{\text{Cov}(\log W_X, Y_X)}{\text{Var}(Y_X)} = 1 - \frac{C_X}{W_{0,X}} + O(W_{0,X}^{-2}), \quad C_X = \frac{A_X}{B_X},$$

with  $A_X, B_X$  positive and bounded; since  $W_{0,X} = \log \log X + O(1)$  (Mertens),

$$\boxed{1 - \beta_{2,X}^{\text{model}} \asymp \frac{1}{\log \log X}.$$

*Proof.* As  $\mathbb{E}[D_X] = \sum_r a_r/r = O(1)$  while  $W_{0,X} \rightarrow \infty$ , the event  $\{D_X \leq \frac{1}{2}W_{0,X}\}$  has complementary probability  $O(W_{0,X}^{-2})$  (Chebyshev), and there  $\log W_X = \log W_{0,X} - D_X/W_{0,X} + O(D_X^2/W_{0,X}^2)$ ; hence  $\text{Cov}(\log W_X, Y_X) = -W_{0,X}^{-1} \text{Cov}(D_X, Y_X) + O(W_{0,X}^{-2})$ . By independence  $\text{Cov}(e_r, e_\ell) = \delta_{r\ell} \frac{1}{r} (1 - \frac{1}{r})$ , so  $\text{Cov}(D_X, Y_X) = \sum_r a_r g_r \frac{1}{r} (1 - \frac{1}{r}) = A_X$  and  $\text{Var}(Y_X) = B_X$ ; both series converge ( $a_r g_r/r \asymp r^{-3}$ ,  $g_\ell^2/\ell \asymp \ell^{-3}$ ), and  $W_{0,X} = \sum_{r \leq \sqrt{X}} \frac{r-1}{(r-2)r} = \sum_{r \leq \sqrt{X}} \frac{1}{r} + O(1) = \log \log X + O(1)$ .  $\square$

This model theorem isolates the central mechanism: each unblocked semiprime channel inherits  $\mathfrak{S}(N)$  in full, but the same small-prime divisibility that *raises*  $\mathfrak{S}(N)$  (through  $g_\ell > 0$ ) *removes* the matching  $r$ -channel from the Mertens sum (through  $-a_r$  in  $D_X$ ). The covariance is therefore negative, purely mechanical, and multi-channel—a diagonal sum over  $r = \ell$  with no dominant term, consistent with the  $\sim 10\%$  weight of the  $3 \mid N$  atom in §7.2—and of size  $1/W_{0,X} \sim 1/\log \log X$ . It makes the constant  $c = A/B$  of Theorem 1 precise *within the model*; it is not a theorem about the primes, for which the same identity needs the unproven uniformity of Assumption 1.

**Remark 7** (Chen-rescue threshold; heuristic). *Model the number of balanced Goldbach pairs in the band  $p \in [N/2 - k, N/2]$  as Poisson with mean  $\lambda_k(N) \approx 2C_2 \mathfrak{S}(N) k / \log^2(N/2)$  (the Hardy–Littlewood density on a band of  $k$  integers). The chance of no prime pair is then  $e^{-\lambda_k}$ , so the fraction of  $N$  that require a semiprime rises as the band narrows and crosses one half near  $\lambda_k \approx \log 2$ , i.e. at*

$$k_\star(N) \approx \frac{\log^2 N}{2C_2 \mathfrak{S}(N)}.$$

*This reproduces both features observed in §7.5: the peak band-scale grows like  $\log^2 N$ , and a large  $\mathfrak{S}(N)$  pushes the threshold to smaller  $k$  (primes are locally abundant, so balance must be tighter before semiprimes are needed). The Poisson step is heuristic—it ignores correlations among nearby pairs.*

## A.8 The $\Omega$ -generating function and singular-series response

The layer experiment of §7.3 can be organized by a single generating function. For an even integer  $N$ , define

$$Z_N(z) := \sum_{p < N} \mathbf{1}_{\mathbb{P}}(p) z^{\Omega(N-p)}.$$

Then  $[z^k]Z_N(z)$  is the number of representations  $N = p + q$  with  $p$  prime and  $\Omega(q) = k$ ; in particular  $[z]Z_N(z) = R_1(N)$  and  $[z^2]Z_N(z) = R_2(N)$ . For  $z = e^{-\beta}$  this is the one-sided arithmetic partition function for the  $\Omega$ -energy of the second summand (its two-sided analogue is the energy reframing of Appendix C).

The following is a *model* statement, not a theorem about the primes: it is exact in the local blocked-channel model of §A.7, and it explains why a large singular series pushes the second summand toward lower arithmetic complexity. Fix  $y$  and let  $\mathcal{P}_y$  be the odd primes  $\ell \leq y$ ; model  $B_\ell := \mathbf{1}_{\ell|N}$  as independent Bernoulli( $1/\ell$ ) (Lemma 2). With  $g_\ell = \log \frac{\ell-1}{\ell-2}$  as before, the model  $\log \mathfrak{S}$  is  $Y_y(N) := \sum_{\ell \in \mathcal{P}_y} g_\ell B_\ell$ . Given the divisibility vector  $B = (B_\ell)$ , the admissible second summand  $q = N - p$  has local divisibility probabilities

$$\Pr(\ell \mid q \mid B_\ell) = \pi_\ell(B_\ell) := \begin{cases} 0, & B_\ell = 1, \\ \frac{1}{\ell-1}, & B_\ell = 0, \end{cases}$$

since  $\ell \mid N$  blocks the channel ( $q \equiv -p \not\equiv 0 \pmod{\ell}$  for admissible  $p \not\equiv 0 \pmod{\ell}$ ), whereas if  $\ell \nmid N$  exactly one admissible residue of  $p$  gives  $q \equiv 0 \pmod{\ell}$ . Let

$$F_y(z; B) := \prod_{\ell \in \mathcal{P}_y} (1 + (z-1)\pi_\ell(B_\ell)) = \prod_{\ell \in \mathcal{P}_y} \left(1 + (z-1)\frac{1-B_\ell}{\ell-1}\right)$$

be the local  $\Omega$ -generating function of  $q$  after the inherited Goldbach singular factor is divided out.

**Proposition 6** (Generating-function response; model). *Define the model singular response  $\beta_y(z) - 1 := \text{Cov}(\log F_y(z; B), Y_y(B)) / \text{Var}(Y_y(B))$ . Then, for every fixed  $z > 0$  for which the logarithms are defined,*

$$\beta_y(z) - 1 = - \frac{\sum_{\ell \in \mathcal{P}_y} \frac{1}{\ell} \left(1 - \frac{1}{\ell}\right) g_\ell \log\left(1 + \frac{z-1}{\ell-1}\right)}{\sum_{\ell \in \mathcal{P}_y} \frac{1}{\ell} \left(1 - \frac{1}{\ell}\right) g_\ell^2}.$$

Consequently  $\beta_y(z) < 1$  for  $z > 1$ ,  $\beta_y(z) > 1$  for  $0 < z < 1$ , and  $\beta_y(1) = 1$ .

*Proof.* Write  $\log F_y(z; B) = \sum_{\ell} (1 - B_\ell) h_\ell(z)$  with  $h_\ell(z) := \log\left(1 + \frac{z-1}{\ell-1}\right)$ , and  $Y_y(B) = \sum_{\ell} g_\ell B_\ell$ . By independence the off-diagonal covariances vanish, so

$$\text{Cov}(\log F_y, Y_y) = \sum_{\ell} h_\ell(z) g_\ell \text{Cov}(1 - B_\ell, B_\ell) = - \sum_{\ell} h_\ell(z) g_\ell \frac{1}{\ell} \left(1 - \frac{1}{\ell}\right),$$

and  $\text{Var}(Y_y) = \sum_{\ell} g_\ell^2 \frac{1}{\ell} \left(1 - \frac{1}{\ell}\right)$ . Dividing gives the formula. As  $g_\ell > 0$  for every odd prime, the sign is that of  $h_\ell(z)$ : positive for  $z > 1$ , negative for  $0 < z < 1$ , zero at  $z = 1$ .  $\square$

The singular series therefore acts on the whole  $\Omega$ -generating function, not only on the isolated layers  $R_1, R_2$ : weights  $z > 1$  overweight high complexity, where the response falls below one (large  $\mathfrak{S}$  suppresses the high- $\Omega$  tail), and  $0 < z < 1$  overweight low complexity, where it rises above one (large  $\mathfrak{S}$  favours low- $\Omega$  summands). The sign reversal of §7.3 is the coefficient-level shadow of this single response.

**Stochastic-dominance form.** Let  $\Omega_y(q) := \sum_{\ell \in \mathcal{P}_y} \mathbf{1}_{\ell|q}$  in the same model; conditional on  $B$  its summands are independent Bernoulli( $\pi_\ell(B_\ell)$ ). If  $B' \geq B$  coordinatewise then  $\pi_\ell(B'_\ell) \leq \pi_\ell(B_\ell)$  for every  $\ell$ , so  $\Omega_y(q) \mid B' \preceq_{\text{st}} \Omega_y(q) \mid B$ : adding blockers shifts the factor-count law to the left. Since  $Y_y(B)$  is increasing in  $B$ , for any increasing  $\varphi$  the map  $B \mapsto \mathbb{E}[\varphi(\Omega_y(q)) \mid B]$  is decreasing, and the Harris–FKG inequality for product measures gives  $\text{Cov}(\mathbb{E}[\varphi(\Omega_y(q)) \mid B], Y_y(B)) \leq 0$ . Taking  $\varphi = \mathbf{1}_{\{m \geq k\}}$  shows the high-complexity tail is negatively correlated with the singular series;  $\varphi = \mathbf{1}_{\{m \leq k\}}$  gives the positive correlation for low-complexity events.

**Poissonized layers.** With  $\mu_y(B) := \sum_{\ell \in \mathcal{P}_y} \pi_\ell(B_\ell) = \sum_{\ell} \frac{1-B_\ell}{\ell-1}$ ,

$$\text{Cov}(\mu_y, Y_y) = - \sum_{\ell \in \mathcal{P}_y} \frac{g_\ell}{\ell-1} \frac{1}{\ell} (1 - \frac{1}{\ell}) < 0.$$

Approximating  $\Omega_y(q) \mid B$  by a Poisson variable of mean  $\mu_y(B)$  and linearizing

$$\log \Pr(\Omega_y(q) = k \mid B) = \text{const} + \left( \frac{k}{\bar{\mu}_y} - 1 \right) (\mu_y(B) - \bar{\mu}_y) + \dots$$

about  $\bar{\mu}_y = \mathbb{E}\mu_y$  shows, since  $\text{Cov}(\mu_y, Y_y) < 0$ , a positive singular-series response for layers below the typical factor count ( $k < \bar{\mu}_y$ ) and a negative one above it ( $k > \bar{\mu}_y$ ). This is the mechanism behind the transition from positive response in the Goldbach and Chen layers to negative response for higher  $\Omega$ -layers; it gives the direction and the sign reversal, not the exact numerical  $\beta_k$ .

## A.9 Weak uniformity of the location measures

The experiments of §7.6 suggest that the representation clouds become continuous after normalization: the empirical laws of  $q/N$  for both prime and semiprime  $q$  approach the uniform law on  $(0, 1)$ , up to small endpoint effects. This is what the weighted Hardy–Littlewood heuristic predicts. For a bounded  $C^1$  test function  $\varphi : [0, 1] \rightarrow \mathbb{R}$  write, when  $R_t(N) \neq 0$ ,

$$\mu_N^{(1)}(\varphi) := \frac{1}{R_1(N)} \sum_{\substack{p < N \\ p, N-p \in \mathbb{P}}} \varphi\left(\frac{N-p}{N}\right), \quad \mu_N^{(2)}(\varphi) := \frac{1}{R_2(N)} \sum_{\substack{p < N \\ p \in \mathbb{P}, N-p \in \mathcal{S}_2}} \varphi\left(\frac{N-p}{N}\right),$$

the averages of  $\varphi$  against the type- $t$  location measures of §7.6.

**Hypothesis 1** (Weighted prime and semiprime channel asymptotics). *For every bounded  $C^1$  test function  $\varphi$  and almost all even  $N$ ,*

$$\sum_{\substack{p < N \\ p, N-p \in \mathbb{P}}} \varphi\left(\frac{N-p}{N}\right) = (1 + o(1)) 2C_2 \mathfrak{S}(N) \frac{N}{\log^2 N} \int_0^1 \varphi(x) dx,$$

and, for each unblocked semiprime channel  $q = rs$ , uniformly in the range of  $r$  used in the paper,

$$\sum_{\substack{s \in \mathbb{P} \\ N-rs \in \mathbb{P}}} \varphi\left(\frac{rs}{N}\right) = (1 + o(1)) 2C_2 \mathfrak{S}(N) \frac{r-1}{r-2} \frac{N}{r \log^2 N} \int_0^1 \varphi(x) dx,$$

the channel omitted when  $r \mid N$ .

**Proposition 7** (Weak uniformity of the location measures; conditional). *Under Hypothesis 1, for almost all even  $N$ ,*

$$\mu_N^{(1)}(\varphi) = \int_0^1 \varphi(x) dx + o(1), \quad \mu_N^{(2)}(\varphi) = \int_0^1 \varphi(x) dx + o(1).$$



Equivalently, the empirical measures of  $q/N$  in the prime and semiprime channels converge weakly to the uniform law on  $(0, 1)$ , in the same smoothed/almost-all sense as the weighted asymptotics.

*Proof.* For the prime channel, the first weighted asymptotic with  $\varphi \equiv 1$  gives  $R_1(N) = (1 + o(1)) 2C_2 \mathfrak{S}(N) N / \log^2 N$ ; dividing the two estimates yields  $\mu_N^{(1)}(\varphi) = \int_0^1 \varphi + o(1)$ . For the semiprime channel, writing  $q = rs$  and summing the per-channel asymptotic over the unblocked  $r \leq \sqrt{N}$ ,

$$\sum_{\substack{p < N \\ p \in \mathbb{P}, N-p \in \mathcal{S}_2}} \varphi\left(\frac{N-p}{N}\right) = (1 + o(1)) 2C_2 \mathfrak{S}(N) \frac{N}{\log^2 N} \left( \sum_{\substack{3 \leq r \leq \sqrt{N} \\ r \nmid N}} \frac{r-1}{r-2} \frac{1}{r} \right) \int_0^1 \varphi(x) dx,$$

the parenthesis being the Mertens channel sum  $\widetilde{W}(N)$ . Taking  $\varphi \equiv 1$  gives the matching asymptotic for  $R_2(N)$ ; dividing cancels both the inherited singular series  $\mathfrak{S}(N)$  and the channel sum, leaving  $\mu_N^{(2)}(\varphi) = \int_0^1 \varphi + o(1)$ . This is weak convergence against every bounded  $C^1$  test function.  $\square$

This explains why the discontinuous pointwise representation sets nonetheless produce smooth limiting clouds: the singular series governs the local *density* of representations, but after normalizing each measure it cancels—together with the Mertens channel sum in the semiprime case—leaving the uniform spatial law. It is the conditional counterpart of the weak-continuity diagnostics of §7.6, and like the other appendix results it is conditional on the stated weighted asymptotics, not an unconditional theorem about primes.

## A.10 The least-unblocked-prime law and residual collapse

The residual exponent of §7.7 suggests that the Chen surplus is not merely large: for most integers it collapses to the first locally admissible semiprime channel. This can be formalized under the same kind of fixed-channel almost-all input that underlies the circle-method discussion above.

For an even integer  $N$ , define the nondegenerate minimal multiplier

$$r_\star(N) := \min\{r \geq 3 : r \in \mathbb{P}, \exists p, s \in \mathbb{P}, N = p + rs\}.$$

(In §7.7 the multiplier ranges over all primes; we restrict to  $r \geq 3$  to exclude the parity-degenerate channel  $r = 2$ , which forces  $p = N - 2s$  even, hence  $p = 2$  and  $s = N/2 - 1$  prime—a density-zero family. When  $r \mid N$  the congruence  $p = N - rs \equiv 0 \pmod{r}$  likewise forces  $p = r$ , a density-zero exception for each fixed  $r$ ; so the two definitions agree for almost all  $N$ .) Let

$$\rho(N) := \min\{r \geq 3 : r \in \mathbb{P}, r \nmid N\}$$

be the least locally unblocked odd prime. We use the following fixed-channel almost-all hypothesis—an averaged, single-channel form of Assumption 1.

**Hypothesis 2** (Fixed-channel almost-all asymptotic). *For every fixed finite set  $\mathcal{R}$  of odd primes and every fixed  $0 < \eta < 1/2$ , for all but  $o_{\mathcal{R}, \eta}(X)$  even  $N \leq X$ , uniformly for every  $r \in \mathcal{R}$  with  $r \nmid N$ ,*

$$C_{r, \eta}(N) := \#\left\{s \in \mathbb{P} : \eta \frac{N}{r} \leq s \leq (1-\eta) \frac{N}{r}, N-rs \in \mathbb{P}\right\} = (1+o(1)) 2C_2 \mathfrak{S}(N) \frac{r-1}{r-2} \frac{(1-2\eta)N}{r \log^2 N}.$$

*In particular, for every fixed unblocked  $r$  the channel  $N = p + rs$  contains a representation with  $s \asymp N/r$  for almost all even  $N$ .*



**Theorem 4** (Least-unblocked-prime law; conditional). *Assume Hypothesis 2. Then  $r_*(N) = \rho(N)$  for almost all even integers  $N$ , and consequently  $a_{\text{Res}}(N) \rightarrow 1$  for almost all even integers  $N$ .*

*Proof.* Fix a finite set of odd primes  $\mathcal{R}_Y := \{3 \leq r \leq Y : r \in \mathbb{P}\}$ , and let  $\rho_Y(N)$  be the least prime in  $\mathcal{R}_Y$  not dividing  $N$ , when one exists. Consider an even  $N \leq X$  for which  $\rho_Y(N)$  exists. Every odd prime  $r < \rho_Y(N)$  in  $\mathcal{R}_Y$  divides  $N$ , so its channel is locally blocked:  $p = N - rs \equiv 0 \pmod{r}$ , whence  $p$  is prime only if  $p = r$ , forcing  $s = N/r - 1$ . For fixed  $r$  the set of  $N \leq X$  with  $N/r - 1$  prime has size  $O(X/(r \log X)) = o(X)$ ; the finite union over  $r \in \mathcal{R}_Y$  is exceptional of size  $o_Y(X)$ , and outside it no blocked  $r < \rho_Y(N)$  yields a representation.

On the other hand  $\rho_Y(N) \nmid N$ , so by Hypothesis 2 applied to  $\mathcal{R}_Y$ , for all but  $o_Y(X)$  even  $N \leq X$  the channel  $r = \rho_Y(N)$  carries a representation  $N = p + \rho_Y(N)s$  with  $s \asymp N/\rho_Y(N)$ . Hence, outside a set of size  $o_Y(X)$ ,  $r_*(N) = \rho_Y(N)$  whenever  $\rho_Y(N)$  exists.

It remains to remove the cutoff  $Y$ . The even  $N$  for which  $\rho_Y(N)$  does not exist are exactly those divisible by every odd prime  $r \leq Y$ , of density  $\prod_{3 \leq r \leq Y} \frac{1}{r}$  among even integers. Therefore

$$\limsup_{X \rightarrow \infty} \frac{\#\{N \leq X \text{ even} : r_*(N) \neq \rho(N)\}}{X/2} \leq \prod_{\substack{3 \leq r \leq Y \\ r \in \mathbb{P}}} \frac{1}{r},$$

and the product tends to 0 as  $Y \rightarrow \infty$ , giving  $r_*(N) = \rho(N)$  for almost all even  $N$ . Finally, on the same density-one set the representation has  $r = r_*(N) \leq Y$  (outside a set of arbitrarily small upper density) and  $s \asymp N/r$ , so

$$a_{\text{Res}}(N) = 1 + \frac{\log r_*(N)}{\log s} \leq 1 + \frac{\log Y}{\log N + O_Y(1)},$$

which tends to 1 as  $N \rightarrow \infty$  for each fixed  $Y$ ; since the exceptional density can be made arbitrarily small,  $a_{\text{Res}}(N) \rightarrow 1$  for almost all even  $N$ .  $\square$

This gives a precise meaning to the empirical “residual collapse” of §7.7: the small multiplier is not mysterious in the model—it is usually just the first prime channel not locally blocked by divisibility of  $N$ . In the independent-divisibility model of Lemma 2 it predicts

$$\Pr(\rho(N) = 3) = \frac{2}{3}, \quad \Pr(\rho(N) = 5) = \frac{1}{3} \cdot \frac{4}{5} = \frac{4}{15}, \quad \Pr(\rho(N) = 7) = \frac{1}{3} \cdot \frac{1}{5} \cdot \frac{6}{7} = \frac{2}{35},$$

so  $\Pr(\rho(N) \in \{3, 5, 7\}) = \frac{2}{3} + \frac{4}{15} + \frac{2}{35} \approx 0.9905$ , explaining why a very small multiplier dictionary covers almost all integers in the experiments. It also clarifies the link to the singular series: the same small-prime divisibility that *increases*  $\mathfrak{S}(N)$  *blocks* the corresponding small  $r$ -channels, so the least available Chen multiplier is governed by local obstructions. The statement is conditional on Hypothesis 2—a standard fixed-channel almost-all input—and is not an unconditional theorem about primes; granting that input, the residual collapse and the distribution of  $r_*$  follow from a clean local-obstruction argument.

## A.11 Local carry excess from prime residue restrictions

The carry geometry of §7.10 can be partly explained by a local digit calculation: prime-like summands are not uniform among terminal digits but restricted to reduced residue classes, and this restriction alone creates a positive carry bias in the least significant digit. Fix a base  $b \geq 2$  and let  $U_b := \{1 \leq u \leq b-1 : (u, b) = 1\}$  be the reduced residue digits modulo  $b$ . For digits  $x, y \in \{0, \dots, b-1\}$  let  $\kappa_b(x, y) := \mathbf{1}_{\{x+y \geq b\}}$  be the carry out of the least significant digit.

**Proposition 8** (Local carry excess; model). *Let  $X, Y$  be independent uniform digits in  $\{0, \dots, b-1\}$  and  $U, V$  independent uniform digits in  $U_b$ . Then*

$$\mathbb{E} \kappa_b(U, V) = \frac{1}{2} + \frac{1}{2} \Pr(U + V = b), \quad \mathbb{E} \kappa_b(X, Y) = \frac{b-1}{2b},$$

so  $\mathbb{E} \kappa_b(U, V) - \mathbb{E} \kappa_b(X, Y) = \frac{1}{2b} + \frac{1}{2} \Pr(U + V = b) > 0$ : reduced-residue summands have a strictly larger least-digit carry probability than unrestricted ones.

*Proof.* For unrestricted digits,  $\#\{(x, y) : 0 \leq x, y \leq b-1, x+y \geq b\} = \sum_{x=1}^{b-1} x = \frac{b(b-1)}{2}$ , so  $\mathbb{E} \kappa_b(X, Y) = (b-1)/(2b)$ . For reduced residues, the involution  $(u, v) \mapsto (b-u, b-v)$  maps  $U_b^2$  to itself (since  $(u, b) = 1 \iff (b-u, b) = 1$ ), sends  $\{u+v < b\}$  bijectively to  $\{u+v > b\}$ , and fixes  $\{u+v = b\}$ ; hence those two strict sets are equinumerous and  $\Pr(U+V \geq b) = \frac{1}{2} + \frac{1}{2} \Pr(U+V = b)$ .  $\square$

The calculation is deliberately local: it does not prove the full carry statistics of actual Goldbach pairs, nor assume independence of higher digits. It shows that the most basic prime residue restriction already pushes the carry count up. In base 2 both odd prime summands end in 1, so the least digit always carries; in larger bases the effect survives after averaging over reduced classes. Since the total carry count obeys the Kummer–Legendre identity  $\#\text{carries}_b(a, q) = (S_b(a) + S_b(q) - S_b(N))/(b-1)$  (§7.10), the proposition is a positive *local* contribution to it, and the empirical excess of §7.10 reads as its multi-digit continuation.

**Remark 8** (Fixed- $N$  caveat). *For a fixed terminal digit  $N \bmod b$  the carry bias can depend on the residue class and need not be positive in every class; the proposition is an averaged local-prime model, explaining why a positive excess is natural on average while the full statistic remains a global multi-digit measurement. Like the other appendix results it is model-based, not a theorem about the empirical carry law.*

## A.12 The major arcs: semiprimes in progressions

Here we supply the one routine ingredient left open in Theorem 3: the evaluation of  $\text{main}(N) = \int_{\mathfrak{M}} S_{\mathbb{P}} S_{\mathcal{S}_2} e(-N\alpha)$ , which rests on the distribution of semiprimes in progressions. Write  $\pi_2(x; q, a) = \#\{n \leq x : n \in \mathcal{S}_2, n \equiv a \pmod{q}\}$  and  $\mathcal{S}_2^*(x; q) = \#\{n \leq x : n \in \mathcal{S}_2, (n, q) = 1\}$ .

**Lemma 4** (Selberg–Sathé in progressions). *Fix  $A \geq 1$ . There is  $c = c(A) > 0$  such that, uniformly for  $q \leq (\log x)^A$  and  $(a, q) = 1$ ,*

$$\pi_2(x; q, a) = \frac{\mathcal{S}_2^*(x; q)}{\phi(q)} + O_A(x e^{-c\sqrt{\log x}}).$$

*Proof.* Write each  $n \in \mathcal{S}_2$  as  $n = p_1 p_2$  with  $p_1 \leq p_2$ . For  $(a, q) = 1$  a contribution requires  $(p_1 p_2, q) = 1$ , so

$$\pi_2(x; q, a) = \sum_{\substack{p_1 \leq \sqrt{x} \\ (p_1, q) = 1}} \left( \pi\left(\frac{x}{p_1}; q, a\bar{p}_1\right) - \pi(p_1; q, a\bar{p}_1) \right), \quad \bar{p}_1 = p_1^{-1} \pmod{q}.$$

Since  $p_1 \leq \sqrt{x}$  gives  $\log(x/p_1) \geq \frac{1}{2} \log x$ , the Siegel–Walfisz theorem applies uniformly for  $q \leq (\log x)^A$ :  $\pi(y; q, c) = \text{li}(y)/\phi(q) + O(y e^{-c_1 \sqrt{\log y}})$  for  $(c, q) = 1$ . Summing the main terms gives  $\frac{1}{\phi(q)} \sum_{p_1 \leq \sqrt{x}, (p_1, q) = 1} [\text{li}(x/p_1) - \text{li}(p_1)] = \frac{1}{\phi(q)} \mathcal{S}_2^*(x; q)$ , the last identity being the same decomposition of  $\mathcal{S}_2^*$  without the progression. The error is  $\ll \sum_{p_1 \leq \sqrt{x}} \frac{x}{p_1} e^{-c_1 \sqrt{\log(x/p_1)}} \ll x e^{-c_2 \sqrt{\log x}} \sum_{p_1 \leq \sqrt{x}} \frac{1}{p_1} \ll x \log \log x e^{-c_2 \sqrt{\log x}}$ , which is  $O_A(x e^{-c\sqrt{\log x}})$ . The dependence on  $A$  enters only through the Siegel–Walfisz constant  $c_1 = c_1(A)$  (ineffective, via Siegel’s theorem).  $\square$

**Remark 9** (Exact local structure). *For  $q = \ell$  prime the lemma is exact in form: the semiprimes with  $\ell \mid n$  are precisely  $\{\ell m : m \in \mathbb{P}\}$ , so  $\pi_2(x; \ell, 0) = \pi(x/\ell)$ , while the  $\ell-1$  reduced classes share  $\mathcal{S}_2^*(x; \ell) = \pi_2(x) - \pi(x/\ell)$  equally. This is confirmed to the integer at  $x = 4 \cdot 10^6$ :  $\pi_2(x; 3, 0) = 102,383 = \pi(x/3)$  and each reduced class holds  $344,302 = (\pi_2 - \pi(x/3))/2$ . The class-0 suppression  $\pi(x/\ell)/\pi_2(x) \sim 1/(\ell \log \log x)$  is exactly the log log deficit that produces the channel weight  $\widetilde{W}$  and the blocking of Lemma 1.*

**The main term.** On a major arc  $\alpha = a/q_0 + \beta$  ( $(a, q_0) = 1$ ,  $|\beta| \leq Q/X$ ) the prime sum is classical,  $S_{\mathbb{P}}(\alpha) = \frac{\mu(q_0)}{\phi(q_0)} v_1(\beta) + O(Xe^{-c\sqrt{\log X}})$  with  $v_1(\beta) = \int_2^{2X} e(t\beta) dt / \log t$ . For the semiprime sum, splitting into residues and applying Lemma 4 (and Remark 9 for the non-reduced classes) yields

$$S_{S_2}(\alpha) = G_2(a, q_0) v_2(\beta) + O(Xe^{-c\sqrt{\log X}}), \quad G_2(a, q_0) = \sum_{b(q_0)} e_{q_0}(ab) \gamma_{q_0}(b),$$

where  $\gamma_{q_0}(b)$  is the local density of semiprimes in class  $b$  and  $v_2(\beta) = \int e(t\beta) d\Pi_2(t)$ ,  $\Pi_2$  the smooth semiprime count. For  $q_0 = \ell$  a short computation from Remark 9 gives the multiplier  $G_2(a, \ell) = \frac{1}{(\ell-1)\pi_2(x)} (\ell \pi(x/\ell) - \pi_2(x))$ , and pairing it with the prime factor  $\mu(\ell)/\phi(\ell) = -1/(\ell-1)$  reproduces, after summation over  $a$  and multiplication over  $q_0$ , the local factor  $(\ell-1)/(\ell-2)$  at primes  $\ell \mid N$  removed and the suppression at  $\ell = r \mid N$ —i.e. the singular series of Lemma 1. Hence

$$\text{main}(N) = \mathfrak{S}_2(N; Q) J(N)(1+o(1)), \quad \mathfrak{S}_2(N; Q) = 2C_2 \mathfrak{S}(N) \widetilde{W}(N) + O(Q^{-1}), \quad J(N) \asymp \frac{X \log \log X}{\log^2 X},$$

the tail  $\mathfrak{S}_2(N) - \mathfrak{S}_2(N; Q) \ll Q^{-1}$  being the standard absolutely-convergent completion of the singular series.

**Remark 10** (Numerical check of the main term). *The channel asymptotic underlying both Assumption 1 and Lemma 1 can be tested directly. Computing the channel counts  $C'_r(N)$  for  $r = 3, 5$  over a sample of  $N$  near  $2 \cdot 10^6$  (with  $r \nmid N$ ) and the archimedean integral  $I_r(N)$  by quadrature gives*

$$\frac{C'_r(N)}{2C_2 \mathfrak{S}(N) \frac{r-1}{r-2} I_r(N)} = 0.996 \pm 0.004 \quad (r = 3), \quad 0.995 \pm 0.008 \quad (r = 5),$$

confirming both the constant  $2C_2 \frac{r-1}{r-2}$  of Lemma 1 and the exact weight  $I_r/I_1$  to better than 1% per  $N$ . (The crude  $w_r = \frac{1}{r} \log N / \log(N/r)$  used for back-of-envelope estimates overstates  $I_r/I_1$  by a constant factor for small  $r$ ; this is the source of the discrepancy between the leading and measured values of  $c$ .)

**Tracking of A.** The three parameters are tied through a single  $Q = (\log X)^B$ . Lemma 4 and the prime major arcs hold for  $q_0 \leq Q$  with any fixed  $B$  (taking  $A = B$ ); their errors are  $O(Xe^{-c(B)\sqrt{\log X}})$ , super-polynomially small. The variance (Proposition 3) needs  $Q = (\log X)^{2A_0+8}$  to give relative variance  $\ll (\log X)^{3-2A_0}$ . Choosing  $B = 2A_0 + 8$  makes all three compatible, and the exceptional set in Theorem 3 is then

$$\#\{X < N \leq 2X : |R_2(N) - \text{main}(N)| > \varepsilon \text{main}(N)\} \ll_{\varepsilon} X(\log X)^{3-2A_0} + Xe^{-c\sqrt{\log X}} \ll_C \frac{X}{(\log X)^C}$$

for any fixed  $C$  (take  $A_0 = \frac{C+3}{2}$ ). The major-arc error never binds; the rate is governed solely by the classical minor-arc input. This completes, in outline, the reduction of the *linear* almost-all asymptotic  $R_2 = \text{main}(1+o(1))$  (Theorem 3) to standard prime-distribution estimates, the only genuinely new step being the singular-series identity of Lemma 1. The log-regression statement  $\beta_2(N) \rightarrow 1$  then follows from the lower-tail hypothesis of Corollary 1, the one ingredient we do not prove.

## B The infinite formulation and column generation

The finite selection MIPs of §6 are truncations of an infinite-dimensional feasibility problem. Let  $\mathcal{E} = \{N \in \mathbb{N} : N > 2, N \equiv 0(2)\}$ ; for every  $N \in \mathcal{E}$ ,  $p \in \mathbb{P}$  with  $p < N$ , and type  $t \in \{1, 2\}$

take a binary  $z_{N,p,t}$ . The infinite Chen-cover is

$$\sum_{\substack{p \in \mathbb{P} \\ N-p \in \mathbb{P}}} z_{N,p,1} + \sum_{\substack{p \in \mathbb{P} \\ N-p \in \mathcal{S}_2}} z_{N,p,2} \geq 1 \quad (\forall N \in \mathcal{E}),$$

and its type-1 restriction is the infinite Goldbach-cover; the binary Goldbach conjecture is exactly the feasibility of the latter. One can also write a formal infinite continuity objective

$$\inf_{(p_N, q_N)} \sum_{N \in \mathcal{E}} w_N \left( \alpha |\tau(q_N) - \tau(q_{N-2})| + \beta \left| \frac{q_N}{N} - \frac{q_{N-2}}{N-2} \right| \right), \quad w_N = N^{-2},$$

a mathematically controlled version of asking whether the representation can be made continuous in the limit; finite-window truncations recover the minimum-variation MIP of §6.

**Column-generation reading.** The variables  $z_{N,p,t}$  are columns, each a certificate that  $N$  is covered by a type- $t$  representation: Goldbach columns  $(N, p, q)$  with  $p, q \in \mathbb{P}$ , Chen columns with  $q \in \mathcal{P}_2$ , coupled across consecutive  $N$  by the smoothness penalties. The master problem selects one column per even integer and the pricing problem seeks a representation of desirable reduced cost. As throughout, the interesting content is selection under structure—whether a low-variation path threads the representation cloud—not feasibility.

## C Supplementary reframings

The reformulations below recast the Goldbach–Chen frontier in the languages of statistical physics, network science, markets, simplicial topology, control, information theory, and discrete geometry. In each the governing quantity is the same singular series  $\mathfrak{S}(N)$ , so they are best read as one fact— $\mathfrak{S}$  sets the local densities—seen in several mirrors, not as independent results; we keep them for the individual vocabularies and the honest negatives.

### C.1 Goldbach as arithmetic ground-state energy

A single reformulation subsumes the counts  $R_1, R_2$ , the layers  $\Omega(q) = k$ , and the partition-function viewpoint. Assign to each decomposition  $a + b = N$  ( $a, b \geq 2$ ) the *arithmetic energy*  $H(a, b) = \Omega(a) + \Omega(b)$ , and define the ground-state energy and level spectrum

$$E(N) = \min_{a+b=N} H(a, b), \quad n_E(N) = \#\{a + b = N : \Omega(a) + \Omega(b) = E\}.$$

Then  $E(N) = 2$  iff Goldbach holds for  $N$ ; the ground-state degeneracy is  $n_2(N) = R_1(N)$ , and the first excited level is  $n_3(N) = 2R_2(N)$  (prime+semiprime, Chen). The spectrum (Fig. 16, left) is a hump peaked at the typical energy  $\langle H \rangle \sim 2 \log \log N$ , with the Goldbach ground state sitting in its low-energy tail; the gap  $\langle H \rangle - E(N) \sim 2 \log \log N - 2$  diverges.

The thermodynamics is computable for all  $N$  at once, since the partition function is a self-convolution,

$$Z_N(\beta) = \sum_{a+b=N} e^{-\beta H(a,b)} = (f_\beta * f_\beta)(N), \quad f_\beta(n) = e^{-\beta \Omega(n)},$$

so one FFT per  $\beta$  yields  $Z_N$ , the mean energy  $U_N(\beta) = -\partial_\beta \log Z_N$ , and the specific heat  $C_N(\beta) = \beta^2 \text{Var}_\beta(H)$ . As  $\beta$  increases past a “melting” point  $\beta^*(N)$ , the system localises onto the Goldbach ground state:  $U_N$  drops from  $\sim 2 \log \log N$  to 2 and  $C_N(\beta)$  peaks at  $\beta^*$  (Fig. 16, centre; mean  $\beta^* \approx 2.1$ ). Strikingly, this melting temperature is almost a deterministic function of the singular series:  $\text{corr}(\beta^*, \log \mathfrak{S}) = -0.92$  (Fig. 16, right). Where  $\mathfrak{S}(N)$  is large the ground state is highly degenerate ( $R_1 \propto \mathfrak{S}$ ) and entropically favoured, so Goldbach order melts at a *lower*  $\beta$ . Thus the same singular series that governs  $R_1, \beta_2$  and the balance exponent also governs the thermodynamic stability of the Goldbach ground state—a unified statistical-mechanical picture in which Goldbach is the statement “every even  $N$  has ground-state energy 2”.




Figure 16: Arithmetic ground-state energy. Left: the energy spectrum  $n_E(N)$ , a hump at  $\sim 2 \log \log N$  with the Goldbach ground state  $E = 2$  (red) in its tail. Centre: mean energy  $U(\beta) \rightarrow 2$  and specific heat  $C(\beta)$ , whose peak  $\beta^* \approx 2.1$  is the melting temperature of Goldbach order. Right:  $\beta^*$  versus  $\log \mathfrak{S}(N)$  (correlation  $-0.92$ ).

## C.2 Adversarial robustness of the Goldbach graph

View the representations as a bipartite graph: even integers  $N$  on one side, primes  $p$  on the other, with an edge  $N \sim p$  when  $N - p$  is prime; the degree of  $N$  is  $R_1(N)/2$ . How robust is Goldbach to deletion of primes? On a window of  $1.5 \cdot 10^4$  even integers near  $10^6$  we compare three adversaries (Fig. 17).

*Random deletion* is almost powerless: deleting each prime independently with probability  $\phi$  leaves  $N$  representable unless all its  $\sim R_1(N)/2$  disjoint pairs are hit, an event of probability  $(1 - (1 - \phi)^2)^{R_1(N)/2}$ . Goldbach survives in the *whole* window until  $\phi \approx 0.97$ ; one may delete 95% of all primes at random and still represent every  $N$ .

*Targeted deletion* is efficient but local: since the pairs  $\{p, N - p\}$  of a fixed  $N$  are disjoint, breaking  $N$  is a vertex cover of a matching, costing exactly  $R_1(N)/2$  primes; the cheapest target is the lowest-degree  $N$ , giving  $\kappa = \min_N R_1(N)/2$ . In the window this is 4.6% of the primes—an order of magnitude less than the random threshold, but aimed at a single  $N$ .

*Modular deletion* is the striking case. Delete the primes in one reduced class  $a \bmod m$ . The survivors lie in  $(\mathbb{Z}/m)^\times \setminus \{a\}$ , and an  $N$  is destroyed when its class is missing from the survivors' sumset. For  $m = 3$ , deleting the class 1 (i.e. *half* of all primes) leaves only the class 2, whose sumset is  $\{2 + 2\} = \{1\}$ : every  $N \equiv 0, 2 \pmod{3}$  loses all its representations, destroying 59% of the window (Fig. 17, right). The same 50% of primes removed at random destroys nothing. Thus the adversary manufactures a *local obstruction*—the exact mechanism the singular series rules out for the primes themselves. The effect is confined to the smallest moduli:  $m = 4$  deleting class 1 kills  $N \equiv 0 \pmod{4}$  (50%), while for  $m \geq 5$  the surviving classes already sum onto everything and no  $N$  is fully obstructed.

In short, the Goldbach graph is *random-robust but structure-fragile*: invulnerable to losing almost all primes by chance, yet shattered by removing one arithmetic progression. This is a network-science counterpart of the local-global theme: Goldbach's redundancy is enormous, but it is arithmetic, not statistical, redundancy.

## C.3 Goldbach as a market: liquidity and the fair-weather substitute

Read each even  $N$  as a market that must clear by hiring two prime inputs  $p + q = N$ ; the liquidity is the number of matches,  $L_G(N) = R_1(N)$ . Chen lets one input be a *minimally composite substitute* (a semiprime), giving  $L_C(N) = R_1(N) + R_2(N)$  and a substitute spread  $S_C(N) = R_2/R_1 \approx 2.3$ : in normal times the Chen market is three times more liquid. We ask whether the substitute also provides *resilience* under a supply shock that removes each prime with probability  $\phi$ . A Goldbach match survives with probability  $(1 - \phi)^2$  (two primes),




Figure 17: Robustness of the Goldbach graph. Left: fraction of window  $N$  destroyed versus fraction of primes deleted—random deletion (curve) only bites near  $\phi = 1$ , whereas the modular attacks at  $\phi = 0.5$  (mod 3, 4) destroy 50–60%, and the targeted cut needs only  $\kappa = 4.6\%$ . Right: under the mod-3 attack,  $N \equiv 0$  and  $N \equiv 2$  are obstructed while  $N \equiv 1$  is untouched.

but a Chen match only with  $(1 - \phi)^3$ , because the semiprime  $q = rs$  needs its two factors  $r, s$  to survive as well. Hence the expected surviving liquidities are  $L_G(\phi) = R_1(1 - \phi)^2$  and  $L_C(\phi) = L_G(\phi) + R_2(1 - \phi)^3$ , and the substitute spread *decays linearly*,

$$S_C(\phi) = \frac{L_C(\phi) - L_G(\phi)}{L_G(\phi)} = S_C(0)(1 - \phi) \rightarrow 0.$$

Solving  $L(\phi) = 1$  for the illiquidity threshold, both markets collapse at essentially the same point,  $\phi_G \approx \phi_C \approx 0.989$ , and the resilience premium  $\phi_C - \phi_G \approx 2 \cdot 10^{-4}$  is negligible (Fig. 18). The economic reading is clean and a little counter-intuitive: the Chen channel is *fair-weather liquidity*—abundant when primes are plentiful, but worth ever less precisely as they become scarce, because the composite good carries a higher assembly cost. The semiprime substitute deepens the market in calm times; it does not hedge the crisis.




Figure 18: Goldbach as a market under a prime supply shock  $\phi$ . Left: expected surviving liquidity of the Goldbach and Chen markets, both crossing the illiquidity threshold near  $\phi = 0.99$ . Right: the Chen substitute spread  $S_C(\phi) = S_C(0)(1 - \phi)$  vanishes linearly—no resilience premium.



## C.4 Persistent topology of additive decompositions (a negative note)

It is tempting to filter additive decompositions of  $N$  by arithmetic complexity rather than geometric distance: form the complex  $K_N^{(k)} = \{\{a, b\} : a + b = N, \Omega(a) + \Omega(b) \leq k\}$  and let  $k$  grow. But for the *binary* problem this is topologically empty: each integer  $a$  lies in the single pair  $\{a, N - a\}$ , so  $K_N^{(k)}$  is a disjoint matching for every  $k$ . Its 0-dimensional persistence is therefore exactly the energy spectrum of §C.1—each pair is a component born at its energy  $\Omega(a) + \Omega(b)$  and never dying—and all higher homology vanishes. There is no topological structure to discover beyond the spectrum.

Genuine higher dimension appears only at the *ternary* level. For odd  $N$ , form the graph on primes  $p < N$  with an edge  $\{p, q\}$  whenever  $N - p - q$  is prime, so that each edge *is* a ternary representation  $(p, q, N - p - q)$ . At  $N = 99,999$  this graph has 9,591 vertices and  $3.9 \cdot 10^6$  edges (density 0.084, minimum degree 2, so it is connected—a graph form of Helfgott’s ternary theorem). Yet it is strongly *anti*-clustered: it contains far fewer triangles than a random graph of equal density, so ternary representations do not aggregate into higher-order simplices. The honest conclusion is that the additive-complexity filtration adds nothing to the spectral and graph-theoretic pictures already developed; we record it as a negative result.

## C.5 Goldbach as control: the value of arithmetic information

Finally, view the even integers as a sequence of markets to be cleared by a controller: at each  $N$  the controller picks a prime  $p$  (the action) and clears if  $q = N - p$  is prime. Goldbach is the statement that a successful policy exists; the computational question is which *simple* policy clears fastest. A naive policy ignores the arithmetic of  $N$ ; a *residue-aware* policy with information budget  $B$  uses the residues  $N \bmod \ell$  for primes  $\ell \leq B$  to prefer  $p$  with  $N - p$  coprime to those  $\ell$ —i.e. it avoids spending tries on partners forced to share a small factor. Over a window near  $10^6$  (Fig. 19), the single-shot clearing probability rises from 18% (naive) to 39% at  $B = 13$ , and the expected number of tries in a sequential search falls from 5.5 to 2.5—a 55% reduction. The marginal value of each prime of information is largest at  $\ell = 3$  (the factor  $\frac{\ell-1}{\ell-2} = 2$ ) and decreases thereafter, exactly tracking the singular series. Unlike the other reframings in this section, this one yields a concrete *algorithm* rather than a restatement: the residue-aware ordering is a drop-in search heuristic—enumerate candidate  $p$  in increasing order of the number of small primes  $\ell \leq B$  that  $N - p$  is forced to share with  $N$ —whose benchmark above (expected tries  $5.5 \rightarrow 2.5$  at  $B = 13$ , a 55% reduction) is a measured speed-up of the per- $N$  Goldbach search—the cost of finding one representation for a given  $N$ , distinct from the global verification  $R_1 > 0$  for all  $N \leq X$ , which the FFT already settles—not an analogy. The same local arithmetic the singular series encodes is thus directly actionable.

## C.6 The information content of the comet

The spectral saturation invites an information-theoretic reading: how compressible is the comet, and what is irreducible? Decomposing the variance of  $\log R_1(N)$  into a smooth trend, the local arithmetic (the conditional mean given  $N \bmod 30030$ , which captures  $\mathfrak{S}$  non-linearly), and a residual, we find (Fig. 20)

$$\underbrace{81.1\%}_{\text{trend } N/\log^2 N} + \underbrace{18.8\%}_{\text{singular series } \mathfrak{S}(N)} + \underbrace{0.11\%}_{\text{irreducible}} .$$

The comet is thus 99.9% compressible *by this particular code*—the conditional mean given  $N \bmod 30030 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 13$ . We stress that this figure is relative to the chosen conditioning set, not a theorem: the “irreducible” 0.11% is what survives after removing structure up to the prime 13, and a richer conditioning (more primes, or non-residue features) would shrink it further. With that caveat, the residual is informative: its power spectrum is *flat*—white noise,






Figure 19: Value of arithmetic information for the Goldbach controller. As the information budget  $B$  grows, the single-shot clearing probability (blue) rises and the sequential clearing cost (red) falls; the gain per prime tracks the singular-series factor  $(\ell - 1)/(\ell - 2)$ .

in contrast to the sharply peaked spectrum of  $R_1$  itself (§7.9)—with lag-1 autocorrelation  $-0.04$  and standard deviation  $2.7\%$ , a few times the Poisson floor  $\langle R_1 \rangle^{-1/2} = 0.9\%$ . So *up to modulus* 30030 the Goldbach comet is almost entirely a singular-series codeword and the residue carries no further *small-modulus* arithmetic signal; we do not claim the absence of all further structure, only that none is visible below this conditioning.




Figure 20: Information budget of the comet *relative to conditioning on  $N \bmod 30030$* . Left: the variance of  $\log R_1$  splits into a smooth trend (81%), the singular series (18.8%), and a residual (0.11%) that is irreducible *by this code* (richer conditioning would shrink it). Right: the residual’s power spectrum is flat—white noise, with none of the small-modulus arithmetic peaks of Fig. 13.

## C.7 Discrete geometry of the Chen surface

Each Chen representation  $N = p + rs$  ( $q = N - p$  semiprime,  $r = \text{spf}(q) \leq s$ ) is a lattice point on the surface  $p + rs = N$ . In normalised logarithmic coordinates  $u = \log r / \log N$ ,  $v = \log s / \log N$  the points fill the triangle  $\{u \leq v, u + v \leq 1\}$ , with the edge  $u = 0$  (i.e.  $r$  bounded) the Goldbach boundary and the diagonal  $u = v$  the balanced semiprimes. Accumulating  $6.6 \cdot 10^7$  solutions over a window near  $10^6$  (Fig. 21, left) shows a cloud organised into discrete vertical *rays*—one

for each small prime  $r = 3, 5, 7, \dots$ , since  $r$  is prime—whose brightness decreases with  $r$ : the mass piles up against the Goldbach edge.

Quantitatively, the Li–Liu exponent of a solution,  $a = 1 + \log r / \log s$ , has the spectrum of Fig. 21 (right): it peaks near  $a \approx 1.1$  and decays toward the balanced value 2, with mean 1.38 and *median* 1.30. Half of all Chen solutions already satisfy  $a < 1.3$ , and essentially all sit far below the uniform threshold  $a = 1.9$  of Li–Liu. So the Chen solutions of a typical  $N$  concentrate near the Goldbach boundary: the semiprime second summand is, far more often than not, an unbalanced product with a small factor—the geometric form of the residual-exponent and factor-balance findings of §7.7 and §7.4.




Figure 21: Discrete geometry of  $p + rs = N$ . Left: density of Chen solutions in  $(u, v) = (\log r / \log N, \log s / \log N)$ , with discrete rays at each prime  $r$  and mass against the Goldbach edge  $u = 0$ . Right: the exponent spectrum  $a = 1 + \log r / \log s$ , median 1.30, far below the Li–Liu threshold 1.9.

## Data and code availability

Every figure, table, and quoted number in this paper is reproduced by the accompanying open-source repository (Python; `numpy/scipy/matplotlib`, with `gurobipy` only for the optional MIP of §7.5). Each `src/` module carries a self-test (`python3 src/<module>.py`) and each figure is regenerated by its `experiments/run_*.py` driver; the central results use `run_counts.py` (FFT counts to  $2 \cdot 10^6$ ), `run_scaling.py` (block-convolution sieve to  $10^9$ ), `run_betalaw.py` (the law (24)), `run_estimator_audit.py` (the estimator reconciliation and the  $3 \mid N$  atom of §7.2), and `run_exceptional.py` (the concentration of Table 5). Derived tables and the sieve seeds are stored in `data/`. The repository is available at <https://github.com/MathiasRodriguezCastro/goldbach-chen-frontier>, and a versioned snapshot is archived on Zenodo, [doi:10.5281/zenodo.20725701](https://doi.org/10.5281/zenodo.20725701) (citation metadata in `CITATION.cff`).

## References

- [1] Chen, J. R. (1973). On the representation of a larger even integer as the sum of a prime and the product of at most two primes. *Scientia Sinica*, 16, 157–176.
- [2] Halberstam, H. and Richert, H.-E. (1974). *Sieve Methods*. Academic Press, London. (London Mathematical Society Monographs, 4.)
- [3] Hardy, G. H. and Littlewood, J. E. (1923). Some problems of ‘Partitio Numerorum’; III: On the expression of a number as a sum of primes. *Acta Mathematica*, 44, 1–70.

- [4] Helfgott, H. A. (2014). The ternary Goldbach problem. *arXiv preprint arXiv:1404.2224*.
- [5] Landau, E. (1909). *Handbuch der Lehre von der Verteilung der Primzahlen*. Teubner.
- [6] Li, J. and Liu, J. (2026). Theorem  $(1 + 1.9)$  on the Goldbach Conjecture. *arXiv preprint arXiv:2606.05224*.
- [7] Montgomery, H. L. and Vaughan, R. C. (1975). The exceptional set in Goldbach's problem. *Acta Arithmetica*, 27, 353–370.
- [8] Nathanson, M. B. (1996). *Additive Number Theory: The Classical Bases*. Springer.
- [9] Oliveira e Silva, T., Herzog, S., and Pardi, S. (2014). Empirical verification of the even Goldbach conjecture and computation of prime gaps up to  $4 \cdot 10^{18}$ . *Mathematics of Computation*, 83(288), 2033–2060.
- [10] Sathé, L. G. (1953). On a problem of Hardy on the distribution of integers having a given number of prime factors. *Journal of the Indian Mathematical Society*, 17, 63–141.
- [11] Selberg, A. (1954). Note on a paper by L. G. Sathé. *Journal of the Indian Mathematical Society*, 18, 83–87.
- [12] Tenenbaum, G. (2015). *Introduction to Analytic and Probabilistic Number Theory*. American Mathematical Society, 3rd edition. (Graduate Studies in Mathematics 163.)
- [13] Vaughan, R. C. (1997). *The Hardy–Littlewood Method*. Cambridge University Press, 2nd edition.