## Supplementary Table 1

Overview of kinetic constants used for the construction of the model.

Enzyme	EC number	Kinetic parameter	References	Rate Law
NADA	3.5.1.19	$K_M:9.6\mu{ m M}$	[1]	Product inhibition
		$K_{iP}$ :120 $\mu$ M		
		$k_{cat}$ : $0.65s^{-1}$		
NADS	6.3.5.1	$K_M$ :190 $\mu$ M	[2]	$_{\mathrm{HMM}}$
		$k_{cat}$ :21 $s^{-1}$		
NMNAT	2.7.7.1/2.7.7.	$18 K_{M_{NaMN}}:67.7 \mu M$	$[3]^{1}$	Substrate Competition
		$k_{cat_{NaMN}}:42.9s^{-1}$		
		$K_{M_{NMN}}:22.3\mu{\rm M}$		
		$k_{cat_{NMN}}:53.8s^{-1}$		
		$K_{M_{NMN}}$ :59 $\mu\mathrm{M}$		
		$k_{cat_{NAD}}:129.1s^{-1}$	$[4]^2$	
		$K_{M_{NaAD}}:502\mu\mathrm{M}$	5. 10	
		$k_{cat_{NaAD}}:103.8s^{-1}$	$[4]^3$	
NMNT	2.1.1.1	$K_M$ :400 $\mu$ M	[5]	Product inhibition
		$K_{iP}$ :60 $\mu$ M		
		$k_{cat}:8.1s^{-1}$	[6]	
NamPRT	6.3.5.1	$K_M$ :5nM	[7]	HMM
		$k_{cat}$ :0.0077 $s^{-1}$		
NAPRT	2.4.2.11	$K_M$ :23 $\mu\mathrm{M}$	[8]	HMM
		$k_{cat}:3.3s^{-1}$		
SIRT1	3.5.1	$K_M$ :29 $\mu$ M	[9]	Product inhibition
		$K_{iP}$ :60 $\mu$ M		
		$k_{cat}$ :0.67 $s^{-1}$		
$NT5^4$	3.1.3.5	$K_M:100\mu{ m M}$	$[10]^5$	HMM
		$k_{cat}:10s^{-16}$		

The total enzyme concentration was set to 10 for all enzymes if not mentioned otherwise. Concentration of potentiantial cosubstrate was assumed to be constant and considered to be not limiting the reaction and thus represented by the maximal velocities given.

 $<sup>^{1}</sup>$ Values for NMNAT1 used

<sup>&</sup>lt;sup>2</sup>Keq used for calculation of turnover rate of reverse reaction

 $<sup>^3</sup>$ Equilibrium constant used for calculation of turnover rate of reverse reaction

 $<sup>^4\</sup>mathrm{As}$  parameter values for NRK and PNP were not available we omited these enzymes and simulated NA and NAM as direct products

<sup>&</sup>lt;sup>5</sup>approx. IC50 value for NAD used

<sup>&</sup>lt;sup>6</sup>Avarage value for pyrimidines used

## Kinetic Rate Laws

**Product Inhibition** 

$$v = \frac{E_T \cdot k_{cat} \cdot S}{K_M + S + \frac{K_M \cdot P}{K_{iP}}} \tag{1}$$

Henry-Michaelis Menten for irreversible reactions (HMM)

$$v = \frac{E_T \cdot k_{cat} \cdot S}{K_M + S} \tag{2}$$

Substrate Competition at NMNAT

$$v = E_T \cdot \frac{\frac{k_{cat_A} \cdot A \cdot B}{K_{M_A}} - \frac{k_{cat_P} \cdot P \cdot Q}{K_{M_P}}}{1 + \frac{A}{K_{M_A}} + \frac{B}{K_{M_B}} + \frac{P}{K_{M_P}} + \frac{Q}{K_{M_O}}}$$
(3)

## References

- [1] Smith BC, Anderson MA, Hoadley KA, Keck JL, Cleland WW, et al. (2012) Structural and kinetic isotope effect studies of nicotinamidase (Pnc1) from saccharomyces cerevisiae. Biochemistry 51: 243–256.
- [2] Yi CK, Dietrich LS (1972) Purification and properties of yeast nicotinamide adenine dinucleotide synthetase. Journal of Biological Chemistry 247: 4794–4802.
- [3] Sorci L, Cimadamore F, Scotti S, Petrelli R, Cappellacci L, et al. (2007) Initial-rate kinetics of human NMN-adenylyltransferases: Substrate and metal ion specificity, inhibition by products and multisubstrate analogues, and isozyme contributions to NAD+ biosynthesis. Biochemistry 46: 4912–4922.
- [4] Berger F, Lau C, Dahlmann M, Ziegler M (2005) Subcellular compartmentation and differential catalytic properties of the three human nicotinamide mononucleotide adenylyltransferase isoforms. Journal of Biological Chemistry 280: 36334–36341.
- [5] Aksoy S, Szumlanski CL, Weinshilboum RM (1994) Human liver nicotinamide N-methyltransferase. cDNA cloning, expression, and biochemical characterization. Journal of Biological Chemistry 269: 14835–14840.
- [6] Alston TA, Abeles RH (1988) Substrate specificity of nicotinamide methyltransferase isolated from porcine liver. Archives of biochemistry and biophysics 260: 601–608.
- [7] Burgos ES, Schramm VL (2008) Weak coupling of ATP hydrolysis to the chemical equilibrium of human nicotinamide phosphoribosyltransferase. Biochemistry 47: 11086–11096.

- [8] Hanna LS, Hess SL, Sloan DL (1983) Kinetic analysis of nicotinate phosphoribosyltransferase from yeast using high pressure liquid chromatography. The Journal of biological chemistry 258: 9745–9754.
- [9] Borra MT, Langer MR, Slama JT, Denu JM (2004) Substrate Specificity and Kinetic Mechanism of the Sir2 Family of NAD + -Dependent Histone/Protein Deacetylases †. Biochemistry 43: 9877–9887.
- [10] Madrid-Marina V, Fox IH (1986) Human placental cytoplasmic 5'nucleotidase. Kinetic properties and inhibition. Journal of Biological Chemistry 261: 444–448.