

Extended Family of Fullerenes

Michel DEZA

Ecole Normale Supérieure, Paris

This is a joint work with Mathieu DUTOUR SIKIRIC and Mikhail SHTOGRIN, presented at

Overview

- 1 8 families of standard $(\{a, b\}, k)$ -spheres
- 2 Listing of $(\{a, b\}, k)$ -spheres with small p_b
- 3 8 standard families: four smallest members
- 4 Symmetry groups of $(\{a, b\}, k)$ -spheres
- 5 Goldberg-Coxeter construction
- 6 Parameterizing $(\{a, b\}, k)$ -spheres
- 7 Railroads and tight $(\{a, b\}, k)$ -spheres
- 8 Tight pure $(\{a, b\}, k)$ -spheres
- 9 Other analogs of fullerenes: c -disks
- 10 Icosahedrites
- 11 Standard $(\{a, b\}, k)$ -maps on surfaces
- 12 Beyond surfaces

Definition of a fullerene

A (geometric) **fullerene** F_n is a **simple** (i.e., 3-valent) **polyhedron** (putative carbon molecule) whose n vertices (carbon atoms) are arranged in **12 pentagons** and $(\frac{n}{2} - 10)$ **hexagons**.

- F_n exist for all even $n \geq 20$ except $n = 22$.
- $1, 1, 1, 2, 5, \dots, 1812, \dots, 214127713, \dots$ **isomers** F_n , for $n = 20, 24, 26, 28, 30, \dots, 60, \dots, 200, \dots$
- Graphite lattice (6^3) as F_∞ : the **"largest fullerene"**
- Thurston, 1998, implies: no. of F_n grows as n^9 .
- $C_{20}(I_h)$, $C_{60}(I_h)$, $C_{80}(I_h)$ are only **icosahedral** (i.e., with highest symmetry I_h or I) fullerenes with $n \leq 80$ vertices.
- **preferable** (or **IP**) fullerenes, C_n , satisfy isolated pentagon rule.

I. 8 families of standard $(\{a, b\}, k)$ -spheres

(R, k) -spheres: **curvature** $\kappa_i = 1 + \frac{i}{k} - \frac{i}{2}$ of i -gons

- Fix $R \subset \mathbb{N}$, an **(R, k) -sphere** is a k -regular, $k \geq 3$, map on \mathbb{S}^2 whose faces are i -gons, $i \in R$. Let $m = \min$ and $M = \max_{i \in R} i$.
- Let v, e and $f = \sum_i p_i$ be the numbers of vertices, edges and faces of S , where p_i is the number of i -gonal faces. Clearly, $kv = 2e = \sum_i ip_i$ and *Euler formula* $v - e + f = 2$ become $2 = \sum_i p_i \kappa_i$, where $\kappa_i = 1 + \frac{i}{k} - \frac{i}{2}$ is the **curvature** of i -gons.
- So, $m < \frac{2k}{k-2}$. For $m \geq 3$, it implies $3 \leq m, k \leq 5$, i.e. 5 Platonic pairs of parameters $(m, k) = (3, 3), (4, 3), (3, 4), (5, 3), (3, 5)$.
- If $M < \frac{2k}{k-2}$ ($\min_{i \in R} \kappa_i > 0$), then either 1) $k = 3, M \leq 5$, or 2) $k \in \{4, 5\}, M \leq 3$.
So, for $m \geq 3$, they are only Octahedron, Icosahedron and 11 $(\{3, 4, 5\}, 3)$ -spheres: 8 dual *deltahedra*, Cube and its truncations on 1 or 2 opposite vertices (*Dürer octahedron*).
In other words: five Platonic and eight $(\{3, 4, 5\}, 3)$ -spheres.

Standard (R, k) -spheres

- An (R, k) -sphere is **standard** if $M = \frac{2k}{k-2}$, i.e. $\min_{i \in R} \kappa_i = 0$.
So, $(M, k) = (6, 3), (4, 4), (3, 6)$ (Euclidean parameter pairs).
Exclusion of i -faces with $\kappa_i < 0$ simplifies enumeration, while number p_M of *flat* ($\kappa_M = 0$) M -faces not being restricted, there is an infinity of such (R, k) -spheres.
- The number of such v -vertex (R, k) -spheres with $|R| = 2$ increases polynomially with v .
Such spheres admit parametrization and description in terms of rings of (*Gaussian* if $k = 4$ and *Eisenstein* if $k = 3, 6$) *integers*.
All eight series of such spheres will be considered in detail.
- Remaining (R, k) -spheres (with $M > \frac{2k}{k-2}$, i.e. $\min_{i \in R} \kappa_i < 0$) do not admit above, in general. We will consider only simplest case: $(\{3, 4\}, 5)$ -spheres. The number of such v -vertex spheres grows at least exponentially with v .

8 families of standard $(\{a, b\}, k)$ -spheres

- An $(\{a, b\}, k)$ -sphere is an (R, k) -sphere with $R = \{a, b\}$, $1 \leq a < b$. It has $v = \frac{1}{k}(ap_a + bp_b)$ vertices.
- Such standard sphere has $b = \frac{2k}{k-2}$; so, $(b, k) = (6, 3), (4, 4), (3, 6)$ and Euler formula become $2 = \kappa_a p_a = (1 + \frac{a}{k} - \frac{a}{2})p_a = (1 - \frac{a}{b})p_a$.
- So, $p_a = \frac{2b}{b-a}$ and all possible (a, p_a) are:
 $(5, 12), (4, 6), (3, 4), (2, 3)$ for $(b, k) = (6, 3)$;
 $(3, 8), (2, 4)$ for $(b, k) = (4, 4)$;
 $(2, 6), (1, 3)$ for $(b, k) = (3, 6)$.
- Those 8 families can be seen as spheric analogs of the regular plane partitions $\{6^3\}, \{4^4\}, \{3^6\}$ with p_a disclinations ("defects") κ_a added to get the curvature 2 of the sphere.

8 families: existence criteria

[Grünbaum-Motzkin, 1963](#): criterion for $k=3 \leq a$; [Grünbaum, 1967](#): for $(\{3, 4\}, 4)$ -spheres; [Grünbaum-Zaks, 1974](#): for $a = 1, 2$.

k	(a, b)	smallest one	it exists if and only if	p_a	v
3	(5, 6)	Dodecahedron	$p_6 \neq 1$	12	$20 + 2p_6$
3	(4, 6)	Cube	$p_6 \neq 1$	6	$8 + 2p_6$
4	(3, 4)	Octahedron	$p_4 \neq 1$	8	$6 + p_4$
6	(2, 3)	$6 \times K_2$	p_3 is even	6	$2 + \frac{p_3}{2}$
3	(3, 6)	Tetrahedron	p_6 is even	4	$4 + 2p_6$
4	(2, 4)	$4 \times K_2$	p_4 is even	4	$2 + p_4$
3	(2, 6)	$3 \times K_2$	$p_6 = (k^2 + kl + l^2) - 1$	3	$2 + 2p_6$
6	(1, 3)	Trifolium	$p_3 = 2(k^2 + kl + l^2) - 1$	3	$\frac{1+p_3}{2}$
5	(3, 4)	Icosahedron	$p_4 \neq 1$	$2p_4 + 20$	$2p_4 + 12$

$(\{3, 6\}, 3)$ - ([Grünbaum-Motzkin, 1963](#)) and $(\{2, 4\}, 4)$ -spheres ([Deza-Shtogrin, 2003](#)) admit a simple 2-parametric description.

8 families of standard $(\{a, b\}, k)$ -spheres

- Let us denote $(\{a, b\}, k)$ -sphere with v vertices by $\{a, b\}_v$.
- $(\{5, 6\}, 3)$ - and $(\{4, 6\}, 3)$ -spheres are (geometric) **fullerenes** and *boron nitrides*. $\{5, 6\}_{60}(I_h)$: a new *carbon allotrope* C_{60} .
- $(\{a, b\}, 4)$ -spheres are minimal projections of **alternating links**, whose components are their *central circuits* (those going only ahead) and crossings are the vertices.
- By smallest member Dodecahedron $\{5, 6\}_{20}$, Cube $\{4, 6\}_8$, Tetrahedron $\{3, 6\}_4$, Octahedron $\{3, 4\}_6$ and $3 \times K_2$ $\{2, 6\}_2$, $4 \times K_2$ $\{2, 4\}_2$, $6 \times K_2$ $\{2, 3\}_2$, Trifolium $\{1, 3\}_1$, we call eight families: dodecahedrites, cubites, tetrahedrites, octahedrites and 3-bundelites, 4-bundelites, 6-bundelites, trifoliumites.
- **b -icosahedrites** ($(\{3, b\}, 5)$ -spheres) are not standard if $b \geq 3$, $p_b \geq 0$, since $p_3 = p_b(3b-10)+20$ and $\kappa_b = \frac{10-3b}{10b} < 0$.

Generation of standard $(\{a, b\}, k)$ -spheres


- $(\{2, 3\}, 6)$ -spheres, except $2 \times K_2$ and $2 \times K_3$, are the duals of $(\{3, 4, 5, 6\}, 3)$ -spheres with six new vertices put on edge(s).
Exp: $(\{5, 6\}, 3)$ -spheres with 5-gons organized in six pairs.
- $(\{1, 3\}, 6)$ -spheres, except $\{1, 3\}_1$ and $\{1, 3\}_3$, are as above but with 3 edges changed into 2-gons enclosing one 1-gon.
- $(\{2, 6\}, 3)$ -spheres are given by the *Goldberg-Coxeter construction* from **Bundle₃** $= 3 \times K_2$ $\{2, 6\}_2$.
- $(\{1, 3\}, 6)$ -spheres come by the *Goldberg-Coxeter construction* (extended below on 6-regular spheres) from **Trifolium** $\{1, 3\}_1$.

III. $(\{a, b\}, k)$ -spheres
with small p_b : listings


$(\{a, b\}, k)$ -spheres with $p_b \leq 2 < a < b$

- Remind: $(a, k) = (3, 3), (4, 3), (3, 4), (5, 3), (3, 5)$ if $k, a \geq 3$.
- The only $(\{a, b\}, k)$ -spheres with $p_b \leq 1$ are 5 **Platonic** (a^k):
Tetrahedron, Cube ($Prism_4$), Octahedron ($APrism_3$),
Dodecahedron (snub $Prism_5$), Icosahedron (snub $APrism_3$).
- There exists unique **trivial** 3-connected $(\{a, b\}, k)$ -sphere with $p_b = 2$ for $(\{4, b\}, 3)$ -, $(\{3, b\}, 4)$ -, $(\{5, b\}, 3)$ -, $(\{3, b\}, 5)$ -:
 D_{bh} **$Prism_b$** and D_{bd} **$APrism_b$** , **snub $Prism_b$** , **snub $APrism_b$** :
two b -gons separated by b -ring of 4-gons, $2b$ -ring of 3-gons,
two b -rings of 5-gons, two $3b$ -rings of 3-gons.
- Also, for $t \geq 2$, 10 **non-trivial** $(\{a, at\}, k)$ -spheres with $p_{at} = 2$:
5 $(\{a, ta\}, k)$ -spheres are (D_{th}) **necklaces** of polycycles $\{a^k\}$ -e,
3 are (D_{th}) **necklaces** of t v-split $\{3^4\}$ and e-split $\{5^3\}$, $\{3^5\}$,
 $(\{3, 3t\}, 5)$ -spheres C_{th} , D_t are **necklaces** of t v-, f -split $\{3^5\}$.


$(\{a, ta\}, k)$ -spheres with $p_{ta} = 2$, $k=3, 4, 5$; case $t=2$




D_{2h} : $a=3$




$a=4$




$a=5$




$a=5$




$a=3$ D_{2h}




$a=3$ D_{2h}




$a=3$ D_{2h}



$a=3$ D_{2h}




$a=3$ C_{2h}



$a=3$ D_2

Proof method: elementary (a, k) -polycycles

- A (a, k) -polycycle is a 2-connected plane graph with faces partitioned in a -gonal proper faces and holes, exterior face among them, so that vertex degrees are in $\{2, \dots, k\}$ and can be $< k$ only for a vertex lying on the boundary of a hole.
- Any (a, k) -polycycle decomposes uniquely along its bridges (non-boundary going hole-to-hole, possibly, same, edges) into elementary ones. Cf. integer factorisation into primes.
- We listed them for $\kappa_a = 1 + \frac{a}{k} - \frac{a}{2} \geq 0$. Otherwise, continuum.











This $(\{5, 15\}, 3)$ -sphere with $p_{15}=3$ is a 3-holes $(\{5\}, 3)$ -polycycle
 It decomposes into five 1-hole elementary $(\{5\}, k)$ -polycycles.

$(\{a, b\}, 3)$ -spheres with $p_b = 3$

- $(\{a, b\}, k)$ -sphere with $p_b = 3$ exists if and only if $b \equiv 2, a, 2a - 2 \pmod{2a}$ and $b \equiv 4, 6 \pmod{10}$ if $a=5$.
- Such sphere are unique if b is not $\equiv a \pmod{2a}$ and then their symmetry is D_{3h} , except when $(a, k) = (3, 5)$ when the symmetry is D_3 .
- There are 7 such spheres with $t = \lfloor \frac{b}{6} \rfloor = 0$ and 3+4+5+17 of them for any $t \geq 1$.









IV. 8 standard families: 4 smallest members

First four $(\{4, 6\}, 3)$ - and $(\{5, 6\}, 3)$ -spheres






 $O_h (6^4)$  $D_{6h} (18^2)$  $D_{3h} (6^2; 30)$  $D_{2d} (24^2)$  $I_h (10^6)$  $D_{6d} (12; 60)$  $D_{3h} (12^3; 42)$  $T_d (12^7)$

First four $(\{2, 6\}, 3)$ - and $(\{3, 6\}, 3)$ -spheres



Number of $(\{2, 6\}_v)$'s is nr. of representations $v=2(k^2 + kl + l^2)$, $0 \leq l \leq k$ ($GC_{k,l}(\{2, 6\}_2)$). It become 2 for $v=7^2=5^2+15+3^2$.


 $D_{3h} (6)$

 $D_{3h} (6^3)$

 $D_{3h} (12^2)$

 $D_3 (42)$

 $T_d (4^3)$

 $D_{2h} (8^2, 4^2)$

 $T_d (12^3)$

 $T_d (8^6)$


First four $(\{2, 4\}, 4)$ - and $(\{3, 4\}, 4)$ -spheres


 $D_{4h} \text{ } 2_1^2 (2^2)$

 $D_{4h} \text{ } 4_1^2 (4^2)$

 $D_{2h} \text{ } 2 \times 2_1^2 (2^2, 4)$

 $D_{2d} \text{ } 6_2^2 (6^2)$

 $O_h \text{ } 6_2^3 (4^3)$

Borr. rings


 $D_{4d} \text{ } 8_{18} (16)$

 $D_{3h} \text{ } 9_{40} (18)$









(Herschel)*


 $D_2 \text{ } 10_{56}^2$

(6; 14)

Above links/knots are given in [Rolfsen, 1976 and 1990](#) notation.
Herschel graph: the smallest non-Hamiltonian polyhedral graph.

First four $(\{2, 3\}, 6)$ - and $(\{1, 3\}, 6)$ -spheres

 $D_{6h} (2^3)$  $D_{3h} (3; 6)$  $D_{2d} (2^2; 8)$  $T_d (3^4)$  $C_{3v} (3)$  $C_{3h} (3; 6)$  $C_{3v} (6^2)$  $C_3 (21)$

Grünbaum-Zaks, 1974: $\{1, 3\}_v$ exists iff $v = k^2 + kl + l^2$ for integers $0 \leq l \leq k$. We show that the number of $\{1, 3\}_v$'s is the number of such representations of v , i.e. found $GC_{k,l}(\{1, 3\}_1)$.

V. Symmetry groups of $(\{a, b\}, k)$ -spheres

Finite isometry groups



All finite groups of isometries of 3-space \mathbb{E}^3 are classified.

In Schoenflies notations, they are:

- C_1 is the **trivial** group
- C_s is the group generated by a **plane reflexion**
- $C_i = \{I_3, -I_3\}$ is the **inversion** group
- C_m is the group generated by a **rotation** of order m of axis Δ
- C_{mv} (\simeq dihedral group) is the group generated by C_m and m **reflexion containing Δ**
- $C_{mh} = C_m \times C_s$ is the group generated by C_m and the **symmetry by the plane orthogonal to Δ**
- S_{2m} is the group of order $2m$ generated by an **antirotation**, i.e. commuting composition of a rotation and a plane symmetry

Finite isometry groups D_m , D_{mh} , D_{md}

- D_m (\simeq dihedral group) is the group generated of C_m and m rotations of order 2 with axis orthogonal to Δ
- D_{mh} is the group generated by D_m and a plane symmetry orthogonal to Δ
- D_{md} is the group generated by D_m and m symmetry planes containing Δ and which does not contain axis of order 2

 D_{2h}  D_{2d}

Remaining 7 finite isometry groups

- $I_h = H_3$ is the group of **isometries** of **Dodecahedron**;
 $I_h \simeq Alt_5 \times C_2$
- $I \simeq Alt_5$ is the group of **rotations** of Dodecahedron
- $O_h = B_3$ is the group of **isometries** of **Cube**
- $O \simeq Sym(4)$ is the group of **rotations** of Cube
- $T_d = A_3 \simeq Sym(4)$ is the group of **isometries** of **Tetrahedron**
- $T \simeq Alt(4)$ is the group of **rotations** of Tetrahedron
- $T_h = T \cup -T$

While (point group) $Isom(P) \subset Aut(G(P))$ (combinatorial group), **Mani, 1971**: for any 3-polytope P , there is a map-isomorphic 3-polytope P' (so, with the same skeleton $G(P') = G(P)$), such that the group $Isom(P')$ of its isometries is isomorphic to $Aut(G)$.

8 families: symmetry groups

- ① 28 for $\{5, 6\}_v$: $C_1, C_s, C_i; C_2, C_{2v}, C_{2h}, S_4; C_3, C_{3v}, C_{3h}, S_6; D_2, D_{2h}, D_{2d}; D_3, D_{3h}, D_{3d}; D_5, D_{5h}, D_{5d}; D_6, D_{6h}, D_{6d}; T, T_d, T_h; I, I_h$ (Fowler-Manolopoulos, 1995)
 - ② 16 for $\{4, 6\}_v$: $C_1, C_s, C_i; C_2, C_{2v}, C_{2h}; D_2, D_{2h}, D_{2d}; D_3, D_{3h}, D_{3d}; D_6, D_{6h}; O, O_h$ (Deza-Dutour, 2005)
 - ③ 5 for $\{3, 6\}_v$: $D_2, D_{2h}, D_{2d}; T, T_d$ (Fowler-Cremona, 1997)
 - ④ 2 for $\{2, 6\}_v$: D_3, D_{3h} (Grünbaum-Zaks, 1974)
 - ⑤ 18 for $\{3, 4\}_v$: $C_1, C_s, C_i; C_2, C_{2v}, C_{2h}, S_4; D_2, D_{2h}, D_{2d}; D_3, D_{3h}, D_{3d}; D_4, D_{4h}, D_{4d}; O, O_h$ (Deza-Dutour-Shtogrin, 2003)
 - ⑥ 5 for $\{2, 4\}_v$: $D_2, D_{2h}, D_{2d}; D_4, D_{4h}$, all in $[D_2, D_{4h}]$ (same)
 - ⑦ 3 for $\{1, 3\}_v$: C_3, C_{3v}, C_{3h} (Deza-Dutour, 2010)
 - ⑧ 22 for $\{2, 3\}_v$: $C_1, C_s, C_i; C_2, C_{2v}, C_{2h}, S_4; C_3, C_{3v}, C_{3h}, S_6; D_2, D_{2h}, D_{2d}; D_3, D_{3h}, D_{3d}; D_6, D_{6h}; T, T_d, T_h$ (same)
- 38 for icosahedrites $(\{3, 4\}, 5)$ - (same, 2011).

8 families: Goldberg-Coxeter construction $GC_{k,l}(\cdot)$

With $\mathbf{T} = \{T, T_d, T_h\}$, $\mathbf{O} = \{O, O_h\}$, $\mathbf{I} = \{I, I_h\}$, $\mathbf{C}_1 = \{C_1, C_s, C_i\}$, $\mathbf{C}_m = \{C_m, C_{mv}, C_{mh}, S_{2m}\}$, $\mathbf{D}_m = \{D_m, D_{mh}, D_{md}\}$, we get

- ① for $(\{5, 6\}, 3)$:- $\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3, \mathbf{D}_2, \mathbf{D}_3, \mathbf{D}_5, \mathbf{D}_6, \mathbf{T}, \mathbf{I}$
 - ② for $(\{2, 3\}, 6)$:- $\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3, \mathbf{D}_2, \mathbf{D}_3, \{D_6, D_{6h}\}, \mathbf{T}$
 - ③ for $(\{4, 6\}, 3)$:- $\mathbf{C}_1, \mathbf{C}_2 \setminus S_4, \mathbf{D}_2, \mathbf{D}_3, \{D_6, D_{6h}\}, \mathbf{O}$
 - ④ for $(\{3, 4\}, 4)$:- $\mathbf{C}_1, \mathbf{C}_2, \mathbf{D}_2, \mathbf{D}_3, \mathbf{D}_4, \mathbf{O}$
 - ⑤ for $(\{3, 6\}, 3)$:- $\mathbf{D}_2, \{T, T_d\}, \{D_3, D_{3h}\}$
 - ⑥ for $(\{2, 4\}, 4)$:- $\mathbf{D}_2, \{D_4, D_{4h}\}$
 - ⑦ for $(\{2, 6\}, 3)$:- $\{D_3, D_{3h}\}$
 - ⑧ for $(\{1, 3\}, 6)$:- $\mathbf{C}_3 \setminus S_6 = \{C_3, C_{3v}, C_{3h}\}$
- if $(\{3, 4\}, 5)$:- $\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3, \mathbf{C}_4, \mathbf{C}_5, \mathbf{D}_2, \mathbf{D}_3, \mathbf{D}_4, \mathbf{D}_5, \mathbf{T}, \mathbf{O}, \mathbf{I}$.

8 families: Goldberg-Coxeter construction $GC_{k,l}(\cdot)$

With $\mathbf{T}=\{T, T_d, T_h\}$, $\mathbf{O}=\{O, O_h\}$, $\mathbf{I}=\{I, I_h\}$, $\mathbf{C}_1=\{C_1, C_s, C_i\}$, $\mathbf{C}_m=\{C_m, C_{mv}, C_{mh}, S_{2m}\}$, $\mathbf{D}_m=\{D_m, D_{mh}, D_{md}\}$, we get

- ① for $(\{5, 6\}, 3)$:- $\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3, \mathbf{D}_2, \mathbf{D}_3, \mathbf{D}_5, \mathbf{D}_6, \mathbf{T}, \mathbf{I}$
- ② for $(\{2, 3\}, 6)$:- $\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3, \mathbf{D}_2, \mathbf{D}_3, \{D_6, D_{6h}\}, \mathbf{T}$
- ③ for $(\{4, 6\}, 3)$:- $\mathbf{C}_1, \mathbf{C}_2 \setminus S_4, \mathbf{D}_2, \mathbf{D}_3, \{D_6, D_{6h}\}, \mathbf{O}$
- ④ for $(\{3, 4\}, 4)$:- $\mathbf{C}_1, \mathbf{C}_2, \mathbf{D}_2, \mathbf{D}_3, \mathbf{D}_4, \mathbf{O}$
- ⑤ for $(\{3, 6\}, 3)$:- $\mathbf{D}_2, \{T, T_d\}, \{D_3, D_{3h}\}$
- ⑥ for $(\{2, 4\}, 4)$:- $\mathbf{D}_2, \{D_4, D_{4h}\}$
- ⑦ for $(\{2, 6\}, 3)$:- $\{D_3, D_{3h}\}$
- ⑧ for $(\{1, 3\}, 6)$:- $\mathbf{C}_3 \setminus S_6 = \{C_3, C_{3v}, C_{3h}\}$

if $(\{3, 4\}, 5)$:- $\mathbf{C}_1, \mathbf{C}_2, \mathbf{C}_3, \mathbf{C}_4, \mathbf{C}_5, \mathbf{D}_2, \mathbf{D}_3, \mathbf{D}_4, \mathbf{D}_5, \mathbf{T}, \mathbf{O}, \mathbf{I}$.



Spheres of blue symmetry are $GC_{a,b}$ from 1st such; so, given by one complex (Gaussian for $k=4$, Eisenstein for $k=3, 6$) parameter.

Goldberg, 1937 and Coxeter, 1971: $\{5, 6\}_v(I, I_h)$, $\{4, 6\}_v(O, O_h)$, $\{3, 6\}_v(T, T_d)$. Dutour-Deza, 2004 and 2010: for other cases.

VI. Goldberg-Coxeter construction


Goldberg-Coxeter construction $GC_{k,l}(\cdot)$

- Take a 3- or 4-regular plane graph G . The faces of dual graph G^* are triangles or squares, respectively.
- Break each face into pieces according to parameter (k, l) .
Master polygons below have area $\mathcal{A}(k^2 + kl + l^2)$ or $\mathcal{A}(k^2 + l^2)$, where \mathcal{A} is the area of a small polygon.



Gluing the pieces together in a coherent way


- Gluing the pieces so that, say, 2 non-triangles, coming from subdivision of neighboring triangles, form a small triangle, we obtain another **triangulation** or **quadrangulation** of the plane.




- The dual is a 3- or 4-regular plane graph, denoted $GC_{k,l}(G)$; we call it **Goldberg-Coxeter construction**.
- It works for **any** 3- or 4-regular map on **oriented surface**.

$GC_{k,l}(Cube)$ for $(k, l) = (1, 0), (1, 1), (2, 0), (2, 1)$


1,0




1,1







2,0




2,1




Goldberg-Coxeter construction from Octahedron



The case $(k, l) = (1, 1)$











3-regular case
 $GC_{1,1}$ is called **leapfrog**
 $(\frac{1}{3}$ -truncation of the dual)
truncated Octahedron



4-regular case
 $GC_{1,1}$ is called **medial**
 $(\frac{1}{2}$ -truncation)
Cuboctahedron

The case $(k, l) = (k, 0)$ of $GC_{k,l}(G)$: k -inflation


Chamfering (*quadrupling*) $GC_{2,0}(G)$ of 8 1st $(\{a, b\}, k)$ -spheres, $(a, b) = (2, 6), (3, 6), (4, 6), (5, 6)$ and $(2, 4), (3, 4), (1, 3), (2, 3)$, are:


 $D_{3h} (12^2)$

 $T_d (8^6)$

 $O_h (12^8)$

 $I_h (20^{12})$

 $D_{4h} (4^4)$

 $O_h (8^6)$

 $C_{3v} (6^2)$

 $D_{6h} (4^3, 6^2)$


For 4-regular G , $GC_{2k^2,0}(G) = GC_{k,k}(GC_{k,k}(G))$ by $(k+ki)^2 = 2k^2i$.

First four $GC_{k,l}(3 \times K_2)$ and $GC_{k,l}(4 \times K_2)$


All $(\{2, 6\}, 3)$ -spheres are $G_{k,l}(3 \times K_2)$: D_{3h} , D_{3h} , D_3 if $l=0, k$, else.




D_{3h} $3 \times K_2$




D_{3h} leapfrog




D_{3h} $G_{2,0}$




D_3 $G_{2,1}$




D_{4h} $4 \times K_2$



D_{4h} medial











D_{4h} $G_{2,0}$



D_4 $G_{2,1}$

First four $GC_{k,l}(6 \times K_2)$ and $GC_{k,l}(\text{Trifolium})$

 D_{6h}  $D_{3d} \ G_{1,1}$  $D_{6h} \ G_{2,0}$  $D_6 \ G_{2,1}$  C_{3v}  $C_{3h} \ G_{1,1}$  $C_{3v} \ G_{2,0}$  $C_3 \ G_{2,1}$

All $(\{2, 3\}, 6)$ -spheres are $G_{k,l}(6 \times K_2)$: C_{3v} , C_{3h} , C_3 if $l=0, k$, else.

Plane tilings $\{4^4\}$, $\{3^6\}$ and complex rings $\mathbb{Z}[i]$, $\mathbb{Z}[w]$

- The vertices of regular plane tilings $\{4^4\}$ and $\{3^6\}$ form each, convenient algebraic structures: lattice and ring. Path-metrics of those graphs are l_1 - 4-metric and hexagonal 6-metric.
- $\{4^4\}$: square lattice \mathbb{Z}^2 and ring $\mathbb{Z}[i]=\{z=k+li : k, l \in \mathbb{Z}\}$ of Gaussian integers with norm $N(z)=z\bar{z}=k^2+l^2=|| (k, l) ||^2$.
- $\{3^6\}$: hexagonal lattice $A^2=\{x \in \mathbb{Z}^3 : x_0+x_1+x_2=0\}$ and ring $\mathbb{Z}[w]=\{z=k+lw : k, l \in \mathbb{Z}\}$, where $w=e^{i\frac{\pi}{3}}=\frac{1}{2}(1+i\sqrt{3})$, of Eisenstein integers with norm $N(z)=z\bar{z}=k^2-k l+l^2$. We identify points $x=(x_0, x_1, x_2) \in A^2$ with $x_0+x_1 w \in \mathbb{Z}[w]$.

Plane tilings $\{4^4\}$, $\{3^6\}$ and complex rings $\mathbb{Z}[i]$, $\mathbb{Z}[w]$

- The vertices of regular plane tilings $\{4^4\}$ and $\{3^6\}$ form each, convenient algebraic structures: lattice and ring. Path-metrics of those graphs are l_1 - 4-metric and hexagonal 6-metric.
- $\{4^4\}$: square lattice \mathbb{Z}^2 and ring $\mathbb{Z}[i]=\{z=k+li : k, l \in \mathbb{Z}\}$ of Gaussian integers with norm $N(z)=z\bar{z}=k^2+l^2=|| (k, l) ||^2$.
- $\{3^6\}$: hexagonal lattice $A^2=\{x \in \mathbb{Z}^3 : x_0+x_1+x_2=0\}$ and ring $\mathbb{Z}[w]=\{z=k+lw : k, l \in \mathbb{Z}\}$, where $w=e^{i\frac{\pi}{3}}=\frac{1}{2}(1+i\sqrt{3})$, of Eisenstein integers with norm $N(z)=z\bar{z}=k^2-kl+l^2$. We identify points $x=(x_0, x_1, x_2) \in A^2$ with $x_0+x_1w \in \mathbb{Z}[w]$.
- A natural number $n = \prod_i p_i^{\alpha_i}$ is of form $n=k^2+l^2$ if and only if any α_i is even, whenever $p_i \equiv 3 \pmod{4}$ (Fermat Theorem). It is of form $n = k^2 + kl + l^2$ if and only if $p_i \equiv 2 \pmod{3}$.
- The first cases of non-unicity with $\gcd(k, l)=\gcd(k_1, l_1)=1$ are $91=9^2+9+1^2=6^2+30+5^2$ and $65=8^2+1^2=7^2+4^2$.
The first cases with $l=0$ are $7^2=5^2+15+3^2$ and $5^2=4^2+3^2$.

The bilattice of vertices of hexagonal plane tiling $\{6^3\}$

- We identify the *hexagonal lattice* A^2 (or *equilateral triangular lattice* of the vertices of the *regular plane tiling* $\{3^6\}$) with *Eisenstein ring* (of Eisenstein integers) $\mathbb{Z}[w]$.
- The hexagon centers of $\{6^3\}$ form $\{3^6\}$. Also, with vertices of $\{6^3\}$, they form $\{3^6\}$, rotated by 90° and scaled by $\frac{1}{3}\sqrt{3}$.
- The complex coordinates of vertices of $\{6^3\}$ are given by vectors $v_1=1$ and $v_2=w$. The lattice $L=\mathbb{Z}v_1+\mathbb{Z}v_2$ is $\mathbb{Z}[w]$.
- The vertices of $\{6^3\}$ form **bilattice** $L_1 \cup L_2$, where the bipartite complements, $L_1=(1+w)L$ and $L_2=1+(1+w)L$, are stable under multiplication. Using this,

$GC_{k,l}(G)$ for 6-regular graph G can be defined similarly to 3- and 4-regular case, **but only for** $k + lw \in L_2$, i.e. $k \equiv l \pm 1 \pmod{3}$.

Ring formalism

$\mathbb{Z}[i]$ ([Gaussian integers](#)) and $\mathbb{Z}[\omega]$ ([Eisenstein integers](#)) are *unique factorization rings*

Dictionary

	3-regular G	4-regular G	6-regular G
the ring	Eisenstein $\mathbb{Z}[\omega]$	Gaussian $\mathbb{Z}[i]$	Eisenstein $\mathbb{Z}[\omega]$
Euler formula	$\sum_i (6 - i)p_i = 12$	$\sum_i (4 - i)p_i = 8$	$\sum_i (3 - i)p_i = 6$
curvature 0	hexagons	squares	triangles
ZC-circuits	zigzags	central circuits	both
$GC_{11}(G)$	leapfrog graph	medial graph	or. tripling

Goldberg-Coxeter operation in ring terms

- Associate $z=k+lw$ (Eisenstein) or $z=k+li$ (Gaussian integer) to the pair (k, l) in 3-,6- or 4-regular case. Operation $GC_z(G)$ correspond to scalar multiplication by $z=k+lw$ or $k+li$.
- Writing $GC_z(G)$, instead of $GC_{k,l}(G)$, one has:

$$GC_z(GC_{z'}(G)) = GC_{zz'}(G)$$

- If G has v vertices, then $GC_{k,l}(G)$ has $vN(z)$ vertices, i.e., $v(k^2+l^2)$ in 4-regular and $v(k^2+kl+l^2)$ in 3- or 6-reg. case.

Goldberg-Coxeter operation in ring terms

- Associate $z=k+lw$ (Eisenstein) or $z=k+li$ (Gaussian integer) to the pair (k, l) in 3-, 6- or 4-regular case. Operation $GC_z(G)$ correspond to scalar multiplication by $z=k+lw$ or $k+li$.
- Writing $GC_z(G)$, instead of $GC_{k,l}(G)$, one has:

$$GC_z(GC_{z'}(G)) = GC_{zz'}(G)$$


- If G has v vertices, then $GC_{k,l}(G)$ has $vN(z)$ vertices, i.e., $v(k^2+l^2)$ in 4-regular and $v(k^2+kl+l^2)$ in 3- or 6-reg. case.
- $GC_z(G)$ has all rotational symmetries of G in 3- and 4-regular case, and all symmetries if $l=0, k$ in general case.
- $GC_z(G)=GC_{\bar{z}}(\bar{G})$ where \bar{G} differs by a plane symmetry only from G . So, if G has a symmetry plane, we reduce to $0 \leq l \leq k$; otherwise, graphs $GC_{k,l}(G)$ and $GC_{l,k}(G)$ are not isomorphic.

$GC_{k,l}(G)$ for 6-regular plane graph G and any k, l

- Bipartition of G^* gives vertex 2-coloring, say, red/blue of G .
- **Truncation** $Tr(G)$ of $\{1, 2, 3\}_v$ is a 3-regular $\{2, 4, 6\}_{6v}$.
- Coloring white vertices of G gives face 3-coloring of $Tr(G)$. White faces in $Tr(G)$ correspond to such in $GC_{k,l}(Tr(G))$.
- For $k \equiv l \pm 1 \pmod{3}$, i.e. $k + lw \in L_2$, define $GC_{k,l}(G)$ as $GC_{k,l}(Tr(G))$ with all white faces shrunk.
- If $k \equiv l \pmod{3}$, faces of $Tr(G)$ are white in $GC_{k,l}(Tr(G))$. Among 3 faces around each vertex, one is white. Coloring other red gives unique 3-coloring of $GC_{k,l}(Tr(G))$. Define $GC_{k,l}(G)$ as pair G_1, G_2 with $Tr(G_1) = Tr(G_2) = GC_{k,l}(Tr(G))$ obtained from it by shrinking all red or blue faces.
- $GC_{1,0}(G) = G$ and $GC_{1,1}(G)$ is **oriented tripling**.

Oriented tripling $GC_{1,1}(G)$ of 6-regular plane graph G


- Let C_1, C_2 be bipartite classes of G^* . For each C_i , **oriented tripling** $GC_{1,1}(G)$ is 6-regular plane graph $Or_{C_i}(G)$ coming by each vertex of $G \rightarrow 3$ vertices and 4 3-gonal faces of $Or_{C_i}(G)$. Symmetries of $Or_{C_i}(G)$ are symmetries of G preserving C_i .
- Orient edges of C_i clockwise. Select 3 of 6 neighbors of each vertex v : $\{2, 4, 6\}$ are those with directed edge going to v ; for $\{1, 5, 5\}$, edges go to them.




- Any $z=k+lw \neq 0$ with $k \equiv l \pmod{3}$ can be written as $(1+w)^s(k'+l'w)w$, where $s \geq 0$ and $k' \equiv l' \pmod{3}$. So, it holds reduction $GC_{k,l}(G) = G_{k',l'}(Or^s(G))$.

Examples of oriented tripling $GC_{1,1}(G)$


Below: $\{2, 3\}_2$ and $\{2, 3\}_4$ have *unique* oriented tripling.




2 D_{6h}



6 D_{3d}




4 T_d




12 T_h

Examples of oriented tripling $GC_{1,1}(G)$


Below: $\{2, 3\}_2$ and $\{2, 3\}_4$ have *unique* oriented tripling.




2 D_{6h}




6 D_{3d}




4 T_d




12 T_h




1 C_{3v}




3 C_{3h}



9 C_{3v}



27 C_{3h}




81 C_{3v}


Above: first 4 *consecutive* oriented triplings of the Trifolium.

VII. Parameterizing $(\{a, b\}, k)$ -spheres


Example: construction of the $(\{3, 6\}, 3)$ -spheres in $Z[\omega]$



In the central triangle
ABC, let A be the origin
of the complex plane



The corresponding
triangulation



All $(\{3, 6\}, 3)$ -spheres
come this way; two
complex parameters
in $Z[\omega]$ defined by
the points B and C

Parameterizing standard $(C_b = 0)$ $(\{a, b\}, k)$ -spheres

Thurston, 1998 implies: $(\{a, b\}, k)$ -spheres have $p_a - 2$ parameters and the number of v -vertex ones is $O(v^{m-1})$ if $m = p_a - 2 > 2$.

Idea: since b -gons are of zero curvature, it suffices to give relative positions of a -gons having curvature $2k - a(k - 2) > 0$.

At most $p_a - 1$ vectors will do, since one position can be taken 0.

But once $p_a - 1$ a -gons are specified, the last one is constrained.

The number of m -parametrized spheres with at most v vertices is $O(v^m)$ by direct integration. The number of such v -vertex spheres is $O(v^{m-1})$ if $m > 1$, by a Tauberian theorem.

Parameterizing standard $(C_b = 0)$ $(\{a, b\}, k)$ -spheres

Thurston, 1998 implies: $(\{a, b\}, k)$ -spheres have $p_a - 2$ parameters and the number of v -vertex ones is $O(v^{m-1})$ if $m = p_a - 2 > 2$.

Idea: since b -gons are of zero curvature, it suffices to give relative positions of a -gons having curvature $2k - a(k - 2) > 0$.

At most $p_a - 1$ vectors will do, since one position can be taken 0.

But once $p_a - 1$ a -gons are specified, the last one is constrained.

The number of m -parametrized spheres with at most v vertices is $O(v^m)$ by direct integration. The number of such v -vertex spheres is $O(v^{m-1})$ if $m > 1$, by a Tauberian theorem.

- Goldberg, 1937: $\{a, 6\}_v$ (highest 2 symmetries): 1 parameter
- Fowler and al., 1988: $\{5, 6\}_v$ (D_5 , D_6 or T): 2 parameters.
- Grünbaum-Motzkin, 1963: $\{3, 6\}_v$: 2 parameters.
- Deza-Shtogrin, 2003: $\{2, 4\}_v$: 2 parameters.
- Thurston, 1998: $\{5, 6\}_v$: 10 (again complex) parameters.
- Graver, 1999: $\{5, 6\}_v$: 20 integer parameters.
- Rivin, 1994: parameter description by dihedral angles.

Parameterizing (R, k) -spheres with $\min_{i \in R} C_i \geq 0$

Thurston, 1998 parametrized (dually, as triangulations) such $(R, 3)$ -spheres, i.e. 19 series of $(\{3, 4, 5, 6\}, 3)$ -spheres.

In general, such (R, k) -spheres are given by $m = \sum_{3 \leq i < \frac{2k}{k-2}} p_i - 2$ complex parameters z_1, \dots, z_m .

The number of vertices is expressed as a non-degenerate Hermitian form $q = q(z_1, \dots, z_m)$ of signature $(1, m - 1)$.

Let H^m be the cone of $z = (z_1, \dots, z_m) \in \mathbb{C}^m$ with $q(z) > 0$.

Given (R, k) -sphere is described by different parameter sets; let $M = M(\{p_3, \dots, p_m\}, k)$ be the discrete linear group preserving q .

For $k=3$, the quotient $H^m / (\mathbb{R}_{>0} \times M)$ is of finite covolume (Thurston, 1998, actually, 1993). Sah, 1994 deduced from it that the number of corresponding spheres grows as $O(v^{m-1})$.

Dutour partially generalized above for other k and surface maps.

8 families: number of complex parameters by groups

- ① $\{5, 6\}_v$ $C_1(10)$, $C_2(6)$, $C_3(4)$, $D_2(4)$, $D_3(3)$, $D_5(2)$, $D_6(2)$, $T(2)$, $\{I, I_h\}(1)$
- ② $\{4, 6\}_v$ $C_1(4)$, $C_2 \setminus S_4(3)$, $D_2(2)$, $D_3(2)$, $\{D_6, D_{6h}\}(1)$, $\{O, O_h\}(1)$
- ③ $\{3, 4\}_v$ $C_1(6)$, $C_2(4)$, $D_2(3)$, $D_3(2)$, $D_4(2)$, $\{O, O_h\}(1)$
- ④ $\{2, 3\}_v$ $C_1(4)$, $C_2(3?)$, $C_3(3?)$, $D_2(2?)$, $D_3(2?)$, $T(1)$, $\{D_6, D_{6h}\}(1)$
- ⑤ $\{3, 6\}_v$ $D_2(2)$, $\{T, T_d\}(1)$
- ⑥ $\{2, 4\}_v$ $D_2(2)$, $\{D_4, D_{4h}\}(1)$
- ⑦ $\{2, 6\}_v$ $\{D_3, D_{3h}\}(1)$
- ⑧ $\{1, 3\}_v$ $\{C_3, C_{3v}, C_{3h}\}(1)$

Thurston, 1998 implies: $(\{a, b\}, k)$ -spheres have $p_a - 2$ parameters and the number of v -vertex ones is $O(v^{m-1})$ if $m = p_a - 2 > 1$.

Number of complex parameters

 $\{5, 6\}_v$

Group	#param.
C_1	10
C_2	6
C_3, D_2	4
D_3	3
D_5, D_6, T	2
I	1

 $\{3, 4\}_v$

Group	#param.
C_1	6
C_2	4
D_2	3
D_3, D_4	2
O	1

 $\{4, 6\}_v$

Group	#param.
C_1	4
C_2	3
D_2, D_3	2
D_6, O	1

 $\{2, 3\}_v$

Group	#param.
C_1	4
C_2, C_3	3?
D_2, D_3	2?
D_6, T	1

$\{3, 6\}_v$ - and $\{2, 4\}_v$: 2 **complex** parameters but 3 **natural** ones will do: *pseudoroad* length, number of circumscribing *railroads*, *shift*.



VIII. Railroads and tight $(\{a, b\}, k)$ -spheres

ZC-circuits

- The edges of any plane graph are doubly covered by **zigzags** (**Petri** or **left-right paths**), i.e., circuits such that any two but not three consecutive edges bound the same face.
- The edges of any *Eulerian* (i.e., even-valent) plane graph are partitioned by its **central circuits** (those going straight ahead).
- A **ZC-circuit** means *zigzag* or *central circuit* as needed.
CC- or **Z-vector** enumerate lengths of above circuits.

ZC-circuits


- The edges of any plane graph are doubly covered by **zigzags** (**Petri** or **left-right paths**), i.e., circuits such that any two but not three consecutive edges bound the same face.
- The edges of any *Eulerian* (i.e., even-valent) plane graph are partitioned by its **central circuits** (those going straight ahead).
- A **ZC-circuit** means *zigzag* or *central circuit* as needed.
CC- or **Z-vector** enumerate lengths of above circuits.
- A **railroad** in a 3-, 4- or 6-regular plane graph is a circuit of 6-, 4- or 3-gons, each adjacent to neighbors on opposite edges.
Any railroad is bound by two "parallel" ZC-circuits. It (any if 4-, simple if 3- or 6-regular) can be collapsed into 1 ZC-circuit.




Railroad in a 6-regular sphere: examples

$APrism_3$ with 2 base 3-gons doubled is the $\{2, 3\}_6$ (D_{3d}) with CC-vector $(3^2, 4^3)$, all five central circuits are simple.


Base 3-gons are separated by a **simple railroad** R of six 3-gons, bounded by two parallel central 3-circuits around them. Collapsing R into one 3-circuit gives the $\{2, 3\}_3$ (D_{3h}) with CC-vector $(3; 6)$.




$D_{3d} (3^2, 4^3)$



$D_{3h} (3; 6)$




$T_d (3^4)$




Above $\{2, 3\}_4$ (T_d) has no railroads but it is not **strictly tight**, i.e. no any central circuit is adjacent to a non-3-gon *on each side*.

Railroads flower: Trifolium $\{1, 3\}_1$

Railroads can be simple or self-intersect, including **triple** if $k = 3$.
First such **Dutour** $(\{a, b\}, k)$ -spheres for $(a, b) = (4, 6), (5, 6)$ are:



$\{4, 6\}_{66}(D_{3h})$ **twice**



$\{5, 6\}_{172}(C_{3v})$

Which plane curves with at most triple self-intersections come so?

Number of ZC-circuits in tight $(\{a, b\}, k)$ -sphere

Call an $(\{a, b\}, k)$ -sphere **tight** if it has no railroads.

- ≤ 15 for $\{5, 6\}_v$ Shtogrin-Deza-Dutour, 2011
- ≤ 9 for $\{4, 6\}_v$ and $\{2, 3\}_v$ Deza-Dutour, 2005 and 2010
- ≤ 3 for $\{2, 6\}_v$ and $\{1, 3\}_v$ same
- ≤ 6 for $\{3, 4\}_v$ Deza-Shtogrin, 2003
- Any $\{3, 6\}_v$ has ≥ 3 zigzags with equality iff it is tight. All $\{3, 6\}_v$ are tight iff $\frac{v}{4}$ is prime > 2 and none iff it is even
- Any $\{2, 4\}_v$ has ≥ 2 central circuits with equality iff it is tight. There is a tight one for any even v .

Number of ZC-circuits in tight $(\{a, b\}, k)$ -sphere

Call an $(\{a, b\}, k)$ -sphere **tight** if it has no railroads.

- ≤ 15 for $\{5, 6\}_v$ Shtogrin-Deza-Dutour, 2011
- ≤ 9 for $\{4, 6\}_v$ and $\{2, 3\}_v$ Deza-Dutour, 2005 and 2010
- ≤ 3 for $\{2, 6\}_v$ and $\{1, 3\}_v$ same
- ≤ 6 for $\{3, 4\}_v$ Deza-Shtogrin, 2003
- Any $\{3, 6\}_v$ has ≥ 3 zigzags with equality iff it is tight. All $\{3, 6\}_v$ are tight iff $\frac{v}{4}$ is prime > 2 and none iff it is even
- Any $\{2, 4\}_v$ has ≥ 2 central circuits with equality iff it is tight. There is a tight one for any even v .






First tight ones with max. of ZC-circuits are $GC_{21}(\{a, b\}_{\min})$: $\{5, 6\}_{140}(I)$, $\{2, 6\}_{14}(D_3)$, $\{3, 4\}_{30}(0)$; and $\{a, b\}_{\min}$: $\{3, 6\}_4(T_d)$, $\{2, 4\}_2(D_{4h})$ with $ZC=(28^{15}), (14^3), (10^6), (4^3), (2^2)$, all simple. $\{4, 6\}_{88}(D_{2h})$ and $\{2, 3\}_{44}(D_{3h})$ are smallest with 8 zigzags.

Maximal number M_v of central circuits in *any* $\{2, 3\}_v$

- $M_v = \frac{v}{2} + 1, \frac{v}{2} + 2$ for $v \equiv 0, 2 \pmod{4}$. It is realized by the series of symmetry D_{2d} with CC-vector $2^{\frac{v}{2}}, 2v_{0,v}$ and of symmetry D_{2h} with CC-vector $2^{\frac{v}{2}}, v_{0, \frac{v-2}{4}}^2$ if $v \equiv 0, 2 \pmod{4}$.
- For odd v , M_v is $\lfloor \frac{v}{3} \rfloor + 3$ if $v \equiv 2, 4, 6 \pmod{9}$ and $\lfloor \frac{v}{3} \rfloor + 1$, otherwise. Define t_v by $\frac{v-t_v}{3} = \lfloor \frac{v}{3} \rfloor$. M_v is realized by the series of symmetry C_{3v} if $v \equiv 1 \pmod{3}$ and D_{3h} , otherwise. CC-vector is $3^{\lfloor \frac{v}{3} \rfloor}, (2\lfloor \frac{v}{3} \rfloor + t_v)_{0, \lfloor \frac{v-2t_v}{9} \rfloor}^3$ if $v \equiv 2, 4, 6 \pmod{9}$ and $3^{\lfloor \frac{v}{3} \rfloor}, (2v + t_v)_{0, v+2t_v}$, otherwise.

Smallest CC-knotted or Z-knotted $\{2, 3\}_v$

- The **minimal number** of central circuits or zigzags, 1, have **CC-knotted** and **Z-knotted** $\{2, 3\}_v$. They correspond to plane curves with only triple self-intersection points. For $v \leq 16$, there are 1, 2, 4, 7, 9, 12 Z-knotted if $v=3, 7, 9, 11, 13, 15$ and 1, 2, 2, 4, 11, 9, 1, 19 CC-knotted if $v=4, 6, 8, 10, 12, 14, 15, 16$.
- Conjecture** (holds if $v \leq 54$): any Z-knotted $\{2, 3\}_v$ has odd v and a CC-knotted $\{2, 3\}_v$ is Z-knotted if and only if v is odd.

4 D_2 6 D_3 6 C_2 8 D_2 15 C_1

IX. Tight pure $(\{a, b\}, k)$ -spheres

Tight $(\{a, b\}, k)$ -spheres with only simple ZC-circuits

- Call $(\{a, b\}, k)$ -sphere **pure** if any of its ZC-circuits is *simple*, i.e. has no self-intersections.
- Any $(\{3, 6\}, 3)$ - or $(\{2, 4\}, 4)$ -sphere is pure. They are tight if and only if have three or, respectively, two ZC-circuits.
- Any ZC-circuit of $\{2, 6\}_v$ or $\{1, 3\}_v$ self-intersects.

Tight $(\{a, b\}, k)$ -spheres with only simple ZC-circuits


- Call $(\{a, b\}, k)$ -sphere **pure** if any of its ZC-circuits is *simple*, i.e. has no self-intersections.
- Any $(\{3, 6\}, 3)$ - or $(\{2, 4\}, 4)$ -sphere is pure. They are tight if and only if have three or, respectively, two ZC-circuits.
- Any ZC-circuit of $\{2, 6\}_v$ or $\{1, 3\}_v$ self-intersects.

The number of tight pure $(\{a, b\}, k)$ -spheres is:


- 1 9? for $\{5, 6\}_v$ computer-checked for $v \leq 200$
- 2 2 for $\{4, 6\}_v$
- 3 8 for $\{3, 4\}_v$
- 4 5 for $\{2, 3\}_v$
- 5 ∞ for $\{2, 4\}_v$: ≥ 1 for any possible (i.e. even) v
- 6 ∞ for $\{3, 6\}_v$: ≥ 1 for any odd $\frac{v}{4}$
(all if it is prime > 2 and none if it is even)
- 7 0 for $\{2, 6\}_v$ and $\{1, 3\}_v$

All tight $(\{4, 6\}, 3)$ -spheres with only simple zigzags

There are exactly two such spheres: **Cube** and its leapfrog $GC_{11}(\text{Cube})$, **truncated Octahedron**.




8 O_h (6^4)




24 O_h (12^6)

All tight $(\{4, 6\}, 3)$ -spheres with only simple zigzags

There are exactly two such spheres: **Cube** and its leapfrog $GC_{11}(\text{Cube})$, **truncated Octahedron**.



8 O_h (6^4)



24 O_h (12^6)


Proof is based on a) The size of intersection of two simple zigzags in any $(\{4, 6\}, 3)$ -sphere is 0, 2, 4 or 6 and

b) Tight $(\{4, 6\}, 3)$ -sphere has at most 9 zigzags.


For $(\{2, 3\}, 6)$ -spheres, a) holds also, implying a similar result.

All tight $(\{3, 4\}, 4)$ -spheres with only simple central circuits


The **medial** of a connected plane graph $G = (X, E)$ is the graph $Med(G)$ of edges of G with two being adjacent if they have a common vertex and bound the same face. $Med(G)$ is a 4-regular plane graph; its central circuits correspond to zizags of G .
1st and 2nd below are the medials of Tetrahedron and Cube.




6 O_h (4^3)
Octahedron




12 O_h (6^4)
 $GC_{11}(Oct.)$




12 D_{3h} (6^4)




14 D_{4h}
($6^2, 8^2$)




20 D_{2d} (8^5)



22 D_{2h}
($8^3, 10^2$)




30 O (10^6)
 $GC_{21}(Oct.)$




32 D_{4h}
($10^4, 12^2$)


8 tight $(\{5, 6\}, 3)$ -spheres with only simple zigzags




20 I_h (10^6)




28 T_d (12^7)




48 D_3 (16^9)




60 I_h (18^{10})




76 D_{2d}
($22^4, 20^7$)



88 T (22^{12})




92 T_h
($24^6, 22^6$)




140 I , (28^{15})

The medials of 1 – 4, 6, 8-th above and of next one form **complete arrangements of pseudocircles** (CAP), i.e. any two intersect twice. Among 9, only 1, 4, 6, 8-th above are **zigzag-transitive**.

Other such 60-vertex $(\{5, 6\}, 3)$ -sphere



60 I_h (18^{10})



60 D_3 (18^{10})


This pair was first answer on a question in [Grünbaum, 1967, 2003](#) *Convex Polytopes* about existence of different *simple* polyhedra with the same p -vector and Z -vector.

Both have 60 vertices of degree 3; 12 5- and 20 6-gonal faces; and 10 (simple) zigzags of length 18 each. But they are different and their groups have, 1 and 3 orbits, respectively, on zigzags.


Pseudocircles arrangements from tight pure spheres

- A simple central circuit can be seen as a **Jordan curve**, i.e. a simple and closed plane curve.
- A **(k, t) -AP** (arrangements of pseudocircles) is a set of k Jordan curves where any two intersect (triple or tangent points excluded) exactly in t points; so, there are $t(k-1)$ points. It is a tight pure 4-regular graph with k central circuits of length $t(k-1)$ intersecting pairwise in t points. It is a projection of a link; **Borromean rings** is $(3, 2)$ -AP.
- For $F_{20}(I_h)$, $F_{28}(T_d)$, $F_{48}(D_3)$, $F_{60}(I_h)$, $F_{60}D_3$, $F_{88}(T)$, $F_{140}(I)$, their medials form $(k, 2)$ -APs with $k = 6, 7, 9, 10, 10, 12, 15$.
- The medials of truncated Tetrahedron, Cube, Icosahedron, Dodecahedron form $(3, 6)$ -, $(4, 6)$ -, $(10, 2)$ -, $(6, 6)$ -APs.
- For $OC_6(O_h)$, $OC_{12}(O_h)$, $OC_{12}(D_{3h})$, $OC_{20}(D_{2d})$, $OC_{30}(0)$, their central circuits form $(k, 2)$ -APs with $k = 3, 4, 4, 5, 6$.


Tight $(\{2, 3\}, 6)$ -spheres with only simple ZC-circuits




2 D_{6h} (2^3)
(6^2)




4 T_d (3^4)
(6^4)




6 D_3 no
($12, 8^3$)




8 D_{2d} ($5^4, 4$)
no



D_{6h} ($4^3, 6^2$)
(8^6) no



12 T_h (6^6)
(12^6)



14 D_6 no
(14^6)

All **CC-pure, tight**: Nrs. 1,2,4,5,6 (Nrs. 3,7 are not CC-pure).

All **Z-pure, tight**: Nrs. 1,2,3,6,7 (4 is not Z-pure, 5 is not Z-tight).

1st, 3rd are **strictly** CC-, Z-**tight**: all ZC-circuits sides touch 2-gons

X. Other fullerene analogs:
 $(\{a, b, c\}, k)$ -disks ($p_c=1$)

Other fullerene-like non-standard $(\min_{i \in R} \kappa_i < 0)$ spheres

Related **non-standard** (R, k) -spheres with $\kappa_{\max\{i \in R\}} < 0$, are:

- **G-fulleroids** (Deza-Delgado, 2000; Jendrol-Trenkler, 2001 and Kardos, 2007): $(\{5, b\}, 3)$ -spheres with $b \geq 7$ and symmetry G .
- **b-Icosahedrites**: $(\{3, b\}, 5)$ -spheres with $b \geq 4$. They have $p_3 = (3b - 10)p_b + 20$ 3-gons and $v = 2(b - 3)p_b + 12$ vertices. Snub Cube and Snub Dodecahedron are the cases $(b, v; \text{group}) = (4, 24; O)$ and $(5, 60; I)$.
- **Haeckel, 1887**: $(\{5, 6, c\}, 3)$ -spheres with $c = 7, 8$ representing skeletons of radiolarian zooplankton **Aulonia hexagona**.
- **$(\{a, b, c\}, k)$ -disk** is an $(\{a, b, c\}, k)$ -sphere with $p_c = 1$; so, its $v = \frac{2}{k-2}(p_a - 1 + p_b) = \frac{2}{2k-a(k-2)}(a + c + p_b(b-a))$ and (setting $b' = \frac{2k}{k-2}$) $p_a = \frac{b'+c}{b'-a} + p_b \frac{b-b'}{b'-a}$. So, $p_a = \frac{b+c}{b-a}$ if $b = b'$ (8 families). An $(\{a, b, c\}, k)$ -disk is non-standard iff $\max\{a, b, c\} > \frac{2k}{k-2}$.
- **Fullerene c-disk** is the case $(a, b, c; k) = (5, 6, c; 3)$ of above. So, they have $p_5 = c + 6$ and $v = 2(p_6 + c + 5)$ vertices.

Fullerene c -disks: big picture

- **Fullerene c -polycycle**: an c -gon partitioned into 5- and 6-gons with vertices of degree 3 inside and 3 or 2 on the c -gon.
- **Fullerene c -disk**: full. c -polycycle without vertices of degree 2; so, $p_5 = p_6 + 6$. If $c \in \{5, 6\}$, it is a **fullerene** without a face.
- **Fullerene c -patch**: full. c -polycycle which is a fullerene's part; so, $p_5 \leq 12$. It is a fullerene c -disk if and only if $c \in \{5, 6\}$.
- **Theorem**: full. c -disk with a face having ≥ 2 common edges with c -gon (so, non-3-connected) exists if and only if $c \geq 8$. So, any fullerene c -disk with $3 \leq c \leq 7$ is polyhedral.

Conjecture (checked for $c \leq 20$):

- 1) minimal fullerene c -disk has $2(c + 11)$ vertices if $c \geq 13$.
 - 2) Only 3 **gap** full. c -disks: $(c, v) = (5, 22), (3, 24), (1, 42)$.
- **Fullerene c -thimble**: a full. c -disk with only 5-gons adjacent to the c -gon. It exists if and only if $c \geq 5$, always polyhedral.
- Conjecture**: minimal fullerene c -thimble has $5c - 5$ or $5c - 6$ vertices for odd or even, respectively, $c \geq 5$.

Reducibility of fullerene c -disks


- In a full. c -disk, a **zigzag** is an edge-circuit alternating left and right turns. The zigzags doubly cover the edges.
- A **belt** is simple circuit of 6-gons, adjacent to their neighbors on opposite faces. It is bounded by 2 disjoint simple zigzags. Call a fullerene c -disk is **reducible** if it has a belt.
- The belts of a full. c -thimble form a cylinder. So, c -thimbles are cuts of full. nanotubes: c -belt \rightarrow two c -rings of 5-gons.
- Any simple zigzag in an irreducible full. c -disk has adjacent 5-gon on each side and intersects any other simple zigzag. So, the number of simple zigzags is at most $\frac{5(c+6)}{2}$.
- Each zigzag of an irreducible **pure** (all zigzags are simple) fullerene, is adjacent to at least two 5-gons on each side. So, their number is $\leq \frac{5(6+6)}{4}=15$. $F_{140}(I)$ has Z -vector 28^{15} .
- **Conjecture:** pure irreducible fullerenes are only 9 fullerenes $F_v(G)$ with $(v, G)=(20, I_h), (28, T_d), (48, D_3), (60, I_h)$ and $(60, D_3), (76, D_{2d}), (88, T), (92, T_h), (140, I)$.

Minimal fullerene c -disks for $1 \leq c \leq 8$


It is 1-vertex-, 1-edge-truncated, usual F_{20} , F_{24} for $c=3, 4$, **5, 6**.

It comes from minimal 4-disk for $c=2$: add edge with 2-gon on it.


Checked for $c \leq 20$: it has $p_6=14, 6, 3, 2, 0, 1, 3, 4, 6, 7, 8, 5$ and $=6$ if $1 \leq c \leq 12$ and $c \geq 13$. Unique unless 2, 3, 10 for $c=9, 10, 11$.




1 40 C_s




2 26 C_{2v}




3 22 C_{3v}




4 22 C_{2v}




5 20 I_h



6 24 D_{6d}




7 30 C_s




8 34 C_{2v}


Minimal fullerene c -disks for $c \geq 9$







9-2 40

 C_{3v} 

10-3 44

 C_{2v} 









11-10 48

 C_s 12 44 C_{2v} 13 48 C_s 14 50 C_2 15 52 C_s 16 54 C_2

Conjecture: for $c \geq 13$, the only minimal c -disk is **c -pentatube**
 $B + \text{Hex}_3 + \text{Pen}_{c-12} + \text{Hex}_3 + B$ (symmetry C_s/C_2 for odd/even c).

Symmetries of fullerene c -disks


- Their groups: C_m, C_{mv} with $m \equiv 0 \pmod{c}$ (since any symmetry should stabilize unique c -gonal face) and $m \in \{1, 2, 3, 5, 6\}$ since the axis pass by a vertex, edge or face.
- The minimal such 8- and 9-disks are given below.

8 34 C_{2v} 8 36 C_s 8 38 C_1 8 38 C_2 9 40 C_{3v} 9 40 C_s 9 42 C_1 9 52 C_3

XI. Icosahedrites: $(\{3, 4\}, 5)$ -spheres

Icosahedrites, i.e., $(\{3, 4\}, 5)$ -spheres

- They have $p_3 = 2p_b + 20$ and $v = 2p_b + 12$ vertices.
- Their number is 1, 0, 1, 1, 5, 12, 63, 246, 1395, 7668, 45460 for $v = 12$, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32. It grows at least exponentially with v .
- p_a is fixed in for standard $(\{a, b\}, k)$ -spheres permitting Goldberg-Coxeter construction and parametrization of graphs which imply the polynomial growth of their number. It does not happen for icosahedrites; no parametrization for them.




A-operation keeps symmetries; *B*-operation: only rotational ones.

Proof for the number of icosahedrites

A **weak zigzag** is a left/right, but never extreme, edge-circuit.


If a v -vertex icosahedrite has a **simple** weak zigzag of length 6, a $(v+6)$ -vertex one come by inserting a **corona** (6-ring of three 4-gons alternated by three pairs of adjacent 3-gons) instead of it. But such spheres exist for $v=18, 20, 22$; so, for $v \equiv 0, 2, 4 \pmod{6}$. There are two options of inserting corona; so, the number of v -vertex icosahedrites grows at least exponentially.



$$12 \ I_h$$

$$wZ = 6^{10}$$


$$Z = 10^6$$



$$18 \ D_3$$

$$6^2, 8^3; 54_{0,9}$$


$$90_{27,18}$$



$$20 \ D_{2d}$$

$$6^4, 20^2; 18^2_{0,3}$$

$$10^4; 30^2_{3,0}$$



$$22 \ D_{5h}$$





$$6^{10}; 50_{15,0}$$

$$10^2; 90_{15,20}$$

An usual (strong) **zigzag** is a left/right, both extreme, edge-circuit.


38 symmetry groups of icosahedrites

- Agregating $\mathbf{C}_1 = \{C_1, C_s, C_i\}$, $\mathbf{C}_m = \{C_m, C_{mv}, C_{mh}, S_{2m}\}$, $\mathbf{D}_m = \{D_m, D_{mh}, D_{md}\}$, $\mathbf{T} = \{T, T_d, T_h\}$, $\mathbf{O} = \{O, O_h\}$, $\mathbf{I} = \{I, I_h\}$, all 38 symmetries of $(\{3, 4\}, 5)$ -spheres are:
 \mathbf{C}_1 , \mathbf{C}_m , \mathbf{D}_m for $2 \leq m \leq 5$ and \mathbf{T} , \mathbf{O} , \mathbf{I} .
- Any group appear an infinite number of times since one gets an infinity by applying A -operation iteratively.
- Group limitations came from k -fold axis only. Is it occurs for all $(\{a, b\}, k)$ -spheres with b -faces of negative curvature?
- Examples (minimal whenever $v \leq 32$) are given below:


22 C_1 22 C_s 32 C_i 72 O_h

Minimal $(\{3, 4\}, 5)$ -spheres of 5-fold symmetry


It exists iff $p_4 \equiv 0 \pmod{5}$, i.e., $v = 2p_4 + 12 \equiv 2 \pmod{10}$.




32 D_5




22 D_{5h}




32 D_{5d}




12 I_h Icosahed.




52 C_5



62 C_{5h}




72 C_{5v}




72 S_{10}

Minimal $(\{3, 4\}, 5)$ -spheres of 4-fold symmetry


It exists iff $p_4 \equiv 2 \pmod{4}$, i.e., $v = 2p_4 + 12 \equiv 0 \pmod{8}$.




32 D_4




16 D_{4d}




40 D_{4h}




24 O Snub cube




40 C_4



40 C_{4v}



32 C_{4h}




32 S_8


Icosahedron, Snub Cube and, with $(b, v; G) = (5, 60; I)$, Snub Dodecahedron are the only vertex-transitive $(\{3, b\}, 5)$ -spheres.

Minimal $(\{3, 4\}, 5)$ -spheres of 3-fold symmetry


It exists iff $p_4 \equiv 0 \pmod{3}$, i.e., $v = 2p_4 + 12 \equiv 0 \pmod{6}$.




18 D_3




24 D_{3d}




30 D_{3h}




36 T_d




30 C_3



30 C_{3v}











24 C_{3h}




24 S_6

Minimal $(\{3, 4\}, 5)$ -spheres of 2-fold symmetry


 $20 D_2$  $20 D_{2d}$  $24 D_{2h}$  $36 T_h$  $20 C_2$  $22 C_{2v}$  $28 C_{2h}$  $28 S_4$

Face-regular $(\{3, b\}, 5)$ -spheres


- A 3-connected map (on sphere or torus) is **pR_i face-regular** if any p -gonal face is adjacent to exactly i p -gons.
- No $(\{3, b\}, 5)$ -sphere, besides Icosahedron $3R_3$, is $3R_i$.
- Clearly, bR_j $(\{3, b\}, 5)$ -sphere has $j \frac{p_b}{2}$ $(b - b)$ -edges. So, bR_j with odd j implies that 4 divides $v = 2p_b(b - 3) + 12$.
- There is infinity of bR_j $(\{3, b\}, 5)$ -spheres for $j = 0, 1, 2$.




20 D_2 $4R_1$
 $G_4 = 2K_2$



36 T_d $4R_1$
 $G_4 = 6K_2$



22 D_{5h} $4R_2$
 $G_4 = C_5$



24 D_{3d} $4R_2$
 $G_4 = C_6$

b -gon-transitive of $(\{3, b\}, 5)$ -spheres

- Icosahedron (snub $APrism_3$) is regular. So, let $p_b > 0$.
- Snub $APrism_b$ has $v = 4b$ vertices (2 orbits of size $2b$), 2 b -gons (1 orbit) and $6b$ 3-gons (2 orbits of size $3b$). Its group G is D_{bd} for $b \geq 4$.
- With $(b, v; G) = (4, 24; O), (5, 60; I)$, Snub Cube and Snub Dodecahedron are only vertex-transitive $(\{3, b\}, 5)$ -spheres. They are also b -gon-transitive and have 2 orbits of triangles.
- Do other **b -gon-transitive** $(\{3, b\}, 5)$ -spheres or $(\{3, b\}, 5)$ -spheres **with at most 3 orbits of faces** exist?

XII. Standard $(\{a, b\}, k)$ -maps on surfaces

Standard (R, k) -maps

- Given $R \subset \mathbb{N}$ and a surface \mathbb{F}^2 , an (R, k) - \mathbb{F}^2 is a k -regular map M on surface \mathbb{F}^2 whose faces have gonality $i \in R$.
- Euler characteristic** $\chi(M)$ is $v - e + f$, where v, e and $f = \sum_i p_i$ are the numbers of vertices, edges and faces of M .
- Since $kv = 2e = \sum_i ip_i$, Euler formula $\chi = v - e + f$ becomes Gauss-Bonnet-like one $\chi(M) = \sum_i p_i \kappa_i$.
- Again, let our maps be **standard**, i.e., $\min_{i \in R} (1 + \frac{i}{k} - \frac{i}{2}) = 0$.
So, $M = \max\{i \in R\} = \frac{2k}{k-2}$ and $(M, k) = (6, 3), (4, 4), (3, 6)$.
- There are infinity of standard maps (R, k) - \mathbb{F}^2 , since the number p_M of flat ($\kappa_M = 0$) faces is not restricted.
- Also, $\chi \geq 0$ with $\chi = 0$ if and only if $R = \{m\}$.
So, \mathbb{F}^2 is $\mathbb{S}^2, \mathbb{T}^2, \mathbb{P}^2, \mathbb{K}^2$ with $\chi = 2, 0, 1, 0$, respectively.
- Such $(\{a, b\}, k)$ - \mathbb{F}^2 map has $b = \frac{2k}{k-2}$, $p_a = \frac{\chi b}{b-a}$, $v = \frac{1}{k}(ap_a + bp_b)$
So, $(a=b, k) = (6, 3), (3, 6), (4, 4)$ if \mathbb{F}^2 is \mathbb{T}^2 or \mathbb{K}^2 .
- But $\chi = \frac{p_3 - 2p_4}{10}$ for icosahedrite maps $(\{3, 4\}, 5)$ (non-standard)
So, $\chi < 0$ is possible and $\chi = 0$ (i.e., $\mathbb{F}^2 = \mathbb{T}^2, \mathbb{K}^2$) iff $p_3 = 2p_4$.

Digression on interesting non-standard $(\{5, 6, c\}, 3)$ -maps

Such maps, generalizing fullerenes, have $c \geq 7$. Examples are:

- **Haeckel, 1887**: $(\{5, 6, c\}, 3)$ -spheres with $c = 7, 8$ representing skeletons of radiolarian zooplankton **Aulonia hexagona**
- **Fullerene c -disks** ($(\{5, 6, c\}, 3)$ -spheres with $p_c = 1$) if $c \geq 7$ (**Deza-Dutour-Shtogrin, 2011-2012**)
- **G-fulleroids** (**Deza-Delgado, 2000; Jendrol-Trenkler, 2001 and Kardos, 2007**): $(\{5, b\}, 3)$ -spheres with $b \geq 7$ and symmetry G
- **Azulenoids**: $(\{5, 6, 7\}, 3)$ -tori; so, $g = 1, p_5 = p_7$ (**Kirby-Diudea, 2003, et al.**)
- **Schwartzits**: $(\{5, 6, c\}, 3)$ -maps on minimal surfaces of constant negative curvature ($g \geq 2$) with $c = 7, 8$ (**Terrones-MacKay, 1997, et al.**)
Knor-Potocnik-Siran-Skrekovski, 2010: such $(\{6, c\}, 3)$ -maps exist for any $g \geq 2, p_6 \geq 0$ and $c = 7, 8, 9, 10, 12$. For $c = 7, 8$ such polyhedral maps exist.

The $(\{a, b\}, k)$ -maps on torus and Klein bottle

The connected *closed* (compact and without boundary) irreducible surfaces are: sphere \mathbb{S}^2 , torus \mathbb{T}^2 (two **orientable**), real projective plane \mathbb{P}^2 and Klein bottle \mathbb{K}^2 with $\chi = 2, 0, 1, 0$, respectively.

The maps $(\{a, b\}, k)$ - \mathbb{T}^2 and $(\{a, b\}, k)$ - \mathbb{K}^2 have $a = b = \frac{2k}{k-2}$; so, $(a = b, k)$ should be $(6, 3)$, $(3, 6)$ or $(4, 4)$.

We consider only **polyhedral** maps, i.e. no loops or multiple edges (1- or 2-gons), and any two faces intersect in edge, point or \emptyset only.

The $(\{a, b\}, k)$ -maps on torus and Klein bottle

The connected *closed* (compact and without boundary) irreducible surfaces are: sphere \mathbb{S}^2 , torus \mathbb{T}^2 (two **orientable**), real projective plane \mathbb{P}^2 and Klein bottle \mathbb{K}^2 with $\chi = 2, 0, 1, 0$, respectively.

The maps $(\{a, b\}, k)$ - \mathbb{T}^2 and $(\{a, b\}, k)$ - \mathbb{K}^2 have $a = b = \frac{2k}{k-2}$; so, $(a = b, k)$ should be $(6, 3)$, $(3, 6)$ or $(4, 4)$.

We consider only **polyhedral** maps, i.e. no loops or multiple edges (1- or 2-gons), and any two faces intersect in edge, point or \emptyset only.

Smallest \mathbb{T}^2 and \mathbb{K}^2 -embeddings for $(a=b, k)=(6, 3), (3, 6), (4, 4)$:


as 6-regular **triangulations**: K_7 and $K_{3,3,3}$ ($p_3 = 14, 18$);

as 3-regular **polyhexes**: **Heawood graph** (dual K_7) and dual $K_{3,3,3}$;


as 4-regular **quadrangulations**: K_5 and $K_{2,2,2}$ ($p_4 = 5, 6$).

K_5 and $K_{2,2,2}$ are also smallest $(\{3, 4\}, 4)$ - \mathbb{P}^2 and $(\{3, 4\}, 4)$ - \mathbb{S}^2 , while K_4 is the smallest $(\{4, 6\}, 3)$ - \mathbb{P}^2 and $(\{3, 6\}, 3)$ - \mathbb{S}^2 .

Smallest 3-regular maps on \mathbb{T}^2 and \mathbb{K}^2 : duals K_7 , $K_{3,3,3}$



Smallest 3-regular maps on \mathbb{T}^2 and \mathbb{K}^2 : duals K_7 , $K_{3,3,3}$



3-regular polyhexes on \mathbb{T}^2 , cylinder, Möbius surface, \mathbb{K}^2 are $\{6^3\}$'s **quotients** by fixed-point-free group of isometries, generated by: two translations, a transl., a glide reflection, transl. *and* glide reflection.

8 families: symmetry groups with inversion


The point symmetry groups with inversion operation are: T_h , O_h , I_h , C_{mh} , D_{mh} with even m and D_{md} , S_{2m} with odd m . So, they are

- ① 9 for $\{5, 6\}_v$: C_i , C_{2h} , D_{2h} , D_{3d} , D_{6h} , S_6 , T_h , D_{5d} , I_h
- ② 7 for $\{2, 3\}_v$: C_i , C_{2h} , D_{2h} , D_{3d} , D_{6h} , S_6 , T_h
- ③ 6 for $\{4, 6\}_v$: C_i , C_{2h} , D_{2h} , D_{3d} , D_{6h} , O_h
- ④ 6 for $\{3, 4\}_v$: C_i , C_{2h} , D_{2h} , D_{3d} , D_{4h} , O_h
- ⑤ 2 for $\{2, 4\}_v$: D_{2h} , D_{4h}
- ⑥ 1 for $\{3, 6\}_v$: D_{2h}
- ⑦ 0 for $\{2, 6\}_v$ and $\{1, 3\}_v$
- ⑧ Cf. 12 for **icosahedrites** ($(\{3, 4\}, 5)$ -spheres):
 C_i , C_{2h} , C_{4h} , D_{2h} , D_{4h} , D_{3d} , D_{5d} , S_6 , S_{10} , T_h , O_h , I_h


(R, k) -maps on the **projective plane** are the antipodal quotients of centrosymmetric (R, k) -spheres; so, halving their p -vector and v .

Smallest $(\{a, b\}, k)$ -maps on the projective plane


- The smallest ones for $(a, b) = (4, 6), (3, 4), (3, 6), (5, 6)$ are:
 K_4 (smallest \mathbb{P}^2 -quadrangulation), K_5 , 2-truncated K_4 , dual K_6 (Petersen graph), i.e., the antipodal quotients of Cube
 $\{4, 6\}_8$, $\{3, 4\}_{10}(D_{4h})$, $\{3, 6\}_{16}(D_{2h})$, Dodecahedron $\{5, 6\}_{20}$.
- The smallest ones for $(a, b) = (2, 4), (2, 3)$ are points with 2, 3 loops; smallest without loops are $4 \times K_2$, $6 \times K_2$ but on \mathbb{P}^2 .




$\{4, 6\}_4$



$\{3, 4\}_5$




$\{3, 6\}_8$



$\{2, 4\}_2$

Smallest $(\{5, 6\}, 3)\text{-}\mathbb{P}^2$ and $(\{3, 4\}, 5)\text{-}\mathbb{P}^2$

The Petersen graph (in positive role) is the smallest \mathbb{P}^2 -fullerene. Its \mathbb{P}^2 -dual, K_6 , is the smallest \mathbb{P}^2 -icosahedrite (half-Icosahedron). K_6 is also the smallest (with 10 triangles) triangulation of \mathbb{P}^2 .



6 families on projective plane: parameterizing

- ① $\{5, 6\}_v$: $C_i, C_{2h}, D_{2h}, S_6, D_{3d}, D_{6h}, T_h, D_{5d}, I_h$
- ② $\{2, 3\}_v$: $C_i, C_{2h}, D_{2h}, S_6, D_{3d}, D_{6h}, T_h$
- ③ $\{4, 6\}_v$: $C_i, C_{2h}, D_{2h}, D_{3d}, D_{6h}, O_h$
- ④ $\{3, 4\}_v$: $C_i, C_{2h}, D_{2h}, D_{3d}, D_{4h}, O_h$
- ⑤ $\{2, 4\}_v$: D_{2h}, D_{4h}
- ⑥ $\{3, 6\}_v$: D_{2h}

6 families on projective plane: parameterizing

- ① $\{5, 6\}_v$: $C_i, C_{2h}, D_{2h}, S_6, D_{3d}, D_{6h}, T_h, D_{5d}, I_h$
- ② $\{2, 3\}_v$: $C_i, C_{2h}, D_{2h}, S_6, D_{3d}, D_{6h}, T_h$
- ③ $\{4, 6\}_v$: $C_i, C_{2h}, D_{2h}, D_{3d}, D_{6h}, O_h$
- ④ $\{3, 4\}_v$: $C_i, C_{2h}, D_{2h}, D_{3d}, D_{4h}, O_h$
- ⑤ $\{2, 4\}_v$: D_{2h}, D_{4h}
- ⑥ $\{3, 6\}_v$: D_{2h}

$(\{2, 3\}, 6)$ -spheres T_h and D_{6h} are $GC_{k,k}(2 \times Tetrahedron)$ and, for $k \equiv 1, 2 \pmod{3}$, $GC_{k,0}(6 \times K_2)$, respectively. Other spheres of blue symmetry are $GC_{k,l}$ with $l = 0, k$ from the first such sphere.

So, each of 7 blue-symmetric families is described by one natural parameter k and contains $O(\sqrt{v})$ spheres with at most v vertices.

$(\{a, b\}, k)$ -maps on Euclidean plane and 3-space

- An $(\{a, b\}, k)$ - \mathbb{E}^2 is a k -regular tiling of \mathbb{E}^2 by a - and b -gons.
- $(\{a, b\}, k)$ - \mathbb{E}^2 have $p_a \leq \frac{b}{b-a}$ and $p_b = \infty$. It follows from [Alexandrov, 1958](#): any metric on \mathbb{E}^2 of non-negative curvature can be realized as a metric of convex surface on \mathbb{E}^3 . In fact, consider plane metric such that all faces became regular in it. Its curvature is 0 on all interior points (faces, edges) and ≥ 0 on vertices. A convex surface is at most half- \mathbb{S}^2 .
- There are ∞ of $(\{a, b\}, k)$ - \mathbb{E}^2 if $2 \leq p_a \leq \frac{b}{b-a}$ and 1 if $p_a = 0, 1$.
- The **plane fullerenes** (or **nanocones**) $(\{5, 6\}, k)$ - \mathbb{E}^2 are classified by [Klein and Balaban, 2007](#): the number of *equivalence* (isomorphism up to a finite induced subgraph) classes is 2, 2, 2, 1 for $p_5 = 2, 3, 4, 5$, respectively.





$(\{a, b\}, k)$ -maps on Euclidean plane and 3-space

- An $(\{a, b\}, k)\text{-}\mathbb{E}^2$ is a k -regular tiling of \mathbb{E}^2 by a - and b -gons.
- $(\{a, b\}, k)\text{-}\mathbb{E}^2$ have $p_a \leq \frac{b}{b-a}$ and $p_b = \infty$. It follows from [Alexandrov, 1958](#): any metric on \mathbb{E}^2 of non-negative curvature can be realized as a metric of convex surface on \mathbb{E}^3 . In fact, consider plane metric such that all faces became regular in it. Its curvature is 0 on all interior points (faces, edges) and ≥ 0 on vertices. A convex surface is at most half- \mathbb{S}^2 .
- There are ∞ of $(\{a, b\}, k)\text{-}\mathbb{E}^2$ if $2 \leq p_a \leq \frac{b}{b-a}$ and 1 if $p_a = 0, 1$.
- The **plane fullerenes** (or **nanocones**) $(\{5, 6\}, k)\text{-}\mathbb{E}^2$ are classified by [Klein and Balaban, 2007](#): the number of *equivalence* (isomorphism up to a finite induced subgraph) classes is 2, 2, 2, 1 for $p_5 = 2, 3, 4, 5$, respectively.
- An $(\{a, b\}, k)\text{-}\mathbb{E}^3$ is a 3-periodic k' -regular face-to-face tiling of the Euclidean 3-space \mathbb{E}^3 by $(\{a, b\}, k)$ -spheres.
- Next, we will mention such tilings by 4 special fullerenes, which are important in Chemistry and Crystallography. Then we consider extension of $(\{a, b\}, k)$ -maps on manifolds.

XIII. Beyond surfaces

Frank-Kasper $(\{a, b\}, k)$ -spheres and tilings


- A $(\{a, b\}, k)$ -sphere is **Frank-Kasper** if no b -gons are adjacent.
- All cases are: smallest ones in 8 families, 3 $(\{5, 6\}, 3)$ -spheres (24-, 26-, 28-vertex fullerenes), $(\{4, 6\}, 3)$ -sphere $Prism_6$, 3 $(\{3, 4\}, 4)$ -spheres ($APrism_4$, $APrism_3^2$, Cuboctahedron), $(\{2, 4\}, 4)$ -sphere doubled square and two $(\{2, 3\}, 6)$ -spheres (tripled triangle and doubled Tetrahedron).

20, I_h 24 D_{6d} 26, D_{3h} 28, T_d

FK space fullerenes

A **FK space fullerene** is a 3-periodic 4-regular face-to-face tiling of 3-space \mathbb{E}^3 by four Frank-Kasper fullerenes $\{5, 6\}_v$.

They appear in crystallography of alloys, clathrate hydrates, zeolites and bubble structures. The most important, A_{15} , is below.



Weaire-Phelan, 1994: best known solution of weak Kelvin problem

Other \mathbb{E}^3 -tilings by $(\{a, b\}, k)$ -spheres

An $(\{a, b\}, k)\text{-}\mathbb{E}^3$ is a 3-periodic k' -regular face-to-face \mathbb{E}^3 -tiling by $(\{a, b\}, k)$ -spheres.

Deza-Shtogrin, 1999: first known non-FK **space fullerene**

$(\{5, 6\}, 3)\text{-}\mathbb{E}^3$: 4-regular \mathbb{E}^3 -tiling by $\{5, 6\}_{20}$, $\{5, 6\}_{24}$ and its elongation $\simeq \{5, 6\}_{36}$ (D_{6h}) in proportion 7:2:1.


Fullerene manifolds

- Given $3 \leq a < b \leq 6$, $\{a, b\}$ -manifold is a $(d-1)$ -dimensional d -valent compact connected *manifold* (locally homeomorphic to \mathbb{R}^{d-1}) whose 2-faces are only a - or b -gonal.
- So, any i -face, $3 \leq i \leq d$, is a polytopal i - $\{a, b\}$ -manifold.
- Most interesting case is $(a, b) = (5, 6)$ (**fullerene manifold**), when $d = 2, 3, 4, 5$ only since (Kalai, 1990) any 5-polytope has a 3- or 4-gonal 2-face.

Fullerene manifolds

- Given $3 \leq a < b \leq 6$, $\{a, b\}$ -manifold is a $(d-1)$ -dimensional d -valent compact connected *manifold* (locally homeomorphic to \mathbb{R}^{d-1}) whose 2-faces are only a - or b -gonal.
- So, any i -face, $3 \leq i \leq d$, is a polytopal i - $\{a, b\}$ -manifold.
- Most interesting case is $(a, b) = (5, 6)$ (**fullerene manifold**), when $d = 2, 3, 4, 5$ only since (Kalai, 1990) any 5-polytope has a 3- or 4-gonal 2-face.
- The smallest polyhex is 6-gon on \mathbb{T}^2 . The "greatest": $\{633\}$, the convex hull of vertices of $\{63\}$, realized on a horosphere.
- Prominent 4-fullerene (600-vertex on \mathbb{S}^3) is **120-cell** ($\{533\}$). The "greatest" polypent: $\{5333\}$, tiling of \mathbb{H}^4 by 120-cells.

Projection of 120-cell in 3-space



$\{533\}$: 600 vertices, 120 dodecahedral facets, $|Aut| = 14,400$

4- and 5-fullerenes

- All known finite 4-fullerenes are "mutations" of 120-cell by interfering in one of ways to construct it: tubes of 120-cells, coronas, inflation-decoration method, etc.
Some putative facets: $F_n(G)$ with $(n, G) = (20, I_h), (24, D_{6h}), (26, D_3), (28, T_d), (30, D_{5h}), (32, D_{3h}), (36, D_{6h})$.
- Space fullerenes $(\{5, 6\}, 3)\text{-}\mathbb{E}^3$: example of infinite 4-fullerenes.

4- and 5-fullerenes

- All known finite 4-fullerenes are "mutations" of 120-cell by interfering in one of ways to construct it: tubes of 120-cells, coronas, inflation-decoration method, etc.
Some putative facets: $F_n(G)$ with $(n, G) = (20, I_h), (24, D_{6h}), (26, D_3), (28, T_d), (30, D_{5h}), (32, D_{3h}), (36, D_{6h})$.
- Space fullerenes $(\{5, 6\}, 3)\text{-}\mathbb{E}^3$: example of infinite 4-fullerenes.
- All known 5-fullerenes come from $\{5333\}$'s by following ways.
With 6-gons also: glue two $\{5333\}$'s on some 120-cells and delete their interiors. If it is done on only one 120-cell, it is $\mathbb{R} \times \mathbb{S}^3$ (so, simply-connected).
Finite compact ones: the quotients of $\{5333\}$ by its symmetry group (partitioned into 120-cells) and gluings of them.

Quotient d -fullerenes

- Selberg, 1960, Borel, 1963: if a discrete group of motions of a symmetric space has a compact fundamental domain, then it has a torsion-free normal subgroup of finite index.
- So, the *quotient* of a d -fullerene by such symmetry group (its points are group orbits) is a **finite d -fullerene**.

Quotient d -fullerenes

- Selberg, 1960, Borel, 1963: if a discrete group of motions of a symmetric space has a compact fundamental domain, then it has a torsion-free normal subgroup of finite index.
- So, the *quotient* of a d -fullerene by such symmetry group (its points are group orbits) is a **finite d -fullerene**.
- Exp. 1: Polyhexes on \mathbb{T}^2 , cylinder, Möbius surface and \mathbb{K}^2 are the quotients of $\{6^3\}$ by discontinuous fixed-point-free group of isometries, generated by: 2 translations, a translation, a glide reflection, translation *and* glide reflection, respectively.

Quotient d -fullerenes

- Selberg, 1960, Borel, 1963: if a discrete group of motions of a symmetric space has a compact fundamental domain, then it has a torsion-free normal subgroup of finite index.
- So, the *quotient* of a d -fullerene by such symmetry group (its points are group orbits) is a **finite d -fullerene**.
- Exp. 1: Polyhexes on \mathbb{T}^2 , cylinder, Möbius surface and \mathbb{K}^2 are the quotients of $\{6^3\}$ by discontinuous fixed-point-free group of isometries, generated by: 2 translations, a translation, a glide reflection, translation *and* glide reflection, respectively.
- Exp 2: **Poincaré dodecahedral space**: the quotient of 120-cell by I_h ; so, its f -vector is $(5, 10, 6, 1) = \frac{1}{120}f(120\text{-cell})$.
- Cf. 6-, 12-regular \mathbb{H}^3 -tilings $\{5, 3, 4\}$, $\{5, 3, 5\}$ by $\{5, 6\}_{20}$ and 6-regular \mathbb{H}^3 -tiling by (right-angled) $\{5, 6\}_{24}$.
Seifert-Weber, 1933 and **Löbell, 1931** spaces are quotients of last 2 with f -vectors $(1, 6, p_5=6, 1)$, $(24, 72, 48+8=p_5+p_6, 8)$.