

# Optimization Models and Algorithms

Applications - Capacitated Vehicle Routing Problem

Mathieu Lerouge  
mathieu.lerouge@unibo.it

29 October 2024



ALMA MATER STUDIORUM  
UNIVERSITÀ DI BOLOGNA

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
- 3 Three-index formulation (Section 3.2)
- 4 Conclusion

# CVRP - Definition

## Capacitated Vehicle Routing Problem (CVRP):

- **Determining routes** for a fleet of identical vehicles, which have the same capacity and are originally located at a central depot, in order **to serve a set of clients** having different demands.
- Routes are such that:
  - ▶ each route **starts and ends at the depot**;
  - ▶ each client is **served by exactly one vehicle**;
  - ▶ the total demand served by each vehicle **does not exceed its capacity**.
- The aim is to **minimize the total cost** of all routes, given that any vehicle displacement has a cost.

# CVRP - Definition

## Capacitated Vehicle Routing Problem (CVRP):

- **Determining routes** for a fleet of identical vehicles, which have the same capacity and are originally located at a central depot, in order **to serve a set of clients** having different demands.
- Routes are such that:
  - ▶ each route **starts and ends at the depot**;
  - ▶ each client is **served by exactly one vehicle**;
  - ▶ the total demand served by each vehicle **does not exceed its capacity**.
- The aim is to **minimize the total cost** of all routes, given that any vehicle displacement has a cost.

# CVRP - Definition

## Capacitated Vehicle Routing Problem (CVRP):

- **Determining routes** for a fleet of identical vehicles, which have the same capacity and are originally located at a central depot, in order **to serve a set of clients** having different demands.
- Routes are such that:
  - ▶ each route **starts and ends at the depot**;
  - ▶ each client is **served by exactly one vehicle**;
  - ▶ the total demand served by each vehicle **does not exceed its capacity**.
- The aim is to **minimize the total cost** of all routes, given that any vehicle displacement has a cost.

# CVRP - Illustration




Figure: Illustration of the CVRP.

(Source: <https://ingenieriaindustrialonline.com>)

# CVRP - Notations and assumptions

## Notations:

- $G = (V, A)$ : directed graph, with nodes  $V = \{0, 1, \dots, n\}$ , such that:
  - ▶ 0 represents the **depot**;
  - ▶  $\{1, \dots, n\}$  represent the **clients**.
- $d_j$ : **demand** of client  $j$  (for  $j \in V \setminus \{0\}$ ).
- $k$ : **number** of identical vehicles.
- $b$ : vehicle **capacity**.
- $c_a$ : **cost** associated with arc  $a$  (for  $a \in A$ ).

## Assumptions:

- $G$  is assumed to be **complete** (without loss of generality).
- **Triangle inequalities** assumed to hold, therefore:  
visiting once client  $j \equiv$  visiting once node  $j$ .

# CVRP - Notations and assumptions

## Notations:

- $G = (V, A)$ : directed graph, with nodes  $V = \{0, 1, \dots, n\}$ , such that:
  - ▶ 0 represents the **depot**;
  - ▶  $\{1, \dots, n\}$  represent the **clients**.
- $d_j$ : **demand** of client  $j$  (for  $j \in V \setminus \{0\}$ ).
- $k$ : **number** of identical vehicles.
- $b$ : vehicle **capacity**.
- $c_a$ : **cost** associated with arc  $a$  (for  $a \in A$ ).

## Assumptions:

- $G$  is assumed to be **complete** (without loss of generality).
- **Triangle inequalities** assumed to hold, therefore:  
visiting once client  $j \equiv$  visiting once node  $j$ .



# CVRP - Notations and assumptions

## Notations:

- $G = (V, A)$ : directed graph, with nodes  $V = \{0, 1, \dots, n\}$ , such that:
  - ▶ 0 represents the **depot**;
  - ▶  $\{1, \dots, n\}$  represent the **clients**.
- $d_j$ : **demand** of client  $j$  (for  $j \in V \setminus \{0\}$ ).
- $k$ : **number** of identical vehicles.
- $b$ : vehicle **capacity**.
- $c_a$ : **cost** associated with arc  $a$  (for  $a \in A$ ).

## Assumptions:

- $G$  is assumed to be **complete** (without loss of generality).
- **Triangle inequalities** assumed to hold, therefore:  
visiting once client  $j \equiv$  visiting once node  $j$ .

# CVRP - Notations and assumptions

## Notations:

- $G = (V, A)$ : directed graph, with nodes  $V = \{0, 1, \dots, n\}$ , such that:
  - ▶ 0 represents the **depot**;
  - ▶  $\{1, \dots, n\}$  represent the **clients**.
- $d_j$ : **demand** of client  $j$  (for  $j \in V \setminus \{0\}$ ).
- $k$ : **number** of identical vehicles.
- $b$ : vehicle **capacity**.
- $c_a$ : **cost** associated with arc  $a$  (for  $a \in A$ ).

## Assumptions:

- $G$  is assumed to be **complete** (without loss of generality).
- **Triangle inequalities** assumed to hold, therefore:  
visiting once client  $j \equiv$  visiting once node  $j$ .

# CVRP - Notations and assumptions

## Notations:

- $G = (V, A)$ : directed graph, with nodes  $V = \{0, 1, \dots, n\}$ , such that:
  - ▶ 0 represents the **depot**;
  - ▶  $\{1, \dots, n\}$  represent the **clients**.
- $d_j$ : **demand** of client  $j$  (for  $j \in V \setminus \{0\}$ ).
- $k$ : **number** of identical vehicles.
- $b$ : vehicle **capacity**.
- $c_a$ : **cost** associated with arc  $a$  (for  $a \in A$ ).

## Assumptions:

- $G$  is assumed to be **complete** (without loss of generality).
- **Triangle inequalities** assumed to hold, therefore:  
visiting once client  $j \equiv$  visiting once node  $j$ .

# CVRP - Notations and assumptions

## Notations:

- $G = (V, A)$ : directed graph, with nodes  $V = \{0, 1, \dots, n\}$ , such that:
  - ▶ 0 represents the **depot**;
  - ▶  $\{1, \dots, n\}$  represent the **clients**.
- $d_j$ : **demand** of client  $j$  (for  $j \in V \setminus \{0\}$ ).
- $k$ : **number** of identical vehicles.
- $b$ : vehicle **capacity**.
- $c_a$ : **cost** associated with arc  $a$  (for  $a \in A$ ).

## Assumptions:

- $G$  is assumed to be **complete** (without loss of generality).
- **Triangle inequalities** assumed to hold, therefore:  
visiting once client  $j \equiv$  visiting once node  $j$ .

# CVRP - Notations and assumptions

## Notations:

- $G = (V, A)$ : directed graph, with nodes  $V = \{0, 1, \dots, n\}$ , such that:
  - ▶ 0 represents the **depot**;
  - ▶  $\{1, \dots, n\}$  represent the **clients**.
- $d_j$ : **demand** of client  $j$  (for  $j \in V \setminus \{0\}$ ).
- $k$ : **number** of identical vehicles.
- $b$ : vehicle **capacity**.
- $c_a$ : **cost** associated with arc  $a$  (for  $a \in A$ ).

## Assumptions:

- $G$  is assumed to be **complete** (without loss of generality).
- **Triangle inequalities** assumed to hold, therefore:  
visiting once client  $j \equiv$  visiting once node  $j$ .

# CVRP - Various formulations

- In this course, we study **three formulations** of the CVRP:
  1. **Two-index** formulation;
  2. **Three-index** formulation;
  3. **Set-partitioning** formulation.
- One formulation might be preferred to another depending on the context / problem:
  - ▶ basic CVRP,
  - ▶ CVRP with heterogeneous capacities,
  - ▶ CVRP with time-windows,
  - ▶ ...
- ⚠ In this course, while we model the CVRP as an **ILP model**, we are “only” interested in the **resolution of its LP-relaxation**.

# CVRP - Various formulations

- In this course, we study **three formulations** of the CVRP:
  1. **Two-index** formulation;
  2. **Three-index** formulation;
  3. **Set-partitioning** formulation.
- One formulation might be preferred to another depending on the context / problem:
  - ▶ basic CVRP,
  - ▶ CVRP with heterogeneous capacities,
  - ▶ CVRP with time-windows,
  - ▶ ...
- ⚠ In this course, while we model the CVRP as an **ILP model**, we are “only” interested in the **resolution of its LP-relaxation**.

# CVRP - Various formulations

- In this course, we study **three formulations** of the CVRP:
  1. **Two-index** formulation;
  2. **Three-index** formulation;
  3. **Set-partitioning** formulation.
- One formulation might be preferred to another depending on the context / problem:
  - ▶ basic CVRP,
  - ▶ CVRP with heterogeneous capacities,
  - ▶ CVRP with time-windows,
  - ▶ ...

⚠ In this course, while we model the CVRP as an **ILP model**, we are “only” interested in the **resolution of its LP-relaxation**.



# CVRP - Various formulations

- In this course, we study **three formulations** of the CVRP:
  1. **Two-index** formulation;
  2. **Three-index** formulation;
  3. **Set-partitioning** formulation.
- One formulation might be preferred to another depending on the context / problem:
  - ▶ basic CVRP,
  - ▶ CVRP with heterogeneous capacities,
  - ▶ CVRP with time-windows,
  - ▶ ...
- ⚠ In this course, while we model the CVRP as an **ILP model**, we are “only” interested in the **resolution of its LP-relaxation**.

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
- 3 Three-index formulation (Section 3.2)
- 4 Conclusion

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
  - Formulation generalities
  - Strong model of challenging constraints
  - Weak model of challenging constraints
- 3 Three-index formulation (Section 3.2)
- 4 Conclusion

# Two-index formulation - Principle

## Principle:

- This model uses, for each arc  $a \in A$ , a binary variable  $x_a$  to indicate whether  $a$  belongs to any vehicle route in the solution, **without specifying to which one**.
- In the literature, this model is often written using as variable indices **nodes instead of arcs**, i.e. using  $x_{ij}$ , with  $(i, j) \in V^2$  such that  $(i, j) \in A$ , instead of  $x_a$ .  
⇒ Hence the name “**two-index**” formulation.

# Two-index formulation - Principle

## Principle:

- This model uses, for each arc  $a \in A$ , a binary variable  $x_a$  to indicate whether  $a$  belongs to any vehicle route in the solution, **without specifying to which one**.
- In the literature, this model is often written using as variable indices **nodes instead of arcs**, i.e. using  $x_{ij}$ , with  $(i, j) \in V^2$  such that  $(i, j) \in A$ , instead of  $x_a$ .
  - ➔ Hence the name “**two-index**” formulation.

# Two-index formulation - Decision variables and objective

## Decision variables:

$$x_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to one of the } k \text{ routes,} \\ 0 & \text{otherwise.} \end{cases}$$

## Objective:

$$\min \sum_{a \in A} c_a x_a$$

# Two-index formulation - Decision variables and objective

## Decision variables:

$$x_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to one of the } k \text{ routes,} \\ 0 & \text{otherwise.} \end{cases}$$

## Objective:

$$\min \sum_{a \in A} c_a x_a$$

# Two-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a = 1, \quad i \in V \setminus \{0\}$$

(Flow at client nodes & single client visit)

$$\sum_{a \in \delta^+(i)} x_a = 1, \quad i \in V \setminus \{0\}$$

$$\sum_{a \in \delta^-(0)} x_a = k,$$

(Flow at depot & fixing nb of vehicles to  $k$ )

$$\sum_{a \in \delta^+(0)} x_a = k,$$

$$??? \quad ??? \quad (\text{Subtour elimination})$$

$$??? \quad ??? \quad (\text{Capacity})$$

$$x_a \in \{0, 1\}, \quad a \in A$$

⇒ “Challenging” constraints.



# Two-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a = 1, \quad i \in V \setminus \{0\} \quad (\text{Flow at client nodes \& single client visit})$$

$$\sum_{a \in \delta^+(i)} x_a = 1, \quad i \in V \setminus \{0\}$$

$$\sum_{a \in \delta^-(0)} x_a = k, \quad (\text{Flow at depot \& fixing nb of vehicles to } k)$$

$$\sum_{a \in \delta^+(0)} x_a = k,$$

$$??? \quad ??? \quad (\text{Subtour elimination})$$

$$??? \quad ??? \quad (\text{Capacity})$$

$$x_a \in \{0, 1\}, \quad a \in A$$

⇒ “Challenging” constraints.

# Two-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a = 1, \quad i \in V \setminus \{0\} \quad (\text{Flow at client nodes \& single client visit})$$

$$\sum_{a \in \delta^+(i)} x_a = 1, \quad i \in V \setminus \{0\}$$

$$\sum_{a \in \delta^-(0)} x_a = k, \quad (\text{Flow at depot \& fixing nb of vehicles to } k)$$

$$\sum_{a \in \delta^+(0)} x_a = k,$$

$$??? \quad ??? \quad (\text{Subtour elimination})$$

$$??? \quad ??? \quad (\text{Capacity})$$

$$x_a \in \{0, 1\}, \quad a \in A$$

⇒ “Challenging” constraints.

# Two-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a = 1, \quad i \in V \setminus \{0\} \quad (\text{Flow at client nodes \& single client visit})$$

$$\sum_{a \in \delta^+(i)} x_a = 1, \quad i \in V \setminus \{0\}$$

$$\sum_{a \in \delta^-(0)} x_a = k, \quad (\text{Flow at depot \& fixing nb of vehicles to } k)$$

$$\sum_{a \in \delta^+(0)} x_a = k,$$

$$??? \quad ??? \quad (\text{Subtour elimination})$$

$$??? \quad ??? \quad (\text{Capacity})$$

$$x_a \in \{0, 1\}, \quad a \in A$$

→ “Challenging” constraints.

# Two-index formulation - Challenging constraints

**Two ways** for modeling these challenging constraints  
(one single set of constraints for both considerations).

1. **Stronger** model, but **hard-to-use** in practice, based on:  
variants of ATSP (alternative) **subtour elimination constraints**  
and the **optimal value** of a BPP;
  2. **Weaker** but sufficient model, **easy-to-use** in practice, based on:  
variants of ATSP (alternative) **subtour elimination constraints**  
and the **trivial lower bound** of the same BPP.
- Let us recall some previous notions and results!

# Two-index formulation - Challenging constraints

**Two ways** for modeling these challenging constraints  
(one single set of constraints for both considerations).

1. **Stronger** model, but **hard-to-use** in practice, based on:  
variants of ATSP (alternative) **subtour elimination constraints**  
and the **optimal value** of a BPP;
  2. **Weaker** but sufficient model, **easy-to-use** in practice, based on:  
variants of ATSP (alternative) **subtour elimination constraints**  
and the **trivial lower bound** of the same BPP.
- Let us recall some previous notions and results!

# Two-index formulation - Challenging constraints

**Two ways** for modeling these challenging constraints  
(one single set of constraints for both considerations).

1. **Stronger** model, but **hard-to-use** in practice, based on:  
variants of ATSP (alternative) **subtour elimination constraints**  
and the **optimal value** of a BPP;
2. **Weaker** but sufficient model, **easy-to-use** in practice, based on:  
variants of ATSP (alternative) **subtour elimination constraints**  
and the **trivial lower bound** of the same BPP.


⇒ Let us recall some previous notions and results!

# Two-index formulation - Challenging constraints

**Two ways** for modeling these challenging constraints  
(one single set of constraints for both considerations).

1. **Stronger** model, but **hard-to-use** in practice, based on:  
variants of ATSP (alternative) **subtour elimination constraints**  
and the **optimal value** of a BPP;
  2. **Weaker** but sufficient model, **easy-to-use** in practice, based on:  
variants of ATSP (alternative) **subtour elimination constraints**  
and the **trivial lower bound** of the same BPP.
- Let us recall some previous notions and results!

# Recalls about ATSP - Illustration



**Figure:** Illustration of the ATSP.  
(Source: <https://en.wikipedia.org/>)



# Recalls about ATSP - Subtour elimination constraints

## Subtour elimination constraints:

Given a graph  $G = (V, A)$  and binary decision variables  $\{x_a\}$ ,  
s.t. for each  $a \in A$ ,  $x_a = 1$  if  $a$  belongs to the solution, 0 otherwise,  
**subtour elimination constraints** are:

$$\sum_{a \in A(S)} x_a \leq |S| - 1, \quad \forall S \subseteq V, 2 \leq |S| \leq |V| - 2$$



$$\sum_{a \in \delta^+(S)} x_a \geq 1, \quad \forall S \subseteq V, 2 \leq |S| \leq |V| - 2.$$

# Recalls about ATSP - Subtour elimination constraints

## Subtour elimination constraints:

Given a graph  $G = (V, A)$  and binary decision variables  $\{x_a\}$ ,  
s.t. for each  $a \in A$ ,  $x_a = 1$  if  $a$  belongs to the solution, 0 otherwise,  
**subtour elimination constraints** are:

$$\sum_{a \in A(S)} x_a \leq |S| - 1, \quad \forall S \subseteq V, 2 \leq |S| \leq |V| - 2$$



$$\sum_{a \in \delta^+(S)} x_a \geq 1, \quad \forall S \subseteq V, 2 \leq |S| \leq |V| - 2.$$

# Recalls about ATSP - Subtour elimination constraints

## Subtour elimination constraints:

Given a graph  $G = (V, A)$  and binary decision variables  $\{x_a\}$ ,  
s.t. for each  $a \in A$ ,  $x_a = 1$  if  $a$  belongs to the solution, 0 otherwise,  
**subtour elimination constraints** are:

$$\sum_{a \in A(S)} x_a \leq |S| - 1, \quad \forall S \subseteq V, 2 \leq |S| \leq |V| - 2$$



$$\sum_{a \in \delta^+(S)} x_a \geq 1, \quad \forall S \subseteq V, 2 \leq |S| \leq |V| - 2.$$

# Recalls about ATSP - Subtour elimination constraints

## Subtour elimination constraints:


Given a graph  $G = (V, A)$  and binary decision variables  $\{x_a\}$ ,  
s.t. for each  $a \in A$ ,  $x_a = 1$  if  $a$  belongs to the solution, 0 otherwise,  
**subtour elimination constraints** are:

$$\sum_{a \in A(S)} x_a \leq |S| - 1, \quad \forall S \subseteq V, 2 \leq |S| \leq |V| - 2$$



$$\sum_{a \in \delta^+(S)} x_a \geq 1, \quad \forall S \subseteq V, 2 \leq |S| \leq |V| - 2.$$

# Recalls about BPP - Illustration



**Figure:** Illustration of the BPP.  
(Source: Chraibi et al. (2022))

# Recalls about BPP - Trivial lower bound

## Trivial lower bound:

Given  $n$  items of weights  $d_1, \dots, d_n$  and identical bins of capacity  $b$ , a **trivial lower bound** on the nb of bins required to pack all items is:

$$\frac{\sum_{i=1}^n d_i}{b}.$$

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
  - Formulation generalities
  - Strong model of challenging constraints
  - Weak model of challenging constraints
- 3 Three-index formulation (Section 3.2)
- 4 Conclusion

# Strong model - Principle and constraints

## Idea:

- Given  $S \subseteq V \setminus \{0\}$ , let  $\sigma(S)$  be the **optimal value** of the **BPP** obtained by considering as:
  - set of **objects**  $\equiv$  subset of **clients**  $S$ ,
  - objects' **weights**  $\equiv$  clients' **demands**  $\{d_j\}_{j \in S}$ ,
  - container **capacity**  $\equiv$  vehicle **capacity**  $b$ .
- "Given a set of clients  $S \subseteq V \setminus \{0\}$ , **at least**  $\sigma(S)$  **vehicles** are needed to serve these clients."

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \sigma(S), \quad S \subseteq V \setminus \{0\}$$

- Subtour elimination** constraints as  $\sigma(S) \geq 1$ .
- Capacity** constraints by definition of  $\sigma(S)$ .



# Strong model - Principle and constraints

## Idea:

- Given  $S \subseteq V \setminus \{0\}$ , let  $\sigma(S)$  be the **optimal value** of the **BPP** obtained by considering as:
  - set of **objects**  $\equiv$  subset of **clients**  $S$ ,
  - objects' **weights**  $\equiv$  clients' **demands**  $\{d_j\}_{j \in S}$ ,
  - container **capacity**  $\equiv$  vehicle **capacity**  $b$ .
- "Given a set of clients  $S \subseteq V \setminus \{0\}$ , **at least**  $\sigma(S)$  **vehicles** are needed to serve these clients."

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \sigma(S), \quad S \subseteq V \setminus \{0\}$$

- Subtour elimination** constraints as  $\sigma(S) \geq 1$ .
- Capacity** constraints by definition of  $\sigma(S)$ .

# Strong model - Principle and constraints

## Idea:

- Given  $S \subseteq V \setminus \{0\}$ , let  $\sigma(S)$  be the **optimal value** of the **BPP** obtained by considering as:
  - set of **objects**  $\equiv$  subset of **clients**  $S$ ,
  - objects' **weights**  $\equiv$  clients' **demands**  $\{d_j\}_{j \in S}$ ,
  - container **capacity**  $\equiv$  vehicle **capacity**  $b$ .
- "Given a set of clients  $S \subseteq V \setminus \{0\}$ , **at least**  $\sigma(S)$  **vehicles** are needed to serve these clients."

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \sigma(S), \quad S \subseteq V \setminus \{0\}$$

- Subtour elimination** constraints as  $\sigma(S) \geq 1$ .
- Capacity** constraints by definition of  $\sigma(S)$ .

# Strong model - Principle and constraints

## Idea:

- Given  $S \subseteq V \setminus \{0\}$ , let  $\sigma(S)$  be the **optimal value** of the **BPP** obtained by considering as:
  - set of **objects**  $\equiv$  subset of **clients**  $S$ ,
  - objects' **weights**  $\equiv$  clients' **demands**  $\{d_j\}_{j \in S}$ ,
  - container **capacity**  $\equiv$  vehicle **capacity**  $b$ .
- "Given a set of clients  $S \subseteq V \setminus \{0\}$ , **at least**  $\sigma(S)$  **vehicles** are needed to serve these clients."

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \sigma(S), \quad S \subseteq V \setminus \{0\}$$

- Subtour elimination** constraints as  $\sigma(S) \geq 1$ .
- Capacity** constraints by definition of  $\sigma(S)$ .

# Strong model - Principle and constraints

## Idea:

- Given  $S \subseteq V \setminus \{0\}$ , let  $\sigma(S)$  be the **optimal value** of the **BPP** obtained by considering as:
  - set of **objects**  $\equiv$  subset of **clients**  $S$ ,
  - objects' **weights**  $\equiv$  clients' **demands**  $\{d_j\}_{j \in S}$ ,
  - container **capacity**  $\equiv$  vehicle **capacity**  $b$ .
- "Given a set of clients  $S \subseteq V \setminus \{0\}$ , **at least**  $\sigma(S)$  **vehicles** are needed to serve these clients."

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \sigma(S), \quad S \subseteq V \setminus \{0\}$$

- Subtour elimination** constraints as  $\sigma(S) \geq 1$ .
- Capacity** constraints by definition of  $\sigma(S)$ .

# Strong model - Principle and constraints

## Idea:

- Given  $S \subseteq V \setminus \{0\}$ , let  $\sigma(S)$  be the **optimal value** of the **BPP** obtained by considering as:
  - set of **objects**  $\equiv$  subset of **clients**  $S$ ,
  - objects' **weights**  $\equiv$  clients' **demands**  $\{d_j\}_{j \in S}$ ,
  - container **capacity**  $\equiv$  vehicle **capacity**  $b$ .
- "Given a set of clients  $S \subseteq V \setminus \{0\}$ , **at least**  $\sigma(S)$  **vehicles** are needed to serve these clients."

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \sigma(S), \quad S \subseteq V \setminus \{0\}$$

- Subtour elimination** constraints as  $\sigma(S) \geq 1$ .
- Capacity** constraints by definition of  $\sigma(S)$ .

# Strong model - Within CVRP two-index formulation

## CVRP two-index formulation:

$$\begin{aligned}
 \min \quad & \sum_{a \in A} c_a x_a \\
 \text{s.t.} \quad & \sum_{a \in \delta^-(i)} x_a = 1, & i \in V \setminus \{0\} & \text{(Flow at client nodes \& single client visit)} \\
 & \sum_{a \in \delta^+(i)} x_a = 1, & i \in V \setminus \{0\} \\
 & \sum_{a \in \delta^-(0)} x_a = k, & \text{(Flow at depot \& fixing nb of vehicles to } k\text{)} \\
 & \sum_{a \in \delta^+(0)} x_a = k, \\
 & \sum_{a \in \delta^+(S)} x_a \geq \sigma(S), & S \subseteq V \setminus \{0\} & \text{(Subtour elimination \& vehicle capacity)} \\
 & x_a \in \{0, 1\}, & a \in A
 \end{aligned}$$

⚠ Exponential nb of constraints  $\rightarrow$  separation with  $\tilde{S} \subseteq 2^{V \setminus \{0\}}$ .

# Strong model - Within CVRP two-index formulation

## CVRP two-index formulation **LP-relaxation**:

$$\begin{aligned}
 \min \quad & \sum_{a \in A} c_a x_a \\
 \text{s.t.} \quad & \sum_{a \in \delta^-(i)} x_a = 1, & i \in V \setminus \{0\} & \text{(Flow at client nodes \& single client visit)} \\
 & \sum_{a \in \delta^+(i)} x_a = 1, & i \in V \setminus \{0\} \\
 & \sum_{a \in \delta^-(0)} x_a = k, & \text{(Flow at depot \& fixing nb of vehicles to } k) \\
 & \sum_{a \in \delta^+(0)} x_a = k, \\
 & \sum_{a \in \delta^+(S)} x_a \geq \sigma(S), & S \subseteq V \setminus \{0\} & \text{(Subtour elimination \& vehicle capacity)} \\
 & x_a \in [0, 1], & a \in A
 \end{aligned}$$

⚠ Exponential nb of constraints → separation with  $\tilde{S} \subseteq 2^{V \setminus \{0\}}$ .

# Strong model - Within CVRP two-index formulation

## CVRP two-index formulation LP-relaxation:

$$\begin{aligned}
 \min \quad & \sum_{a \in A} c_a x_a \\
 \text{s.t.} \quad & \sum_{a \in \delta^-(i)} x_a = 1, & i \in V \setminus \{0\} & \text{(Flow at client nodes \& single client visit)} \\
 & \sum_{a \in \delta^+(i)} x_a = 1, & i \in V \setminus \{0\} \\
 & \sum_{a \in \delta^-(0)} x_a = k, & \text{(Flow at depot \& fixing nb of vehicles to } k\text{)} \\
 & \sum_{a \in \delta^+(0)} x_a = k, \\
 & \sum_{a \in \delta^+(S)} x_a \geq \sigma(S), & S \subseteq V \setminus \{0\} & \text{(Subtour elimination \& vehicle capacity)} \\
 & x_a \in [0, 1], & a \in A
 \end{aligned}$$

⚠ **Exponential nb of constraints** → separation with  $\tilde{S} \subseteq 2^{V \setminus \{0\}}$ .



# Strong model - Within CVRP two-index formulation

## CVRP two-index formulation LP-relaxation:

$$\begin{aligned}
 \min \quad & \sum_{a \in A} c_a x_a \\
 \text{s.t.} \quad & \sum_{a \in \delta^-(i)} x_a = 1, & i \in V \setminus \{0\} & \text{(Flow at client nodes \& single client visit)} \\
 & \sum_{a \in \delta^+(i)} x_a = 1, & i \in V \setminus \{0\} \\
 & \sum_{a \in \delta^-(0)} x_a = k, & \text{(Flow at depot \& fixing nb of vehicles to } k\text{)} \\
 & \sum_{a \in \delta^+(0)} x_a = k, \\
 & \sum_{a \in \delta^+(S)} x_a \geq \sigma(S), & S \in \tilde{\mathcal{S}} & \text{(Subtour elimination \& vehicle capacity)} \\
 & x_a \in [0, 1], & a \in A
 \end{aligned}$$

⚠ **Exponential nb** of constraints → **separation** with  $\tilde{\mathcal{S}} \subseteq 2^{V \setminus \{0\}}$ .

# Strong model - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{\mathcal{S}}$ ). Let  $x^*$  be its **optimal solution**.

## Separation:

- Identify **if there exists** a subset  $S^* \subseteq V \setminus \{0\}$  such that:

$$\sum_{a \in \delta^+(S^*)} x_a^* < \sigma(S^*)$$

- Use **(M)ILP** to model this problem into a **separation problem**?

⚠ **Cannot be formulated** as a reasonable (M)ILP because the model would have to both select vertices to form  $S^*$  and compute the BPP optimal value  $\sigma(S^*)$ .

→ Use heuristics to find  $S^*$  or change of separation problem.

# Strong model - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{\mathcal{S}}$ ). Let  $x^*$  be its **optimal solution**.

## Separation:

- Identify **if there exists** a subset  $S^* \subseteq V \setminus \{0\}$  such that:

$$\sum_{a \in \delta^+(S^*)} x_a^* < \sigma(S^*)$$

- Use (M)ILP to model this problem into a **separation problem**?

⚠ **Cannot be formulated** as a reasonable (M)ILP because the model would have to both select vertices to form  $S^*$  and compute the BPP optimal value  $\sigma(S^*)$ .

→ Use heuristics to find  $S^*$  or change of separation problem.

# Strong model - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{\mathcal{S}}$ ). Let  $x^*$  be its **optimal solution**.

## Separation:

- Identify **if there exists** a subset  $S^* \subseteq V \setminus \{0\}$  such that:

$$\sum_{a \in \delta^+(S^*)} x_a^* < \sigma(S^*)$$

- Use **(M)ILP** to model this problem into a **separation problem**?

⚠ **Cannot be formulated** as a reasonable (M)ILP because the model would have to both select vertices to form  $S^*$  and compute the BPP optimal value  $\sigma(S^*)$ .

→ Use heuristics to find  $S^*$  or change of separation problem.

# Strong model - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{\mathcal{S}}$ ). Let  $x^*$  be its **optimal solution**.

## Separation:

- Identify **if there exists** a subset  $S^* \subseteq V \setminus \{0\}$  such that:

$$\sum_{a \in \delta^+(S^*)} x_a^* < \sigma(S^*)$$

- Use **(M)ILP** to model this problem into a **separation problem**?

**⚠ Cannot be formulated** as a reasonable (M)ILP because the model would have to both select vertices to form  $S^*$  and compute the BPP optimal value  $\sigma(S^*)$ .

⇒ Use heuristics to find  $S^*$  or change of separation problem.

# Strong model - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{\mathcal{S}}$ ). Let  $x^*$  be its **optimal solution**.

## Separation:

- Identify **if there exists** a subset  $S^* \subseteq V \setminus \{0\}$  such that:

$$\sum_{a \in \delta^+(S^*)} x_a^* < \sigma(S^*)$$

- Use **(M)ILP** to model this problem into a **separation problem**?

⚠ **Cannot be formulated** as a reasonable (M)ILP because the model would have to both select vertices to form  $S^*$  and compute the BPP optimal value  $\sigma(S^*)$ .

→ Use heuristics to find  $S^*$  or change of separation problem.

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
  - Formulation generalities
  - Strong model of challenging constraints
  - Weak model of challenging constraints
- 3 Three-index formulation (Section 3.2)
- 4 Conclusion

# Weak model - Principle and constraints

## Principle:

- Same BPP problem as before, but rather than making use of the optimal value  $\sigma(S)$ , we use its **trivial lower bound**:

$$\sigma(S) \geq \frac{\sum_{j \in S} d_j}{b}$$

- “Given a set of clients  $S \subseteq V \setminus \{0\}$ , at least  $\frac{\sum_{j \in S} d_j}{b}$  vehicles are needed to serve these clients.”

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \frac{\sum_{j \in S} d_j}{b}, \quad S \subseteq V \setminus \{0\}$$



# Weak model - Principle and constraints

## Principle:

- Same BPP problem as before, but rather than making use of the optimal value  $\sigma(S)$ , we use its **trivial lower bound**:

$$\sigma(S) \geq \frac{\sum_{j \in S} d_j}{b}$$

- “Given a set of clients  $S \subseteq V \setminus \{0\}$ , **at least**  $\frac{\sum_{j \in S} d_j}{b}$  **vehicles** are needed to serve these clients.”

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \frac{\sum_{j \in S} d_j}{b}, \quad S \subseteq V \setminus \{0\}$$

# Weak model - Principle and constraints

## Principle:

- Same BPP problem as before, but rather than making use of the optimal value  $\sigma(S)$ , we use its **trivial lower bound**:

$$\sigma(S) \geq \frac{\sum_{j \in S} d_j}{b}$$

- “Given a set of clients  $S \subseteq V \setminus \{0\}$ , **at least**  $\frac{\sum_{j \in S} d_j}{b}$  **vehicles** are needed to serve these clients.”

## Challenging constraints:

$$\sum_{a \in \delta^+(S)} x_a \geq \frac{\sum_{j \in S} d_j}{b}, \quad S \subseteq V \setminus \{0\}$$

# Weak model - Within CVRP two-index formulation

## CVRP two-index formulation:

$$\begin{aligned}
 \min \quad & \sum_{a \in A} c_a x_a \\
 \text{s.t.} \quad & \sum_{a \in \delta^-(i)} x_a = 1, & i \in V \setminus \{0\} & \text{(Flow at client nodes \& single client visit)} \\
 & \sum_{a \in \delta^+(i)} x_a = 1, & i \in V \setminus \{0\} \\
 & \sum_{a \in \delta^-(0)} x_a = k, & \text{(Flow at depot \& fixing nb of vehicles to } k\text{)} \\
 & \sum_{a \in \delta^+(0)} x_a = k, \\
 & \sum_{a \in \delta^+(S)} x_a \geq \frac{\sum_{j \in S} d_j}{b}, & S \subseteq V \setminus \{0\} & \text{(Subtour elimination \& vehicle capacity)} \\
 & x_a \in \{0, 1\}, & a \in A
 \end{aligned}$$

⚠ Exponential nb of constraints → separation with  $\tilde{S} \subseteq 2^{V \setminus \{0\}}$ .

# Weak model - Within CVRP two-index formulation

## CVRP two-index formulation LP-relaxation:

$$\begin{aligned}
 \min \quad & \sum_{a \in A} c_a x_a \\
 \text{s.t.} \quad & \sum_{a \in \delta^-(i)} x_a = 1, & i \in V \setminus \{0\} & \text{(Flow at client nodes \& single client visit)} \\
 & \sum_{a \in \delta^+(i)} x_a = 1, & i \in V \setminus \{0\} \\
 & \sum_{a \in \delta^-(0)} x_a = k, & \text{(Flow at depot \& fixing nb of vehicles to } k\text{)} \\
 & \sum_{a \in \delta^+(0)} x_a = k, \\
 & \sum_{a \in \delta^+(S)} x_a \geq \frac{\sum_{j \in S} d_j}{b}, & S \subseteq V \setminus \{0\} & \text{(Subtour elimination \& vehicle capacity)} \\
 & x_a \in [0, 1], & a \in A
 \end{aligned}$$

⚠ Exponential nb of constraints → separation with  $\tilde{S} \subseteq 2^{V \setminus \{0\}}$ .

# Weak model - Within CVRP two-index formulation

## CVRP two-index formulation LP-relaxation:

$$\begin{aligned}
 \min \quad & \sum_{a \in A} c_a x_a \\
 \text{s.t.} \quad & \sum_{a \in \delta^-(i)} x_a = 1, & i \in V \setminus \{0\} & \text{(Flow at client nodes \& single client visit)} \\
 & \sum_{a \in \delta^+(i)} x_a = 1, & i \in V \setminus \{0\} & \\
 & \sum_{a \in \delta^-(0)} x_a = k, & & \text{(Flow at depot \& fixing nb of vehicles to } k) \\
 & \sum_{a \in \delta^+(0)} x_a = k, & & \\
 & \sum_{a \in \delta^+(S)} x_a \geq \frac{\sum_{j \in S} d_j}{b}, & S \subseteq V \setminus \{0\} & \text{(Subtour elimination \& vehicle capacity)} \\
 & x_a \in [0, 1], & a \in A &
 \end{aligned}$$

⚠ **Exponential nb of constraints** → separation with  $\tilde{S} \subseteq 2^{V \setminus \{0\}}$ .

# Weak model - Within CVRP two-index formulation

## CVRP two-index formulation LP-relaxation:

$$\begin{aligned}
 \min \quad & \sum_{a \in A} c_a x_a \\
 \text{s.t.} \quad & \sum_{a \in \delta^-(i)} x_a = 1, & i \in V \setminus \{0\} & \text{(Flow at client nodes \& single client visit)} \\
 & \sum_{a \in \delta^+(i)} x_a = 1, & i \in V \setminus \{0\} & \\
 & \sum_{a \in \delta^-(0)} x_a = k, & & \text{(Flow at depot \& fixing nb of vehicles to } k\text{)} \\
 & \sum_{a \in \delta^+(0)} x_a = k, & & \\
 & \sum_{a \in \delta^+(S)} x_a \geq \frac{\sum_{j \in S} d_j}{b}, & S \in \tilde{\mathcal{S}} & \text{(Subtour elimination \& vehicle capacity)} \\
 & x_a \in [0, 1], & a \in A &
 \end{aligned}$$

⚠ **Exponential nb** of constraints → **separation** with  $\tilde{\mathcal{S}} \subseteq 2^{V \setminus \{0\}}$ .

# Weak formulation - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{S}$ ). Let  $x^*$  be its **optimal solution**.

## Separation:

- Identify if there exists a subset  $S^* \subseteq V \setminus \{0\}$  such that:

$$\sum_{a \in \delta^+(S^*)} x_a^* < \frac{\sum_{j \in S^*} d_j}{b} \quad \text{i.e.} \quad \sum_{j \in S^*} d_j - b \sum_{a \in \delta^+(S^*)} x_a^* > 0$$

- Use **ILP** to model this problem into a **separation problem**.

# Weak formulation - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{S}$ ). Let  $x^*$  be its **optimal solution**.

## Separation:

- Identify **if there exists** a subset  $S^* \subseteq V \setminus \{0\}$  such that:

$$\sum_{a \in \delta^+(S^*)} x_a^* < \frac{\sum_{j \in S^*} d_j}{b} \quad \text{i.e.} \quad \sum_{j \in S^*} d_j - b \sum_{a \in \delta^+(S^*)} x_a^* > 0$$

- Use **ILP** to model this problem into a **separation problem**.



# Weak formulation - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{S}$ ). Let  $x^*$  be its **optimal solution**.

## Separation:

- Identify **if there exists** a subset  $S^* \subseteq V \setminus \{0\}$  such that:

$$\sum_{a \in \delta^+(S^*)} x_a^* < \frac{\sum_{j \in S^*} d_j}{b} \quad \text{i.e.} \quad \sum_{j \in S^*} d_j - b \sum_{a \in \delta^+(S^*)} x_a^* > 0$$

- Use **ILP** to model this problem into a **separation problem**.

# Weak formulation - Separation problem

## Decision variables:

- $y_i = \begin{cases} 1 & \text{if client } i \text{ belongs to } S^*, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(S^*), \\ 0 & \text{otherwise.} \end{cases}$

## Objective:

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{j \in S^*} d_j - b \sum_{a \in \delta^+(S^*)} x_a^* > 0 \quad ?$$

↓

$$\max \quad \sum_{i \in V} d_i y_i - b \sum_{a \in A} x_a^* z_a > 0 \quad ?$$

# Weak formulation - Separation problem

## Decision variables:

- $y_i = \begin{cases} 1 & \text{if client } i \text{ belongs to } S^*, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(S^*), \\ 0 & \text{otherwise.} \end{cases}$

## Objective:

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{j \in S^*} d_j - b \sum_{a \in \delta^+(S^*)} x_a^* > 0 \quad ?$$

↓

$$\max \sum_{i \in V} d_i y_i - b \sum_{a \in A} x_a^* z_a > 0 \quad ?$$

# Weak formulation - Separation problem

## Decision variables:

- $y_i = \begin{cases} 1 & \text{if client } i \text{ belongs to } S^*, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(S^*), \\ 0 & \text{otherwise.} \end{cases}$

## Objective:

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{j \in S^*} d_j - b \sum_{a \in \delta^+(S^*)} x_a^* > 0 \quad ?$$

↓

$$\max \quad \sum_{i \in V} d_i y_i - b \sum_{a \in A} x_a^* z_a > 0 \quad ?$$

# Weak formulation - Separation problem

## Decision variables:

- $y_i = \begin{cases} 1 & \text{if client } i \text{ belongs to } S^*, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(S^*), \\ 0 & \text{otherwise.} \end{cases}$

## Objective:

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{j \in S^*} d_j - b \sum_{a \in \delta^+(S^*)} x_a^* > 0 \quad ?$$

↓

$$\max \quad \sum_{i \in V} d_i y_i - b \sum_{a \in A} x_a^* z_a > 0 \quad ?$$

# Weak formulation - Separation problem

## Separation problem:

$$\begin{aligned} \max \quad & \sum_{i \in V} d_i y_i - b \sum_{a \in A} x_a^* z_a \\ \text{s.t.} \quad & y_0 = 0 \\ & z_{(i,j)} \leq y_i, \quad (i,j) \in A \\ & z_{(i,j)} \leq 1 - y_j, \quad (i,j) \in A \\ & z_{(i,j)} \geq y_i - y_j, \quad (i,j) \in A \\ & y_i \in \{0, 1\}, \quad i \in V \\ & z_a \in \{0, 1\}, \quad a \in A \end{aligned}$$

→ If the optimal value is positive,  
**add**  $S^* = \{i \in V \setminus \{0\}, y_i = 1\}$  to  $\tilde{S}$ .

# Weak formulation - Separation problem

## Separation problem:

$$\begin{aligned} \max \quad & \sum_{i \in V} d_i y_i - b \sum_{a \in A} x_a^* z_a \\ \text{s.t.} \quad & y_0 = 0 \\ & z_{(i,j)} \leq y_i, \quad (i,j) \in A \\ & z_{(i,j)} \leq 1 - y_j, \quad (i,j) \in A \\ & z_{(i,j)} \geq y_i - y_j, \quad (i,j) \in A \\ & y_i \in \{0, 1\}, \quad i \in V \\ & z_a \in \{0, 1\}, \quad a \in A \end{aligned}$$

→ If the optimal value is positive,  
**add**  $S^* = \{i \in V \setminus \{0\}, y_i = 1\}$  to  $\tilde{S}$ .

# Weak formulation - Separation problem

## Separation problem:

$$\begin{aligned} \max \quad & \sum_{i \in V} d_i y_i - b \sum_{a \in A} x_a^* z_a \\ \text{s.t.} \quad & y_0 = 0 \\ & z_{(i,j)} \leq y_i, \quad (i,j) \in A \\ & z_{(i,j)} \leq 1 - y_j, \quad (i,j) \in A \\ & z_{(i,j)} \geq y_i - y_j, \quad (i,j) \in A \\ & y_i \in \{0, 1\}, \quad i \in V \\ & z_a \in \{0, 1\}, \quad a \in A \end{aligned}$$

→ If the optimal value is positive,  
add  $S^* = \{i \in V \setminus \{0\}, y_i = 1\}$  to  $\tilde{S}$ .



# Weak formulation - Separation problem

## Separation problem:

$$\begin{aligned} \max \quad & \sum_{i \in V} d_i y_i - b \sum_{a \in A} x_a^* z_a \\ \text{s.t.} \quad & y_0 = 0 \\ & z_{(i,j)} \leq y_i, \quad (i,j) \in A \\ & z_{(i,j)} \leq 1 - y_j, \quad (i,j) \in A \\ & z_{(i,j)} \geq y_i - y_j, \quad (i,j) \in A \\ & y_i \in \{0, 1\}, \quad i \in V \\ & z_a \in \{0, 1\}, \quad a \in A \end{aligned}$$

- If the optimal value is positive,  
**add**  $S^* = \{i \in V \setminus \{0\}, y_i = 1\}$  to  $\tilde{S}$ .

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
- 3 Three-index formulation (Section 3.2)
- 4 Conclusion

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
- 3 Three-index formulation (Section 3.2)
  - Formulation generalities
  - Extension to CVRP with heterogenous capacities
  - Extension to CVRP with time windows
- 4 Conclusion

# Three-index formulation - Principle

## Principle:

- This model uses, for each arc  $a \in A$  and each vehicle  $h \in \{1, \dots, k\}$ , a binary variable  $x_a^h$  to indicate whether arc  $a$  belongs to **vehicle  $h$ 's route**.
- From a resolution perspective, this three-index formulation does not offer any advantages over the two-index one.  
However, it is considerably **easier to adapt it** to integrate additional characteristics of routing problems.

# Three-index formulation - Principle

## Principle:

- This model uses, for each arc  $a \in A$  and each vehicle  $h \in \{1, \dots, k\}$ , a binary variable  $x_a^h$  to indicate whether arc  $a$  belongs to **vehicle  $h$ 's route**.
- From a resolution perspective, this three-index formulation does not offer any advantages over the two-index one.  
However, it is considerably **easier to adapt it** to integrate additional characteristics of routing problems.

# Three-index formulation - Decision variables and objective

## Decision variables:

- $x_a^h = \begin{cases} 1 & \text{if arc } a \text{ belongs to route } h, \\ 0 & \text{otherwise.} \end{cases}$
- $y_i^h = \begin{cases} 1 & \text{if client } i \text{ is served by route } h, \\ 0 & \text{otherwise.} \end{cases}$

⇒ Variables  $\{y_i^h\}$  are introduced for readability purpose: they correspond to sum over part of variables  $\{x_a\}$ .

## Objective:

$$\min \sum_{h=1}^k \sum_{a \in A} c_a x_a^h$$

# Three-index formulation - Decision variables and objective

## Decision variables:

- $x_a^h = \begin{cases} 1 & \text{if arc } a \text{ belongs to route } h, \\ 0 & \text{otherwise.} \end{cases}$
- $y_i^h = \begin{cases} 1 & \text{if client } i \text{ is served by route } h, \\ 0 & \text{otherwise.} \end{cases}$

⇒ Variables  $\{y_i^h\}$  are introduced for readability purpose: they correspond to sum over part of variables  $\{x_a\}$ .

## Objective:

$$\min \sum_{h=1}^k \sum_{a \in A} c_a x_a^h$$

# Three-index formulation - Decision variables and objective

## Decision variables:

- $x_a^h = \begin{cases} 1 & \text{if arc } a \text{ belongs to route } h, \\ 0 & \text{otherwise.} \end{cases}$
  - $y_i^h = \begin{cases} 1 & \text{if client } i \text{ is served by route } h, \\ 0 & \text{otherwise.} \end{cases}$
- Variables  $\{y_i^h\}$  are introduced for readability purpose: they correspond to sum over part of variables  $\{x_a\}$ .

## Objective:

$$\min \sum_{h=1}^k \sum_{a \in A} c_a x_a^h$$



# Three-index formulation - Decision variables and objective

## Decision variables:

- $x_a^h = \begin{cases} 1 & \text{if arc } a \text{ belongs to route } h, \\ 0 & \text{otherwise.} \end{cases}$
- $y_i^h = \begin{cases} 1 & \text{if client } i \text{ is served by route } h, \\ 0 & \text{otherwise.} \end{cases}$

→ Variables  $\{y_i^h\}$  are introduced for readability purpose:  
they correspond to sum over part of variables  $\{x_a\}$ .

## Objective:

$$\min \sum_{h=1}^k \sum_{a \in A} c_a x_a^h$$

# Three-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Constraints

## Constraints of LP-relaxation:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in [0, 1], \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in [0, 1], \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$



# Three-index formulation - Constraints

## Constraints of LP-relaxation:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in [0, 1], \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in [0, 1], \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Constraints

## Constraints of LP-relaxation:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in [0, 1], \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in [0, 1], \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Constraints

## Constraints of LP-relaxation with separation:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \in \tilde{S}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in [0, 1], \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in [0, 1], \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Three-index formulation - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{S}$ ). Let  $\bar{x}$ ,  $\bar{y}$  be its **optimal solution**.

## Separation:

- Identify if there exist a route  $h$ , a subset  $\bar{S} \subseteq V \setminus \{0\}$  and a client  $i \in \bar{S}$  such that:

$$\sum_{a \in \delta^+(\bar{S})} \bar{x}_a^h < \bar{y}_i^h$$

- Use **ILP** to model this as  $\mathcal{O}(kn)$  separation problems:
  - by considering each route  $h \in \{1, \dots, k\}$  and each client  $i \in V \setminus \{0\}$  such that  $\bar{y}_i^h > 0$ ,
  - and by searching for  $\bar{S}$  (with  $i \in \bar{S}$ ) as above.

# Three-index formulation - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{S}$ ). Let  $\bar{x}$ ,  $\bar{y}$  be its **optimal solution**.

## Separation:

- Identify **if there exist** a route  $h$ , a subset  $\bar{S} \subseteq V \setminus \{0\}$  and a client  $i \in \bar{S}$  such that:

$$\sum_{a \in \delta^+(\bar{S})} \bar{x}_a^h < \bar{y}_i^h$$

- Use **ILP** to model this as  $\mathcal{O}(kn)$  separation problems:
  - ▶ by considering each route  $h \in \{1, \dots, k\}$  and each client  $i \in V \setminus \{0\}$  such that  $\bar{y}_i^h > 0$ ,
  - ▶ and by searching for  $\bar{S}$  (with  $i \in \bar{S}$ ) as above.

# Three-index formulation - Separation

Consider the CVRP **LP-relaxation** (initialized with a “small” subset of challenging constraints  $\tilde{S}$ ). Let  $\bar{x}$ ,  $\bar{y}$  be its **optimal solution**.

## Separation:

- Identify **if there exist** a route  $h$ , a subset  $\bar{S} \subseteq V \setminus \{0\}$  and a client  $i \in \bar{S}$  such that:

$$\sum_{a \in \delta^+(\bar{S})} \bar{x}_a^h < \bar{y}_i^h$$

- Use **ILP** to model this as  $\mathcal{O}(kn)$  **separation problems**:
  - ▶ by considering each route  $h \in \{1, \dots, k\}$  and each client  $i \in V \setminus \{0\}$  such that  $\bar{y}_i^h > 0$ ,
  - ▶ and by searching for  $\bar{S}$  (with  $i \in \bar{S}$ ) as above.

# Three-index formulation - Separation problem

Let  $h \in \{1, \dots, k\}$  be a route and  $i \in V \setminus \{0\}$  a client such that  $\bar{y}_i^h > 0$ .

Decision variables:

- $w_j = \begin{cases} 1 & \text{if client } i \text{ belongs to } \bar{S}, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(\bar{S}), \\ 0 & \text{otherwise.} \end{cases}$

Objective:

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{a \in \delta^+(\bar{S})} \bar{x}_a^h < \bar{y}_i^h \quad ?$$

↓

$$\max \sum_{a \in A} \bar{x}_a^h z_a < \bar{y}_i^h \quad ?$$

# Three-index formulation - Separation problem

Let  $h \in \{1, \dots, k\}$  be a route and  $i \in V \setminus \{0\}$  a client such that  $\bar{y}_i^h > 0$ .

**Decision variables:**

- $w_j = \begin{cases} 1 & \text{if client } i \text{ belongs to } \bar{S}, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(\bar{S}), \\ 0 & \text{otherwise.} \end{cases}$

**Objective:**

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{a \in \delta^+(\bar{S})} \bar{x}_a^h < \bar{y}_i^h \quad ?$$

↓

$$\max \sum_{a \in A} \bar{x}_a^h z_a < \bar{y}_i^h \quad ?$$



# Three-index formulation - Separation problem

Let  $h \in \{1, \dots, k\}$  be a route and  $i \in V \setminus \{0\}$  a client such that  $\bar{y}_i^h > 0$ .

**Decision variables:**

- $w_j = \begin{cases} 1 & \text{if client } i \text{ belongs to } \bar{S}, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(\bar{S}), \\ 0 & \text{otherwise.} \end{cases}$

**Objective:**

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{a \in \delta^+(\bar{S})} \bar{x}_a^h < \bar{y}_i^h \quad ?$$

↓

$$\max \sum_{a \in A} \bar{x}_a^h z_a < \bar{y}_i^h \quad ?$$

# Three-index formulation - Separation problem

Let  $h \in \{1, \dots, k\}$  be a route and  $i \in V \setminus \{0\}$  a client such that  $\bar{y}_i^h > 0$ .

**Decision variables:**

- $w_j = \begin{cases} 1 & \text{if client } i \text{ belongs to } \bar{S}, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(\bar{S}), \\ 0 & \text{otherwise.} \end{cases}$

**Objective:**

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{a \in \delta^+(\bar{S})} \bar{x}_a^h < \bar{y}_i^h \quad ?$$

$$\max \quad \sum_{a \in A} \bar{x}_a^h z_a < \bar{y}_i^h \quad ?$$

# Three-index formulation - Separation problem

Let  $h \in \{1, \dots, k\}$  be a route and  $i \in V \setminus \{0\}$  a client such that  $\bar{y}_i^h > 0$ .

**Decision variables:**

- $w_j = \begin{cases} 1 & \text{if client } i \text{ belongs to } \bar{S}, \\ 0 & \text{otherwise.} \end{cases}$
- $z_a = \begin{cases} 1 & \text{if arc } a \text{ belongs to } \delta^+(\bar{S}), \\ 0 & \text{otherwise.} \end{cases}$

**Objective:**

$$\exists S^* \subseteq V \setminus \{0\}, \quad \sum_{a \in \delta^+(\bar{S})} \bar{x}_a^h < \bar{y}_i^h \quad ?$$

$$\downarrow$$
$$\max \sum_{a \in A} \bar{x}_a^h z_a < \bar{y}_i^h \quad ?$$

# Three-index formulation - Separation problem

## Separation problem:

$$\begin{aligned}
 \max \quad & \sum_{a \in A} \bar{x}_a^h z_a \\
 \text{s.t.} \quad & w_0 = 0 \\
 & w_i = 1 \\
 & z_{(i,j)} \leq w_i, \quad (i,j) \in A \\
 & z_{(i,j)} \leq 1 - w_j, \quad (i,j) \in A \\
 & z_{(i,j)} \geq w_i - w_j, \quad (i,j) \in A \\
 & w_i \in \{0, 1\}, \quad i \in V \\
 & z_a \in \{0, 1\}, \quad a \in A
 \end{aligned}$$

➔ If the optimal value is smaller than  $\bar{y}_i^h$ ,  
 add  $S^* = \{i \in V \setminus \{0\}, w_i = 1\}$  to  $\tilde{S}$ .

# Three-index formulation - Separation problem

## Separation problem:

$$\begin{aligned}
 \max \quad & \sum_{a \in A} \bar{x}_a^h z_a \\
 \text{s.t.} \quad & w_0 = 0 \\
 & w_i = 1 \\
 & z_{(i,j)} \leq w_i, \quad (i,j) \in A \\
 & z_{(i,j)} \leq 1 - w_j, \quad (i,j) \in A \\
 & z_{(i,j)} \geq w_i - w_j, \quad (i,j) \in A \\
 & w_i \in \{0, 1\}, \quad i \in V \\
 & z_a \in \{0, 1\}, \quad a \in A
 \end{aligned}$$

➔ If the optimal value is smaller than  $\bar{y}_i^h$ ,  
 add  $S^* = \{i \in V \setminus \{0\}, w_i = 1\}$  to  $\tilde{S}$ .

# Three-index formulation - Separation problem

## Separation problem:

$$\begin{aligned}
 \max \quad & \sum_{a \in A} \bar{x}_a^h z_a \\
 \text{s.t.} \quad & w_0 = 0 \\
 & w_i = 1 \\
 & z_{(i,j)} \leq w_i, \quad (i,j) \in A \\
 & z_{(i,j)} \leq 1 - w_j, \quad (i,j) \in A \\
 & z_{(i,j)} \geq w_i - w_j, \quad (i,j) \in A \\
 & w_i \in \{0, 1\}, \quad i \in V \\
 & z_a \in \{0, 1\}, \quad a \in A
 \end{aligned}$$

➔ If the optimal value is smaller than  $\bar{y}_i^h$ ,  
 add  $S^* = \{i \in V \setminus \{0\}, w_i = 1\}$  to  $\tilde{S}$ .

# Three-index formulation - Separation problem

## Separation problem:

$$\begin{aligned}
 \max \quad & \sum_{a \in A} \bar{x}_a^h z_a \\
 \text{s.t.} \quad & w_0 = 0 \\
 & w_i = 1 \\
 & z_{(i,j)} \leq w_i, \quad (i,j) \in A \\
 & z_{(i,j)} \leq 1 - w_j, \quad (i,j) \in A \\
 & z_{(i,j)} \geq w_i - w_j, \quad (i,j) \in A \\
 & w_i \in \{0, 1\}, \quad i \in V \\
 & z_a \in \{0, 1\}, \quad a \in A
 \end{aligned}$$

→ If the optimal value is smaller than  $\bar{y}_i^h$ ,  
**add**  $\mathbf{S}^* = \{i \in V \setminus \{0\}, w_i = 1\}$  to  $\tilde{\mathcal{S}}$ .

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
- 3 Three-index formulation (Section 3.2)
  - Formulation generalities
  - Extension to CVRP with heterogenous capacities
  - Extension to CVRP with time windows
- 4 Conclusion



# CVRP with heterog. cap. - Definition and notations

**Same** as for the basic CVRP with **a few changes**

## Definition:

- **Determining routes** for a fleet of ~~identical~~ **heterogeneous** vehicles, which have ~~the same capacity~~ **different capacities** and are originally located at a central depot, in order **to serve a set of clients** having different demands.
- ...

## Notations:

- $G = (V, A), \{d_j\}, \dots$  ;
- ~~$b$ : vehicle capacity~~  $b_h$ : **capacity** of vehicle  $h$ .

# CVRP with heterog. cap. - Definition and notations

**Same** as for the basic CVRP with **a few changes**

## Definition:

- **Determining routes** for a fleet of ~~identical~~ **heterogeneous** vehicles, which have ~~the same capacity~~ **different capacities** and are originally located at a central depot, in order **to serve a set of clients** having different demands.
- ...

## Notations:

- $G = (V, A), \{d_j\}, \dots$  ;
- ~~$b$ : vehicle capacity~~  $b_h$ : **capacity of vehicle  $h$ .**

# CVRP with heterog. cap. - Definition and notations

**Same** as for the basic CVRP with **a few changes**

## Definition:

- **Determining routes** for a fleet of ~~identical~~ **heterogeneous** vehicles, which have ~~the same capacity~~ **different capacities** and are originally located at a central depot, in order **to serve a set of clients** having different demands.
- ...

## Notations:

- $G = (V, A), \{d_j\}, \dots$  ;
- ~~$b$ : vehicle capacity~~  $b_h$ : **capacity of vehicle  $h$ .**

# CVRP with heterog. cap. - Definition and notations

**Same** as for the basic CVRP with **a few changes**

## Definition:

- **Determining routes** for a fleet of ~~identical~~ **heterogeneous** vehicles, which have ~~the same capacity~~ **different capacities** and are originally located at a central depot, in order **to serve a set of clients** having different demands.
- ...

## Notations:

- $G = (V, A), \{d_j\}, \dots$  ;
- ~~$b$ : vehicle capacity~~  $b_h$ : **capacity of vehicle  $h$** .

# CVRP with heterog. cap. - Decision var. and objective

## Decision variables:

- $x_a^h = \begin{cases} 1 & \text{if arc } a \text{ belongs to route } h, \\ 0 & \text{otherwise.} \end{cases}$
- $y_i^h = \begin{cases} 1 & \text{if client } i \text{ is served by route } h, \\ 0 & \text{otherwise.} \end{cases}$

## Objective:

$$\min \sum_{h=1}^k \sum_{a \in A} c_a x_a^h$$

# CVRP with heterog. cap. - Constraints

## Constraints:

$$\sum_{a \in \delta^-(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at client node})$$

$$\sum_{a \in \delta^+(i)} x_a^h = y_i^h, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{h=1}^k y_i^h = 1, \quad i \in V \setminus \{0\} \quad (\text{Each client visited once})$$

$$\sum_{a \in \delta^-(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\} \quad (\text{Flow at depot})$$

$$\sum_{a \in \delta^+(0)} x_a^h = 1, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

$$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b^h, \quad h \in \{1, \dots, k\} \quad (\text{Capacity})$$

$$\sum_{a \in \delta^+(S)} x_a^h \leq y_i^h, \quad h \in \{1, \dots, k\}, S \subseteq V \setminus \{0\}, i \in S \quad (\text{Subtour elimination})$$

$$x_a^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, a \in A$$

$$y_i^h \in \{0, 1\}, \quad h \in \{1, \dots, k\}, i \in V \setminus \{0\}$$

# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
- 3 Three-index formulation (Section 3.2)
  - Formulation generalities
  - Extension to CVRP with heterogenous capacities
  - Extension to CVRP with time windows
- 4 Conclusion

# CVRP with time windows - Definition

Same as for the basic CVRP with **additional considerations**

## Definition:

- **Determining routes** for a fleet of identical vehicles, which have the same capacity and are originally located at a central depot, in order **to serve a set of clients** having different demands and different visiting time windows.
- Routes are such that:
  - ▶ each route **starts at the depot at time 0**, goes from client to client **without idle time**, and **ends at the depot**;
  - ▶ each client is **served by exactly one vehicle**, during their **time window**, given that the service time is null;
  - ▶ the total demand served by each vehicle **does not exceed its capacity**.
- ...



# CVRP with time windows - Definition

Same as for the basic CVRP with **additional considerations**

## Definition:

- **Determining routes** for a fleet of identical vehicles, which have the same capacity and are originally located at a central depot, in order **to serve a set of clients** having different demands and different visiting time windows.
- Routes are such that:
  - ▶ each route **starts at the depot at time 0**, goes from client to client **without idle time**, and **ends at the depot**;
  - ▶ each client is **served by exactly one vehicle**, during their **time window**, given that the service time is null;
  - ▶ the total demand served by each vehicle **does not exceed its capacity**.
- ...

# CVRP with time windows - Definition

Same as for the basic CVRP with **additional considerations**

## Definition:

- **Determining routes** for a fleet of identical vehicles, which have the same capacity and are originally located at a central depot, in order **to serve a set of clients** having different demands and different visiting time windows.
- Routes are such that:
  - ▶ each route **starts at the depot at time 0**, goes from client to client **without idle time**, and **ends at the depot**;
  - ▶ each client is **served by exactly one vehicle**, during their **time window**, given that the service time is null;
  - ▶ the total demand served by each vehicle **does not exceed its capacity**.

# CVRP with time windows - Definition

Same as for the basic CVRP with **additional considerations**

## Definition:

- **Determining routes** for a fleet of identical vehicles, which have the same capacity and are originally located at a central depot, in order **to serve a set of clients** having different demands and different visiting time windows.
- Routes are such that:
  - ▶ each route **starts at the depot at time 0**, goes from client to client **without idle time**, and **ends at the depot**;
  - ▶ each client is **served by exactly one vehicle**, during their **time window**, given that the service time is null;
  - ▶ the total demand served by each vehicle **does not exceed its capacity**.
- ...

# CVRP with time windows - Notations and assumptions

## Notations:

- $G = (V, A), \{d_j\}, \dots$  ;
- $[e_i, l_i]$ : **time window** of client  $i$ .

## Assumptions:

- $G$  complete graph and triangle inequalities hold.
- $c_a$  is not only the cost of the arc  $a$  but also the **travel time** (and the time spent between two nodes must be exactly  $c_a$ ).

# CVRP with time windows - Notations and assumptions

## Notations:

- $G = (V, A), \{d_j\}, \dots$  ;
- $[e_i, l_i]$ : **time window** of client  $i$ .

## Assumptions:

- $G$  complete graph and triangle inequalities hold.
- $c_a$  is not only the cost of the arc  $a$  but also the **travel time** (and the time spent between two nodes must be exactly  $c_a$ ).

# CVRP with time windows - Notations and assumptions

## Notations:

- $G = (V, A), \{d_j\}, \dots$  ;
- $[e_i, l_i]$ : **time window** of client  $i$ .

## Assumptions:

- $G$  complete graph and triangle inequalities hold.
- $c_a$  is not only the cost of the arc  $a$  but also the **travel time** (and the time spent between two nodes must be exactly  $c_a$ ).

# CVRP with time windows - Notations and assumptions

## Notations:

- $G = (V, A), \{d_j\}, \dots$  ;
- $[e_i, l_i]$ : **time window** of client  $i$ .

## Assumptions:

- $G$  complete graph and triangle inequalities hold.
- $c_a$  is not only the cost of the arc  $a$  but also the **travel time** (and the time spent between two nodes must be exactly  $c_a$ ).

# CVRP with time windows - Decision var. and objective

## Decision variables:

- $\{x_a^h\}$  and  $\{y_i^h\}$
- $s_i^h \in \mathbb{R}^+$ : time at which client  $i$  is served, if served by vehicle  $h$ ,  
( $s_i^h$  equals 0 if not served by vehicle  $h$ )

## Objective:

$$\min \sum_{h=1}^k \sum_{a \in A} c_a x_a^h$$



# CVRP with time windows - Decision var. and objective

## Decision variables:

- $\{x_a^h\}$  and  $\{y_i^h\}$
- $s_i^h \in \mathbb{R}^+$ : time at which client  $i$  is served, if served by vehicle  $h$ ,  
( $s_i^h$  equals 0 if not served by vehicle  $h$ )

## Objective:

$$\min \sum_{h=1}^k \sum_{a \in A} c_a x_a^h$$

# CVRP with time windows - Decision var. and objective

## Decision variables:

- $\{x_a^h\}$  and  $\{y_i^h\}$
- $s_i^h \in \mathbb{R}^+$ : time at which client  $i$  is served, if served by vehicle  $h$ ,  
( $s_i^h$  equals 0 if not served by vehicle  $h$ )

## Objective:

$$\min \sum_{h=1}^k \sum_{a \in A} c_a x_a^h$$

# CVRP with time windows - Travel time constraints

## Travel time constraints:

- **Logical (non-linear)** formulation:

$$x_{(i,j)}^h = 1 \Rightarrow s_j^h = s_i^h + c_{(i,j)}$$

- **Linear** formulation using “**big-M**” techniques:

$$\begin{aligned} s_j^h &\geq s_i^h + c_{(i,j)} - M(1 - x_{(i,j)}^h) \\ s_j^h &\leq s_i^h + c_{(i,j)} + M(1 - x_{(i,j)}^h) \end{aligned}$$

with  $M$  chosen “sufficiently” large.

- ⚠ Such big-M formulations often cause **weak LP relaxation**:

- ▶ if possible, **try to avoid** big-M formulation;
- ▶ if not, use **smallest possible** (but still “safe”) values of  $M$ .

# CVRP with time windows - Travel time constraints

## Travel time constraints:

- **Logical (non-linear)** formulation:

$$x_{(i,j)}^h = 1 \Rightarrow s_j^h = s_i^h + c_{(i,j)}$$

- **Linear** formulation using “**big-M**” techniques:

$$\begin{aligned} s_j^h &\geq s_i^h + c_{(i,j)} - M(1 - x_{(i,j)}^h) \\ s_j^h &\leq s_i^h + c_{(i,j)} + M(1 - x_{(i,j)}^h) \end{aligned}$$

with  $M$  chosen “sufficiently” large.

- ⚠ Such big-M formulations often cause **weak LP relaxation**:

- ▶ if possible, **try to avoid** big-M formulation;
- ▶ if not, use **smallest possible** (but still “safe”) values of  $M$ .

# CVRP with time windows - Travel time constraints

## Travel time constraints:

- **Logical (non-linear)** formulation:

$$x_{(i,j)}^h = 1 \Rightarrow s_j^h = s_i^h + c_{(i,j)}$$

- **Linear** formulation using “**big-M**” techniques:

$$\begin{aligned} s_j^h &\geq s_i^h + c_{(i,j)} - M(1 - x_{(i,j)}^h) \\ s_j^h &\leq s_i^h + c_{(i,j)} + M(1 - x_{(i,j)}^h) \end{aligned}$$

with  $M$  chosen “sufficiently” large.

⚠ Such big-M formulations often cause **weak LP relaxation**:

- ▶ if possible, **try to avoid** big-M formulation;
- ▶ if not, use **smallest possible** (but still “safe”) values of  $M$ .

# CVRP with time windows - Travel time constraints

## Travel time constraints:

- **Logical (non-linear)** formulation:

$$x_{(i,j)}^h = 1 \Rightarrow s_j^h = s_i^h + c_{(i,j)}$$

- **Linear** formulation using “**big-M**” techniques:

$$\begin{aligned} s_j^h &\geq s_i^h + c_{(i,j)} - M(1 - x_{(i,j)}^h) \\ s_j^h &\leq s_i^h + c_{(i,j)} + M(1 - x_{(i,j)}^h) \end{aligned}$$

with  $M$  chosen “sufficiently” large.

- ⚠ Such big-M formulations often cause **weak LP relaxation**:

- ▶ if possible, **try to avoid** big-M formulation;
- ▶ if not, use **smallest possible** (but still “safe”) values of  $M$ .

# CVRP with time windows - Travel time constraints

## Travel time constraints:

- **Logical (non-linear)** formulation:

$$x_{(i,j)}^h = 1 \Rightarrow s_j^h = s_i^h + c_{(i,j)}$$

- **Linear** formulation using “**big-M**” techniques:

$$\begin{aligned} s_j^h &\geq s_i^h + c_{(i,j)} - M(1 - x_{(i,j)}^h) \\ s_j^h &\leq s_i^h + c_{(i,j)} + M(1 - x_{(i,j)}^h) \end{aligned}$$

with  $M$  chosen “sufficiently” large.

- ⚠ Such big-M formulations often cause **weak LP relaxation**:

- ▶ if possible, **try to avoid** big-M formulation;
- ▶ if not, use **smallest possible** (but still “safe”) values of  $M$ .

# CVRP with time windows - Constraints

## Constraints:

$\sum_{a \in \delta^-(i)} x_a^h = y_i^h,$	$h \in \{1, \dots, k\}, i \in V \setminus \{0\}$	(Flow at client)
$\vdots$	$\vdots$	$\vdots$
$\sum_{i \in V \setminus \{0\}} d_i y_i^h \leq b,$	$h \in \{1, \dots, k\}$	(Capacity)
$s_0^h = 0,$	$h \in \{1, \dots, k\}$	(Time 0)
$s_i^h \geq e_i y_i^h,$	$h \in \{1, \dots, k\}, i \in V \setminus \{0\}$	(Time window)
$s_i^h \leq l_i y_i^h,$	$h \in \{1, \dots, k\}, i \in V \setminus \{0\}$	
$s_j^h \geq s_i^h + c_{(i,j)} - M(1 - x_{(i,j)}^h),$	$h \in \{1, \dots, k\}, i \in V, j \in V \setminus \{0\}$	(Travel time)
$s_j^h \leq s_i^h + c_{(i,j)} + M(1 - x_{(i,j)}^h),$	$h \in \{1, \dots, k\}, i \in V, j \in V \setminus \{0\}$	
$x_a^h \in \{0, 1\},$	$h \in \{1, \dots, k\}, a \in A$	
$y_i^h \in \{0, 1\},$	$h \in \{1, \dots, k\}, i \in V \setminus \{0\}$	
$s_j^h \geq 0,$	$h \in \{1, \dots, k\}, i \in V$	



# Plan

- 1 Introduction to CVRP
- 2 Two-index formulation (Section 3.1)
- 3 Three-index formulation (Section 3.2)
- 4 Conclusion

# Conclusion

- The **CVRP** serves as a **core model** for routing problems, commonly applied to freight transportation. Despite being a basic problem, it already presents the **combined complexities of standard routing (TSP) and loading (BPP) problems**, leading to the use of **separation** techniques, or other sophisticated techniques.
- We studied two different formulations:
  - ▶ the **two-index formulation** that is the **simplest** one;
  - ▶ the **three-index formulation** that is **easier to adapt** for incorporating additional characteristics (e.g. time windows or heterogeneous capacities).
- Another formulation to study: the **set partitioning formulation** that leverages **column generation** techniques.

# Conclusion

- The **CVRP** serves as a **core model** for routing problems, commonly applied to freight transportation. Despite being a basic problem, it already presents the **combined complexities of standard routing (TSP) and loading (BPP) problems**, leading to the use of **separation** techniques, or other sophisticated techniques.
- We studied two different formulations:
  - ▶ the **two-index formulation** that is the **simplest** one;
  - ▶ the **three-index formulation** that is **easier to adapt** for incorporating additional characteristics (e.g. time windows or heterogeneous capacities).
- Another formulation to study: the **set partitioning formulation** that leverages **column generation** techniques.

# Conclusion

- The **CVRP** serves as a **core model** for routing problems, commonly applied to freight transportation. Despite being a basic problem, it already presents the **combined complexities of standard routing (TSP) and loading (BPP) problems**, leading to the use of **separation** techniques, or other sophisticated techniques.
- We studied two different formulations:
  - ▶ the **two-index formulation** that is the **simplest** one;
  - ▶ the **three-index formulation** that is **easier to adapt** for incorporating additional characteristics (e.g. time windows or heterogeneous capacities).
- Another formulation to study: the **set partitioning formulation** that leverages **column generation** techniques.