

# Local Gap Colorings from Edge Labelings

Axel Brandt\*, Brent Moran†, Kapil Nepal‡, Florian Pfender§, Devon Sigler¶

## Abstract

We study a local version of gap vertex–distinguishing edge coloring. From an edge labeling  $f: E \rightarrow \{1, \dots, k\}$  of a graph  $G$ , an induced vertex coloring  $c$  is obtained by coloring the vertices with the greatest difference between incident edge labels. The local gap chromatic number  $\chi_{\Delta}^e(G)$  is the minimum  $k$  for which there exists an edge coloring such that  $c(u) \neq c(v)$  for all edges  $uv$ . We prove that  $\chi(G) \leq \chi_{\Delta}^e(G) \leq \chi(G) + 1$  for all graphs  $G$ , where  $\chi(G)$  denotes the chromatic number of  $G$ . Further we find graph classes attaining both bounds.

## 1 Introduction

Unless otherwise stated, a graph  $G$  is simple, finite, and undirected with no isolated vertex. Standard graph theory notation ([13]) is used throughout.

Derived graph colorings, typically obtained from graph labelings, have been widely studied. In 1988, Chartrand et al. [2] introduced the *irregularity strength* of a graph  $G$ , which is the smallest positive integer  $k$  such that each edge can have a label from  $[k] := \{1, \dots, k\}$  so that the sum of labels of edges incident to each vertex is distinct. This topic was further studied in [1, 3, 8], among others. In 2008, Gyori et al. [4] introduced a variation that seeks the smallest positive integer  $k$  such that each edge can have a label from  $[k]$  so that the sets of the weights on edges incident to vertices are distinct.

In addition to the global constraints described above, local constraints have also been studied. The most active problem in this area, the 1–2–3 Conjecture, was proposed in 2004 by Karoński, Łuczak, and Thomason [6]. For a graph  $G$ , let  $\chi_{\Sigma}^e(G)$  be the smallest positive integer  $k$  such that  $G$  has a labeling  $\ell: E(G) \rightarrow [k]$  such that, for every edge  $uv \in E(G)$ ,  $\sum_{e \ni u} \ell(e) \neq \sum_{e \ni v} \ell(e)$ .

**1–2–3 Conjecture** (Karoński, Łuczak, Thomason [6]). *If  $G$  has no component isomorphic to  $K_2$ , then  $\chi_{\Sigma}^e(G) \leq 3$ .*

Under the same assumptions, Kalkowski, Karoński, Pfender [5] showed that  $\chi_{\Sigma}^e(G) \leq 5$ . For a survey of work on the 1–2–3 Conjecture and derived colorings, we direct the reader to [11].

---

\*Dept. of Math. and Stat. Sciences, University of Colorado Denver, Axel.Brandt@ucdenver.edu, partially funded by NSF GK-12 Transforming Experiences Grant DGE-0742434

†Dept. of Math. and Stat. Sciences, University of Colorado Denver, Brent.Moran@ucdenver.edu

‡Dept. of Math. and Stat. Sciences, University of Colorado Denver, Kapil.Nepal@ucdenver.edu

§Dept. of Math. and Stat. Sciences, University of Colorado Denver, Florian.Pfender@ucdenver.edu, partially funded by Simons Foundation grant #276726

¶Dept. of Math. and Stat. Sciences, University of Colorado Denver, Devon.Sigler@ucdenver.edu

In this paper we are interested in a particular derived vertex coloring called *gap vertex-distinguishing edge coloring*. Here the derived coloring  $c_\ell$  of a vertex is

$$g_\ell(v) = \begin{cases} \ell(e)_{e \ni v} & \text{if } d(v) = 1 \\ \max_{e \ni v} \ell(e) - \min_{e \ni v} \ell(e) & \text{otherwise,} \end{cases}$$

where  $\ell : E(G) \rightarrow [k]$ . An edge coloring is called *gap vertex distinguishing* when all vertices have distinct colors. The minimum  $k$  such that a gap vertex-distinguishing edge coloring exists is called the *gap chromatic number* of  $G$  and denoted  $\text{gap}(G)$ . Introduced by Tahraoui, Duchêne, and Kheddouci in [12], they conjectured that  $\text{gap}(G) \leq n(G) + 1$ . In [10], Scheidweiler and Triesch prove this conjecture for connected graphs, but disprove it in general by finding a class of graphs with  $\text{gap}(G) = n(G) + 2$ .

Tahraoui et al. [12] introduce a similar derived coloring that distinguished adjacent vertices. The *gap-adjacent-chromatic number* of a graph  $G$ ,  $\text{gap}_{\text{ad}}(G)$ , is the minimum  $k$  for which a labeling  $\ell : E(G) \rightarrow [k]$  exists so that  $g_\ell$  induces a proper vertex coloring. Scheidweiler and Triesch [9] prove the following:

**Theorem 1** ([9]). *If  $\chi(G) \in \{2, 3\}$ , then  $\text{gap}_{\text{ad}}(G) \leq \chi(G) + 1$ .*

**Theorem 2** ([9]). *If  $G$  is a graph without isolated edges, then  $\chi(G) - 1 \leq \text{gap}_{\text{ad}}(G) \leq \chi(G) + 5$ .*

In this paper, we improve Theorem 2 by giving sharp bounds.

**Theorem 3.** *If  $G$  is a graph without isolated edges, then  $\chi(G) \leq \text{gap}_{\text{ad}}(G) \leq \chi(G) + 1$  unless  $G$  is a star, in which case  $\text{gap}_{\text{ad}}(G) = 1 = \chi(G) - 1$ .*

In order to define  $\text{gap}(G)$  for graphs with more than one leaf, the special treatment of leaves in the definition of  $g_\ell$  is necessary. In the local version, leaves are permitted to have the same color. As such, it is natural to consider the following simpler definition for the gap color of vertices

$$c_\ell(v) = \max_{e \ni v} \ell(e) - \min_{e \ni v} \ell(e),$$

where  $\ell : E(G) \rightarrow [k]$  and leaves do not receive special treatment. An edge coloring is a *local gap  $k$ -coloring* when adjacent vertices have distinct colors. The minimum  $k$  for which a local gap  $k$ -coloring exists is called the *local gap chromatic number* of  $G$  and denoted by  $\chi_\Delta^e(G)$ . In this paper we prove the following:

**Theorem 4.** *If  $G$  has no isolated edges, then  $\chi_\Delta^e(G) \in \{\chi(G), \chi(G) + 1\}$ .*

Despite the difference between  $c_\ell$  and  $g_\ell$ , we are able to use Theorem 4 to prove Theorem 3. All bounds in Theorem 3 and 4 are sharp. as we will show in the following section.

## 2 Sharpness Examples for Theorems 3 and 4

Assigning edge labels from  $[k]$  allows for  $k$  vertex colors under  $c_\ell$ , namely  $0, \dots, k - 1$ . Therefore  $\chi_\Delta^e(G) \geq \chi(G)$ . Equality is achieved by many graphs, some of which we discuss in Section 5. A similar argument implies  $\text{gap}_{\text{ad}}(G) \geq \chi(G) - 1$ . However not all graphs have  $\chi_\Delta^e(G) = \chi(G)$ . For

example, consider a complete graph with a pendant edge added to each vertex. It is easy to see that  $\chi_\Delta^e(G) = \chi(G) + 1$  and  $\text{gap}_{\text{ad}}(G) = \chi(G)$ .

For an example with  $\chi_\Delta^e(G) = \text{gap}_{\text{ad}}(G) = \chi(G) + 1$ , consider the following graph. Let  $s$  and  $r$  be positive integers such that  $s \geq r + 1 \geq 3$ , and let  $K_s^r$  be the complete  $r$ -partite graph with  $s$  vertices in each partite set  $X_1, \dots, X_r$ . For all  $i$  and all  $u, v \in X_i$ , add a new vertex that is adjacent to  $u$  and  $v$  and call this graph  $G$ . Since  $G$  has no leaves,  $\chi_\Delta^e(G) = \text{gap}_{\text{ad}}(G)$ . We claim that  $\chi_\Delta^e(G) = r + 1 = \chi(G) + 1$ .

To see this, suppose that  $\chi_\Delta^e(G) = r$ . Let  $c_\ell$  be a local gap  $r$ -coloring of  $G$ . Since  $K_s^r \subseteq G$  and there exists exactly one partition of  $K_s^r$  into  $r$  independent sets, there is some partite set of  $K_s^r$ , say  $X_1$ , such that  $c_\ell(v) = 0$  for all  $v \in X_1$ . For every  $v \in X_1$ , all edges incident to  $v$  have the same label. This partitions  $X_1$  into at most  $r$  classes depending on the label of the incident edges. Since  $s \geq r + 1$ , there are two vertices  $u, v \in X_1$  that have the same label on all incident edges. Thus the vertex  $x$  outside  $K_s^r$  adjacent to  $u$  and  $v$  has  $c_\ell(x) = 0$ , a contradiction.

One can generalize this construction by taking a large enough blow-up of any graph  $G$  and joining new vertices to every  $t$ -tuple from the independent sets of the blow-up corresponding to vertices of  $G$ .

### 3 An Upper Bound for $\chi_\Delta^e(G)$

We turn our attention to proving an upper bound for  $\chi_\Delta^e(G)$ . For this purpose, we define the following sets based on distance to any  $X \subseteq V(G)$  (see Figure 1):

$$\begin{aligned} V_i(X) &= \{x \in V(G) : d(x, X) = i\}, \\ U_i(X) &= \{x \in V_i(X) : N(x) \subseteq V_{i-1}(X)\}, \\ E_i(X) &= \{xy \in E(G) : x \in V_{i-1}(X), y \in V_i(X)\}, \\ F_i(X) &= \{xy \in E_i(G) : y \in U_i(X)\}, \end{aligned}$$

where  $i \in \mathbb{N}$ ,  $V_0 = U_0 = X$ . For  $i \geq 2$ , let

$$\begin{aligned} V'_{i-1}(X) &= \{x \in V_{i-1}(X) : \exists xy \in F_i(X)\} \\ F'_{i-1}(X) &= \{wx \in E_{i-1}(G) : x \in V'_{i-1}(X)\}. \end{aligned}$$




Figure 1: Sets defined for proofs.

For brevity we write  $v$  when  $X = \{v\}$  and say that  $v$  has *gap color*  $m$  when  $c_\ell(v) = m$ . To prove Theorem 4 we cover three cases based on chromatic number in the following lemmas.

**Lemma 1.** *Let  $G$  be a connected bipartite graph not isomorphic to  $K_2$ . Then  $\chi_\Delta^e(G) \leq 3$ .*

*Proof.* Let  $v$  be a vertex in  $G$  that is not adjacent to a leaf. Define an edge labeling  $\ell$  as follows for edge  $e \in E_i(v)$  (see Figure 2):

$$\ell(e) = \begin{cases} 3 & \text{if } i \equiv 1 \pmod{4}, \text{ or both } i \equiv 0 \pmod{4} \text{ and } e \in F_i(v), \\ 1 & \text{if } i \equiv 2 \pmod{4}, \text{ or both } i \equiv 3 \pmod{4} \text{ and } e \in F'_i(v), \\ 2 & \text{otherwise.} \end{cases}$$




Figure 2: An edge labeling for connected bipartite graphs including derived gap colors.

Notice that each  $V_i(v)$  is an independent set since  $G$  is bipartite. When  $i$  is even, the gap color of vertices in  $V_i(v)$  is either 0 or 1. When  $i$  is odd, the gap color of vertices in  $V_i(v)$  is either 0 or 2. Toward  $\ell$  being a local gap 3-coloring, we show vertices with gap color 0 are independent. Since  $v$  is not adjacent to a leaf,  $c_\ell(x) = 2$  for all  $x \in V_1(v)$ . Notice that every vertex in some  $U_i(v)$  has gap color 0, and all of their neighbors have nonzero gap color. When  $i \equiv 2 \pmod{4}$ , every neighbor of a vertex in  $V_i$  with gap color 0 has gap color 2. Vertices in  $V'_i(v)$  when  $i \equiv 3 \pmod{4}$  have gap color 2 and are adjacent to vertices whose gap color is either 0 or 1. Thus,  $\ell$  is a local gap 3-coloring.  $\square$

**Lemma 2.** *Let  $G$  be a connected tripartite graph. Then  $\chi_\Delta^e(G) \leq 4$ .*

*Proof.* Let  $C_1, C_2, C_3$  be the color classes of a proper coloring of  $V(G)$  in which every vertex in  $C_i$  has a neighbor in  $C_j$  for  $1 \leq j < i \leq 3$ . Let  $Y_1 = \{v \in C_1 : N(v) \subseteq C_2\}$ ,  $Y_2 = C_1 \setminus Y_1$ ,  $X_1 = \{v \in C_2 : N(v) \subseteq C_1\}$ , and  $X_2 = C_2 \setminus X_1$ . We refine  $Y_1$  and  $X_1$  by defining  $Y_{11} = \{v \in Y_1 : N(v) \subseteq X_1\}$ ,  $Y_{12} = Y_1 \setminus Y_{11}$ ,  $X_{11} = \{v \in X_1 : N(v) \subseteq Y_1\}$ , and  $X_{12} = X_1 \setminus X_{11}$ . Define a partial edge labeling  $\ell$  as follows (see Figure 3).

$$\ell(uv) = \begin{cases} 4 & \text{if } u \in C_3 \text{ and } v \in C_1, \\ 3 & \text{if } u \in C_3 \text{ and } v \in C_2, \\ 1 & \text{if } u \in X_2 \text{ and } v \in Y_2. \end{cases}$$

Notice that vertices in  $C_3$ ,  $X_2$ , and  $Y_2$  have gap colors 1, 2, and 3 respectively. Also notice that the remaining edges all live in a bipartite graph. We attempt to mimic Lemma 1 without changing the existing gap colors in  $C_3$ ,  $X_2$ , and  $Y_2$ . We examine two completions of this coloring dependent upon  $Y'$ .

First assume that  $Y_{12} = \emptyset$ . Since  $G$  is connected,  $X_{12} \neq \emptyset$ . For all leaves  $u \in X_{12}$  incident to some  $v$ , define  $\ell(uv) = 1$ . Complete the partial edge labeling for all remaining edges  $uv \in E_i(Y_2)$




Figure 3: A partial edge labeling for connected tripartite graphs including derived gap colors.

with  $u \in C_2$  and  $v \in C_1$  as follows (see Figure 4).

$$\ell(uv) = \begin{cases} 4 & \text{if } i \equiv 1 \pmod{4}, \\ 2 & \text{if } i \equiv 2 \pmod{4}, \\ 3 & \text{if } i \equiv 3 \pmod{4}, \\ 1 & \text{if } i \equiv 0 \pmod{4}. \end{cases}$$




Figure 4: Completing the partial tripartite labeling from Figure 3 when  $Y_{12} = \emptyset$ .

The gap color of vertices in  $X_1$  is 0 or 2, and the gap color of vertices in  $Y_1$  is 0 or 1. Hence, the vertices in  $C_2$  have gap color 0 or 2, and vertices in  $C_1$  have gap color 0, 1, or 3. Again we must show the vertices with gap color 0 are independent. Notice that if a vertex in  $(X_1 \cup Y_1) \cap V_i(Y_2)$  has a neighbor in  $(X_1 \cup Y_1) \cap V_{i+1}(Y_2)$  then it has nonzero gap color. Let  $x \in (X \cup Y) \cap V_i(Y_2)$  with  $c_\ell(x) = 0$ . Then every neighbor of  $x$  is in  $V_{i-1}(Y_2)$  and has a neighbor in  $V_i(Y_2)$ . Hence every neighbor of  $x$  has nonzero gap color. Therefore  $\ell$  is a local gap 4-coloring.

Otherwise,  $Y_{12} \neq \emptyset$ . Complete the edge labeling for all remaining edges  $uv \in E_i(X_2)$  with

$u \in C_2$  and  $v \in C_1$  as follows (see Figure 5).

$$\ell(uv) = \begin{cases} 1 & \text{if } i \equiv 1 \pmod{4}, u \in X_{12}, \text{ and } v \in Y_2, \\ 3 & \text{if } i \equiv 1 \pmod{4}, \text{ and } u \in C_2 \setminus X_{12} \text{ or } v \in Y_1, \\ 2 & \text{if } i \equiv 2 \pmod{4}, \\ 4 & \text{if } i \equiv 3 \pmod{4}, \\ 1 & \text{if } i \equiv 0 \pmod{4}. \end{cases}$$




Figure 5: Completing the partial tripartite labeling from Figure 3 when  $Y_{12} \neq \emptyset$ .

The gap color of vertices in  $X_1$  is either 0 or 2, and the gap color of vertices in  $Y_1$  are 0, 1, or 3. As before, the neighbors of any vertex with gap color 0 in  $X_1 \cup Y_1$  have nonzero gap color. Thus,  $\ell$  is a local gap 4-coloring.  $\square$

**Lemma 3.** *Let  $G$  be a connected graph with  $\chi(G) \geq 4$ . Then  $\chi_{\Delta}^e(G) \leq \chi(G) + 1$ .*

*Proof.* Let  $\chi = \chi(G)$ , and let  $C_1, \dots, C_{\chi}$  be the color classes of a proper coloring on  $V(G)$  in which every vertex in  $C_i$  has a neighbor in  $C_j$  for  $1 \leq j < i \leq \chi$ . Let  $X_{11}, X_{12}, X_2, Y_{11}, Y_{12}$ , and  $Y_2$  be defined as in the proof of Lemma 2. Define a partial edge labeling for  $uv \in E(G)$  as follows (see Figure 6).

$$\ell(uv) = \begin{cases} \chi - i + 2 & \text{if } u \in C_i \text{ and } v \in C_{i-1} \text{ for some } i = 4, \dots, \chi \\ 1 & \text{if } u \in C_2 \text{ and } v \in C_i \text{ for some } i = 4, \dots, \chi \\ 2 & \text{if } u \in C_i \text{ and } v \in C_j \text{ for } 4 \leq i < j - 1 \leq \chi - 1 \\ \chi & \text{if } u \in C_3 \text{ and } v \in C_2 \\ 2 & \text{if } u \in C_3 \text{ and } v \in C_1 \\ \chi + 1 & \text{if } v \in Y_2 \end{cases}$$

Notice that for  $i = 3, \dots, \chi$  vertices in  $C_i$  have gap color  $\chi - i + 1$ , vertices in  $Y_2$  with a neighbor in  $C_2$  have gap color  $\chi - 1$ , and vertices in  $Y_2$  with no neighbor in  $C_2$  have gap color 0. As in Lemma 2, every remaining edge lives in a bipartite graph and we attempt to mimic Lemma 1 without changing the existing gap colors. We further refine  $X_2$  and  $Y_{12}$  by defining  $Y_{121} = \{v \in Y_{12} : N(v) \subseteq X_2\}$ ,  $X_{21} = \{v \in X_2 : N(v) \subseteq C_3 \cup (Y_{12} \setminus Y_{121})\}$ ,  $Y_{122} = \{v \in Y_{12} \setminus Y_{121} : N(v) \cap X_{21} \neq \emptyset, N(v) \cap X_{12} \neq \emptyset\}$ ,




Figure 6: A partial edge labeling of a  $\chi$  partite graph with  $\chi \geq 4$ .

$X_{22} = X_2 \setminus X_{21}$ , and  $Y_{123} = Y_{12} \setminus \{Y_{121}, Y_{122}\}$ . Label edges  $uv \in E(G)$  as follows:

$$\ell(uv) = \begin{cases} \chi + 1 & \text{if } u \in Y_{121}, \text{ or } u \in Y_{122} \text{ and } v \in X_1, \\ \chi - 1 & \text{if } u \in Y_{122} \text{ and } v \in X_{21}. \end{cases}$$

Let  $X'$  be the set of vertices in  $C_2$  that have an incident edge already labeled. Complete the partial edge labeling for all remaining edges  $uv \in E_i(X')$  with  $u \in C_2$  and  $v \in C_1$  as follows (see Figure 7).

$$\ell(uv) = \begin{cases} \chi & \text{if } i \equiv 1 \pmod{4}, \\ \chi - 2 & \text{if } i \equiv 2 \pmod{4}, \\ \chi - 3 & \text{if } i \equiv 3 \pmod{4}, \\ \chi - 1 & \text{if } i \equiv 0 \pmod{4}. \end{cases}$$




Figure 7: Completing the partial edge labeling from Figure 6.

Every edge is labeled under this labelling even if  $Y_{12} = \emptyset$  since  $G$  is connected. Notice that vertices in  $X_2$  adjacent to  $Y_{121} \cup Y_2$  have gap color 1 or  $\chi$ , and the vertices with gap color 1 in  $X_2$  have neighbors strictly in  $C_1 \cup C_3$ . Since vertices in  $X_2$  with gap color 0 don't have neighbors in  $Y_{121}$ , vertices with gap color 0 in  $X_1 \cup Y_1$  are independent. Notice that vertices in  $Y_{122}$  may have an incident edge labeled  $\chi$  from  $X_{22}$  but do not have an incident edge labeled  $\chi - 2$ . Thus vertices in  $Y_{122}$  have gap color 2. Vertices with gap color 0 in  $X_{12}$  have neighborhoods strictly in  $Y_2 \cup Y_{122}$  and thus are independent from vertices with gap color 0. The remaining parity of colors guarantees  $\ell$  is a local gap  $(\chi + 1)$ -coloring.  $\square$

*Proof of Theorem 4.* Apply Lemmas 1, 2, and 3 to appropriate components of  $G$ .  $\square$

## 4 Improved Bounds for $\text{gap}_{\text{ad}}(G)$

Recall that for a graph  $G$ ,  $g_\ell$  is the gap coloring associated with  $\text{gap}_{\text{ad}}(G)$ . In this section, we improve the bounds on the gap-adjacent-chromatic number of graphs. We begin by showing  $\text{gap}_{\text{ad}}(G) \geq \chi(G)$  when  $G$  is not a star, and then provide a sharp upper bound for all graphs.

**Lemma 4.** *Let  $G$  be a connected graph. Then  $\text{gap}_{\text{ad}}(G) = \chi(G) - 1$  if and only if  $G$  is a star.*

*Proof.* If  $G$  is a star, then  $\ell: E(G) \rightarrow \{1\}$  gives  $\text{gap}_{\text{ad}}(G) = \chi(G) - 1$ . Assume  $G$  is not a star. Let  $L$  be the leaves of  $G$  and  $\ell: E(G) \rightarrow [k]$ . Since a proper coloring of  $G - L$  can be extended to  $G$  without using an additional color,  $\chi(G - L) = \chi(G)$ . From the definition,  $g_\ell(v) \in \{0, \dots, k - 1\}$  for all  $v \in V(G - L)$ . Thus, in order to properly color  $G - L$ ,  $k \geq \chi(G - L) = \chi(G)$ .  $\square$

**Lemma 5.** *For any connected graph  $G$  not isomorphic to  $K_2$ ,  $\text{gap}_{\text{ad}}(G) \leq \chi(G) + 1$ .*

*Proof.* Theorem 1 gives the desired result when  $\chi(G) \in \{2, 3\}$ . Thus we may assume  $\chi(G) \geq 4$  and, in particular, that  $G$  is not a path. If  $G$  has no leaves, then  $\text{gap}_{\text{ad}}(G) = \chi_\Delta^e(G)$  since the two colorings associated with these parameters differ only on leaves. Lemma 4 completes the base case. We proceed by induction on the number of leaves.

Assume  $\text{gap}_{\text{ad}}(H) \leq \chi(H) + 1$  for all graphs  $H$  with  $k$  leaves and let  $G$  be a graph with  $k + 1$  leaves. Let  $P = v_0 v_1 \dots v_p u$  be a minimum length path in  $G$  with  $d(v_0) = 1$  and  $d(u) \geq 3$ . Let  $G' = G - \{v_0, \dots, v_p\}$ . By the induction hypothesis,  $\text{gap}_{\text{ad}}(G') \leq \chi(G') + 1 = \chi(G) + 1$ . Let  $\ell: E(G') \rightarrow [\chi(G) + 1]$  be a labeling such that  $g_\ell$  induces a proper vertex coloring of  $G'$ . We extend  $\ell$  to  $E(G)$ .

**Case 1:  $p = 0$ .** If  $g_\ell(u) = 0$ , then define  $\ell(uv_0) = \ell(uw)$  for some edge  $uw$  in  $E(G')$ . Otherwise there are at least two distinct labels on edges incident to  $u$  in  $E(G')$ . Thus we can label  $uv_0$  from among those distinct labels in such a way that  $g_\ell(v_0) \neq g_\ell(u)$ .

**Case 2:  $p > 0$ .** Define  $\ell(uv_p) = \ell(uw)$  for some edge  $uw$  in  $E'(G)$ . It is straightforward to iteratively label the remaining edges  $v_p v_{p-1}, \dots, v_1 v_0$  with values from  $\{1, 2, 3\}$  so that  $g_\ell$  induces a proper vertex coloring of  $G$ .

In either case,  $g_\ell$  induces a proper vertex coloring of  $G$  for some  $\ell: E(G) \rightarrow [\chi(G) + 1]$ .  $\square$

## 5 Further Pursuits

We determine  $\chi_\Delta^e(G)$  exactly for cliques, cycles, and trees in the following propositions. It is easy to see that  $\chi_\Delta^e(K_3) = 4$ .

**Proposition 1.** *Let  $n \geq 4$ . Then  $\chi_\Delta^e(K_n) = n$ .*

*Proof.* Let  $v_1 \dots v_{n-1}$  be a cycle on all but one vertex,  $v_0$ , of  $K_n$ . Define  $\ell(v_{n-1}v_1) = 1$ ,  $\ell(v_i v_{i+1}) = i + 2$  for  $i = 2, \dots, n - 2$ , and  $\ell(e) = 2$  for all remaining edges  $e$ . Note that  $c_\ell(v_i) = i$  for  $i = 0, \dots, n - 1$ .  $\square$

**Proposition 2.** *Let  $C_n$  be a cycle on  $n \geq 4$  vertices. Then*

$$\chi_\Delta^e(C_n) = \begin{cases} 2 & \text{if } n \equiv 0 \pmod{4}, \\ 3 & \text{otherwise.} \end{cases}$$



*Proof.* A local gap 2-coloring must alternate gap colors between 0 and 1 along vertices. This is possible precisely when  $n \equiv 0 \pmod{4}$ . For  $n \equiv 2 \pmod{4}$ , Theorem 4 then implies  $\chi_{\Delta}^e(C_n) = 3$ . For odd  $n \geq 5$ , let  $C_n = v_1 \dots v_n$ . If  $n \equiv 1 \pmod{4}$ , define  $\ell(v_n v_1) = 3$  and, for  $i \in \{1, \dots, n-1\}$ ,

$$\ell(v_i v_{i+1}) = \begin{cases} 1 & \text{if } i \equiv 1, 2 \pmod{4}, \\ 2 & \text{if } i \equiv 0, 3 \pmod{4}. \end{cases}$$

If  $n \equiv 3 \pmod{4}$ , define  $\ell(v_{n-1} v_n) = \ell(v_n v_1) = 3$  and, for  $i \in \{1, n-2\}$ ,

$$\ell(v_i v_{i+1}) = \begin{cases} 1 & \text{if } i \equiv 1, 2 \pmod{4}, \\ 2 & \text{if } i \equiv 0, 3 \pmod{4}. \end{cases}$$

This completes the proof.  $\square$

**Proposition 3.** *Let  $T$  be a tree on  $n$  vertices,  $n \geq 3$ . Then  $\chi_{\Delta}^e(T) = 2$  if all the leaves of  $T$  are in the same partite set of a bipartition of  $T$ ; otherwise  $\chi_{\Delta}^e(T) = 3$ .*

*Proof.* Lemma 1 implies that  $\chi_{\Delta}^e(T) \leq 3$ . Since all leaves have gap color 0, a local gap 2-coloring is not possible if leaves appear in both partite sets of  $T$ .

Now, let all leaves be in the same partite set of  $T$ . Let  $v$  be a leaf of  $T$ . For each  $e \in E_i(v)$ , define  $\ell(e) = 1$  if  $i \equiv 0, 1 \pmod{4}$ , and  $\ell(e) = 2$  otherwise. Since  $T$  has no cycles,  $\ell$  is a local gap 2-coloring of  $T$ .  $\square$

Since determining  $\chi(G)$  is APX-hard in general [7] and  $\chi_{\Delta}^e(G)$  is within an additive constant of  $\chi(G)$ , determining  $\chi_{\Delta}^e(G)$  is APX-hard as well. However, it would be interesting to investigate when  $\chi_{\Delta}^e(G)$  can be determined in polynomial time if  $\chi(G)$  is given as part of the input.

## References

- [1] T. Bohman and D. Kravitz. On the irregularity strength of trees. *J. Graph Theory*, 45(4):241–254, 2004.
- [2] G. Chartrand, M. S. Jacobson, J. Lehel, O. R. Oellermann, S. Ruiz, and F. Saba. Irregular networks. *Congr. Numer.*, 64(197-210):250th, 1988.
- [3] A. Frieze, R. J. Gould, M. Karoński, and F. Pfender. On graph irregularity strength. *J. Graph Theory*, 41(2):120–137, 2002.
- [4] E. Győri, M. Horňák, C. Palmer, and M. Woźniak. General neighbour-distinguishing index of a graph. *Discrete Math.*, 308(5-6):827–831, 2008.
- [5] M. Kalkowski, M. Karoński, and F. Pfender. Vertex-coloring edge-weightings: towards the 1-2-3-conjecture. *J. Combin. Theory Ser. B*, 100(3):347–349, 2010.
- [6] M. Karoński, T. Łuczak, and A. Thomason. Edge weights and vertex colours. *J. Combin. Theory Ser. B*, 91(1):151–157, 2004.
- [7] C. Lund and M. Yannakakis. On the hardness of approximating minimization problems. *J. Assoc. Comput. Mach.*, 41(5):960–981, 1994.

- [8] T. Nierhoff. A tight bound on the irregularity strength of graphs. *SIAM J. Discrete Math.*, 13(3):313–323 (electronic), 2000.
- [9] R. Scheidweiler and E. Triesch. Gap-neighbour-distinguishing colourings. *J. Combin. Math. Combin. Comput.*, (to appear).
- [10] R. Scheidweiler and E. Triesch. New estimates for the gap chromatic number. *Discrete Math.*, 328:42–53, 2014.
- [11] B. Seamone. The 1–2–3 conjecture and related problems: a survey. arXiv:1211.5122.
- [12] M. A. Tahraoui, E. Duchêne, and H. Kheddouci. Gap vertex-distinguishing edge colorings of graphs. *Discrete Math.*, 312(20):3011–3025, 2012.
- [13] D. B. West. *Introduction to graph theory*. Prentice Hall, Inc., Upper Saddle River, NJ, 1996.