

Assignment 2

Matish Singh Tanwar
AI20MTECH11005

Abstract—This document finds the equation of a plane which is at a distance of 7 units from origin and normal to $\begin{pmatrix} 3 \\ 5 \\ -6 \end{pmatrix}$

Download all python codes from

https://github.com/Matish007/Matrix-Theory-EE5609-/tree/master/Assignment_2/codes

and latex-tikz codes from

https://github.com/Matish007/Matrix-Theory-EE5609-/tree/master/Assignment_2

1 PROBLEM

Find the equation of a plane which is at a distance of 7 units from origin and normal to $\begin{pmatrix} 3 \\ 5 \\ -6 \end{pmatrix}$

$$\mathbf{n} = \begin{pmatrix} 3 \\ 5 \\ -6 \end{pmatrix} \quad (1)$$

2 EXPLANATION

Let the equation of plane be:-

$$\mathbf{n}^T \mathbf{x} = c \quad (2)$$

where \mathbf{n} =normal vector to the plane

The distance from the origin is given by:-

$$\frac{|c|}{\|\mathbf{n}\|} = 7 \quad (3)$$

$$\|\mathbf{n}\| = \sqrt{3^2 + 5^2 + (-6)^2} = \sqrt{70} \quad (4)$$

Substituting equation (4) in (3) we get,

$$\frac{|c|}{\sqrt{70}} = 7 \quad (5)$$

$$c = \pm 7 \sqrt{70} \quad (6)$$

Substituting equation (1),(6) in (2) we get two equation of planes,

$$(3 \ 5 \ -6) \mathbf{x} = 7 \sqrt{70} \quad (7)$$

$$(3 \ 5 \ -6) \mathbf{x} = -7 \sqrt{70} \quad (8)$$

Equation (7) and (8) gives us the equation of two planes which are at a distance of 7 units from origin and normal to $\begin{pmatrix} 3 \\ 5 \\ -6 \end{pmatrix}$




Fig. 1: Plane1 with Normal vector




Fig. 2: Plane2 with Normal vector