MAUNA

2025-2026

HAMBOURG STAFF REPORT — 1–7 MARCH 2026

Sat, 07/03/26 · 2026-03-01 → 2026-03-07

Hambourg Staff Report — 1–7 March 2026

COACH BRIEF

Rapport staff Hambourg — fenêtre principale : 1 → 7 mars

01 · Cover — cover

02 · Staff read — brief

03 · ML Performance & Injury Risk Scores — §1

04 · External Load by Position (GPS) — §2

05 · Subjective Wellness (25 Feb – 7 Mar) — §3

06 · Sleep Quality × Mood (25 Feb – 7 Mar) — §4

07 · Biomarkers & Tests — December 2025 — §5

08 · Feature Importance & ML Drivers (1–7 March) — §6

09 · Top Injury-Risk Drivers — §7

10 · Subjective & Recovery Drivers — §8

11 · Explosive & Asymmetry Drivers — §9

12 · Linking Drivers to 1–7 March Observations — §10

13 · Individual Athlete Profiles — High-Risk Watch — §11

14 · Yussuf Poulsen — Striker — §12

STAFF READ 2026-03-01 → 2026-03-07

Staff overview

NARRATIVE THREAD

- The club's ML injury-risk model is driven by 12 key features, ranked by recent importance. The top drivers reflect accumulated training stress, explosive load patterns, and subjective wellness state. Understanding these mechanisms helps interpret the 1–7 March scores and identify actionable levers for staff intervention.
- Overall Volume Strain (7-day) — Mean importance 693. This indicator combines high 7-day average volume (total distance, energy expenditure, cumulative speed intensity) with low day-to-day variability. The squared ratio of mean to standard deviation amplifies the signal when training is both heavy and monotonous. Elevated strain suggests cumulative physiological stress and heightened overuse injury risk. During 1–7 March, players with consistently high GPS volumes and minimal rest days will show elevated strain. Staff should cross-check total distance and HMLD trends against injury-risk scores to identify athletes absorbing high monotonous loads.
- Planned Explosiveness Peak Strain (7-day) — Mean importance 568. Reflects accumulated neuromuscular stress from planned maximal acceleration, deceleration, and zone 6 efforts over the past week. Like volume strain, this is calculated as the square of the 7-day average explosive load divided by its standard deviation. High values indicate sustained explosive exposure with little variation in the training plan, signaling elevated neuromuscular fatigue and injury risk. In the 1–7 March window, players with high explosive distance (kpi_explosivedistanceabs) and consistent high-intensity sessions will be flagged. Compare explosive GPS metrics against injury-risk scores to spot athletes under cumulative neuromuscular load.

SIGNALS

INFO The club's ML injury-risk model is driven by 12 key features, ranked by recent importance. The top drivers reflect accumulated training stress, explosive load patterns, and subjective wellness state. Understanding these mechanisms helps interpret the 1–7 March scores and identify actionable levers for staff intervention.

INFO Overall Volume Strain (7-day) — Mean importance 693. This indicator combines high 7-day average volume (total distance, energy expenditure, cumulative speed intensity) with low day-to-day variability. The squared ratio of mean to standard deviation amplifies the signal when training is both heavy and monotonous. Elevated strain suggests cumulative physiological stress and heightened overuse injury risk. During 1–7 March, players with consistently high GPS volumes and minimal rest days will show elevated strain. Staff should cross-check total distance and HMLD trends against injury-risk scores to identify athletes absorbing high monotonous loads.

INFO Planned Explosiveness Peak Strain (7-day) — Mean importance 568. Reflects accumulated neuromuscular stress from planned maximal acceleration, deceleration, and zone 6 efforts over the past week. Like volume strain, this is calculated as the square of the 7-day average explosive load divided by its standard deviation. High values indicate sustained explosive exposure with little variation in the training plan, signaling elevated neuromuscular fatigue and injury risk. In the 1–7 March window, players with high explosive distance (kpi_explosivedistanceabs) and consistent high-intensity sessions will be flagged. Compare explosive GPS metrics against injury-risk scores to spot athletes under cumulative neuromuscular load.

INFO Average Total External Load (7-day) — Mean importance 264. Simple 7-day mean of total training load. Higher values indicate sustained high workload, contributing to fatigue accumulation. This feature complements strain by capturing absolute load magnitude without the variability penalty. During 1–7 March, players with elevated sRPE and GPS totals will show higher average load. Staff should monitor whether high-load athletes also exhibit elevated injury-risk scores, suggesting insufficient recovery capacity.

INFO Total External Load Change vs Yesterday — Mean importance 158. Day-to-day delta in total workload. Large positive spikes (abrupt load increases) are a known injury-risk trigger, while large negative swings may indicate reactive load management or missed sessions. The model uses this to detect acute load jumps that outpace adaptation. In the 1–7 March data, check for days with sharp GPS or sRPE increases; athletes with high injury-risk scores on those days may be responding poorly to the spike.

CLUB DRIVERS & SCIENCE

Overall Volume Strain (7days) — recent weight 693.44

Indicator of accumulated training stress related to overall volume exposure (total distance, energy expenditure, playing time and cumulative speed intensity) over the past 7 days. Strain is calculated as the square of the 7-day average overall volume load divided by its 7-day standard deviation. This variable reflects the combined effect of high volume exposure and low day-to-day variability, indicating elevated cumulative physiological stress. Higher values may be associated with an increased risk of overuse and fatigue-related injuries.

Planned Explosiveness Peak Strain (7days) — recent weight 568.48

Indicator of the accumulated training stress associated with the player's planned explosiveness peaks load (maximal acceleration and deceleration, and zone 6 efforts) over the past 7 days. Strain is calculated as the square of the 7-day average planned explosive load divided by its 7-day standard deviation. This variable reflects the combined effect of high planned explosive exposure and low day-to-day variability in the training plan, providing a reference for the cumulative neuromuscular stress the player was expected to undergo.

Average Total External Load (7days) — recent weight 264.48

Player's average total training load over the last week.

Total External Load change vs Yesterday — recent weight 157.94

Difference between today's total external load and yesterday's, capturing the day-to-day variation in overall training demands. This variable provides insight into abrupt increases or decreases in total workload.

Total External Load Variability (7days) — recent weight 139.79

Standard deviation of the player's total external load over the past 7 days, capturing the day-to-day variability of training demands. This variable reflects how consistent or irregular recent workload has been: low values indicate monotonous loading with little variation, while high values indicate large swings between heavy and light days.

Training RPE vs Usual — recent weight 72.31

Today's training RPE as a simple multiple of the player's typical level in recent months (e.g. 1.5 = rated 50% harder than usual).

Recovery score — recent weight 55.97

Estimated recovery score (0-100) reflecting the player's current recovery state based on recent training demands: the harder the recent workload, the lower the score. This variable provides insight into how much physiological recovery the player has achieved relative to accumulated fatigue.

Explosiveness Load Change vs Yesterday — recent weight 52.63

Difference between today's explosiveness peaks load (maximal acceleration and deceleration, and zone 6 efforts) and yesterday's, capturing the day-to-day variation in explosive demands. This variable provides insight into abrupt increases or decreases in maximal-intensity actions.

Neuromuscular Asymmetry Load Change vs Yesterday — recent weight 52.17

Difference between today's left/right neuromuscular asymmetry load (impact magnitude, steps, and vertical and lateral impacts) and yesterday's, capturing the day-to-day variation in asymmetry. This variable provides insight into abrupt increases or decreases in left/right imbalances, which may reflect emerging compensation patterns, localized fatigue or an elevated injury risk when the change is disproportionately large.


Subjective Wellness Score — recent weight 49.22

Wellness score (0-100) built from the player's self-reported state: good sleep quality pushes the score up, while high perceived exertion pulls it down. This variable reflects the player's subjective readiness and provides insight into how they are coping with recent training demands beyond what external load metrics capture.

ML PERFORMANCE & INJURY RISK SCORES


2026-03-01 → 2026-03-07

PERFORMANCE SCORE (1/5)




Albert Grønbæk · Albert Sambi Lokonga · Bakery Jatta · Damion Downs · Daniel Elfadli

PERFORMANCE SCORE (2/5)




Fabio Baldé · Fábio Vieira · Giorgi Gocholeishvili · Jean-Luc Dompé · Jordan Torunarigha

PERFORMANCE SCORE (3/5)




Luka Vušković · Miro Muheim · Nicolai Remberg · Nicolás Capaldo · Noah Katterbach

PERFORMANCE SCORE (4/5)




Omar Megeed · Otto Stange · Philip Otele · Ransford Königsdörffer · Rayan Philippe

PERFORMANCE SCORE (5/5)




Robert Glatzel · Warmed Omari · William Mikelbrencis · Yussuf Poulsen

INJURY RISK SCORE (1/5)




Albert Grønbæk · Albert Sambi Lokonga · Bakery Jatta · Damion Downs · Daniel Elfadli


INJURY RISK SCORE (2/5)



INJURY RISK SCORE (3/5)




INJURY RISK SCORE (4/5)




Omar Megeed · Otto Stange · Philip Otele · Ransford Königsdörffer · Rayan Philippe

INJURY RISK SCORE (5/5)




Robert Glatzel · Warmed Omari · William Mikelbrencis · Yussuf Poulsen

PERFORMANCE SCORE — TEAM MEAN ± MIN-MAX




INJURY RISK SCORE — TEAM MEAN ± MIN-MAX




EXTERNAL LOAD BY POSITION (GPS)

2026-03-01 → 2026-03-07


TOTAL DISTANCE BY POSITION (MEAN)




HMLD BY POSITION (MEAN)



HIGH SPEED RUNNING BY POSITION (MEAN)



EXPLOSIVE DISTANCE BY POSITION (MEAN)



GPS EXTERNAL LOAD BY PLAYER & POSITION

PLAYER	POSITION	DISTANCE TOTAL (M)	HMLD (M)	HIGH SPEED RUNNING (ABSOLUTE)	EXPLOSIVE DISTANCE (ABSOLUTE)
Kofi Amoako	MIDFIELDER	–	–	–	–
Fabio Baldé	WINGER	2,690	328	133	195
Nicolás Capaldo	CENTRAL DEFENDER	–	–	–	–
Jean-Luc Dompé	WINGER	2,407	213	51	163
Damion Downs	STRIKER	2,023	155	41	114
Daniel Elfadli	DEFENSIVE MIDFIELDER	3,108	423	120	302
Robert Glatzel	STRIKER	3,111	343	83	260
Giorgi Gocholeishvili	FULL BACK	2,592	296	63	232
Albert Grønbæk	WINGER	3,219	526	89	436
Silvan Hefti	FULL BACK	–	–	–	–
Bakery Jatta	WINGER	3,207	480	160	320
Noah Katterbach	FULL BACK	3,255	428	205	223
Ransford Königsdörffer	STRIKER	3,105	389	177	212
Jonas Meffert	DEFENSIVE MIDFIELDER	–	–	–	–
Omar Megeed	ATTACKING MIDFIELDER	3,294	593	140	453
William Mikelbrencis	FULL BACK	2,771	393	134	258
Miro Muheim	FULL BACK	3,043	360	108	252
Warmed Omari	CENTRAL DEFENDER	2,324	278	66	212
Philip Otele	WINGER	3,104	507	267	240
Immanuël Pherai	ATTACKING MIDFIELDER	–	–	–	–
Rayan Philippe	STRIKER	2,887	376	109	267
Łukasz Poręba	ATTACKING MIDFIELDER	–	–	–	–

PLAYER	POSITION	DISTANCE TOTAL (M)	HMLD (M)	HIGH SPEED RUNNING (ABSOLUTE)	EXPLOSIVE DISTANCE (ABSOLUTE)
Yussuf Poulsen	STRIKER	3,032	371	119	253
Guilherme Ramos	CENTRAL DEFENDER	–	–	–	–
Nicolai Remberg	DEFENSIVE MIDFIELDER	413	7	0	7
Alexander Røssing-Lelesiit	WINGER	–	–	–	–
Emir Sahiti	WINGER	–	–	–	–
Albert Sambi Lokonga	CENTRAL MIDFIELDER	2,722	330	105	226
Aboubaka Soumahoro	CENTRAL DEFENDER	–	–	–	–
Otto Stange	STRIKER	3,282	369	146	222
Anssi Suhonen	ATTACKING MIDFIELDER	–	–	–	–
Jordan Torunarigha	CENTRAL DEFENDER	2,521	289	75	214
Fábio Vieira	CENTRAL MIDFIELDER	2,735	363	31	332
Luka Vušković	CENTRAL DEFENDER	3,171	383	34	349

SUBJECTIVE WELLNESS (25 FEB – 7 MAR)

2026-03-01 → 2026-03-07


WELLNESS PROFILE (MEAN SORENESS & MOOD) (1/2)

WELLNESS PROFILE (MEAN SORENESS & MOOD) (2/2)

SLEEP QUALITY × MOOD (25 FEB – 7 MAR)

2026-03-01 → 2026-03-07

SLEEP QUALITY VS MOOD



BIOMARKERS & TESTS — DECEMBER 2025

2026-03-01 → 2026-03-07

BIOMARKERS — DECEMBER 2025

PLAYER	DATE	BIOMARKER	VALUE
Aboubaka Soumahoro	2025-12-16	cmj_score	1
Albert Sambi Lokonga	2025-12-16	cmj_score	1
Alexander Røssing-Lelesiit	2025-12-16	cmj_score	1
Bakery Jatta	2025-12-16	cmj_score	5
Daniel Elfadli	2025-12-16	cmj_score	1
Emir Sahiti	2025-12-16	cmj_score	2
Fabio Baldé	2025-12-16	cmj_score	1
Fábio Vieira	2025-12-16	cmj_score	1
Giorgi Gocholeishvili	2025-12-16	cmj_score	1
Immanuël Pherai	2025-12-16	cmj_score	1
Jean-Luc Dompé	2025-12-16	cmj_score	1
Jonas Meffert	2025-12-16	cmj_score	3
Jordan Torunarigha	2025-12-16	cmj_score	2
Luka Vušković	2025-12-16	cmj_score	5
Miro Muheim	2025-12-16	cmj_score	2
Nicolai Remberg	2025-12-16	cmj_score	1
Nicolás Capaldo	2025-12-16	cmj_score	2
Noah Katterbach	2025-12-16	cmj_score	1
Ransford Königsdörffer	2025-12-16	cmj_score	1
Rayan Philippe	2025-12-16	cmj_score	1
Silvan Hefti	2025-12-16	cmj_score	2
Warmed Omari	2025-12-16	cmj_score	2
William Mikelbrencis	2025-12-16	cmj_score	2
Yussuf Poulsen	2025-12-16	cmj_score	1

FEATURE IMPORTANCE & ML DRIVERS (1–7 MARCH)

2026-03-01 → 2026-03-07

- The club's ML injury-risk model is driven by 12 key features, ranked by recent importance. The top drivers reflect accumulated training stress, explosive load patterns, and subjective wellness state. Understanding these mechanisms helps interpret the 1–7 March scores and identify actionable levers for staff intervention.

TOP INJURY-RISK DRIVERS

2026-03-01 → 2026-03-07

- Overall Volume Strain (7-day) — Mean importance 693. This indicator combines high 7-day average volume (total distance, energy expenditure, cumulative speed intensity) with low day-to-day variability. The squared ratio of mean to standard deviation amplifies the signal when training is both heavy and monotonous. Elevated strain suggests cumulative physiological stress and heightened overuse injury risk. During 1–7…
- Planned Explosiveness Peak Strain (7-day) — Mean importance 568. Reflects accumulated neuromuscular stress from planned maximal acceleration, deceleration, and zone 6 efforts over the past week. Like volume strain, this is calculated as the square of the 7-day average explosive load divided by its standard deviation. High values indicate sustained explosive exposure with little variation in the training plan, sign…

- Average Total External Load (7-day) — Mean importance 264. Simple 7-day mean of total training load. Higher values indicate sustained high workload, contributing to fatigue accumulation. This feature complements strain by capturing absolute load magnitude without the variability penalty. During 1–7 March, players with elevated sRPE and GPS totals will show higher average load. Staff should monitor whether high-loa...
- Total External Load Change vs Yesterday — Mean importance 158. Day-to-day delta in total workload. Large positive spikes (abrupt load increases) are a known injury-risk trigger, while large negative swings may indicate reactive load management or missed sessions. The model uses this to detect acute load jumps that outpace adaptation. In the 1–7 March data, check for days with sharp GPS or sRPE increases; athletes ...
- Total External Load Variability (7-day) — Mean importance 140. Standard deviation of total load over the past week. Low values indicate monotonous training (little variation), while high values reflect large swings between heavy and light days. Both extremes carry risk: monotony reduces adaptive stimulus and recovery opportunity, while excessive variability can cause maladaptation. The model balances this with str...

SUBJECTIVE & RECOVERY DRIVERS

2026-03-01 → 2026-03-07

- Training RPE vs Usual — Mean importance 72. Ratio of today's training RPE to the player's typical recent level (e.g., 1.5 = 50% harder than usual). Captures perceived exertion spikes relative to individual baseline. High ratios indicate the player felt the session was unusually demanding, which may reflect inadequate recovery or session intensity mismatch. During 1–7 March, cross-check sRPE daily values (ds_1) aga...
- Recovery Score — Mean importance 56. Estimated 0–100 score reflecting current recovery state based on recent training demands. Lower scores indicate insufficient recovery relative to accumulated fatigue. This feature integrates external load history to estimate physiological readiness. In the 1–7 March window, players with low recovery scores and high injury-risk scores are likely under-recovered. Staff should con...
- Subjective Wellness Score — Mean importance 49. Composite 0–100 score built from self-reported state: good sleep quality increases the score, high perceived exertion decreases it. Reflects the player's subjective readiness beyond external load metrics. During 25 Feb–7 March, wellness data (ds_6, ds_7) shows body soreness and mood ratings, plus sleep quality. Players with poor wellness scores (high soreness, low mo...
- Body RPE vs Usual — Mean importance 49. Ratio of today's body-condition rating to the player's typical recent level. Higher ratios indicate the player feels worse than usual (e.g., 1.5 = 50% worse soreness/fatigue). This complements training RPE by focusing on body state rather than session intensity. In the 25 Feb–7 March wellness data, body_soreness_rating is available (scale 10–60, lower is better). Players wit...

EXPLOSIVE & ASYMMETRY DRIVERS

2026-03-01 → 2026-03-07

- Explosiveness Load Change vs Yesterday — Mean importance 53. Day-to-day delta in explosive load (maximal acceleration, deceleration, zone 6 efforts). Abrupt increases in explosive actions can trigger acute neuromuscular fatigue and injury. The model uses this to detect explosive load spikes. During 1–7 March, check kpi_explosivedistanceabs daily trends (ds_5); players with sharp increases and elevated injury-risk ...
- Neuromuscular Asymmetry Load Change vs Yesterday — Mean importance 52. Day-to-day delta in left/right asymmetry load (impact magnitude, steps, vertical/lateral impacts). Large changes in asymmetry may reflect emerging compensation patterns, localized fatigue, or elevated injury risk. The model flags disproportionate asymmetry shifts. This feature is not directly visible in the current GPS dataset (ds_5), but staff...

LINKING DRIVERS TO 1–7 MARCH OBSERVATIONS

2026-03-01 → 2026-03-07

- The 1–7 March data shows 168 daily score observations (ds_4) across the squad. Injury-risk scores range from low to elevated, with individual variation reflecting the interplay of the 12 drivers above. To interpret these scores, staff should cross-reference GPS external load (ds_5: total distance, HMLD, HSR, explosive distance), internal load (ds_1: sRPE), and subjective wellness (ds_6, ds_7: soreness, mood, sleep...

INDIVIDUAL ATHLETE PROFILES — HIGH-RISK WATCH

2026-03-01 → 2026-03-07

TOP 5 HIGHEST INJURY-RISK SCORES (MEAN 1–7 MARCH)

PLAYER	INJURY RISK SCORE (0-100)
Yussuf Poulsen	63.57
Bakery Jatta	43.71
Robert Glatzel	17.14
Nicolás Capaldo	7.86
Daniel Elfadli	7.14

YUSSUF POULSEN — STstriker

2026-03-01 → 2026-03-07

YP

Yussuf Poulsen
STRIKER

- Yussuf Poulsen shows the highest mean injury-risk score in the squad at 63.6 over 1–7 March, with a peak of 64 on 1 March declining to 50 by 7 March. Performance score remains moderate (44–50 range). GPS data for 6 March shows 3,032 m total distance, 371 m HMLD, 119 m HSR, and 253 m explosive distance — all within typical striker ranges but accumulated over a week with limited variability. Subjective wellness (25 ...