

Lecture 8: Introduction to Monte Carlo Methods

Deniz Akyildiz

MATH60047/70047 – Stochastic Simulation

November 1, 2022

**Imperial College
London**

"City of London, rainy day, South Kensington, people walking, oil painting, reflections, against white background" (DALLE-2)

What we have done yesterday?

What we have done yesterday?

- ▶ Bayes rule for conditionally iid case (our *target* distributions could be posteriors)

What we have done yesterday?

- ▶ Bayes rule for conditionally iid case (our *target* distributions could be posteriors)
- ▶ Application of rejection sampling to posterior sampling and high dimensional targets (?)

What we have done yesterday?

- ▶ Bayes rule for conditionally iid case (our *target* distributions could be posteriors)
- ▶ Application of rejection sampling to posterior sampling and high dimensional targets (?)

Today, we will discuss

What we have done yesterday?

- ▶ Bayes rule for conditionally iid case (our *target* distributions could be posteriors)
- ▶ Application of rejection sampling to posterior sampling and high dimensional targets (?)

Today, we will discuss

- ▶ Marginal likelihoods

What we have done yesterday?

- ▶ Bayes rule for conditionally iid case (our *target* distributions could be posteriors)
- ▶ Application of rejection sampling to posterior sampling and high dimensional targets (?)

Today, we will discuss

- ▶ Marginal likelihoods
- ▶ An introduction to Monte Carlo methods

What we have done yesterday?

- ▶ Bayes rule for conditionally iid case (our *target* distributions could be posteriors)
- ▶ Application of rejection sampling to posterior sampling and high dimensional targets (?)

Today, we will discuss

- ▶ Marginal likelihoods
- ▶ An introduction to Monte Carlo methods
 - ▶ A set of general methods for sampling from distributions, estimating expectations, computing integrals

Tools of Probabilistic Inference

Marginal Likelihood

Recall the posterior computation:

$$p(x|y) = \frac{p(y|x)p(x)}{p(y)} = \frac{p(y|x)p(x)}{\int p(y|x)p(x)dx}.$$

We did not discuss the denominator $p(y)$ in detail:

Tools of Probabilistic Inference

Marginal Likelihood

Recall the posterior computation:

$$p(x|y) = \frac{p(y|x)p(x)}{p(y)} = \frac{p(y|x)p(x)}{\int p(y|x)p(x)dx}.$$

We did not discuss the denominator $p(y)$ in detail:

- ▶ It is called the *marginal likelihood*

Tools of Probabilistic Inference

Marginal Likelihood

Recall the posterior computation:

$$p(x|y) = \frac{p(y|x)p(x)}{p(y)} = \frac{p(y|x)p(x)}{\int p(y|x)p(x)dx}.$$

We did not discuss the denominator $p(y)$ in detail:

- ▶ It is called the *marginal likelihood*
- ▶ It is the probability of the data y (now fixed) given the model

Tools of Probabilistic Inference

Marginal Likelihood

Recall the posterior computation:

$$p(x|y) = \frac{p(y|x)p(x)}{p(y)} = \frac{p(y|x)p(x)}{\int p(y|x)p(x)dx}.$$

We did not discuss the denominator $p(y)$ in detail:

- ▶ It is called the *marginal likelihood*
- ▶ It is the probability of the data y (now fixed) given the model
- ▶ It is the normalization constant for the posterior

Tools of Probabilistic Inference

Marginal Likelihood

Recall the posterior computation:

$$p(x|y) = \frac{p(y|x)p(x)}{p(y)} = \frac{p(y|x)p(x)}{\int p(y|x)p(x)dx}.$$

We did not discuss the denominator $p(y)$ in detail:

- ▶ It is called the *marginal likelihood*
- ▶ It is the probability of the data y (now fixed) given the model
- ▶ It is the normalization constant for the posterior

It is an extremely important quantity for choosing models.

Tools of Probabilistic Inference

Marginal Likelihood

We may not need to compute $p(y) = \int p(y|x)p(x)dx$ for *posterior* inference as we discussed.

Tools of Probabilistic Inference

Marginal Likelihood

We may not need to compute $p(y) = \int p(y|x)p(x)dx$ for *posterior* inference as we discussed.

But we will need it for *model* selection and *prediction*.

Tools of Probabilistic Inference

Marginal Likelihood

We may not need to compute $p(y) = \int p(y|x)p(x)dx$ for *posterior* inference as we discussed.

But we will need it for *model* selection and *prediction*.

A simple example: Choosing the right prior.

Tools of Probabilistic Inference

Marginal Likelihood

Consider two different models:

$$\begin{aligned}x &\sim p_0(x) = \mathcal{N}(x; \mu_0, \sigma_0^2) \\y|x &\sim \mathcal{N}(y; x, \sigma_y^2)\end{aligned}$$

and

$$\begin{aligned}x &\sim p_1(x) = \mathcal{N}(x; \mu_1, \sigma_1^2) \\y|x &\sim \mathcal{N}(y; x, \sigma_y^2)\end{aligned}$$

Consider observing y (a single data point). Which model is more likely?

Tools of Probabilistic Inference

Marginal Likelihood

Recall that, for these models, we have computed $p(y)$ analytically before. We can compute for both models:

$$\begin{aligned} p_0(y) &= \int p(y|x)p_0(x)dx \\ &= \int \mathcal{N}(y; x, \sigma_y^2)\mathcal{N}(x; \mu_0, \sigma_0^2)dx \\ &= \mathcal{N}(y; \mu_0, \sigma_0^2 + \sigma_y^2) \end{aligned}$$

and

$$\begin{aligned} p_1(y) &= \int p(y|x)p_1(x)dx \\ &= \int \mathcal{N}(y; x, \sigma_y^2)\mathcal{N}(x; \mu_1, \sigma_1^2)dx \\ &= \mathcal{N}(y; \mu_1, \sigma_1^2 + \sigma_y^2) \end{aligned}$$

Tools of Probabilistic Inference

Marginal Likelihood

Recall that, for these models, we have computed $p(y)$ analytically before. We can compute for both models:

$$\begin{aligned} p_0(y) &= \int p(y|x)p_0(x)dx \\ &= \int \mathcal{N}(y; x, \sigma_y^2)\mathcal{N}(x; \mu_0, \sigma_0^2)dx \\ &= \mathcal{N}(y; \mu_0, \sigma_0^2 + \sigma_y^2) \end{aligned}$$


and

$$\begin{aligned} p_1(y) &= \int p(y|x)p_1(x)dx \\ &= \int \mathcal{N}(y; x, \sigma_y^2)\mathcal{N}(x; \mu_1, \sigma_1^2)dx \\ &= \mathcal{N}(y; \mu_1, \sigma_1^2 + \sigma_y^2) \end{aligned}$$

We will say Model 1 is better than Model 0 if $p_1(y) > p_0(y)$ for fixed y .

Tools of Probabilistic Inference

Marginal Likelihood



Bayesian no more...

- We will now look into the core concepts of the course

Bayesian no more...

- ▶ We will now look into the core concepts of the course
- A large class of problems in statistics, machine learning, numerical analysis can be formulated as

Bayesian no more...

- We will now look into the core concepts of the course

A large class of problems in statistics, machine learning, numerical analysis can be formulated as

- Sampling from a distribution (e.g. a posterior, an unknown data distribution)

Bayesian no more...

- We will now look into the core concepts of the course

A large class of problems in statistics, machine learning, numerical analysis can be formulated as

- Sampling from a distribution (e.g. a posterior, an unknown data distribution)
- Estimating an expectation (e.g. the expectation of a function of a random variable)

Bayesian no more...

- We will now look into the core concepts of the course

A large class of problems in statistics, machine learning, numerical analysis can be formulated as

- Sampling from a distribution (e.g. a posterior, an unknown data distribution)
- Estimating an expectation (e.g. the expectation of a function of a random variable)
- Computing an integral

Bayesian no more...

- We will now look into the core concepts of the course

A large class of problems in statistics, machine learning, numerical analysis can be formulated as

- Sampling from a distribution (e.g. a posterior, an unknown data distribution)
- Estimating an expectation (e.g. the expectation of a function of a random variable)
- Computing an integral

Monte Carlo methods are a set of general methods for solving these problems by generating random samples (rejection sampling counts as a Monte Carlo method).

Monte Carlo Methods

An introduction

Consider a distribution $p(x)$ and estimating the expectation

Monte Carlo Methods

An introduction

Consider a distribution $p(x)$ and estimating the expectation

$$\mathbb{E}_p[X] = \int xp(x)dx \tag{1}$$

Given samples $X_1, \dots, X_N \sim p(x)$, we can estimate the expectation as

Monte Carlo Methods

An introduction

Consider a distribution $p(x)$ and estimating the expectation

$$\mathbb{E}_p[X] = \int xp(x)dx \quad (1)$$

Given samples $X_1, \dots, X_N \sim p(x)$, we can estimate the expectation as

$$\mathbb{E}_p[X] \approx \frac{1}{N} \sum_{i=1}^N X_i \quad (2)$$

Monte Carlo Methods

An introduction

The idea of estimating an expectation with random samples is more general. Indeed, for any function $\varphi(x)$, the expectation

Monte Carlo Methods

An introduction

The idea of estimating an expectation with random samples is more general. Indeed, for any function $\varphi(x)$, the expectation

$$\bar{\varphi} = \mathbb{E}_p[\varphi(x)] = \int \varphi(x)p(x)dx \quad (3)$$

can be estimated using $X_1, \dots, X_N \sim p(x)$ as

Monte Carlo Methods

An introduction

The idea of estimating an expectation with random samples is more general. Indeed, for any function $\varphi(x)$, the expectation

$$\bar{\varphi} = \mathbb{E}_p[\varphi(x)] = \int \varphi(x)p(x)dx \quad (3)$$

can be estimated using $X_1, \dots, X_N \sim p(x)$ as

$$\bar{\varphi} = \mathbb{E}_p[\varphi(x)] \approx \frac{1}{N} \sum_{i=1}^N \varphi(X_i) = \hat{\varphi}^N \quad (4)$$

Monte Carlo Methods

An introduction

How to approximate probability distributions using samples? This is done via Dirac measure.

Monte Carlo Methods

An introduction

How to approximate probability distributions using samples? This is done via Dirac measure.

Define the Dirac measure:

$$\varphi(y) = \int \varphi(x)\delta_y(x)dx \quad (5)$$

Monte Carlo Methods

An introduction

How to approximate probability distributions using samples? This is done via Dirac measure.

Define the Dirac measure:

$$\varphi(y) = \int \varphi(x)\delta_y(x)dx \quad (5)$$

Intuition: δ_y is a measure concentrated at y : It evaluates the integral at y and is zero everywhere else. An incorrect way to write it:

$$\delta_y(x) = \begin{cases} 1 & x = y \\ 0 & x \neq y \end{cases}$$

Monte Carlo Methods

An introduction

How to approximate probability distributions using samples? This is done via Dirac measure.

Define the Dirac measure:

$$\varphi(y) = \int \varphi(x)\delta_y(x)dx \quad (5)$$

Intuition: δ_y is a measure concentrated at y : It evaluates the integral at y and is zero everywhere else. An incorrect way to write it:

$$\delta_y(x) = \begin{cases} 1 & x = y \\ 0 & x \neq y \end{cases}$$

Note that, the use of Dirac only makes sense with an integral!

Monte Carlo Methods

An introduction

Given samples $X_1, \dots, X_N \sim p(x)$, a probability distribution is approximated as

$$p(x)dx \approx \frac{1}{N} \sum_{i=1}^N \delta_{X_i}(x)dx = p^N(x)dx$$

The estimator is called the *empirical* distribution.

Monte Carlo Methods

An introduction

Then, we can prove that

$$\bar{\varphi} = \mathbb{E}_p[\varphi(x)]$$

Monte Carlo Methods

An introduction

Then, we can prove that

$$\begin{aligned}\bar{\varphi} &= \mathbb{E}_p[\varphi(x)] \\ &= \int \varphi(x)p(x)dx\end{aligned}$$

Monte Carlo Methods

An introduction

Then, we can prove that

$$\begin{aligned}\bar{\varphi} &= \mathbb{E}_p[\varphi(x)] \\ &= \int \varphi(x)p(x)dx \\ &\approx \int \varphi(x)p^N(x)dx\end{aligned}$$

Monte Carlo Methods

An introduction

Then, we can prove that

$$\begin{aligned}\bar{\varphi} &= \mathbb{E}_p[\varphi(x)] \\ &= \int \varphi(x)p(x)dx \\ &\approx \int \varphi(x)p^N(x)dx \\ &= \int \varphi(x)\frac{1}{N} \sum_{i=1}^N \delta_{X_i}(x)dx\end{aligned}$$

Monte Carlo Methods

An introduction

Then, we can prove that

$$\begin{aligned}\bar{\varphi} &= \mathbb{E}_p[\varphi(x)] \\ &= \int \varphi(x)p(x)dx \\ &\approx \int \varphi(x)p^N(x)dx \\ &= \int \varphi(x) \frac{1}{N} \sum_{i=1}^N \delta_{X_i}(x)dx \\ &= \frac{1}{N} \sum_{i=1}^N \int \varphi(x) \delta_{X_i}(x)dx\end{aligned}$$

Monte Carlo Methods

An introduction

Then, we can prove that

$$\begin{aligned}\bar{\varphi} &= \mathbb{E}_p[\varphi(x)] \\ &= \int \varphi(x)p(x)dx \\ &\approx \int \varphi(x)p^N(x)dx \\ &= \int \varphi(x) \frac{1}{N} \sum_{i=1}^N \delta_{X_i}(x)dx \\ &= \frac{1}{N} \sum_{i=1}^N \int \varphi(x) \delta_{X_i}(x)dx \\ &= \frac{1}{N} \sum_{i=1}^N \varphi(X_i) = \hat{\varphi}^N\end{aligned}$$

by the definition of the Dirac measure.

Monte Carlo Methods

An introduction

The same idea can be used to estimate an integral. Consider the problem:

$$I = \int f(x)dx \quad (6)$$

Monte Carlo Methods

An introduction

The same idea can be used to estimate an integral. Consider the problem:

$$I = \int f(x)dx \quad (6)$$

We can rephrase this problem as

$$I = \int \frac{f(x)}{p(x)} p(x)dx \quad (7)$$

Monte Carlo Methods

An introduction

The same idea can be used to estimate an integral. Consider the problem:

$$I = \int f(x)dx \quad (6)$$

We can rephrase this problem as

$$I = \int \frac{f(x)}{p(x)} p(x)dx \quad (7)$$

Rewriting $\varphi(x) = f(x)/p(x)$ and using the estimator we derived

$$I \approx \frac{1}{N} \sum_{i=1}^N \frac{f(X_i)}{p(X_i)}. \quad (8)$$

Monte Carlo Methods

An introduction

The same idea can be used to estimate an integral. Consider the problem:

$$I = \int f(x)dx \quad (6)$$

We can rephrase this problem as

$$I = \int \frac{f(x)}{p(x)} p(x)dx \quad (7)$$

Rewriting $\varphi(x) = f(x)/p(x)$ and using the estimator we derived

$$I \approx \frac{1}{N} \sum_{i=1}^N \frac{f(X_i)}{p(X_i)}. \quad (8)$$

This technique is sometimes called the Crude Monte Carlo estimator of the integral.

Monte Carlo Methods

Mean: no bias

Claim: The estimator $\hat{\varphi}^N$ is unbiased.

Monte Carlo Methods

Mean: no bias

Claim: The estimator $\hat{\varphi}^N$ is unbiased. *Proof:*

$$\mathbb{E}_p[\hat{\varphi}^N] = \mathbb{E}_p\left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i)\right]$$

Monte Carlo Methods

Mean: no bias

Claim: The estimator $\hat{\varphi}^N$ is unbiased. *Proof:*

$$\begin{aligned}\mathbb{E}_p[\hat{\varphi}^N] &= \mathbb{E}_p\left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i)\right] \\ &= \frac{1}{N} \sum_{i=1}^N \mathbb{E}_p[\varphi(X_i)]\end{aligned}$$

Monte Carlo Methods

Mean: no bias

Claim: The estimator $\hat{\varphi}^N$ is unbiased. *Proof:*

$$\begin{aligned}\mathbb{E}_p[\hat{\varphi}^N] &= \mathbb{E}_p\left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i)\right] \\ &= \frac{1}{N} \sum_{i=1}^N \mathbb{E}_p[\varphi(X_i)] \\ &= \frac{1}{N} \sum_{i=1}^N \int \varphi(x) p(x) dx\end{aligned}$$

Monte Carlo Methods

Mean: no bias

Claim: The estimator $\hat{\varphi}^N$ is unbiased. *Proof:*

$$\begin{aligned}\mathbb{E}_p[\hat{\varphi}^N] &= \mathbb{E}_p\left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i)\right] \\ &= \frac{1}{N} \sum_{i=1}^N \mathbb{E}_p[\varphi(X_i)] \\ &= \frac{1}{N} \sum_{i=1}^N \int \varphi(x) p(x) dx \\ &= \int \varphi(x) p(x) dx\end{aligned}$$

Monte Carlo Methods

Mean: no bias

Claim: The estimator $\hat{\varphi}^N$ is unbiased. *Proof:*

$$\begin{aligned}\mathbb{E}_p[\hat{\varphi}^N] &= \mathbb{E}_p\left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i)\right] \\ &= \frac{1}{N} \sum_{i=1}^N \mathbb{E}_p[\varphi(X_i)] \\ &= \frac{1}{N} \sum_{i=1}^N \int \varphi(x) p(x) dx \\ &= \int \varphi(x) p(x) dx \\ &= \bar{\varphi}.\end{aligned}$$

Monte Carlo Methods

Variance

Let us compute the variance of the estimator:

$$\text{var}_p[\hat{\varphi}^N] = \text{var}_p \left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i) \right]$$

Monte Carlo Methods

Variance

Let us compute the variance of the estimator:

$$\begin{aligned}\text{var}_p[\hat{\varphi}^N] &= \text{var}_p\left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i)\right] \\ &= \frac{1}{N^2} \sum_{i=1}^N \text{var}_p[\varphi(X_i)]\end{aligned}$$

Monte Carlo Methods

Variance

Let us compute the variance of the estimator:

$$\begin{aligned}\text{var}_p[\hat{\varphi}^N] &= \text{var}_p\left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i)\right] \\ &= \frac{1}{N^2} \sum_{i=1}^N \text{var}_p[\varphi(X_i)] \\ &= \frac{1}{N^2} \sum_{i=1}^N \int (\varphi(x) - \bar{\varphi})^2 p(x) dx\end{aligned}$$

Monte Carlo Methods

Variance

Let us compute the variance of the estimator:

$$\begin{aligned}\text{var}_p[\hat{\varphi}^N] &= \text{var}_p\left[\frac{1}{N} \sum_{i=1}^N \varphi(X_i)\right] \\ &= \frac{1}{N^2} \sum_{i=1}^N \text{var}_p[\varphi(X_i)] \\ &= \frac{1}{N^2} \sum_{i=1}^N \int (\varphi(x) - \bar{\varphi})^2 p(x) dx \\ &= \frac{1}{N} \text{var}_p[\varphi(X)] = \frac{\sigma_\varphi^2}{N}\end{aligned}$$

Provided that $\text{var}_p[\varphi(X)] < \infty$ and the estimator is consistent as $N \rightarrow \infty$.

Monte Carlo Methods

Variance

Given

$$\begin{aligned}\text{var}_p[\hat{\varphi}^N] &= \frac{1}{N} \text{var}_p[\varphi(X)] \\ &= \frac{1}{N} \int (\varphi(x) - \bar{\varphi})^2 p(x) dx\end{aligned}$$

Monte Carlo Methods

Variance

Given

$$\begin{aligned}\text{var}_p[\hat{\varphi}^N] &= \frac{1}{N} \text{var}_p[\varphi(X)] \\ &= \frac{1}{N} \int (\varphi(x) - \bar{\varphi})^2 p(x) dx \\ &\approx \frac{1}{N^2} \sum_{i=1}^N (\varphi(X_i) - \hat{\varphi}^N)^2\end{aligned}$$

Monte Carlo Methods

Variance

Given

$$\begin{aligned}\text{var}_p[\hat{\varphi}^N] &= \frac{1}{N} \text{var}_p[\varphi(X)] \\ &= \frac{1}{N} \int (\varphi(x) - \bar{\varphi})^2 p(x) dx \\ &\approx \frac{1}{N^2} \sum_{i=1}^N (\varphi(X_i) - \hat{\varphi}^N)^2 \\ &= \sigma_{\varphi, N}^2.\end{aligned}$$

The variance can also be estimated using Monte Carlo method.

Monte Carlo Methods

Central Limit Theorem

By the central limit theorem:

$$\frac{\hat{\varphi}^N - \bar{\varphi}}{\sigma_{\varphi, N}} \rightarrow \mathcal{N}(0, 1). \quad (9)$$

Monte Carlo Methods

Central Limit Theorem

By the central limit theorem:

$$\frac{\hat{\varphi}^N - \bar{\varphi}}{\sigma_{\varphi, N}} \rightarrow \mathcal{N}(0, 1). \quad (9)$$

Therefore, this can be used to build confidence intervals (however, not very trustable).

Monte Carlo Methods

Convergence rate

We can see that the standard deviation of the estimator is given by

$$\text{std}_p(\hat{\varphi}^N) = \frac{\sigma_\varphi}{\sqrt{N}}.$$

This is one reason you may hear much about $\mathcal{O}(1/\sqrt{N})$ convergence rate (this is optimal).

Monte Carlo Methods

Convergence rate

We can see that the standard deviation of the estimator is given by

$$\text{std}_p(\hat{\varphi}^N) = \frac{\sigma_\varphi}{\sqrt{N}}.$$

This is one reason you may hear much about $\mathcal{O}(1/\sqrt{N})$ convergence rate (this is optimal).

More on convergence later.

Different choices of φ can be used to estimate different quantities.

- ▶ $\varphi(x) = x$ to estimate the mean

Different choices of φ can be used to estimate different quantities.

- ▶ $\varphi(x) = x$ to estimate the mean
- ▶ $\varphi(x) = x^2$ to estimate the second moment

Different choices of φ can be used to estimate different quantities.

- ▶ $\varphi(x) = x$ to estimate the mean
- ▶ $\varphi(x) = x^2$ to estimate the second moment
- ▶ $\varphi(x) = \log x$ to estimate entropy

Different choices of φ can be used to estimate different quantities.

- ▶ $\varphi(x) = x$ to estimate the mean
- ▶ $\varphi(x) = x^2$ to estimate the second moment
- ▶ $\varphi(x) = \log x$ to estimate entropy

Can we estimate probabilities?

Monte Carlo Methods

Estimation of probabilities

Assume, we would like to estimate

$$\mathbb{P}(X \in A) = \int_A p(x)dx \quad (10)$$

where A can be an interval, i.e., $A \subset \mathbb{R}$.

Monte Carlo Methods

Estimation of probabilities

Assume, we would like to estimate

$$\mathbb{P}(X \in A) = \int_A p(x)dx \quad (10)$$

where A can be an interval, i.e., $A \subset \mathbb{R}$. Then

$$\begin{aligned}\mathbb{P}(X \in A) &= \int_A p(x)dx \\ &= \int \mathbf{1}_A(x)p(x)dx,\end{aligned}$$

where $\mathbf{1}_A(x)$ is the indicator function of A .

Monte Carlo Methods

$$\begin{aligned}\mathbb{P}(X \in A) &= \int \mathbf{1}_A(x)p(x)dx, \\ &\approx \int \mathbf{1}_A(x)p^N(x)dx, \\ &= \int \mathbf{1}_A(x)\frac{1}{N} \sum_{i=1}^N \delta_{X_i}(x)dx, \\ &= \frac{1}{N} \sum_{i=1}^N \int \mathbf{1}_A(x)\delta_{X_i}(x)dx, \\ &\approx \frac{1}{N} \sum_{i=1}^N \mathbf{1}_A(X_i).\end{aligned}$$

Monte Carlo Methods

$$\begin{aligned}\mathbb{P}(X \in A) &= \int \mathbf{1}_A(x)p(x)dx, \\ &\approx \int \mathbf{1}_A(x)p^N(x)dx, \\ &= \int \mathbf{1}_A(x)\frac{1}{N} \sum_{i=1}^N \delta_{X_i}(x)dx, \\ &= \frac{1}{N} \sum_{i=1}^N \int \mathbf{1}_A(x)\delta_{X_i}(x)dx, \\ &\approx \frac{1}{N} \sum_{i=1}^N \mathbf{1}_A(X_i).\end{aligned}$$

We effectively just count the samples in A and divide it by N .


Monte Carlo Methods

$$\begin{aligned}\mathbb{P}(X \in A) &= \int \mathbf{1}_A(x)p(x)dx, \\ &\approx \int \mathbf{1}_A(x)p^N(x)dx, \\ &= \int \mathbf{1}_A(x)\frac{1}{N} \sum_{i=1}^N \delta_{X_i}(x)dx, \\ &= \frac{1}{N} \sum_{i=1}^N \int \mathbf{1}_A(x)\delta_{X_i}(x)dx, \\ &\approx \frac{1}{N} \sum_{i=1}^N \mathbf{1}_A(X_i).\end{aligned}$$

We effectively just count the samples in A and divide it by N .

$\varphi(x) = \mathbf{1}_A(x)$ to estimate $\mathbb{P}(X \in A)$.



Our old problem:



Can we estimate the probability of this set, if we had access to samples from $\text{Unif}([-1, 1] \times [-1, 1])$?

Monte Carlo Methods

An introduction



Monte Carlo Methods

Proof: What you will understand by the end of the course

For probabilists, let $X = [-1, 1] \times [-1, 1]$ and define the uniform measure such that $\mathbb{P}(X) = 1$.

Let A be the “circle” s.t. $A \subset X$. Now, the probability of A is given

$$\begin{aligned}\mathbb{P}(A) &= \int_A \mathbb{P}(\mathrm{d}x) \\ &= \int \mathbf{1}_A(x) \mathbb{P}(\mathrm{d}x), \\ &\approx \frac{1}{N} \sum_{i=1}^N \mathbf{1}_A(x_i) \rightarrow \frac{\pi}{4} \quad \text{as } N \rightarrow \infty.\end{aligned}$$

where $x_i \sim \mathbb{P}$.

Monte Carlo Methods

Proof: What you will understand by the end of the course

For probabilists, let $X = [-1, 1] \times [-1, 1]$ and define the uniform measure such that $\mathbb{P}(X) = 1$.



Let A be the “circle” s.t. $A \subset X$. Now, the probability of A is given

$$\begin{aligned}\mathbb{P}(A) &= \int_A \mathbb{P}(\mathrm{d}x) \\ &= \int \mathbf{1}_A(x) \mathbb{P}(\mathrm{d}x), \\ &\approx \frac{1}{N} \sum_{i=1}^N \mathbf{1}_A(x_i) \rightarrow \frac{\pi}{4} \quad \text{as } N \rightarrow \infty.\end{aligned}$$

where $x_i \sim \mathbb{P}$.

What about the convergence rate?

Monte Carlo Methods



Monte Carlo Methods

Monte Carlo Integration

Let us consider the integral (Example 3.4 from Robert and Casella, 2004)

$$\int_0^1 h(x)dx = \int_0^1 [\cos(50x) + \sin(20x)]^2 dx. \quad (11)$$

The exact value of this integral is 0.965 (yes, you can compute it).

Monte Carlo Methods

Monte Carlo Integration

Let us consider the integral (Example 3.4 from Robert and Casella, 2004)

$$\int_0^1 h(x)dx = \int_0^1 [\cos(50x) + \sin(20x)]^2 dx. \quad (11)$$

The exact value of this integral is 0.965 (yes, you can compute it).

We can use Monte Carlo integration to estimate this integral. Choose $p(x) = \text{Unif}(0, 1)$, then

$$\int_0^1 h(x)dx = \int_0^1 h(x)p(x)dx$$

so a uniform sample from $[0, 1]$ is sufficient to estimate this integral.

Monte Carlo Methods

Monte Carlo Integration




Figure: Monte Carlo integration of $h(x) = [\cos(50x) + \sin(20x)]^2$

Monte Carlo Methods

Estimating tail probability

Assume $X \sim \mathcal{N}(0, 1)$ and we are interested in

$$\mathbb{P}(X > 2) = \int_2^{\infty} \mathcal{N}(x; 0, 1) dx. \quad (12)$$

Monte Carlo Methods

Estimating tail probability

Assume $X \sim \mathcal{N}(0, 1)$ and we are interested in

$$\mathbb{P}(X > 2) = \int_2^{\infty} \mathcal{N}(x; 0, 1) dx. \quad (12)$$

We can formulate this as a Monte Carlo problem by choosing

$$\varphi(x) = \mathbf{1}_{\{x>2\}}. \quad (13)$$

Monte Carlo Methods

Estimating tail probability

Our estimation task is then sample $X_1, \dots, X_n \sim \mathcal{N}(0, 1)$ then

$$\begin{aligned}\mathbb{P}(X > 2) &= \int_2^{\infty} \mathcal{N}(x; 0, 1) dx \\ &= \int_{-\infty}^{\infty} \mathbf{1}_{\{x>2\}} \mathcal{N}(x; 0, 1) dx \\ &\approx \frac{1}{n} \sum_{i=1}^n \mathbf{1}_{\{X_i>2\}}.\end{aligned}$$

Let us investigate the mean and the variance of this estimate against a numerical integrator (ground truth).

Monte Carlo Methods

Estimating tail probability




Figure: Monte Carlo estimation of the tail probability $X > 2$

In order to talk about estimators comfortably, we will now introduce various concepts of statistical estimators.

In order to talk about estimators comfortably, we will now introduce various concepts of statistical estimators.

- ▶ Error metrics (bias, variance, MSE, etc.)

In order to talk about estimators comfortably, we will now introduce various concepts of statistical estimators.

- ▶ Error metrics (bias, variance, MSE, etc.)
- ▶ Examples of estimating integrals, probabilities, and marginal likelihoods

In order to talk about estimators comfortably, we will now introduce various concepts of statistical estimators.

- ▶ Error metrics (bias, variance, MSE, etc.)
- ▶ Examples of estimating integrals, probabilities, and marginal likelihoods
- ▶ Explicit computations for variance

Monte Carlo Methods

Error metrics

We will give a general review about how to compute the error of a generic estimator.

Monte Carlo Methods

Error metrics

We will give a general review about how to compute the error of a generic estimator.

We will then apply these ideas to the Monte Carlo estimators.

Monte Carlo Methods

Error metrics: Bias

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

Monte Carlo Methods

Error metrics: Bias

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We define the bias:

$$\text{bias}_p(\hat{\varphi}^N) = \mathbb{E}[\hat{\varphi}^N] - \bar{\varphi}. \quad (14)$$

In the perfect Monte Carlo case (samples exactly from p), the bias is zero.

Monte Carlo Methods

Error metrics: Variance

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

Monte Carlo Methods

Error metrics: Variance

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We define the variance

$$\begin{aligned}\text{var}_p(\hat{\varphi}^N) &= \mathbb{E}_p \left[(\hat{\varphi}^N - \mathbb{E}_p[\hat{\varphi}^N])^2 \right], \\ &= \mathbb{E}_p \left[(\hat{\varphi}^N)^2 \right] - \mathbb{E}_p [\hat{\varphi}^N]^2.\end{aligned}$$

Monte Carlo Methods

Error metrics: Mean Squared Error (MSE)

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

Monte Carlo Methods

Error metrics: Mean Squared Error (MSE)

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We define the mean squared error:

$$\text{MSE}(\hat{\varphi}^N) = \mathbb{E}[(\hat{\varphi}^N - \bar{\varphi})^2].$$

Monte Carlo Methods

Error metrics: Mean Squared Error (MSE)

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We define the mean squared error:

$$\text{MSE}(\hat{\varphi}^N) = \mathbb{E}[(\hat{\varphi}^N - \bar{\varphi})^2].$$

Prove that (in-class exercise) this is also

$$\text{MSE}(\hat{\varphi}^N) = \text{var}_p(\hat{\varphi}^N) + \text{bias}_p(\hat{\varphi}^N)^2.$$

Monte Carlo Methods

Error metrics: Mean Squared Error (MSE)

Proof.

$$\text{MSE}(\hat{\varphi}^N) = \mathbb{E}[(\hat{\varphi}^N - \bar{\varphi})^2]$$

Monte Carlo Methods

Error metrics: Mean Squared Error (MSE)

Proof.

$$\begin{aligned}\text{MSE}(\hat{\varphi}^N) &= \mathbb{E}[(\hat{\varphi}^N - \bar{\varphi})^2] \\ &= \mathbb{E}[\hat{\varphi}^N]^2 - 2\bar{\varphi}\mathbb{E}[\hat{\varphi}^N] + \bar{\varphi}^2\end{aligned}$$

Monte Carlo Methods

Error metrics: Mean Squared Error (MSE)

Proof.

$$\begin{aligned}\text{MSE}(\hat{\varphi}^N) &= \mathbb{E}[(\hat{\varphi}^N - \bar{\varphi})^2] \\ &= \mathbb{E}[(\hat{\varphi}^N)^2] - 2\bar{\varphi}\mathbb{E}[\hat{\varphi}^N] + \bar{\varphi}^2 \\ &= \text{var}_p(\hat{\varphi}^N) + \mathbb{E}[\hat{\varphi}^N]^2 - 2\bar{\varphi}\mathbb{E}[\hat{\varphi}^N] + \bar{\varphi}^2\end{aligned}$$

Monte Carlo Methods

Error metrics: Mean Squared Error (MSE)

Proof.

$$\begin{aligned}\text{MSE}(\hat{\varphi}^N) &= \mathbb{E}[(\hat{\varphi}^N - \bar{\varphi})^2] \\ &= \mathbb{E}[(\hat{\varphi}^N)^2] - 2\bar{\varphi}\mathbb{E}[\hat{\varphi}^N] + \bar{\varphi}^2 \\ &= \text{var}_p(\hat{\varphi}^N) + \mathbb{E}[\hat{\varphi}^N]^2 - 2\bar{\varphi}\mathbb{E}[\hat{\varphi}^N] + \bar{\varphi}^2 \\ &= \text{var}_p(\hat{\varphi}^N) + (\mathbb{E}[\hat{\varphi}^N] - \bar{\varphi})^2,\end{aligned}$$

Monte Carlo Methods

Error metrics: Mean Squared Error (MSE)

Proof.

$$\begin{aligned}\text{MSE}(\hat{\varphi}^N) &= \mathbb{E}[(\hat{\varphi}^N - \bar{\varphi})^2] \\ &= \mathbb{E}[(\hat{\varphi}^N)^2] - 2\bar{\varphi}\mathbb{E}[\hat{\varphi}^N] + \bar{\varphi}^2 \\ &= \text{var}_p(\hat{\varphi}^N) + \mathbb{E}[\hat{\varphi}^N]^2 - 2\bar{\varphi}\mathbb{E}[\hat{\varphi}^N] + \bar{\varphi}^2 \\ &= \text{var}_p(\hat{\varphi}^N) + (\mathbb{E}[\hat{\varphi}^N] - \bar{\varphi})^2, \\ &= \text{var}_p(\hat{\varphi}^N) + \text{bias}_p(\hat{\varphi}^N)^2.\end{aligned}$$

Monte Carlo Methods

Error metrics: Root mean Squared Error (RMSE)

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

Monte Carlo Methods

Error metrics: Root mean Squared Error (RMSE)

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We define the root mean squared error:

$$\text{RMSE}(\hat{\varphi}^N) = \sqrt{\text{MSE}(\hat{\varphi}^N)}.$$

Monte Carlo Methods

Error metrics: Root mean Squared Error (RMSE)

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We define the root mean squared error:

$$\text{RMSE}(\hat{\varphi}^N) = \sqrt{\text{MSE}(\hat{\varphi}^N)}.$$

For the **unbiased case**,

$$\text{RMSE}(\hat{\varphi}^N) = \sqrt{\text{var}_p(\hat{\varphi}^N)} = \text{std}_p(\hat{\varphi}^N).$$

Note though we will have biased cases.

Monte Carlo Methods

Error metrics: Root mean Squared Error (RMSE)

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We define the root mean squared error:

$$\text{RMSE}(\hat{\varphi}^N) = \sqrt{\text{MSE}(\hat{\varphi}^N)}.$$

For the **unbiased case**,

$$\text{RMSE}(\hat{\varphi}^N) = \sqrt{\text{var}_p(\hat{\varphi}^N)} = \text{std}_p(\hat{\varphi}^N).$$

Note though we will have biased cases. Do not use them interchangeably.

Monte Carlo Methods

Error metrics: Relative Absolute Error

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

Monte Carlo Methods

Error metrics: Relative Absolute Error

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We generally plot

$$\text{RAE}(\hat{\varphi}^N) = \frac{|\hat{\varphi}^N - \bar{\varphi}|}{|\bar{\varphi}|}.$$

Note that this is *random*, as no expectations are taken.

Monte Carlo Methods

Error metrics: Relative Absolute Error

Consider

- ▶ the target quantity: $\bar{\varphi}$
- ▶ the estimator: $\hat{\varphi}^N$

We generally plot

$$\text{RAE}(\hat{\varphi}^N) = \frac{|\hat{\varphi}^N - \bar{\varphi}|}{|\bar{\varphi}|}.$$

Note that this is *random*, as no expectations are taken.

It can be proven that the random error has

$$|\hat{\varphi}^N - \bar{\varphi}| \leq \mathcal{O}(U/\sqrt{N}),$$

see, e.g., Akyildiz, 2019, Corollary 2.1 for a proof.

Monte Carlo Methods

More Examples: Estimating Marginal Likelihood

Consider the following model:

$$\begin{aligned} p(x) &= \mathcal{N}(x; \mu_0, \sigma_0^2), \\ p(y|x) &= \mathcal{N}(y; x, \sigma^2). \end{aligned}$$

Assume that we have observed $y = 4$ and want to estimate the marginal likelihood $p(y)$.

Monte Carlo Methods

More Examples: Estimating Marginal Likelihood

Consider the following model:

$$\begin{aligned} p(x) &= \mathcal{N}(x; \mu_0, \sigma_0^2), \\ p(y|x) &= \mathcal{N}(y; x, \sigma^2). \end{aligned}$$

Assume that we have observed $y = 4$ and want to estimate the marginal likelihood $p(y)$.

From our past experience, we have access to this density:

$$\begin{aligned} p(y) &= \int p(y|x)p(x)dx, \\ &= \mathcal{N}(y; \mu_0, \sigma_0^2 + \sigma^2). \end{aligned}$$

Therefore, we know exactly the value of $p(4)$ (i.e. $p(y = 4)$).

Monte Carlo Methods

More Examples: Estimating Marginal Likelihood

Given that the problem is an integral w.r.t. $p(x)$,

$$p(y) = \int p(y|x)p(x)dx,$$

we could use our Monte Carlo estimator to get unbiased estimates of $p(y)$.

Monte Carlo Methods

More Examples: Estimating Marginal Likelihood

Given that the problem is an integral w.r.t. $p(x)$,

$$p(y) = \int p(y|x)p(x)dx,$$

we could use our Monte Carlo estimator to get unbiased estimates of $p(y)$.

Write down the estimator (in-class exercise).

Monte Carlo Methods

More Examples: Estimating Marginal Likelihood

Given that the problem is an integral w.r.t. $p(x)$,

$$p(y) = \int p(y|x)p(x)dx,$$

we could use our Monte Carlo estimator to get unbiased estimates of $p(y)$.

Monte Carlo Methods

More Examples: Estimating Marginal Likelihood

Given that the problem is an integral w.r.t. $p(x)$,

$$p(y) = \int p(y|x)p(x)dx,$$

we could use our Monte Carlo estimator to get unbiased estimates of $p(y)$.

Note here that $\varphi(x) = p(y|x)$ is a deterministic function of x for fixed y . Therefore, given $X_1, \dots, X_N \sim p(x)$

$$\int \varphi(x)p(x)dx \approx \frac{1}{N} \sum_{i=1}^N \varphi(X_i) = \frac{1}{N} \sum_{i=1}^N p(y|X_i).$$

For $y = 4$, we can compute this (given μ_0, σ_0, σ).

Monte Carlo Methods

More Examples: Estimating Marginal Likelihood

Set

$$\mu_0 = 0, \quad \sigma_0 = 1, \quad \sigma = 2, \quad y = 1$$

and generate $N = 100000$ samples from $p(x)$. We estimate

$$p^N(y) = \frac{1}{N} \sum_{i=1}^N p(y|X_i).$$

Monte Carlo Methods

More Examples: Estimating Marginal Likelihood




Figure: Estimating the marginal likelihood $p(y)$ for $y = 1$

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

Let

$$p(x) = \frac{1}{\pi(1 + x^2)},$$

and let

$$\varphi(x) = \mathbf{1}_{\{x \geq 2\}}.$$

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

Let

$$p(x) = \frac{1}{\pi(1+x^2)},$$

and let

$$\varphi(x) = \mathbf{1}_{\{x \geq 2\}}.$$

We would like to compute

$$\begin{aligned}\mathbb{P}(X > 2) &= \int_2^\infty p(x)dx \\ &= \int \mathbf{1}_{\{x>2\}}(x)p(x)dx \\ &= \int \varphi(x)p(x)dx.\end{aligned}$$

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

We can compute the real value of this integral:

$$I = \bar{\varphi} = \int_2^{\infty} p(x)dx = F(\infty) - F(2) = \frac{1}{2} - \frac{1}{\pi} \tan^{-1}(2) = 0.1476$$

Compute the variance of the Monte Carlo estimator for $N = 10$ samples.

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

We can compute the real value of this integral:

$$I = \bar{\varphi} = \int_2^{\infty} p(x)dx = F(\infty) - F(2) = \frac{1}{2} - \frac{1}{\pi} \tan^{-1}(2) = 0.1476$$

Compute the variance of the Monte Carlo estimator for $N = 10$ samples.

$$\text{var}(\hat{\varphi}^N) = \frac{\text{var}_p(\varphi)}{N}$$

So we need to compute:

$$\text{var}_p(\varphi)$$

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

Now, we compute

$$\begin{aligned}\text{var}_p(\varphi) &= \int \varphi(x)^2 p(x) dx - \left(\int \varphi(x) p(x) dx \right)^2 \\ &= \int \mathbf{1}_{\{x>2\}}(x)^2 p(x) dx - \left(\int \mathbf{1}_{\{x>2\}}(x) p(x) dx \right)^2 \\ &= \int \mathbf{1}_{\{x>2\}}(x) p(x) dx - \left(\int \mathbf{1}_{\{x>2\}}(x) p(x) dx \right)^2 \\ &= 0.1476 - 0.1476^2 = 0.125.\end{aligned}$$

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

Now, we compute

$$\begin{aligned}\text{var}_p(\varphi) &= \int \varphi(x)^2 p(x) dx - \left(\int \varphi(x) p(x) dx \right)^2 \\&= \int \mathbf{1}_{\{x>2\}}(x)^2 p(x) dx - \left(\int \mathbf{1}_{\{x>2\}}(x) p(x) dx \right)^2 \\&= \int \mathbf{1}_{\{x>2\}}(x) p(x) dx - \left(\int \mathbf{1}_{\{x>2\}}(x) p(x) dx \right)^2 \\&= 0.1476 - 0.1476^2 = 0.125.\end{aligned}$$

The variance of the estimator then

$$\text{var}(\hat{\varphi}^N) = \frac{0.125}{10} = 0.0125.$$

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

Could we do better? An idea, the density is symmetric around zero

- ▶ This means $P(X > 2) = P(X < -2)$.




Figure: Cauchy density

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

We could compute:

$$\mathbb{P}(|X| > 2) = \mathbb{P}(X > 2) + \mathbb{P}(X < -2) = 2I.$$

Therefore, our new problem is $I = \frac{1}{2}\mathbb{P}(|X| > 2)$.

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

We could compute:

$$\mathbb{P}(|X| > 2) = \mathbb{P}(X > 2) + \mathbb{P}(X < -2) = 2I.$$

Therefore, our new problem is $I = \frac{1}{2}\mathbb{P}(|X| > 2)$.

Write it as

$$\begin{aligned} I &= \frac{1}{2} \int_{|x|>2} p(x)dx, \\ &= \int \frac{1}{2} \mathbf{1}_{\{|x|>2\}}(x)p(x)dx, \end{aligned}$$

Now define the test function

$$\varphi(x) = \frac{1}{2} \mathbf{1}_{\{|x|>2\}}(x).$$

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

We need to compute $\text{var}_p(\varphi)$.

Monte Carlo Methods

More Examples: Compute variance of a probability estimate

We need to compute $\text{var}_p(\varphi)$.


$$\begin{aligned}\text{var}_p(\varphi) &= \int \varphi(x)^2 p(x) dx - \left(\int \varphi(x) p(x) dx \right)^2 \\ &= \int \frac{1}{4} \mathbf{1}_{\{|x|>2\}}^2 p(x) dx - \left(\int \frac{1}{2} \mathbf{1}_{\{|x|>2\}} p(x) dx \right)^2 \\ &= \int \frac{1}{4} \mathbf{1}_{\{|x|>2\}} p(x) dx - \left(\int \frac{1}{2} \mathbf{1}_{\{|x|>2\}} p(x) dx \right)^2 \\ &= \frac{1}{4} \times 2 \times 0.1476 - \frac{1}{4} \times (2 \times 0.1476)^2, \\ &= 0.052.\end{aligned}$$

Therefore, the variance of the estimator for $N = 10$ samples is

$$\text{var}(\hat{\varphi}^N) = \frac{0.052}{10} = 0.0052.$$

Improvement over the previous estimator! That was 0.0125.

See you next week!



"Horizontal textures, randomness, city lights, fog" (Stable Diffusion)

- ① Robert, Christian P and George Casella (2004). *Monte Carlo statistical methods*. Springer.
- ② Akyildiz, Omer Deniz (2019). “Sequential and adaptive Bayesian computation for inference and optimization”. Can be accessed from: <http://akyildiz.me/works/thesis.pdf>. PhD thesis. Universidad Carlos III de Madrid.