

Mathematical Logic (MATH6/70132; P65), Problem Sheet 1

[1] Let p denote 'I will pass this course,' q denote 'I do my homework regularly' and r denote 'I am lucky.' Express in symbolic form each of the following propositions:

- I will pass this course only if I do my homework regularly.
- Doing homework regularly is a necessary condition for me to pass this course.
- If I do my homework regularly and I do not pass this course then I am unlucky.
- If I do not do my homework regularly and I pass this course then I am lucky.

[2] Show that ϕ and ψ are logically equivalent if and only if $(\phi \leftrightarrow \psi)$ is a tautology.

[3] Suppose ϕ and ψ are logically equivalent propositional formulas whose propositional variables are amongst p_1, \dots, p_n . Let η_1, \dots, η_n be formulas and let θ and χ be the formulas obtained by replacing each occurrence of p_i by η_i in ϕ and ψ respectively. Show that θ and χ are logically equivalent.

[4] Suppose p, q, r are propositional variables. For each of the following formulas find a formula in disjunctive normal form (– see 1.1.9 in notes) which is logically equivalent to it.

- $((p \rightarrow q) \rightarrow ((\neg p) \wedge q))$.
- $(\neg((p \rightarrow q) \rightarrow r))$.

ϕ	ψ	$(\phi \mid \psi)$
T	T	F
T	F	T
F	T	T
F	F	T

[5] The connective \mid has truth table:

- Show that this connective is adequate.
- Show that \mid and the NOR connective \downarrow are the only binary connectives which are adequate.

[6] How many truth functions f of n propositional variables are there with the property that $f(T, T, \dots, T) = T$? Can all of these be expressed as the truth function of a formula of n variables using only the connectives \wedge, \vee ? Explain your answer.

[7] (i) For which values of the propositional variables p_1, p_2, p_3 does the following propositional formula θ have truth value F :

$$((p_1 \rightarrow ((\neg p_2) \rightarrow p_3)) \rightarrow ((p_3 \rightarrow p_2) \rightarrow p_1))?$$

Find a formula in disjunctive normal form which is logically equivalent to $(\neg\theta)$.

(ii) Find a formula χ with three propositional variables p_1, p_2, p_3 whose truth value is T if and only if p_1, p_2, p_3 have truth values T, F, T (respectively). Justify your answer.

[8] Below is the start of a proof that if ϕ, ψ are propositional formulas, then $((\neg\psi) \rightarrow (\psi \rightarrow \phi))$ is a theorem (of L). Write this out again, but at each stage give the reasoning, and then complete the proof:

- $((\neg\psi) \rightarrow ((\neg\phi) \rightarrow (\neg\psi)))$
- $((((\neg\phi) \rightarrow (\neg\psi)) \rightarrow (\psi \rightarrow \phi)))$

Denote this formula by χ

- $(\chi \rightarrow ((\neg\psi) \rightarrow \chi))$