

# Notes on NEWUOA

Zaikun Zhang \*

June 13, 2021 12:02pm

We define  $\bar{\mathcal{X}}_k$  to be a point in  $\mathcal{X}_k$  such that

$$f(\bar{\mathcal{X}}_k) = \min\{f(x) : x \in \mathcal{X}_k\}. \quad (0.1) \text{?eq:xopt?}$$

If multiple points attain the minimum, then we take the earliest one visited by the algorithm.

---

## Algorithm 0.1 OPTimization based on Interpolation Models (OPTIM)

---

Input  $\Delta_0 \in (0, +\infty)$ ,  $\tau > 0$ ,  $m \in \{n+2, n+3, \dots, (n+1)(n+2)/2\}$ , and  $\mathcal{X}_0 \subset \mathbb{R}^n$  with  $|\mathcal{X}_0| = m$  and  $\kappa(\mathcal{X}_0) \leq \kappa_0$ . Set  $k = 0$ .

1. **Model construction.** Pick  $Q_k \in \{Q : Q(x) = f(x) \text{ for all } x \in \mathcal{X}_k\}$ .
2. **Trust-region step evaluation.** Calculate

$$x_k^+ \approx \operatorname{argmin}\{Q_k(x) : \|x - \bar{\mathcal{X}}_k\| \leq \Delta_k\}. \quad (0.2) \text{?eq:xget?}$$

If  $\|x_k^+ - \bar{\mathcal{X}}_k\| < \alpha\Delta_k$ , then set  $\Delta_{k+1} = \theta\Delta_k$ . Otherwise, update  $\Delta_k$  to  $\Delta_{k+1}$  according to  $r_k = [f(\bar{\mathcal{X}}_k) - f(x_k^+)]/[Q_k(\bar{\mathcal{X}}_k) - Q_k(x_k^+)]$ .

3. **Interpolation set update.** If  $\|x_k^+ - \bar{\mathcal{X}}_k\| \geq \alpha\Delta_k$ , then calculate

$$x_k^- \approx \operatorname{argmin}\{\kappa(\mathcal{X}_k \cup x_k^+ \setminus x) : x \in \mathcal{X}_k\}, \quad (0.3) \text{?eq:xdrop?}$$

and set  $\mathcal{X}_{k+1} = \mathcal{X}_k \cup x_k^+ \setminus x_k^-$  if  $r_k > 0$  or  $\kappa(\mathcal{X}_k \cup x_k^+ \setminus x_k^-) \leq \kappa_0$ .

4. **Geometry improvement.** If  $\|x_k^+ - \bar{\mathcal{X}}_k\| < \alpha\Delta_k$ , or  $\|x_k^+ - \bar{\mathcal{X}}_k\| > \alpha\Delta_k$  but  $r_k \leq 0$  and  $\kappa(\mathcal{X}_k \cup x_k^+ \setminus x_k^-) > \kappa_0$ , then calculate

$$y_k^- = \operatorname{argmax}\{\|y - \bar{\mathcal{X}}_k\| : y \in \mathcal{X}_k\}, \quad (0.4) \text{?eq:ydrop?}$$

$$y_k^+ \approx \operatorname{argmin}\{\kappa(\mathcal{X}_k \cup y \setminus y_k^-) : \|y - \bar{\mathcal{X}}_k\| \leq \Delta_k\}, \quad (0.5) \text{?eq:yget?}$$

and set  $\mathcal{X}_{k+1} = \mathcal{X}_k \cup y_k^+ \setminus y_k^-$ .

---



---

\*Hong Kong Polytechnic University, [zaikun.zhang@polyu.edu.hk](mailto:zaikun.zhang@polyu.edu.hk)

---

**Algorithm 0.2** NEWUOA

Input  $\Delta_0 \in (0, +\infty)$ ,  $\tau > 0$ ,  $m \in \{n+2, n+3, \dots, (n+1)(n+2)/2\}$ , and  $\mathcal{X}_0 \subset \mathbb{R}^n$  with  $|\mathcal{X}_0| = m$  and  $\kappa(\mathcal{X}_0) \leq \kappa_0$ . Define  $Q_0 = \operatorname{argmin}\{\|\nabla Q\|_F : Q \in \mathcal{Q} \text{ and } Q(x) = f(x) \text{ for all } x \in \mathcal{X}_0\}$ . Set  $k = 0$ .

1. **Trust-region step evaluation.** Set

$$x_k^+ \approx \operatorname{argmin}\{Q_k(x) : \|x - \bar{\mathcal{X}}_k\| \leq \Delta_k\}. \quad (0.6) \text{ ?eq:xgetn?}$$

$$S = \mathbb{1}(\|x_k^+ - \bar{\mathcal{X}}_k\| < \rho_k/2), \quad (0.7) \text{ ?eq:shortd?}$$

$$R = \mathbb{1}(S = 1 \text{ and the errors in recent models are small}). \quad (0.8) \text{ ?eq:redrho?}$$

If  $R = 1$ , then set  $\mathcal{X}_{k+1} = \mathcal{X}_k$ ,  $Q_{k+1} = Q_k$ , and go to step 4. If  $S = 1$  and  $R = 0$ , then set  $\Delta_{k+1} = \max\{\Delta_k/10, \rho_k\}$ . If  $S = 0$ , then evaluate  $r_k = [f(\bar{\mathcal{X}}_k) - f(x_k^+)]/[Q_k(\bar{\mathcal{X}}_k) - Q_k(x_k^+)]$  and update  $\Delta_k$  to  $\Delta_{k+1}$  according to  $r_k$ .

2. **Interpolation set and model update.** If  $S = 0$ , then calculate

$$x_k^- \approx \operatorname{argmin}\{\kappa(\mathcal{X}_k \cup x_k^+ \setminus x) : x \in \mathcal{X}_k\}, \quad (0.9) \text{ ?eq:xdropn?}$$

and let  $\mathcal{X}'_k = \mathcal{X}_k \cup x_k^+ \setminus x_k^-$  if  $r_k > 0$  or  $\kappa(\mathcal{X}_k \cup x_k^+ \setminus x_k^-) \leq \kappa_0$ . In any other case,  $\mathcal{X}'_k = \mathcal{X}_k$ . Set

$$Q'_k = \operatorname{argmin}\{\|Q - Q_k\|_F : Q \in \mathcal{Q} \text{ and } Q(x) = f(x) \text{ for all } x \in \hat{\mathcal{X}}_k\}. \quad (0.10) \text{ ?eq:updateeq1?}$$

3. **Geometry improvement.** If  $S = 1$  or  $r_k < 1/10$ , then set

$$y_k^- = \operatorname{argmax}\{\|y - \bar{\mathcal{X}}'_k\| : y \in \mathcal{X}'_k\}. \quad (0.11) \text{ ?eq:ydropn?}$$

If  $\|y_k^- - \bar{\mathcal{X}}'_k\| \geq 2\Delta_{k+1}$ , then define  $\Delta'_k = \max\{\rho_k, \min\{\|y_k^- - \bar{\mathcal{X}}'_k\|/10, \Delta_{k+1}/2\}\}$ , calculate

$$y_k^+ \approx \operatorname{argmin}\{\kappa(\mathcal{X}'_k \cup y \setminus y_k^-) : \|y - \bar{\mathcal{X}}'_k\| \leq \Delta'_k\}, \quad (0.12) \text{ ?eq:ygetn?}$$

and set  $\mathcal{X}_{k+1} = \mathcal{X}'_k \cup y_k^+ \setminus y_k^-$ .

4. **Resolution enhancement.** If  $R = 1$ , then reduce  $\rho_k$  by about a factor of 10 to obtain  $\rho_{k+1}$ , and set  $\Delta_{k+1} = \max\{\rho_{k+1}/2, \rho_k\}$ . If  $R = 0$ , then set  $\rho_{k+1} = \rho_k$ .

---