

# ACLIM2 CMIP6 ROMSNPZ Indices quick start guide

K. Holsman

## Contents

<b>Download the ACLIM2 repo &amp; data</b>	<b>2</b>
Clone the ACLIM2 repo . . . . .	2
Option 1: Use R . . . . .	2
Option 2: Download the zipped repo . . . . .	2
Option 3: Use git commandline . . . . .	3
Get the data . . . . .	3
Set up the Workspace . . . . .	4
<b>Read this before you start</b>	<b>5</b>
Overview . . . . .	5
ROMSNPZ versions . . . . .	5
ROMSNPZ variables . . . . .	6
Data outputs . . . . .	8
<b>Indices &amp; bias correction</b>	<b>9</b>
Weekly indices . . . . .	10
Monthly indices . . . . .	11
Seasonal indices . . . . .	11
Annual indices . . . . .	12
Annual survey rep. indices . . . . .	12
<b>Plot &amp; concat Indices</b>	<b>12</b>
NRS indices (André) . . . . .	13
monthly indices (Andy) . . . . .	19
weekly indices (Jon) . . . . .	21
<b>Output to .dat file (ADMB/ TMB users)</b>	<b>23</b>
Use R to make .dat file . . . . .	23
<b>APPENDIX A: Create &amp; bias correct ACLIM2 indices</b>	<b>28</b>

## Download the ACLIM2 repo & data

### Clone the ACLIM2 repo

To run this tutorial first clone the ACLIM2 repository to your local drive:

#### Option 1: Use R

This set of commands, run within R, downloads the ACLIM2 repository and unpacks it, with the ACLIM2 directory structure being located in the specified `download_path`. This also performs the folder renaming mentioned in Option 2.

```
# Specify the download directory
main_nm      <- "ACLIM2"

# Note: Edit download_path for preference
download_path <- path.expand("~/")
dest_fldr    <- file.path(download_path,main_nm)

url          <- "https://github.com/kholsman/ACLIM2/archive/main.zip"
dest_file    <- file.path(download_path,paste0(main_nm,".zip"))
download.file(url=url, destfile=dest_file)

# unzip the .zip file (manually unzip if this doesn't work)
setwd(download_path)
unzip (dest_file, exdir = download_path,overwrite = T)

#rename the unzipped folder from ACLIM2-main to ACLIM2
file.rename(paste0(main_nm,"-main"), main_nm)
setwd(main_nm)
```

#### Option 2: Download the zipped repo

Download the full zip archive directly from the **ACLIM2 Repo** using this link: <https://github.com/kholsman/ACLIM2> and unzip its contents while preserving directory structure.

**Important!** If downloading from zip, please **rename the root folder** from ACLIM2-main (in the zipfile) to ACLIM2 (name used in cloned copies) after unzipping, for consistency in the following examples.

Your final folder structure should look like this:

Name	Date Modified	Size	Kind
ACLIM2.Rproj	Today at 10:02 AM	205 bytes	R Project
> Data	Today at 10:08 AM	--	Folder
> Docs	Oct 26, 2021 at 4:12 PM	--	Folder
> Figs	Oct 26, 2021 at 3:42 PM	--	Folder
GettingStarted_Beri...10K_ROMSNPZ.html	Today at 11:55 AM	10.9 MB	HTML text
GettingStarted_Bering10K_ROMSNPZ.md	Today at 11:55 AM	84 KB	Markdown File
GettingStarted_Bering10K_ROMSNPZ.pdf	Today at 11:55 AM	7.2 MB	PDF Document
GettingStarted_Beri...10K_ROMSNPZ.Rmd	Today at 11:55 AM	87 KB	R Markdown File
GettingStarted_Bering10K_ROMSNPZ.tex	Today at 11:55 AM	171 KB	TeX File
plot_ts.R	Oct 26, 2021 at 11:14 AM	5 KB	R Source File
> R	Today at 2:46 PM	--	Folder
README.md	Today at 11:55 AM	84 KB	Markdown File
trash	Oct 26, 2021 at 2:09 PM	--	Folder
> untitled folder	Apr 5, 2022 at 11:46 AM	--	Folder
> Vignettes	Today at 3:03 PM	--	Folder

### Option 3: Use git commandline

If you have git installed and can work with it, this is the preferred method as it preserves all directory structure and can aid in future updating. Use this from a **terminal command line, not in R**, to clone the full ACLIM2 directory and sub-directories:

```
git clone https://github.com/kholsman/ACLIM2.git
```

## Get the data

**Step 1)** \* Go to the google drive and download the zipped file with the R data 2022\_03\_07\_Rdata.zip:

- 00\_ACLIM\_shared > 02\_Data > Newest\_Data(use this) > 2022\_03\_07\_Rdata.zip
- Unzip the folder and move the contents of the zipped folder to your local folder ACLIM2/Data/in.

**Step 2)** \* Go to the google drive and download the zipped file with the R ACLIM2 indices ACLIM2\_indices.zip:

- 00\_ACLIM\_shared > 02\_Data > Newest\_Data(use this) > 2022\_05\_16\_unzip\_and\_put\_contents\_in\_the\_Data\_o
- Unzip the folder and move the contents of the zipped folder to your local folder ACLIM2/Data/out.

## Set up the Workspace

Open R() and used 'setwd()' to navigate to the root ACLIM2 folder (.e.g, ~/mydocuments/ACLIM2)

```
# set the workspace to your local ACLIM2 folder
# e.g., "/Users/kholsman/Documents/GitHub/ACLIM2"
# setwd( path.expand("~/Documents/GitHub/ACLIM2") )

# -----
# SETUP WORKSPACE
tmstp <- format(Sys.time(), "%Y_%m_%d")
main  <- getwd()  #"~/GitHub_new/ACLIM2"

# loads packages, data, setup, etc.
suppressWarnings(source("R/make.R"))
```

```
## -----
## ALIM2/R/setup.R settings
## -----
## data_path          : /Volumes/LaCie/romsnpz/roms_for_public
## Rdata_path         : /Volumes/LaCie/romsnpz/2022_05_16_Rdata/roms_for_public
## redownload_level3_mox: FALSE
## update.figs        : FALSE
## load_gis           : FALSE
## update.outputs     : TRUE
## update.figs        : FALSE
## dpiIN              : 150
## update.figs        : FALSE
## -----
## -----
##
## The following datasets are public, please cite as Hermann et al. 2019 (v.H16) and Kearney et al. 2020
## B10K-H16_CMIP5_CESM_BIO_rcp85
## B10K-H16_CMIP5_CESM_rcp45
## B10K-H16_CMIP5_CESM_rcp85
## B10K-H16_CMIP5_GFDL_BIO_rcp85
## B10K-H16_CMIP5_GFDL_rcp45
## B10K-H16_CMIP5_GFDL_rcp85
## B10K-H16_CMIP5_MIROC_rcp45
## B10K-H16_CMIP5_MIROC_rcp85
## B10K-H16_CORECFS
## B10K-K20_CORECFS
##
## The following datasets are still under embargo, please do not share outside of ACLIM:
## B10K-K20P19_CMIP6_cesm_historical
## B10K-K20P19_CMIP6_cesm_ssp126
## B10K-K20P19_CMIP6_cesm_ssp585
## B10K-K20P19_CMIP6_gfdl_historical
## B10K-K20P19_CMIP6_gfdl_ssp126
## B10K-K20P19_CMIP6_gfdl_ssp585
## B10K-K20P19_CMIP6_miroc_historical
## B10K-K20P19_CMIP6_miroc_ssp126
## B10K-K20P19_CMIP6_miroc_ssp585
```

---

## Read this before you start

### Overview

The **ACLIM2 github repository** contains R code and Rdata files for working with netcdf-format data generated from the **downscaled ROMSNPZ modeling** of the ROMSNPZ Bering Sea Ocean Modeling team; Drs. Hermann, Cheng, Kearney, Pilcher, Ortiz, and Aydin. The code and R resources described in this tutorial are maintained by Kirstin Holsman as part of NOAA's **ACLIM project** for the Bering Sea. *See Hollowed et al. 2020 for more information about the ACLIM project.*

---

This document provides an overview of accessing, plotting, and creating bias corrected indices for ACLIM2 based on CMIP6 (embargoed for ACLIM2 users until 2023) and CMIP5 (publicly available) simulations. This guide assumes analyses will take place in R() and that users have access to the data folder within the ACLIM2 shared drive. For more information also see the full tutorial ("GettingStarted\_Bering10K\_ROMSNPZ" available at the bottom of **this repo page**).

**Important!** A few key things to know before getting started are detailed below. Please review this information before getting started.

### ROMSNPZ versions

**Important!** ACLIM1 CMIP5 and ACLIM2 CMIP5 and CMIP6 datasets use different base models.

There are two versions of the ROMSNPZ model:

1. ACLIM1 an older 10-depth layer model used for CMIP5 ("H-16")
2. ACLIM2 a new 30-depth layer model used for CMIP6 ("K20" or "K20P19")

The models are not directly comparable, therefore the projections should be bias corrected and recentered to baselines of hindcasts of each model (forced by "observed" climate conditions). i.e. CMIP5 and CMIP6 have corresponding hindcasts:

1. Hindcast for CMIP5 "H19" -> H16\_CORECFS
2. Hindcast for CMIP5 "K20P19" -> H16\_CORECFS
3. Hindcast for CMIP6 "K20P19" -> K20\_CORECFS

In addition for CMIP6 "historical" runs are available for bias correcting. We will use those below.

For a list of the available simulations for ACLIM enter the following in R():

```
# list of the climate scenarios
data.frame(sim_list)
```

```
##                                sim_list
## 1                      B10K-K20_CORECFS
## 2      B10K-H16_CMIP5_CESM_BIO_rcp85
## 3      B10K-H16_CMIP5_CESM_rcp45
```

```

## 4      B10K-H16_CMIP5_CESM_rcp85
## 5      B10K-H16_CMIP5_GFDL_BIO_rcp85
## 6      B10K-H16_CMIP5_GFDL_rcp45
## 7      B10K-H16_CMIP5_GFDL_rcp85
## 8      B10K-H16_CMIP5_MIROC_rcp45
## 9      B10K-H16_CMIP5_MIROC_rcp85
## 10     B10K-H16_CORECFS
## 11     B10K-K20P19_CMIP5_CESM_rcp45
## 12     B10K-K20P19_CMIP5_CESM_rcp85
## 13     B10K-K20P19_CMIP5_GFDL_rcp45
## 14     B10K-K20P19_CMIP5_GFDL_rcp85
## 15     B10K-K20P19_CMIP5_MIROC_rcp45
## 16     B10K-K20P19_CMIP5_MIROC_rcp85
## 17     B10K-K20P19_CMIP6_cesm_historical
## 18     B10K-K20P19_CMIP6_cesm_ssp126
## 19     B10K-K20P19_CMIP6_cesm_ssp585
## 20     B10K-K20P19_CMIP6_gfdl_historical
## 21     B10K-K20P19_CMIP6_gfdl_ssp126
## 22     B10K-K20P19_CMIP6_gfdl_ssp585
## 23     B10K-K20P19_CMIP6_miroc_historical
## 24     B10K-K20P19_CMIP6_miroc_ssp126
## 25     B10K-K20P19_CMIP6_miroc_ssp585

```

## ROMSNPZ variables

For a list of the available variables from the ROMSNPZ:

```

# Metadata for variables
(srvy_var_def[-(1:5),])

```

```

##          name          units
## 6          Ben      mg C m^-2
## 7        DetBen      mg C m^-2
## 8          Hsbl      meter
## 9        IceNH4    mmol N m^-3
## 10       IceNO3    mmol N m^-3
## 11       IcePhL      mg C m^-3
## 12         aice
## 13         hice      meter
## 14       shflux    watt meter-2
## 15       ssflux    meter second-1
## 16   Cop_integrated    (mg C m^-3)*m
## 17   Cop_surface5m      mg C m^-3
## 18   Eup0_integrated    (mg C m^-3)*m
## 19   Eup0_surface5m      mg C m^-3
## 20   EupS_integrated    (mg C m^-3)*m
## 21   EupS_surface5m      mg C m^-3
## 22   Iron_bottom5m    micromol Fe m-3
## 23   Iron_integrated    (micromol Fe m-3)*m
## 24   Iron_surface5m    micromol Fe m-3
## 25   Jel_integrated    (mg C m^-3)*m
## 26   Jel_surface5m      mg C m^-3
## 27   MZL_integrated    (mg C m^-3)*m

```

```

## 28      MZL_surface5m      mg C m-3
## 29      NCaO_integrated      (mg C m-3)*m
## 30      NCaO_surface5m      mg C m-3
## 31      NCaS_integrated      (mg C m-3)*m
## 32      NCaS_surface5m      mg C m-3
## 33      NH4_bottom5m      mmol N m-3
## 34      NH4_integrated      (mmol N m-3)*m
## 35      NH4_surface5m      mmol N m-3
## 36      NO3_bottom5m      mmol N m-3
## 37      NO3_integrated      (mmol N m-3)*m
## 38      NO3_surface5m      mmol N m-3
## 39      PhL_integrated      (mg C m-3)*m
## 40      PhL_surface5m      mg C m-3
## 41      PhS_integrated      (mg C m-3)*m
## 42      PhS_surface5m      mg C m-3
## 43      prod_Cop_integrated      mg C m-2 d-1
## 44      prod_Eup0_integrated      mg C m-2 d-1
## 45      prod_EupS_integrated      mg C m-2 d-1
## 46      prod_Eup_integrated      (milligram carbon meter-3 d-1)*m
## 47      prod_Jel_integrated      mg C m-2 d-1
## 48      prod_MZL_integrated      mg C m-2 d-1
## 49      prod_NCaO_integrated      mg C m-2 d-1
## 50      prod_NCaS_integrated      mg C m-2 d-1
## 51      prod_NCa_integrated      (milligram carbon meter-3 d-1)*m
## 52      prod_PhL_integrated      mg C m-2 d-1
## 53      prod_PhS_integrated      mg C m-2 d-1
## 54      salt_surface5m
## 55      temp_bottom5m      Celsius
## 56      temp_integrated      (Celsius)*m
## 57      temp_surface5m      Celsius
## 58      uEast_bottom5m      meter second-1
## 59      uEast_surface5m      meter second-1
## 60      vNorth_bottom5m      meter second-1
## 61      vNorth_surface5m      meter second-1
##                                     longname
## 6      Benthic infauna concentration
## 7      Benthic detritus concentration
## 8      depth of oceanic surface boundary layer
## 9      Ice ammonium concentration
## 10     Ice nitrate concentration
## 11     Ice algae concentration
## 12     fraction of cell covered by ice
## 13     average ice thickness in cell
## 14     surface net heat flux
## 15     surface net salt flux, (E-P)*SALT
## 16     Small copepod concentration, integrated over depth
## 17     Small copepod concentration, surface 5m mean
## 18     Offshore euphausiid concentration, integrated over depth
## 19     Offshore euphausiid concentration, surface 5m mean
## 20     On-shelf euphausiid concentration, integrated over depth
## 21     On-shelf euphausiid concentration, surface 5m mean
## 22     iron concentration, bottom 5m mean
## 23     iron concentration, integrated over depth
## 24     iron concentration, surface 5m mean

```

```

## 25         Jellyfish concentration, integrated over depth
## 26         Jellyfish concentration, surface 5m mean
## 27         Microzooplankton concentration, integrated over depth
## 28         Microzooplankton concentration, surface 5m mean
## 29 Offshore large copepod concentration, integrated over depth
## 30         Offshore large copepod concentration, surface 5m mean
## 31 On-shelf large copepod concentration, integrated over depth
## 32         On-shelf large copepod concentration, surface 5m mean
## 33         Ammonium concentration, bottom 5m mean
## 34         Ammonium concentration, integrated over depth
## 35         Ammonium concentration, surface 5m mean
## 36         Nitrate concentration, bottom 5m mean
## 37         Nitrate concentration, integrated over depth
## 38         Nitrate concentration, surface 5m mean
## 39 Large phytoplankton concentration, integrated over depth
## 40         Large phytoplankton concentration, surface 5m mean
## 41 Small phytoplankton concentration, integrated over depth
## 42         Small phytoplankton concentration, surface 5m mean
## 43         Cop net production rate, summed over depth
## 44         Eup0 net production rate, summed over depth
## 45         EupS net production rate, summed over depth
## 46 secondary production Euphausiids, integrated over depth
## 47         Jel net production rate, summed over depth
## 48         MZL net production rate, summed over depth
## 49         NCaO net production rate, summed over depth
## 50         NCaS net production rate, summed over depth
## 51 secondary production Neocalanus, integrated over depth
## 52         PhL net production rate, summed over depth
## 53         PhS net production rate, summed over depth
## 54         salinity, surface 5m mean
## 55         potential temperature, bottom 5m mean
## 56         potential temperature, integrated over depth
## 57         potential temperature, surface 5m mean
## 58         u-momentum component, geo-rotated, bottom 5m mean
## 59         u-momentum component, geo-rotated, surface 5m mean
## 60         v-momentum component, geo-rotated, bottom 5m mean
## 61         v-momentum component, geo-rotated, surface 5m mean

```

## Data outputs

**Important!** There are 2 types of post-processed data available for use in ACLIM.

The ROMSNPZ team has developed a process to provide standardized post-processed outputs from the large (and non-intuitive) ROMSNPZ grid. These have been characterized as:

1. Level 1 (original ROMSNPZ U,V, grid, not rotated or corrected)
2. Level 2 (lat long bi-weekly high res versions, shouldn't be needed and are difficult to work with)
3. **Level 3 indices (depth corrected and area weighted means for each model variable; i.e., what we will mostly use)**
  - a. "ACLIMsurveyrep\_": groundfish survey replicated (replicated in space and time)
  - b. "ACLIMregion\_": weekly strata based averages



To get more information about each of these level 3 datasets enter this in R:

```
# Metadata for Weekly ("ACLIMregion...") indices
head(all_info1)
```

```
##              name                                     Type B10KVersion  CMIP  GCM
## 1 B10K-H16_CMIP5_CESM_BIO_rcp85 Weekly regional indices          H16 CMIP5 CESM
## 2   B10K-H16_CMIP5_CESM_rcp45 Weekly regional indices          H16 CMIP5 CESM
## 3   B10K-H16_CMIP5_CESM_rcp85 Weekly regional indices          H16 CMIP5 CESM
## 4 B10K-H16_CMIP5_GFDL_BIO_rcp85 Weekly regional indices          H16 CMIP5 GFDL
## 5   B10K-H16_CMIP5_GFDL_rcp45 Weekly regional indices          H16 CMIP5 GFDL
## 6   B10K-H16_CMIP5_GFDL_rcp85 Weekly regional indices          H16 CMIP5 GFDL
##      BIO Carbon_scenario      Start      End nvars
## 1  TRUE                rcp85 2006-01-22 12:00:00 2099-12-27 12:00:00    59
## 2 FALSE                rcp45 2006-01-22 12:00:00 2081-02-16 12:00:00    59
## 3 FALSE                rcp85 2006-01-22 12:00:00 2099-12-27 12:00:00    59
## 4  TRUE                rcp85 2006-01-22 12:00:00 2099-12-27 12:00:00    59
## 5 FALSE                rcp45 2006-01-22 12:00:00 2099-12-27 12:00:00    59
## 6 FALSE                rcp85 2006-01-22 12:00:00 2099-12-27 12:00:00    59
```

```
# Metadata for Weekly ("ACLIMsurveyrep...") indices
head(all_info2)
```

```
##              name                                     Type B10KVersion  CMIP  GCM  BIO
## 1 B10K-H16_CMIP5_CESM_BIO_rcp85 Survey replicated          H16 CMIP5 CESM  TRUE
## 2   B10K-H16_CMIP5_CESM_rcp45 Survey replicated          H16 CMIP5 CESM FALSE
## 3   B10K-H16_CMIP5_CESM_rcp85 Survey replicated          H16 CMIP5 CESM FALSE
## 4 B10K-H16_CMIP5_GFDL_BIO_rcp85 Survey replicated          H16 CMIP5 GFDL  TRUE
## 5   B10K-H16_CMIP5_GFDL_rcp45 Survey replicated          H16 CMIP5 GFDL FALSE
## 6   B10K-H16_CMIP5_GFDL_rcp85 Survey replicated          H16 CMIP5 GFDL FALSE
##      Carbon_scenario Start  End nvars
## 1      rcp85    1970 2100    60
## 2      rcp45    1970 2100    60
## 3      rcp85    1970 2100    60
## 4      rcp85    1970 2100    60
## 5      rcp45    1970 2100    60
## 6      rcp85    1970 2100    60
```

## Indices & bias correction

### *ACLIM2 Indices*

The next step creates ACLIM2 indices (i.e., Level4) based on the Level3 output for each hindcast, historical run, and CMIP6 projection. The script below then bias corrects each index using the historical run and recenters the projection on the corresponding hindcast (such that projections are  $\Delta$  from historical mean values for the reference period `deltays <- 1970:2000`).

*NESB & SEBS averaged indices* The average water column values for each variable from the ROMSNPZ model strata x weekly Level2 outputs ('ACLIMregion') was calculated and used to calculate the strata-area weighted mean value for the NEBS and SEBS weekly, monthly, seasonally, and annually. Similarly, for survey replicated ('ACLIMsurveyrep') Level2 outputs the average water column value for each variable at each station was calculated used to calculate the strata-area weighted mean value for the NEBS and SEBS

annually. These indices were calculate for hindcast, historical, and projection scenarios, and used to bias correct the projections. More information on the methods for each can be found in the tabs below and the code immediately following this section will re-generate the bias corrected indices. All of the bias corrected outputs can be found in the “Data/out/CMIP6” folder.




Figure 1: Raw (top row) and bias corrected (bottom row)bottom temperature indices based on survey replicated Level3 outputs for the SEBS

**Important!** Note that for projections the ‘mn\_val’ represents raw mean values, while ‘val\_bias-corrected’ is the bias corrected mn\_val (should be used instead of the raw values). In all cases, for variables that are log-normally distributed (cannot be < 0), the  $\ln(\text{mn\_val})$  were used to bias correct and were then back transformed to non-log space after correction:

For normally distributed variables ( $Y$ ):

$$Y_{t,k}^{fut'} = \bar{Y}_{k,T}^{hind} + \left( \frac{\sigma_{k,T}^{hind}}{\sigma_{k,T}^{hist}} * (Y_{t,k}^{fut} - \bar{Y}_{k,T}^{hist}) \right)$$

where  $\bar{Y}_{y,k}^{fut'}$  is the bias corrected variable  $k$  value for time-step  $t$  (e.g., year, month, or season),  $\bar{Y}_{k,T}^{hind}$  is the mean value of the variable  $k$  during the reference period  $\bar{T} = [1980, 2013]$  from the hindcast model,  $\sigma_{k,T}^{hind}$  is the standard deviation of the hindcast during the reference period  $\bar{T}$ ,  $\sigma_{k,T}^{hist}$  is the standard deviation of the historical run during the reference period,  $Y_{t,k}^{fut}$  is the value of the variable from the projection at time-step  $t$  and  $\bar{Y}_{k,T}^{hist}$  is the average value from the historical run during reference period  $\bar{T}$ .

For log-normally distributed variables( $Y$ ):

$$Y_{y,k}^{fut'} = e^{\ln \bar{Y}_{k,T}^{hind} + \left( \frac{\hat{\sigma}_{k,T}^{hind}}{\hat{\sigma}_{k,T}^{hist}} * (\ln Y_{t,k}^{fut} - \ln \bar{Y}_{k,T}^{hist}) \right)}$$

, where  $\hat{\sigma}_{k,T}^{hist}$  and  $\hat{\sigma}_{k,T}^{hind}$  are the standard deviation of the  $\ln \bar{Y}_{k,T}^{hist}$  and  $\ln \bar{Y}_{k,T}^{hind}$  during the reference period  $\bar{T}$  (respectively).

## Weekly indices

Uses the strata x weekly data (‘ACLIMregion’) to generate strata-specific averages in order to generate the strata area-weighted averages for each week  $w$  each year  $y$ .

$$\bar{Y}_{w,y,k} = \frac{\sum_l^{n_s} \left( \frac{1}{n_i} \sum_t^{n_t} Y_{k,w,y,s,t} \right) * A_s}{\sum_s^{n_s} A_s}$$

, where  $Y_{k,w,y,s,t}$  is the value of the variable  $k$  in strata  $s$  at time  $t$  in year  $y$ ,  $A_s$  is the area of strata  $s$ ,  $n_i$  is the number of stations in strata  $s$ , and  $n_s$  is the number of strata  $s$  in each basin (NEBS or SEBS).

$\bar{Y}_{w,y,k}$  was calculated for the hindcast, historical run, and projection time-series. For projections  $\bar{Y}_{w,y,k}$  was bias corrected using the corresponding historical and hindcast values such that:

$$\bar{Y}_{w,y,k}^{fut'} = \bar{Y}_{w,k}^{hind} + \left( \frac{\sigma_{w,k}^{hind}}{\sigma_{w,k}^{hist}} * (\bar{Y}_{w,y,k}^{fut} - \bar{Y}_{w,k}^{hist}) \right)$$

, where  $\bar{Y}_{w,k}^{hist}$  and  $\bar{Y}_{w,k}^{hind}$  are the average historical weekly values across years in the period (1980 to 2012 ; adjustable in `R/setup.R`).

## Monthly indices

Uses the strata x weekly data ('ACCLIMregion') to generate strata-specific averages in order to generate the strata area-weighted averages for each month  $m$  each year  $y$ .

$$\bar{Y}_{m,y,k} = \frac{1}{n_w} \sum_w^{n_w} \bar{Y}_{w,y,k}$$

, where  $\bar{Y}_{w,y,k}$  are the weekly average indices for variable  $k$  in year  $y$  from the previous step,  $n_w$  is the number of weeks in each month  $m$ .

$\bar{Y}_{m,y,k}$  was calculated for the hindcast, historical run, and projection time-series. For projections  $\bar{Y}_{m,y,k}$  was bias corrected using the corresponding historical and hindcast values such that:

$$\bar{Y}_{m,y,k}^{fut'} = \bar{Y}_{m,k}^{hind} + \left( \frac{\sigma_{m,k}^{hind}}{\sigma_{m,k}^{hist}} * (\bar{Y}_{m,y,k}^{fut} - \bar{Y}_{m,k}^{hist}) \right)$$

, where  $\bar{Y}_{m,k}^{hist}$  and  $\bar{Y}_{m,k}^{hind}$  are the average historical monthly values across years in the period (1980 to 2012 ; adjustable in `R/setup.R`).

## Seasonal indices

Uses the strata x weekly data ('ACCLIMregion') to generate strata-specific averages in order to generate the strata area-weighted averages for each season  $l$  each year  $y$ .

$$\bar{Y}_{l,y,k} = \frac{1}{n_w} \sum_w^{n_w} \bar{Y}_{w,y,k}$$

, where  $\bar{Y}_{w,y,k}$  are the weekly average indices for variable  $k$  in year  $y$  from the previous step,  $n_w$  is the number of weeks in each season  $l$ .

$\bar{Y}_{l,y,k}$  was calculated for the hindcast, historical run, and projection time-series. For projections  $\bar{Y}_{l,y,k}$  was bias corrected using the corresponding historical and hindcast values such that:

$$\bar{Y}_{l,y,k}^{fut'} = \bar{Y}_{l,k}^{hind} + \left( \frac{\sigma_{l,k}^{hind}}{\sigma_{l,k}^{hist}} * (\bar{Y}_{l,y,k}^{fut} - \bar{Y}_{l,k}^{hist}) \right)$$

, where  $\bar{Y}_{l,k}^{hist}$  and  $\bar{Y}_{l,k}^{hind}$  are the average historical seasonal values across years in the reference period (1980 to 2012 ; adjustable in `R/setup.R`).

## Annual indices

Uses the strata x weekly data ('ACLIMregion') to generate strata-specific averages in order to generate the strata area-weighted averages for each season  $l$  each year  $y$ .

$$\bar{Y}_{y,k} = \frac{1}{n_w} \sum_w^{n_w} \bar{Y}_{w,y,k}$$

, where  $\bar{Y}_{w,y,k}$  are the weekly average indices for variable  $k$  in year  $y$  from the previous step,  $n_w$  is the number of weeks in each year  $y$ .

$\bar{Y}_{y,k}$  was calculated for the hindcast, historical run, and projection time-series. For projections  $\bar{Y}_{y,k}$  was bias corrected using the corresponding historical and hindcast values such that:

$$\bar{Y}_{y,k}^{fut'} = \bar{Y}_k^{hind} + \left( \frac{\sigma_k^{hind}}{\sigma_k^{hist}} * (\bar{Y}_{y,k}^{fut} - \bar{Y}_k^{hist}) \right)$$

, where  $\bar{Y}_k^{hind}$  and  $\bar{Y}_k^{hist}$  are the average historical values across years in the reference period (1980 to 2012 ; adjustable in R/setup.R).

## Annual survey rep. indices

Uses the station specific survey replicated (in time and space) data ('ACLIMsurveyrep') to generate strata-specific averages in order to generate the strata area-weighted averages for each year  $y$ .

$$\bar{Y}_{y,k} = \frac{\sum_l^{n_s} (\frac{1}{n_i} \sum_i^{n_i} Y_{k,y,s,i}) * A_s}{\sum_s^{n_s} A_s}$$

, where  $Y_{k,y,s,i}$  is the value of the variable  $k$  at station  $i$  in strata  $s$  in year  $y$ ,  $A_s$  is the area of strata  $s$ ,  $n_i$  is the number of stations in strata  $s$ , and  $n_s$  is the number of strata  $s$  in each basin (NEBS or SEBS).

$\bar{Y}_{y,k}$  was calculated for the hindcast, historical run, and projection time-series. For projections  $\bar{Y}_{y,k}$  was bias corrected using the corresponding historical and hindcast values such that:

$$\bar{Y}_{y,k}^{fut'} = \bar{Y}_k^{hind} + \left( \frac{\sigma_k^{hind}}{\sigma_k^{hist}} * (\bar{Y}_{y,k}^{fut} - \bar{Y}_k^{hist}) \right)$$

, where  $\bar{Y}_k^{hind}$  and  $\bar{Y}_k^{hist}$  are the average historical values across years in the reference period (1980 to 2012 ; adjustable in R/setup.R).

Appendix A includes the code used to generate the ACLIM2 indices and bias correct them. That code can be run to re-make the indices if you like but takes approx 30 mins a CMIP to run.

## Plot & concat Indices

The following code will open an interactive shiny() app for exploring the indices. You can also view this online at (kkh2022.shinyapps.io/ACLIM2\_indices)[https://kkh2022.shinyapps.io/ACLIM2\_indices/].

```
shiny::runApp("/Users/kholsman/Documents/GitHub/ACLIM2/R/shiny_aclim/ACLIM2_indices/app.R")  
  
# alternatively you can extract the data you want using the get_var()function
```

```
df <- get_var(typeIN = "annual",
              plotvar = "temp_bottom5m", plothist = F)

df$plot
head(df$dat)
```




Figure 2: “Raw (top row) and bias corrected (bottom row) bottom temperature indices based on survey replicated Level3 outputs for the SEBS”

## NRS indices (André)

```
suppressMessages(source("R/make.R"))

# preview possible variables

load(file = "Data/out/weekly_vars_C.Rdata")
load(file = "Data/out/weekly_vars.Rdata")
load(file = "Data/out/srvy_vars_C.Rdata")
load(file = "Data/out/srvy_vars.Rdata")

load(paste0("Data/out/K20P19_CMIP6/allEBS_means/ACLIM_annual_hind_mn.Rdata"))
varall <- unique(ACLIM_annual_hind$var)
varall

scens <- c("ssp126", "ssp585")
GCMs <- c("miroc", "gfdl", "cesm")
```


```

varlist <- c("temp_bottom5m","fracbelow2","uEast_surface5m")

# get the variable you want:
df <- get_var( typeIN    = "annual",
               plotvar   = varlist[1],
               bcIN      = "raw",
               CMIPIN    = "K20P19_CMIP6",
               plothist  = F, # ignore the hist runs
               removeyr1 = T) # "Remove first year of projection ( burn in)"

df$plot+coord_cartesian(ylim = c(0, 7))

```



```

head(df$dat)
#
# # get the variable you want:
# df <- get_var( typeIN    = "annual",
#               plotvar   = varlist[1],
#               bcIN      = "raw",
#               CMIPIN    = "K20P19_CMIP6_C",
#               plothist  = F, # ignore the hist runs
#               removeyr1 = T) # "Remove first year of projection ( burn in)"
#
# df$plot+coord_cartesian(ylim = c(0, 7))
#
# head(df$dat)
#
# # get the variable you want:

```


```

# df <- get_var( typeIN    = "annual",
#               plotvar    = "uEast_surface5m",
#               bcIN       = "bias corrected",
#               CMIPIN     = "K20P19_CMIP6_C",
#               plothist    = F, # ignore the hist runs
#               removeyr1  = T) #"Remove first year of projection ( burn in)"
#
# df$plot
# head(df$dat)

# get the variable you want:
df <- get_var( typeIN    = "annual",
               plotvar    = "pH_depthavg",
               bcIN       = "bias corrected",
               CMIPIN     = "K20P19_CMIP6_C",
               plothist    = F, # ignore the hist runs
               removeyr1  = T) #"Remove first year of projection ( burn in)"

df$plot

```



```
head(df$dat)
```

```

# get the variable you want:
df <- get_var( typeIN    = "annual",
               plotvar    = "pH_depthavg",
               bcIN       = "bias corrected",
               CMIPIN     = "K20P19_CMIP6_C",


```

```

plothist = F, # ignore the hist runs
removeyr1 = T) #"Remove first year of projection ( burn in)"

```

```
df$plot
```



```
head(df$dat)
```

```

##      var basin year   jday   mnDate  mn_val
## 1 pH_depthavg SEBS 1970 189.5000 1970-07-09 8.002964
## 2 pH_depthavg SEBS 1971 181.5000 1971-07-01 8.008074
## 3 pH_depthavg SEBS 1972 184.0000 1972-07-03 7.996702
## 4 pH_depthavg SEBS 1973 182.0000 1973-07-02 8.022209
## 5 pH_depthavg SEBS 1974 181.1373 1974-07-01 8.023676
## 6 pH_depthavg SEBS 1975 183.5000 1975-07-03 8.030164
##
##      sim gcmcmip GCM  scen sim_type units
## 1 ACLIMregion_C_B10K-K20P19_CORECFS hind hind ssp126 hind
## 2 ACLIMregion_C_B10K-K20P19_CORECFS hind hind ssp126 hind
## 3 ACLIMregion_C_B10K-K20P19_CORECFS hind hind ssp126 hind
## 4 ACLIMregion_C_B10K-K20P19_CORECFS hind hind ssp126 hind
## 5 ACLIMregion_C_B10K-K20P19_CORECFS hind hind ssp126 hind
## 6 ACLIMregion_C_B10K-K20P19_CORECFS hind hind ssp126 hind
##
##      bc GCM_scen GCM_scen_sim CMIP type
## 1 bias corrected hind_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 2 bias corrected hind_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 3 bias corrected hind_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 4 bias corrected hind_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 5 bias corrected hind_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 6 bias corrected hind_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual

```



```

# concat the hind and fut runs by removing years from projection
maxDin <- max(as.vector(df$dat%>%dplyr::filter(sim_type=="hind")%>%dplyr::select(mnDate))[[1]])

newdat <- stitchTS(dat = df$dat,
                   maxD = maxDin)

# newdat has the full set of data
# select miroc_ssp126
head(newdat%>%dplyr::filter(GCM_scen==paste0(GCMs[1], "_", scens[1])))

```

```

##           var basin year    jday    mnDate    mn_val
## 1 pH_depthavg  SEBS 1970 189.5000 1970-07-09 8.002964
## 2 pH_depthavg  SEBS 1971 181.5000 1971-07-01 8.008074
## 3 pH_depthavg  SEBS 1972 184.0000 1972-07-03 7.996702
## 4 pH_depthavg  SEBS 1973 182.0000 1973-07-02 8.022209
## 5 pH_depthavg  SEBS 1974 181.1373 1974-07-01 8.023676
## 6 pH_depthavg  SEBS 1975 183.5000 1975-07-03 8.030164
##
##               sim gcmcmip  GCM    scen sim_type units
## 1 ACLIMregion_C_B10K-K20P19_CORECFS    hind hind ssp126    hind
## 2 ACLIMregion_C_B10K-K20P19_CORECFS    hind hind ssp126    hind
## 3 ACLIMregion_C_B10K-K20P19_CORECFS    hind hind ssp126    hind
## 4 ACLIMregion_C_B10K-K20P19_CORECFS    hind hind ssp126    hind
## 5 ACLIMregion_C_B10K-K20P19_CORECFS    hind hind ssp126    hind
## 6 ACLIMregion_C_B10K-K20P19_CORECFS    hind hind ssp126    hind
##           bc      GCM_scen      GCM_scen_sim      CMIP      type
## 1 bias corrected miroc_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 2 bias corrected miroc_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 3 bias corrected miroc_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 4 bias corrected miroc_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 5 bias corrected miroc_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual
## 6 bias corrected miroc_ssp126 hind_ssp126_hind K20P19_CMIP6_C annual

```

```

tail(newdat%>%dplyr::filter(GCM_scen==paste0(GCMs[1], "_", scens[1])))

```

```


##           var basin year    jday    mnDate    mn_val
## 126 pH_depthavg  SEBS 2094 181.50 2094-07-01 7.879315
## 127 pH_depthavg  SEBS 2095 180.50 2095-06-30 7.886838
## 128 pH_depthavg  SEBS 2096 183.00 2096-07-02 7.858582
## 129 pH_depthavg  SEBS 2097 184.50 2097-07-04 7.904151
## 130 pH_depthavg  SEBS 2098 183.50 2098-07-03 7.933660
## 131 pH_depthavg  SEBS 2099 175.92 2099-06-25 7.883800
##
##               sim               gcmcmip      GCM
## 126 ACLIMregion_C_B10K-K20P19_CMIP6_miroc_ssp126 B10K-K20P19_CMIP6_miroc miroc
## 127 ACLIMregion_C_B10K-K20P19_CMIP6_miroc_ssp126 B10K-K20P19_CMIP6_miroc miroc
## 128 ACLIMregion_C_B10K-K20P19_CMIP6_miroc_ssp126 B10K-K20P19_CMIP6_miroc miroc
## 129 ACLIMregion_C_B10K-K20P19_CMIP6_miroc_ssp126 B10K-K20P19_CMIP6_miroc miroc
## 130 ACLIMregion_C_B10K-K20P19_CMIP6_miroc_ssp126 B10K-K20P19_CMIP6_miroc miroc
## 131 ACLIMregion_C_B10K-K20P19_CMIP6_miroc_ssp126 B10K-K20P19_CMIP6_miroc miroc
##           scen sim_type units           bc      GCM_scen      GCM_scen_sim
## 126 ssp126      proj      bias corrected miroc_ssp126 miroc_ssp126_proj

```

```
## 127 ssp126      proj      bias corrected miroc_ssp126 miroc_ssp126_proj
## 128 ssp126      proj      bias corrected miroc_ssp126 miroc_ssp126_proj
## 129 ssp126      proj      bias corrected miroc_ssp126 miroc_ssp126_proj
## 130 ssp126      proj      bias corrected miroc_ssp126 miroc_ssp126_proj
## 131 ssp126      proj      bias corrected miroc_ssp126 miroc_ssp126_proj
##              CMIP      type
## 126 K20P19_CMIP6_C annual
## 127 K20P19_CMIP6_C annual
## 128 K20P19_CMIP6_C annual
## 129 K20P19_CMIP6_C annual
## 130 K20P19_CMIP6_C annual
## 131 K20P19_CMIP6_C annual
```

```
pp <- ggplot(newdat)+
  geom_line(aes(x=mnDate,y=mn_val,color= GCM_scen, linetype = basin),
            alpha = 0.6,show.legend = FALSE)+
  geom_smooth(aes(x=mnDate,y=mn_val,color= GCM_scen,
                 fill=GCM_scen,linetype = basin),alpha=0.1,
             method="loess",formula='y ~ x',span = .5,show.legend=T)+
  theme_minimal() +
  labs(x="Date",
       y=paste(newdat$var[1], "(" ,newdat$units[1], ")"),
       subtitle = "",
       legend = "",
       title = paste(newdat$var[1], "(" ,newdat$basin[1], "," ,newdat$type[1], ")"))+
  scale_color_discrete()+
  facet_grid(scen~.)
# plot it
pp
```

### pH\_depthavg ( SEBS , annual )



```
# plot it interactively
plotly::ggplotly(pp)
```

## monthly indices (Andy)


```
suppressMessages(source("R/make.R"))

# preview possible variables
load(paste0("Data/out/K20P19_CMIP6/allEBS_means/ACLIM_monthly_hind_mn.Rdata"))
varall <- unique(ACLIM_monthly_hind$var)
varall

scens <- c("ssp126","ssp585")
GCMs <- c("miroc","gfdl", "cesm" )
varlist <- c("temp_bottom5m","fracbelow2","uEast_surface5m")

# get the variable you want:
df <- get_var( typeIN = "monthly",
               plotvar = "temp_bottom5m",
               bcIN = "bias corrected",
               plothist = F, # ignore the hist runs
               removeyr1 = T) # "Remove first year of projection ( burn in)"

head(df$dat)
df$plot
```



```

# concat the hind and fut runs by removing years from projection
maxDin <- max(as.vector(df$dat)%>%dplyr::filter(sim_type=="hind")%>%dplyr::select(mnDate))[[1]]


newdat <- stitchTS(dat = df$dat,
                  maxD = maxDin)

# newdat has the full set of data
# select miroc_ssp126
head(newdat%>%dplyr::filter(GCM_scen==paste0(GCMs[1],"_",scens[1])))
tail(newdat%>%dplyr::filter(GCM_scen==paste0(GCMs[1],"_",scens[1])))

pp <- ggplot(newdat)+
  geom_line(aes(x=mnDate,y=mn_val,color= GCM_scen, linetype = basin),
            alpha = 0.6,show.legend = FALSE)+
  geom_smooth(aes(x=mnDate,y=mn_val,color= GCM_scen,
                 fill=GCM_scen,linetype = basin),alpha=0.1,
             method="loess",formula='y ~ x',span = .5,show.legend=T)+
  theme_minimal() +
  labs(x="Date",
       y=paste(newdat$var[1],"(",newdat$units[1],")"),
       subtitle = "",
       legend = "",
       title = paste(newdat$var[1],"(",newdat$basin[1],",",newdat$type[1],")"))+
  scale_color_discrete()+
  facet_grid(scen~.)
# plot it
pp

```

temp\_bottom5m ( SEBS , monthly )



```
# plot it interactively
plotly::ggplotly(pp)
```

## weekly indices (Jon)


```
suppressMessages(source("R/make.R"))

# preview possible variables
load(paste0("Data/out/K20P19_CMIP6/allEBS_means/ACLIM_weekly_hind_mn.Rdata"))
varall <- unique(ACLIM_weekly_hind$var)
varall

scens <- c("ssp126","ssp585")
GCMs <- c("miroc","gfdl", "cesm" )
varlist <- c("temp_bottom5m","fracbelow2","uEast_surface5m")

# get the variable you want:
df <- get_var( typeIN = "weekly",
               plotvar = "temp_bottom5m",
               bcIN = "bias corrected",
               plothist = F, # ignore the hist runs
               removeyr1 = T) #"Remove first year of projection ( burn in)")

df$plot
```



```

head(df$dat)

# concat the hind and fut runs by removing years from projection
maxDin <- max(as.vector(df$dat%>%dplyr::filter(sim_type=="hind")%>%dplyr::select(mnDate))[[1]])

newdat <- stitchTS(dat = df$dat,
                   maxD = maxDin)


# newdat has the full set of data
# select miroc_ssp126
head(newdat%>%dplyr::filter(GCM_scen==paste0(GCMs[1],"_",scens[1])))

pp <- ggplot(newdat)+
  geom_line(aes(x=mnDate,y=mn_val,color= GCM_scen, linetype = basin),
            alpha = 0.6,show.legend = FALSE)+
  geom_smooth(aes(x=mnDate,y=mn_val,color= GCM_scen,
                 fill=GCM_scen,linetype = basin),alpha=0.1,
             method="loess",formula='y ~ x',span = .5,show.legend=T)+
  theme_minimal() +
  labs(x="Date",
       y=paste(newdat$var[1],"(",newdat$units[1],")"),
       subtitle = "",
       legend = "",
       title = paste(newdat$var[1],"(",newdat$basin[1],",",newdat$type[1],")"))+
  scale_color_discrete()+
  facet_grid(scen~.)

# plot it
pp

```

## temp\_bottom5m ( SEBS , weekly )



```
# plot it interactively
plotly::ggplotly(pp)
```

## Output to .dat file (ADMB/ TMB users)

For CEATTLE I create a .dat file that is read into the ADMB script. That .dat file includes the bias corrected values (e.g., bottom temperature in deg C) used for the bioenergetics and temperature-dependent growth functions as well as Z-score (scaled) values used as covariates on the recruitment function. The section below will step through that .dat file creation for a subset of variables as well as demo chunks of ADMB code for reading that into a ADMB based model.

### Use R to make .dat file

```
# 1 -- create .dat filename & path
# 2 -- rescale (Z-score) data and get variables
# 3 -- write data to hind .dat file
# 3 -- write data to fut .dat file

# 1 -- create .dat filename & path
# -----

# Define the name for the .dat file
file.name <- "ACLIM2_CMIP6_short"
```

```

fn          <- paste(file.name, "_bcs.dat", sep="")
# switches
hind_yrs    <- 1989:2021  # define the years of your estimation model
fut_yrs     <- 2022:2100  # define the years of your projections
archive_old <- T          # Archive the older version of the .dat file?

outpath     <- "Data/out/ADMB_datfiles"
if(!dir.exists(outpath)) dir.create(outpath)

# define hind and fut data files
fndat_hind <- file.path(outpath, paste("KKHhind_", fn, sep=""))
fndat_fut  <- file.path(outpath, paste("KKHfut_", fn, sep=""))
fndat_hind2 <- file.path(outpath, paste("hind_", fn, sep=""))
fndat_fut2  <- file.path(outpath, paste("fut_", fn, sep=""))

# create and archive .dat files
outfile     <- fndat_fut
if(file.exists(outfile)&archive_old){
  # archive older version
  archivefl <- paste0(substr(outfile, start=1, stop=nchar(outfile)-4),
    format(Sys.time(), "%Y%m%d"), ".dat")
  file.rename(outfile, archivefl)
  file.remove(outfile)
}

file.create(outfile)
outfile <- outfile.hind
if(file.exists(outfile)&archive_old){
  # archive older version
  archivefl <- paste0(substr(outfile, start=1, stop=nchar(outfile)-4),
    format(Sys.time(), "%Y%m%d"), ".dat")
  file.rename(outfile, archivefl)
  file.remove(outfile)
}

file.create(outfile)

# 2 -- rescale (Z-score) data and get variables

# preview possible variables
load(paste0("Data/out/K20P19_CMIP6/allEBS_means/ACLIM_annual_hind_mn.Rdata"))
varall <- unique(ACLIM_annual_hind$var)
varall

# get each variable, convert to TS and rbind

plotbasin <- "SEBS"

CMIPS <- c("K20P19_CMIP6", "K20P19_CMIP5")
CMIPS <- c("K20P19_CMIP6")

```



```

for(c in 1:length(CMIPS)){

  # first for annual mean values:
  varlist <- c("largeZoop_integrated", "fracbelow2", "temp_bottom5m", "temp_surface5m")
  typeIN <- "annual"

  load(paste0("Data/out/", CMIPS[c], "/allEBS_means/ACLIM_", typeIN, "_hind_mn.Rdata"))
  load(paste0("Data/out/", CMIPS[c], "/allEBS_means/ACLIM_", typeIN, "_fut_mn.Rdata"))
  eval(parse(text = paste0("dhind <- ACLIM_", typeIN, "_hind")))
  eval(parse(text = paste0("dfut <- ACLIM_", typeIN, "_fut")))

  # rescale the data using mean of the hind
  tmphind <- dhind%>%
    dplyr::filter(var%in%varlist, basin==plotbasin, year%in%hind_yrs)%>%
    dplyr::select(var, basin, year, jday, mnDate, mn_val,
                  mnVal_hind, sdVal_hind, sim, gcmcmip, CMIP, GCM, scen, sim_type, units, long_name)%>%
    dplyr::mutate(bc = "bias corrected",
                  GCM_scen = paste0(GCM, "_", scen),
                  mn_val_scaled = (mn_val-mnVal_hind)/sqrt(sdVal_hind))

  tmpfut <- dfut%>%
    dplyr::filter(var%in%varlist, basin==plotbasin, year%in%fut_yrs)%>%
    dplyr::select(var, basin, year, jday, mnDate, val_biascorrected,
                  mnVal_hind, sdVal_hind, sim, gcmcmip, CMIP, GCM, scen, sim_type, units, long_name)%>%
    dplyr::rename(mn_val = val_biascorrected)%>%
    dplyr::mutate(bc = "bias corrected",
                  GCM_scen = paste0(GCM, "_", scen),
                  mn_val_scaled = (mn_val-mnVal_hind)/sqrt(sdVal_hind))

  # now for seasonal mean values:
  typeIN <- "seasonal"
  varlist <- c("largeZoop_integrated")

  seasonsIN <- unique(seasons$season)
  load(paste0("Data/out/", CMIPS[c], "/allEBS_means/ACLIM_", typeIN, "_hind_mn.Rdata"))
  load(paste0("Data/out/", CMIPS[c], "/allEBS_means/ACLIM_", typeIN, "_fut_mn.Rdata"))
  eval(parse(text = paste0("dhind <- ACLIM_", typeIN, "_hind")))
  eval(parse(text = paste0("dfut <- ACLIM_", typeIN, "_fut")))

  # rescale the data using mean of the hind
  tmphind2 <- dhind%>%
    dplyr::filter(var%in%varlist, basin==plotbasin, year%in%hind_yrs, season%in%seasonsIN)%>%
    dplyr::mutate(var = paste0(var, "_", season))%>%
    dplyr::select(var, basin, year, jday, mnDate, mn_val,
                  mnVal_hind, sdVal_hind, sim, gcmcmip, CMIP, GCM, scen, sim_type, units, long_name)%>%
    dplyr::mutate(bc = "bias corrected",
                  GCM_scen = paste0(GCM, "_", scen),
                  mn_val_scaled = (mn_val-mnVal_hind)/sqrt(sdVal_hind))

  tmpfut2 <- dfut%>%

```

```

dplyr::filter(var%in%varlist,basin==plotbasin,year%in%fut_yrs,season%in%seasonsIN)%>%
dplyr::mutate(var = paste0(var,"_",season))%>%
dplyr::select(var,basin,year, jday,mnDate,val_biascorrected,
              mnVal_hind,sdVal_hind, sim,gcmcmip,CMIP,GCM,scen,sim_type ,units,long_name)%>%
dplyr::rename(mn_val = val_biascorrected)%>%
dplyr::mutate(bc = "bias corrected",
              GCM_scen = paste0(GCM,"_",scen),
              mn_val_scaled = (mn_val-mnVal_hind )/sqrt(sdVal_hind))

if(c ==1){
  hind <- rbind(tmphind,tmphind2)
  fut <- rbind(tmpfut,tmpfut2)
}else{
  hind <- rbind(hind,tmphind,tmphind2)
  fut <- rbind(fut,tmpfut,tmpfut2)
}
}

# plot the data
p <- ggplot(hind)+
  geom_line(aes(x=mnDate,y=mn_val,color=GCM_scen))+
  geom_line(data=fut,aes(x=mnDate,y=mn_val,color=GCM_scen))+
  facet_wrap(~var,scales="free_y")+
  theme_minimal()
p

# plot the data
p_scaled <- ggplot(hind)+
  geom_line(aes(x=mnDate,y=mn_val_scaled,color=GCM_scen))+
  geom_line(data=fut,aes(x=mnDate,y=mn_val_scaled,color=GCM_scen))+
  facet_wrap(~var,scales="free_y")+
  theme_minimal()
p_scaled

d_wide <- reshape2::dcast(hind%>%dplyr::filter(year!=2021),year~c(var),value.var = "mn_val")
corr <- cor(d_wide[,-1])
# remove those where cov is high (temp by season and cold pool by season)
long_dat <- reshape2::melt(corr,variable.name = "variable") %>%
as.data.frame()

# plot covariation between variables
long_dat %>%arrange(value)%>%
ggplot(aes(x=Var1, y=Var2, fill=value)) +
geom_raster() +
scale_fill_viridis_c()+
theme_minimal()+
theme(axis.text.x = element_text(angle = 90))

# 3 -- write data to hind .dat file
# -----

# CEATTLE uses a spp overlap index - you can skip this

```

```

overlapdat <- data.frame(
  atf_OL=c(0.9391937,0.8167094,0.808367,0.5926875,0.7804481,0.5559549,
    0.4006931,0.5881404,0.7856776,0.511565,0.6352048,0.5583476,
    0.5792738,0.5417657,0.8212887,0.6287613,0.4536608,0.6587292,
    0.4884194,0.8289379,0.4399257,0.5950167,0.6388434,0.6111834,
    0.8742649,0.7868746,0.8024257,0.6227457,0.4956742,0.4347917,
    0.4791108,0.4369006,0.5613625,0.4353015),
  south_OL=c(0.9980249,0.9390368,0.9959974,0.6130846,0.951234,0.5851891,
    0.4934879,0.641471,0.9809618,0.5596813,0.7196964,0.6754698,
    0.5774808,0.6041351,0.9406521,0.7949525,0.5306435,0.7977694,
    0.5345031,0.9879945,0.5079171,0.7148121,0.8997132,0.7340859,
    0.9962068,0.9627235,0.998043,0.8111,0.6087638,0.513057,0.5492621,
    0.4971361,0.665453,0.5969653)
)

includeOverlap <- F
overlap <- matrix(1,3,length(sort(unique(hind$year))))
overlap_fut <- array(1,c(3,length(unique(fut$GCM_scen))+1,length(sort(unique(fut$year))))
if(includeOverlap){
  overlap[3,] <- overlapIN
  overlap[3,][overlap[3,]>1]<-1 #covs$BT2to6/covs$BT0to6
}

# Pick up here

# Kir's .dat file
covars <- unique(hind$var)
makeDat_hind(datIN = hind,
  outfile = fndat_hind,
  nsppIN = 3,
  overlapIN = overlap,
  nonScaled_covlist = c("temp_bottom5m","temp_surface5m" ),
  Scaled_covlist = covars)

# generic .dat file
covars <- unique(hind$var)
makeDat_fut( datIN = fut,
  hinddatIN = hind,
  outfile = fndat_fut,
  nsppIN = 3,
  last_nyrs_avg = 10,
  overlapIN = overlap_fut, #(nspp,nsim+1,nyrs_fut)
  overlap_hind=overlap,
  nonScaled_covlist = c("temp_bottom5m","temp_surface5m" ),
  Scaled_covlist = covars)

### Here's a generic version that doesn't include nspp and overla[]
# generic .dat file
covars <- unique(hind$var)
makeDat_hind(datIN = hind,
  outfile = fndat_hind2,
  nsppIN = NULL,

```

```

overlapIN = NULL,
nonScaled_covlist = c("temp_bottom5m", "temp_surface5m" ),
Scaled_covlist = covars)

# generic .dat file
covars <- unique(hind$var)
makeDat_fut( datIN = fut,
             hinddatIN = hind,
             outfile = fndat_fut2,
             nsppIN = NULL,
             last_nyrs_avg = 10,
             overlapIN = NULL, #(nspp, nsim+1, nyrs_fut)
             overlap_hind = NULL,
             nonScaled_covlist = c("temp_bottom5m", "temp_surface5m" ),
             Scaled_covlist = covars)

```

## APPENDIX A: Create & bias correct ACLIM2 indices

The following code shows how the ACLIM2 indices and bias correction was done. You do not need to re-run this (it is included so you can if you want to). To explore the indices skip to the next section.

```

# -----
# SETUP WORKSPACE
# rm(list=ls())
# setwd("Documents/GitHub/ACLIM2")
tmstp <- format(Sys.time(), "%Y_%m_%d")
main <- getwd() # "~/GitHub_new/ACLIM2"

# loads packages, data, setup, etc.
suppressMessages(source("R/make.R"))

tmstamp1 <- format(Sys.time(), "%Y%m%d")
# tmstamp1 <- "20220428"

update_hind <- TRUE # set to true to update hind and hindS; needed annually
update_proj <- TRUE # set to true to update fut; not needed
update_hist <- TRUE # set to true to update fut; not needed

# the reference years for bias correcting in R/setup.R
ref_years

# the year to z-score scale / delta in R/setup.R
deltayrs
# the year to z-score scale / delta in R/setup.R
deltayrs
data_path
load(file.path(Rdata_path, "../weekly_vars_C.Rdata"))
load(file.path(Rdata_path, "../weekly_vars.Rdata"))
load(file.path(Rdata_path, "../srvy_vars_C.Rdata"))
load(file.path(Rdata_path, "../srvy_vars.Rdata"))
rm_wk_list <- c(
  "Iron_bottom5m", "Iron_integrated", "Iron_surface5m",

```

```

    "prod_Eup_integrated", "prod_NCa_integrated",
    "region_area")
rm_wk_list_C <- c( "Iron_bottom5m",
    "prod_Cop_integrated", "prod_Eup0_integrated", "prod_EupS_integrated",
    "prod_Jel_integrated", "prod_MZL_integrated",
    "prod_NCa0_integrated", "prod_NCaS_integrated", "Fe_depthavg",
    "region_area")
rm_var_list <- c(
    "station_id", "latitude",
    "longitude", "stratum",
    "Iron_integrated", "Iron_surface5m",
    "Iron_bottom5m", "prod_Eup_integrated", "prod_NCa_integrated",
    "doy")
rm_var_list_C <- c(
    "station_id", "latitude",
    "longitude", "stratum",
    "prod_Cop_integrated", "prod_Eup0_integrated", "prod_EupS_integrated",
    "prod_Jel_integrated", "prod_MZL_integrated", "Fe_depthavg",
    "prod_NCa0_integrated", "prod_NCaS_integrated",
    # "Iron_integrated", "Iron_surface5m",
    # "Iron_bottom5m", "prod_Eup_integrated", "prod_NCa_integrated"
    "doy")
# remove these variables:
l3srvy_varlist <- srvy_vars[!srvy_vars%in%rm_var_list]
l3wk_varlist    <- weekly_vars[!weekly_vars%in%rm_wk_list]

l3wk_varlist_C  <- weekly_vars_C[!weekly_vars_C%in%rm_wk_list_C]
l3srvy_varlist_C <- srvy_vars_C[!srvy_vars_C%in%rm_var_list_C]
l2_vars         <- c("temp_bottom5m", "temp_surface5m", "average_oxygen_bottom5m")
# remove these variables:
v11 <- srvy_vars[!srvy_vars%in%rm_var_list]
v11_c <- srvy_vars_C[!srvy_vars_C%in%rm_var_list_C]

v12 <- weekly_vars[!weekly_vars%in%rm_wk_list]
v12_c <- weekly_vars_C[!weekly_vars_C%in%rm_wk_list_C]

v1<-unique(c(v11,v12))
v1_c<-unique(c(v11_c,v12_c))
# add in largeZoop (gets generated in make_indices_region_new.R)
v1 <- c(v1,"largeZoop_integrated")
v1_c <- c(v1_c,"largeZoop_integrated")

# Identify which variables would be normally
# distributed (i.e., can have negative values)
normv1 <- c("shflux", "ssflux", "temp_bottom5m",
    "temp_integrated", "temp_surface5m",
    "uEast_bottom5m", "uEast_surface5m",
    "vNorth_bottom5m", "vNorth_surface5m")

normlist <- data.frame(var = v1, lognorm = TRUE)
normlist$lognorm[normlist$var%in%normv1] <- FALSE

normlist_c <- data.frame(var = v1_c, lognorm = TRUE)

```

```

normlist$lognorm[normlist$var%in%normvl] <- FALSE

save(weekly_vars_C,file = "Data/out/weekly_vars_C.Rdata")
save(weekly_vars,file = "Data/out/weekly_vars.Rdata")
save(srvy_vars_C,file = "Data/out/srvy_vars_C.Rdata")
save(srvy_vars,file = "Data/out/srvy_vars.Rdata")

# generate indices and bias corrected projections
# This takes approx 30 mins each

gcmcmipL <- c("B10K-K20P19_CMIP6_miroc",
             "B10K-K20P19_CMIP6_gfdl",
             "B10K-K20P19_CMIP6_cesm")

CMIP6_K20P19_Indices_C <- suppressMessages(
  makeACLIM2_Indices(
    BC_target = "mn_val",
    hind_sim  = "B10K-K20P19_CORECFS",
    histLIST  = paste0(gcmcmipL,"_historical"),
    gcmcmipLIST = gcmcmipL,
    Rdata_pathIN = Rdata_path_C,
    regnm      = "ACLIMregion_C",
    srvynm     = "ACLIMsurveyrep_C",

    normlist_IN = normlist_c,
    sim_listIN  = sim_list[-grep("historical",sim_list)]))

if("CMIP6_K20P19_Indices_C"%in%ls()){
  saved <- FALSE
  saved <- save_indices(flIN = CMIP6_K20P19_Indices_C,
    subfl = "allEBS_means",
    post_txt = "_mn",
    CMIP_fdlr ="K20P19_CMIP6_C")
  fl <- "Data/out/CMIP6_K20P19_Indices_list_C.Rdata"

  if(file.exists(fl)) file.remove(fl)
  save(CMIP6_K20P19_Indices_C, file = fl)
  if(saved){
    rm(CMIP6_K20P19_Indices_C)}else{
    stop("Indices not saved!")
  }
  gc()
}

CMIP6_K20_Indices <- suppressMessages(
  makeACLIM2_Indices(
    BC_target = "mn_val",
    hind_sim  = "B10K-K20_CORECFS",
    histLIST  = paste0(gcmcmipL,"_historical"),
    gcmcmipLIST = gcmcmipL,
    normlist_IN = normlist,
    sim_listIN  = sim_list[-grep("historical",sim_list)]))

```

```

if("CMIP6_K20_Indices"%in%ls()){
  saved <- FALSE
  saved <- save_indices(flIN = CMIP6_K20_Indices,
    subfl = "allEBS_means",
    post_txt = "_mn",
    CMIP_fdlr ="K20P19_CMIP6")
  fl <- "Data/out/CMIP6_K20_Indices_list.Rdata"

  if(file.exists(fl)) file.remove(fl)
  save(CMIP6_K20_Indices, file = fl)
  if(saved){
    rm(CMIP6_K20_Indices)}else{
    stop("Indices not saved!")
  }
  gc()
}

# CMIP5 K20P19
gcmcmipL2 <- c("B10K-K20P19_CMIP5_MIROC","B10K-K20P19_CMIP5_GFDL","B10K-K20P19_CMIP5_CESM")
CMIP5_K20P19_Indices <- suppressMessages(
  makeACLIM2_Indices(
    BC_target = "mn_val",
    hind_sim = "B10K-K20_CORECFS",
    histLIST = rep("B10K-K20_CORECFS",3),
    gcmcmipLIST = gcmcmipL2,
    normlist_IN = normlist,
    regnm = "ACLIMregion",
    srvyntm = "ACLIMsurveyrep",
    sim_listIN = sim_list[-grep("historical",sim_list)]))

if("CMIP5_K20P19_Indices"%in%ls()){
  saved <- FALSE
  saved <- save_indices(flIN = CMIP5_K20P19_Indices,
    subfl = "allEBS_means",
    post_txt = "_mn",
    CMIP_fdlr ="K20P19_CMIP5")

  fl <- "Data/out/CMIP5_K20P19_Indices_list.Rdata"
  if(file.exists(fl)) file.remove(fl)
  save(CMIP5_K20P19_Indices, file = fl)

  if(saved){
    rm(CMIP5_K20P19_Indices)}else{
    stop("Indices not saved!")
  }
  gc()
}

CMIP5_K20P19_Indices_C <- suppressMessages(
  makeACLIM2_Indices(
    BC_target = "mn_val",
    hind_sim = "B10K-K20P19_CORECFS",
    histLIST = rep("B10K-K20P19_CORECFS",3),

```

```

        gcmcmipLIST = gcmcmipL2,
        Rdata_pathIN = Rdata_path_C,
        regnm      = "ACLIMregion_C",
        srvynm     = "ACLIMsurveyrep_C",

        normlist_IN = normlist_c,
        sim_listIN = sim_list[-grep("historical",sim_list)])

if("CMIP5_K20P19_Indices_C"%in%ls()){
  saved <- FALSE
  saved <- save_indices(flIN = CMIP5_K20P19_Indices_C,
    subfl = "allEBS_means",
    post_txt = "_mn",
    CMIP_fdlr ="K20P19_CMIP5_C")

  fl <- "Data/out/CMIP5_K20P19_Indices_list_C.Rdata"
  if(file.exists(fl)) file.remove(fl)
  save(CMIP5_K20P19_Indices_C, file = fl)
  if(saved){
    rm(CMIP5_K20P19_Indices_C)}else{
    stop("Indices not saved!")
  }
  gc()
}

# CMIP5 H16
gcmcmipL2 <- c("B10K-H16_CMIP5_MIROC","B10K-H16_CMIP5_GFDL","B10K-H16_CMIP5_CESM")
CMIP5_H16_Indices <- suppressMessages(
  makeACLIM2_Indices(
    BC_target = "mn_val",
    hind_sim  = "B10K-H16_CORECFS",
    histLIST  = rep("B10K-H16_CORECFS",3),
    gcmcmipLIST = gcmcmipL2,
    sim_listIN = sim_list[-grep("historical",sim_list)])

if("CMIP5_H16_Indices"%in%ls()){
  saved <- FALSE
  saved <- save_indices(flIN = CMIP5_H16_Indices,
    subfl = "allEBS_means",
    post_txt = "_mn",
    CMIP_fdlr ="H16_CMIP5")

  fl <- "Data/out/CMIP5_H16_Indices_list.Rdata"
  if(file.exists(fl)) file.remove(fl)
  save(CMIP5_H16_Indices, file = fl)
  if(saved){
    rm(CMIP5_H16_Indices)}else{
    stop("Indices not saved!")
  }
  gc()
}

if(1==10){
  save(CMIP6_Indices, file = "Data/out/CMIP6_Indices_List.Rdata")
  save(CMIP5_K20_Indices, file = "Data/out/CMIP5_K20_Indices_List.Rdata")
  save(CMIP5_H16_Indices, file = "Data/out/CMIP5_H16_Indices_List.Rdata")
}

```



}




Figure 3: September large zooplankton integrated concentration

## misc

$$B0_{input}^k = \bar{B0}_{(2004:2014)}^k \left( \frac{B0_{2015}^a}{\bar{B0}_{(2004:2014)}^a} \right)$$

Where  $B0_{input}$  is the unfished biomass used for setting inputs of (e.g.,  $B0_{target} = 0.4B0_{input}$ ) and is determined by re-scaling the spawning stock biomass from the status quo assessment in 2015 ( $B0_{2015}^a$ ) to the average model spawning stock biomass for your model between 2004-2014 (i.e.,  $B0_k$ ) using the average unfished biomass from the stock assessment model during the same period ( $B0_a$ ).