

# Homework 1

Matthew Cocci

February 11, 2014

1. (a) We want to show that, for all  $P$ ,

$$U_\alpha(f, P) - L_\alpha(f, P) \leq w_f(||P||) (\alpha(b) - \alpha(a))$$

By the definition of the Upper and Lower Darboux sums,

$$\begin{aligned} U_\alpha(f, P) - L_\alpha(f, P) &= \sum_{i=1}^n M_i(f) \Delta\alpha_i - \sum_{i=1}^n m_i(f) \Delta\alpha_i \\ &= \sum_{i=1}^n [M_i(f) - m_i(f)] \Delta\alpha_i \end{aligned} \tag{1}$$

Now let  $\bar{x}_i$  and  $\underline{x}_i$  be the points such that

$$\begin{aligned} f(\underline{x}_i) &= m_i(f) = \inf_{x \in [x_i, x_{i-1}]} f(x) \\ f(\bar{x}_i) &= M_i(f) = \sup_{x \in [x_i, x_{i-1}]} f(x) \end{aligned}$$

Now since  $\underline{x}_i$  and  $\bar{x}_i$  are both in  $[x_{i-1}, x_i]$ , we can combine this with the definition of  $||P||$  to get

$$|\underline{x}_i - \bar{x}_i| \leq |x_i - x_{i-1}| \leq ||P||$$

By the definition of the modulus of continuity,

$$\begin{aligned} |\underline{x}_i - \bar{x}_i| \leq ||P|| &\Rightarrow |f(\underline{x}_i) - f(\bar{x}_i)| \leq w_f(||P||) \\ &\Leftrightarrow |M_i(f) - m_i(f)| \leq w_f(||P||) \end{aligned}$$

Substituting this fact back into 1, and using the properties of telescoping sums, we get that

$$\begin{aligned} \sum_{i=1}^n [M_i(f) - m_i(f)] \Delta\alpha_i &\leq \sum_{i=1}^n w_f(||P||) \Delta\alpha_i = w_f(||P||) \sum_{i=1}^n \Delta\alpha_i \\ &\leq w_f(||P||) (\alpha(b) - \alpha(a)) \end{aligned}$$

- (b) We know  $f$  is monotone on  $[a, b]$ . Suppose that  $f$  is monotone increasing.<sup>1</sup> Then given any partition,  $P$ ,

$$f(x_{i-1}) \leq f(c) \leq f(x_i) \quad \forall i, c \in [x_{i-1}, x_i]$$

This implies that

$$M_i = f(x_i) \quad m_i = f(x_{i-1}) \quad \forall i \quad (2)$$

Writing  $U_\alpha - L_\alpha$  as above in Equation 1, we see that the expression reduces to

$$\sum_{i=1}^n [M_i(f) - m_i(f)] \Delta\alpha_i = \sum_{i=1}^n [f(x_i) - f(x_{i-1})] \Delta\alpha_i \quad (3)$$

Now let's consider  $w_\alpha(||P||)$ . By definition, it is

$$w_\alpha(||P||) := \sup_{|x-y| \leq ||P||} |\alpha(x) - \alpha(y)|$$

It's clear that for each sub-interval,  $[x_{i-1}, x_i]$ , we have

$$|x_i - x_{i-1}| \leq ||P|| \quad \forall i$$

This implies that

$$|\Delta\alpha_i| = |\alpha(x_i) - \alpha(x_{i-1})| \leq w_\alpha(||P||) \quad \forall i$$

Subbing this into the right-hand side of equation 3, we can simplify using the properties of telescoping sums to get

$$\begin{aligned} \sum_{i=1}^n [f(x_i) - f(x_{i-1})] \Delta\alpha_i &\leq \sum_{i=1}^n [f(x_i) - f(x_{i-1})] w_\alpha(||P||) \\ &\leq w_\alpha(||P||) \sum_{i=1}^n [f(x_i) - f(x_{i-1})] \\ &\Leftrightarrow U_\alpha(f, P) - L_\alpha(f, P) \leq w_\alpha(||P||) [f(b) - f(a)] \end{aligned} \quad (4)$$

Finally, since  $\alpha$  is assumed continuous on the compact interval  $[a, b]$ ,  $\alpha$  is uniformly continuous on  $[a, b]$ . That means

$$\forall \epsilon > 0 \quad \exists \delta > 0 \quad \text{s.t.} \quad |x - y| \leq \delta \quad \Rightarrow \quad |\alpha(x) - \alpha(y)| \leq \epsilon$$

---

<sup>1</sup> Note that I'll lose no generality in assuming that the function  $f$  was monotone increasing. If it's monotone decreasing instead, swap the values of  $M_i$  and  $m_i$  in Equation 2, and proceed in exactly the same way.

So if we can make  $w_\alpha(||P||) < \epsilon$ , by choosing our partition  $P$  such that  $||P|| \leq \delta$ . With Equation 4, we therefore ensure that

$$U_\alpha(f, P) - L_\alpha(f, P) \leq \epsilon[f(b) - f(a)] \quad (5)$$

for any arbitrary  $\epsilon$ . And so by Riemann's Condition,  $f \in \mathcal{R}_\alpha([a, b])$  because for any  $\epsilon > 0$ , we can ensure the upper and lower sums are within that distance of each other by choosing a sufficiently fine partition.

2. **Exercise 51.15:** First, let's establish some basic building blocks for the proof. We have  $\alpha$  increasing on  $[a, b]$  and  $f, g \in \mathcal{R}_\alpha([a, b])$ . By Riemann's condition, it's clear that for all  $\delta > 0$ , there exist partitions  $P_1$  and  $P_2$  such that

$$U_\alpha(f, P_1) - L_\alpha(f, P_1) < \delta \quad (6)$$

$$U_\alpha(g, P_2) - L_\alpha(g, P_2) < \delta \quad (7)$$

Now take the common refinement,  $P^* = P_1 \cup P_2$ . By Lemma 51.5 and Corollary 51.6 (in FoMA), we know that

$$L_\alpha(f, P_1) \leq L_\alpha(f, P^*) \leq U_\alpha(f, P^*) \leq U_\alpha(f, P_1)$$

$$L_\alpha(g, P_2) \leq L_\alpha(g, P^*) \leq U_\alpha(g, P^*) \leq U_\alpha(g, P_2)$$

Combining this with the result from Riemann's condition above (and rewriting as in Question 1), we see that we must also have

$$\begin{aligned} \sum_{i=1}^n [M_i(f) - m_i(f)] \Delta\alpha_i &= U_\alpha(f, P^*) - L_\alpha(f, P^*) \leq U_\alpha(f, P_1) - L_\alpha(f, P_1) < \delta \\ \sum_{i=1}^n [M_i(g) - m_i(g)] \Delta\alpha_i &= U_\alpha(g, P^*) - L_\alpha(g, P^*) \leq U_\alpha(g, P_2) - L_\alpha(g, P_2) < \delta \end{aligned}$$

Now since  $\alpha$  is assumed increasing, we know that  $\Delta\alpha_i \geq 0$  for all  $i$ . Also,  $M_i(\cdot) \geq m_i(\cdot)$  for all  $i$ . Therefore, we can conclude

$$0 \leq M_i(f) - m_i(f) < \delta \quad \forall i \quad (8)$$

$$0 \leq M_i(g) - m_i(g) < \delta \quad \forall i \quad (9)$$

Now, using all of this as groundwork, let's show the main result.

- (a) We'll start by showing  $h(x) = \max\{f, g\} \in \mathcal{R}_\alpha([a, b])$  using Riemann's Condition. So for all  $\epsilon > 0$ , we need to find a partition  $P$  such that

$$U_\alpha(h, P) - L_\alpha(h, P) < \epsilon \quad (10)$$

To do so, take  $\delta = \epsilon/[\alpha(b) - \alpha(a)]$  and use Riemann's condition to find  $P_1$  and  $P_2$  as above, in Equations 6 and 7. Then take their common refinement to find  $P^*$ . **This will be our partition  $P$  such that Equation 10 holds.**

To formally show this, consider any arbitrary interval defined by the partition  $P^*$ . Over any interval  $[x_{i-1}, x_i]$  in  $P^*$ ,

$$\begin{aligned} M_i(h) &= \sup_{x \in [x_{i-1}, x_i]} \max\{f(x), g(x)\} \\ m_i(h) &= \inf_{x \in [x_{i-1}, x_i]} \max\{f(x), g(x)\} \end{aligned}$$

It's clear that we can narrow the list of candidates for  $M_i(h)$  and  $m_i(h)$ :

$$\begin{aligned} M_i(h) &= \max\{M_i(f), M_i(g)\} \\ m_i(h) &= \max\{m_i(f), m_i(g)\} \end{aligned}$$

Let's consider the cases:

- i. Suppose  $M_i(h) = M_i(f)$  and  $m_i(h) = m_i(f)$ . Then by Equation 8 and our choice of  $\delta$ , we have

$$0 \leq M_i(h) - m_i(h) = M_i(f) - m_i(f) \leq \frac{\epsilon}{\alpha(b) - \alpha(a)}$$

- ii. Similarly, if  $M_i(h) = M_i(g)$  and  $m_i(h) = m_i(g)$ . Then by Equation 9 and our choice of  $\delta$ , we have

$$0 \leq M_i(h) - m_i(h) = M_i(g) - m_i(g) \leq \frac{\epsilon}{\alpha(b) - \alpha(a)}$$

- iii. Next, suppose that  $M_i(h) = M_i(f)$  and  $m_i(h) = m_i(g)$ . In that case,

$$m_i(g) > m_i(f) \quad \Rightarrow \quad M_i(f) - m_i(g) < M_i(f) - m_i(f)$$

We also know that  $M_i(f) - m_i(g)$  is bounded below by zero because if not, then  $m_i(g) > M_i(f)$ , implying that we didn't choose  $M_i(h)$  correctly, as  $M_i(g)$  would certainly have been larger than  $m_i(g)$  and, thus also  $M_i(f)$ .

So by this fact, Equation 8, and our choice of  $\delta$ :

$$0 \leq M_i(h) - m_i(h) = M_i(f) - m_i(g) \leq M_i(f) - m_i(f) \leq \frac{\epsilon}{\alpha(b) - \alpha(a)}$$

iv. Finally, suppose that  $M_i(h) = M_i(g)$  and  $m_i(h) = m_i(f)$ . In that case,

$$m_i(f) > m_i(g) \quad \Rightarrow \quad M_i(g) - m_i(f) < M_i(g) - m_i(g)$$

We also know that  $M_i(g) - m_i(f)$  is bounded below by zero because if not, then  $m_i(f) > M_i(g)$ , implying that we didn't choose  $M_i(h)$  correctly, as  $M_i(f)$  would certainly have been larger than  $m_i(f)$  and, thus also  $M_i(g)$ .

So by this fact, Equation 9, and our choice of  $\delta$ :

$$0 \leq M_i(h) - m_i(h) = M_i(g) - m_i(f) \leq M_i(g) - m_i(g) \leq \frac{\epsilon}{\alpha(b) - \alpha(a)}$$

Putting it all together, we managed to bound

$$M_i(h) - m_i(h) \leq \frac{\epsilon}{\alpha(b) - \alpha(a)} \quad \forall i$$

This implies that

$$\begin{aligned} U_\alpha(h, P^*) - L_\alpha(h, P^*) &= \sum_{i=1}^n [M_i(h) - m_i(h)] \Delta\alpha_i \leq \sum_{i=1}^n \frac{\epsilon}{\alpha(b) - \alpha(a)} \Delta\alpha_i \\ &\leq \frac{\epsilon}{\alpha(b) - \alpha(a)} \sum_{i=1}^n \Delta\alpha_i \\ &\leq \frac{\epsilon}{\alpha(b) - \alpha(a)} \cdot (\alpha(b) - \alpha(a)) = \epsilon \end{aligned}$$

So we have a process to get the partition to satisfy Riemann's Condition for integrability for any  $\epsilon$  implying  $h \in \mathcal{R}_\alpha([a, b])$ .

- (b) Next, we show that  $h(x) = \min\{f, g\} \in \mathcal{R}_\alpha([a, b])$  using Riemann's Condition. So for all  $\epsilon > 0$ , we need to again find a sufficient partition  $P$ .

Also again, take  $\delta = \epsilon/[\alpha(b) - \alpha(a)]$  and use Riemann's condition to find  $P_1$  and  $P_2$  as above, in Equations 6 and 7. Then take their common refinement to find  $P^*$ . Again, **this will be our partition  $P$  such that Equation 10 holds.**

To formally show this, consider any arbitrary interval defined by the partition  $P^*$ . Over any interval  $[x_{i-1}, x_i]$  in  $P^*$ ,

$$\begin{aligned} M_i(h) &= \sup_{x \in [x_{i-1}, x_i]} \min\{f(x), g(x)\} \\ m_i(h) &= \inf_{x \in [x_{i-1}, x_i]} \min\{f(x), g(x)\} \end{aligned}$$

Narrowing down the list of candidates for  $M_i(h)$  and  $m_i(h)$ :

$$\begin{aligned} M_i(h) &= \min\{M_i(f), M_i(g)\} \\ m_i(h) &= \min\{m_i(f), m_i(g)\} \end{aligned}$$

The cases are then *exactly analogous* to what we saw for the case of the max. The result is that we again managed to bound

$$M_i(h) - m_i(h) \leq \frac{\epsilon}{\alpha(b) - \alpha(a)} \quad \forall i$$

This implies that

$$\begin{aligned} U_\alpha(h, P^*) - L_\alpha(h, P^*) &= \sum_{i=1}^n [M_i(h) - m_i(h)] \Delta\alpha_i \leq \sum_{i=1}^n \frac{\epsilon}{\alpha(b) - \alpha(a)} \Delta\alpha_i \\ &\leq \frac{\epsilon}{\alpha(b) - \alpha(a)} \sum_{i=1}^n \Delta\alpha_i \\ &\leq \frac{\epsilon}{\alpha(b) - \alpha(a)} \cdot (\alpha(b) - \alpha(a)) = \epsilon \end{aligned}$$

So we have a process to get the partition to satisfy Riemann's Condition for integrability for any  $\epsilon$  implying  $h \in \mathcal{R}_\alpha([a, b])$ .

3. **Exercise 51.18:** We want an example of an increasing  $\alpha$  on  $[a, b]$  and a bounded function  $f$  such that  $|f| \in \mathcal{R}_\alpha([a, b])$  but  $f \notin \mathcal{R}_\alpha([a, b])$ .

Consider the interval  $[-1, 1]$  and the functions

$$f(x) = \begin{cases} -1 & x < 0 \\ 1 & x \geq 0 \end{cases} \quad \alpha(x) = \begin{cases} 0 & x < 0 \\ 1 & x \geq 0 \end{cases} \quad (11)$$

First, for  $|f| \in \mathcal{R}_\alpha([a, b])$ , we have a situation similar to Example 51.9 in the book, as

$$|f(x)| = 1$$

a constant. Then for any partition  $P$ , we have

$$M_i(|f|) = m_i(|f|) = 1$$

which implies (after telescoping the sum) that

$$\begin{aligned} U_\alpha(|f|, P) &= \sum_{i=1}^n M_i(|f|) \Delta\alpha(i) = \sum_{i=1}^n 1 \cdot \Delta\alpha(i) \\ &= \alpha(b) - \alpha(a) = 1 - 0 = 1 \\ L_\alpha(|f|, P) &= \sum_{i=1}^n m_i(|f|) \Delta\alpha(i) = \sum_{i=1}^n 1 \cdot \Delta\alpha(i) \\ &= \alpha(b) - \alpha(a) = 1 - 0 = 1 \end{aligned}$$

And so we satisfy Riemann's Condition as

$$U_\alpha(|f|, P) - L_\alpha(|f|, P) = 0 \leq \epsilon$$

for all  $\epsilon > 0$ , which implies  $|f| \in \mathcal{R}_\alpha([a, b])$

**Show**  $f \notin \mathcal{R}_\alpha([a, b])$

4.

5.