

# Tidy Time Series & Forecasting in R

## 3. Transformations



# Outline


- 1 Per capita adjustments
- 2 Lab Session 6
- 3 Inflation adjustments
- 4 Mathematical transformations
- 5 Lab Session 7

# Outline

- 1 Per capita adjustments
- 2 Lab Session 6
- 3 Inflation adjustments
- 4 Mathematical transformations
- 5 Lab Session 7


# Per capita adjustments

```
global_economy %>%  
  filter(Country == "Australia") %>%  
  autoplot(GDP)
```



# Per capita adjustments

```
global_economy %>%  
  filter(Country == "Australia") %>%  
  autoplot(GDP / Population)
```



# Outline

- 1 Per capita adjustments
- 2 Lab Session 6
- 3 Inflation adjustments
- 4 Mathematical transformations
- 5 Lab Session 7

## Lab Session 6

Consider the GDP information in `global_economy`. Plot the GDP per capita for each country over time. Which country has the highest GDP per capita? How has this changed over time?

# Outline


- 1 Per capita adjustments
- 2 Lab Session 6
- 3 Inflation adjustments**
- 4 Mathematical transformations
- 5 Lab Session 7



# Inflation adjustments

```
print_retail <- aus_retail %>%  
  filter(Industry == "Newspaper and book retailing") %>%  
  group_by(Industry) %>%  
  index_by(Year = year(Month)) %>%  
  summarise(Turnover = sum(Turnover))  
aus_economy <- filter(global_economy, Code == "AUS")  
print_retail %>%  
  left_join(aus_economy, by = "Year") %>%  
  mutate(Adj_turnover = Turnover / CPI) %>%  
  pivot_longer(c(Turnover, Adj_turnover),  
    names_to = "Type", values_to = "Turnover"  
  ) %>%  
  ggplot(aes(x = Year, y = Turnover)) +  
  geom_line() +  
  facet_grid(vars(Type), scales = "free_y") +  
  xlab("Years") + ylab(NULL) +  
  ggtitle("Turnover: Australian print media industry")
```

# Inflation adjustments



# Outline

- 1 Per capita adjustments
- 2 Lab Session 6
- 3 Inflation adjustments
- 4 Mathematical transformations
- 5 Lab Session 7

# Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

# Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as  $y_1, \dots, y_n$  and transformed observations as  $w_1, \dots, w_n$ .

# Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as  $y_1, \dots, y_n$  and transformed observations as  $w_1, \dots, w_n$ .

## Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	$\downarrow$
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

# Variance stabilization

If the data show different variation at different levels of the series, then a transformation can be useful.

Denote original observations as  $y_1, \dots, y_n$  and transformed observations as  $w_1, \dots, w_n$ .


## Mathematical transformations for stabilizing variation

Square root	$w_t = \sqrt{y_t}$	$\downarrow$
Cube root	$w_t = \sqrt[3]{y_t}$	Increasing
Logarithm	$w_t = \log(y_t)$	strength

Logarithms, in particular, are useful because they are more interpretable: changes in a log value are **relative (percent) changes on the original scale**.

# Variance stabilization


```
food <- aus_retail %>%  
  filter(Industry == "Food retailing") %>%  
  summarise(Turnover = sum(Turnover))
```






# Variance stabilization

```
food %>% autoplot(sqrt(Turnover)) +  
  labs(y = "Square root turnover")
```




# Variance stabilization

```
food %>% autoplot(Turnover^(1 / 3)) +  
  labs(y = "Cube root turnover")
```




# Variance stabilization

```
food %>% autoplot(log(Turnover)) +  
  labs(y = "Log turnover")
```



# Variance stabilization

```
food %>% autoplot(-1 / Turnover) +  
  labs(y = "Inverse turnover")
```



# Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

# Box-Cox transformations

Each of these transformations is close to a member of the family of **Box-Cox transformations**:

$$w_t = \begin{cases} \log(y_t), & \lambda = 0; \\ (y_t^\lambda - 1)/\lambda, & \lambda \neq 0. \end{cases}$$

- $\lambda = 1$ : (No substantive transformation)
- $\lambda = \frac{1}{2}$ : (Square root plus linear transformation)
- $\lambda = 0$ : (Natural logarithm)
- $\lambda = -1$ : (Inverse plus 1)

# Box-Cox transformations

# Box-Cox transformations

```
food %>%  
  features(Turnover, features = guerrero)
```

```
## # A tibble: 1 x 1  
##   lambda_guerrero  
##               <dbl>  
## 1              0.0895
```



# Box-Cox transformations


```
food %>%  
  features(Turnover, features = guerrero)
```

```
## # A tibble: 1 x 1  
##   lambda_guerrero  
##               <dbl>  
## 1              0.0895
```

- This attempts to balance the seasonal fluctuations and random variation across the series.
- Always check the results.
- A low value of  $\lambda$  can give extremely large prediction intervals.

# Box-Cox transformations

```
food %>% autoplot(box_cox(Turnover, 0.0524)) +  
  labs(y = "Box-Cox transformed turnover")
```



# Transformations

- Often no transformation needed.
- Simple transformations are easier to explain and work well enough.
- Transformations can have very large effect on PI.
- If the data contains zeros, then don't take logs.
- `logp1()` can be useful for data with zeros.
- If some data are negative, no power transformation is possible unless a constant is added to all values.
- Choosing logs is a simple way to force forecasts to be positive
- Transformations must be reversed to obtain forecasts on the original scale. (Handled automatically by `fable`.)

# Outline

- 1 Per capita adjustments
- 2 Lab Session 6
- 3 Inflation adjustments
- 4 Mathematical transformations
- 5 Lab Session 7

# Lab Session 7

- 1 For the following series, find an appropriate transformation in order to stabilise the variance.
  - ▶ United States GDP from `global_economy`
  - ▶ Slaughter of Victorian “Bulls, bullocks and steers” in `aus_livestock`
  - ▶ Victorian Electricity Demand from `vic_elec`.
  - ▶ Gas production from `aus_production`
- 2 Why is a Box-Cox transformation unhelpful for the `canadian_gas` data?