


Probability and Statistics

Probability: Given Model, how likely is Data? → Well-formed since these are Mathematical questions.

Statistics: Given Data, how likely is Model? → Ill-formed since many Models can generate the same data!



Probability Space

Probability Space: a triple (Ω, F, P) consisting of:

1. Ω the **Sample Space**
2. $F \subseteq 2^\Omega$ a σ -**algebra**¹ on Ω i.e.
 - (a) $\Omega \in F$
 - (b) $E \in F \Rightarrow E^c \in F$
 - (c) $E_1, E_2, \dots \in F \Rightarrow \bigcup_{i=1}^\infty E_i \in F$
3. $P: F \mapsto [0, 1]$ a **Probability Measure** i.e.
 - (a) $P(E) \geq 0$ for $E \in F$
 - (b) $P(\Omega) = 1$
 - (c) $P(\bigcup_{i=1}^\infty E_i) \Rightarrow \sum_{i=1}^\infty P(E_i)$ for $E_i \in F$

Given **Events** $E_i, E \in F$, P also satisfies:

1. **Upward and Downward continuity** of P :
 - (a) $E_i \uparrow E \Rightarrow \lim_{n \rightarrow \infty} P(E_n) = P(E)$
 - (b) $E_i \downarrow E \Rightarrow \lim_{n \rightarrow \infty} P(E_n) = P(E)$
2. **Monotonicity** of P :
 - (a) $E_i \subseteq E_j \Rightarrow P(E_i) \leq P(E_j)$

Conditional Probability

We can compute Probabilities of Events Conditioned on other Events.

Conditional Probability of event A on event B with $P(B) > 0$ is:

$$P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{P(B | A)P(A)}{P(B)}$$

A set of events $\{A_i\}$ are **Mutually Independent** if, for any n element subset of $\{A_i\}$:

$$P\left(\bigcap_{i=1}^n A_i\right) = \prod_{i=1}^n P(A_i)$$

Law of Total Probability: Given Events A and **Partition** $\{B_i\}$ (i.e. where $\bigcup_{i=1}^\infty B_i = \Omega$)


$$P(A) = \sum_{i=1}^\infty P(A | B_i)P(B_i)$$

Random Variables

A **Random Variable** is a \mathbb{B} -Measurable function

$X: (\Omega, F) \mapsto (\mathbb{R}, \mathbb{B})^2$

1. For $A \in \mathbb{B}$ we can compute $P(X \in A)$
2. $P(X \in A) := P(X^{-1}(A)) = P(\{\omega \in \Omega : X(\omega) \in A\})$
3. $P(X^{-1}(\cdot)) := P_X(\cdot)$ which is called the **Push-Forward Measure** of P by X on \mathbb{R}
4. Hence X induces a new Probability Space $(\mathbb{R}, \mathbb{B}, P_X)$ from the original (Ω, F, P)



Random Variables can be uniquely determined by their CDF:
 $F_X(t) := P_X((-\infty, t]) = P(X \leq t)$

1. Right Continuous
2. Non-Negative
3. $\lim_{t \rightarrow \infty} F_X(t) = 1, \lim_{t \rightarrow -\infty} F_X(t) = 0$
4. Any function satisfying above properties is the CDF for some random variable.

Random Variables studied are usually either **Continuous** or **Discrete** (although they can be neither.)

1. If F_X **Absolutely Continuous** then X is a Continuous RV.³
 - (a) **Absolutely Continuous:** F differentiable a.e. and $\exists f(x)$ s.t. $F_X(x) = \int_{-\infty}^x f(u)du$
 - (b) $\Rightarrow \frac{d}{dx} F_X(x) = f(x)$ wherever F is differentiable
 - (c) f is called the **PDF**
 - (d) f is unique a.e. (may not be everywhere!)
 - (e) If X also Non-Negative then **Hazard** of X is $\lambda(t) = \frac{f(t)}{1-F(t)}$
 - i. $1 - F(t) = \exp\left(-\int_0^t \lambda(x) dx\right)$
 - ii. $\lambda(t)$ interpreted as instantaneous survival rate at time t .
 - iii. $\lambda(t) = c \forall t \iff X \sim \text{Exp}(c)$

2. If $\text{Range}(X)$ is countable then X is Discrete.

- (a) $f(x) := P(\{X = x\})$
- (b) analogously, $F_X(t) = \sum_{i=0}^t f(i)$ ⁴
- (c) f is called the **PMF**

Random Vectors

Joint CDF for $\vec{X} = (X_1, X_2, \dots, X_n)$ is

$$F(t_1, \dots, t_n) = P(X_1 \leq t_1, \dots, X_n \leq t_n)$$

1. **Marginal PDF** of $\vec{X}_{1:p} = (X_1, \dots, X_p)$ is

$$f_{\vec{X}_{1:p}}(\vec{u}_{1:p}) = \int_{\vec{X}_{(p+1):n}} f_{\vec{X}}(\vec{u}_{1:p}, \vec{X}_{(p+1):n}) d\vec{X}_{(p+1):n}$$

2. **Conditional PDF** on $\vec{X}_{1:p}$ given $\vec{X}_{(p+1):n}$ is

$$f_{\vec{X}_{1:p}|\vec{X}_{(p+1):n}}(\vec{u}_{1:p}, \vec{u}_{(p+1):n}) = \frac{f_{\vec{X}}(\vec{u}_{1:p}, \vec{u}_{(p+1):n})}{f_{\vec{X}_{(p+1):n}}(\vec{u}_{(p+1):n})}$$

3. **kth Order Statistic** $X_{(k)}$ of \vec{X} is the kth smallest value

$$(a) f_{X_{(1)}}(u) = \sum_{i=1}^n f_{X_i}(u) \prod_{j \neq i} (1 - F_{X_j}(u))$$

$$(b) f_{X_{(n)}}(u) = \sum_{i=1}^n f_{X_i}(u) \prod_{j \neq i} F_{X_j}(u)$$

Moments of a Random Variable

$\mathbb{E}_X(X^r) := \mathbb{E}(X^r)$ is the **rth Moment** of X under the distribution of X

1. $\mathbb{E}(X) = \int_0^\infty 1 - F_X(t) dt - \int_{-\infty}^0 F_X(t) dt$
2. If X Continuous, $\mathbb{E}(X) = \int_{-\infty}^\infty t \cdot f_X(t) dt$
3. **LOTUS**: $\mathbb{E}(g(X)) = \int_{-\infty}^\infty g(t) \cdot f_X(t) dt$
4. Moments need not exist! (i.e. $E(|X^r|) = \pm\infty$)

Can generate Moments using the **MGF** of X :

$M_X(t) = \mathbb{E}(\exp(Xt))$, if $\exists \epsilon > 0$ s.t. $\forall |t| < \epsilon, M_X(t) < \infty$

1. $\exists \epsilon > 0$ s.t. $\forall |t| < \epsilon, M_X(t) = M_Y(t) \Rightarrow X$ and Y have same distribution
2. $\mathbb{E}(|X^r|) = \frac{\partial^r}{\partial t^r} M_X(t) \Big|_{t=0}$, if M_X exists.
3. If $\{X_i\}$ independent RVs, then $M_{\sum X_i}(t) = \prod M_{X_i}(t)$

Moments most commonly analyzed are:

1. **Mean** of X : $\mathbb{E}(X) := \mu_X$
2. **Variance** of X : $Var(X) = \mathbb{E}((X - \mu_X)^2) = \sigma_X^2$

For random vectors \vec{X} we have:

1. $\mathbb{E}(\vec{X}) = [\mathbb{E}(X_1), \dots, \mathbb{E}(X_n)] = \vec{\mu}$
2. $Cov(\vec{X}) = \mathbb{E}[(\vec{X} - \vec{\mu})(\vec{X} - \vec{\mu})^T] = \Sigma$

If X and Y are Random Variables on the same Probability Space

1. **Law of Total Expectation**: If $\mathbb{E}(|X|) < \infty$
 $\mathbb{E}(X) = \mathbb{E}_Y(\mathbb{E}_{X|Y}(X | Y))$
2. **Law of Total Variance**: If $Var(X) < \infty$
 $Var(X) = \mathbb{E}(Var(X | Y)) + Var(\mathbb{E}(X | Y))$

Parametric Family

A **parametric model** is a family of distributions that is defined by a fixed finite number of parameters.

1. Formally, $\mathcal{P}_\Theta = \{p_\theta(\cdot; \theta) \mid \theta \in \Theta\}$ where $p_\theta(\cdot; \theta)$ is a possible density depending on the **Parameter** θ and Θ is the **Parameter Space**
- 2.

Asymptotic Results

Notes

1. For some Ω we cannot use 2^Ω as a σ -algebra since this may contain sets which do not satisfy all of the axioms. See [1] for an example.
2. Where \mathbb{B} is the **Borel σ -algebra**: the smallest σ -algebra containing all the open intervals. This must contain all intervals of the form $(-\infty, x]$, and since X measurable $\Rightarrow F_X$ guaranteed to exist.
3. Continuity of X as a function on Ω has nothing to do with its continuity as a Random Variable (which depends on the absolute continuity of its CDF) [2]
4. Taking P to be the **Counting Measure**, it can be shown that the Lebesgue integral is this sum.

References

- [1] J. S. Rosenthal, *A first look at rigorous probability theory*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, second ed., 2006.
- [2] pidgeot, “On clarifying the relationship between distribution functions in measure theory and probability theory.” Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/976739> (version: 2014-10-16).