


Probability and Statistics

Matthew Scicluna
matthew.scicluna@umontreal.ca

Probability Vs Statistics

Probability: Given Model, how likely is Data? → Well-formed since these are Mathematical questions.

Statistics: Given Data, how likely is Model? → Ill-formed since many Models can create the same data!



Probability Space

Probability Space: a triple (Ω, F, P) consisting of:

1. Ω the **Sample Space**
2. $F \subseteq 2^\Omega$ a σ -algebra on Ω i.e.
 - (a) $\Omega \in F$
 - (b) $E \in F \Rightarrow E^c \in F$
 - (c) $E_1, E_2, \dots \in F \Rightarrow \bigcup_{i=1}^\infty E_i \in F$
3. $P: F \mapsto [0, 1]$ a **Probability Measure** i.e.
 - (a) $P(E) \geq 0$ for $E \in F$
 - (b) $P(\Omega) = 1$
 - (c) $P(\bigcup_{i=1}^\infty E_i) \Rightarrow \sum_{i=1}^\infty P(E_i)$ for $E_i \in F$

Given **Events** $E_i, E \in F$, P also satisfies:

1. **Upward and Downward continuity** of P :
 - (a) $E_i \uparrow E \Rightarrow \lim_{n \rightarrow \infty} P(E_n) = P(E)$
 - (b) $E_i \downarrow E \Rightarrow \lim_{n \rightarrow \infty} P(E_n) = P(E)$
2. **Monotonicity** of P :
 - (a) $E_i \subseteq E_j \Rightarrow P(E_i) \leq P(E_j)$

Conditional Probability

We can compute Probabilities of Events Conditioned on other Events.

Conditional Probability of event A on event B with $P(B) > 0$ is:

$$P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{P(B | A)P(A)}{P(B)}$$

A set of events $\{A_i\}$ are **Mutually Independent** if, for any n element subset of $\{A_i\}$:

$$P\left(\bigcap_{i=1}^n A_i\right) = \prod_{i=1}^n P(A_i)$$

Law of Total Probability: Given Events A and **Partition** $\{B_i\}$ (i.e. where $\bigsqcup_{i=1}^\infty B_i = \Omega$)


$$P(A) = \sum_{i=1}^\infty P(A | B_i)P(B_i)$$

Random Variables

A **Random Variable** is a \mathbb{B} -Measurable function

$X: (\Omega, F) \mapsto (\mathbb{R}, \mathbb{B})$

1. For $A \in \mathbb{B}$ we can compute $P(X \in A)$
2. $P(X \in A) := P(X^{-1}(A)) = P(\{\omega \in \Omega : X(\omega) \in A\})$
3. $P(X^{-1}(\cdot)) := P_X(\cdot)$ which is called the **Push-Forward Measure** of P by X on \mathbb{R}
4. Hence X induces a new Probability Space $(\mathbb{R}, \mathbb{B}, P_X)$ from the original (Ω, F, P)



Continuous Random Variables test

Moments of a Random Variable