

# MXB103 Exam Notes

## Ordinary Differential Equations

### Euler's Method

what's given: function, where you want to approximate, step value (h), and initial condition (when  $n = 0$ )

follow the formula:  $y_{n+1} = y_n + hf'(x_n)$

n	$x_n$	$y_n$
0	initial x	initial y
1	initial x + h	follow fomula
2	$x_1 + h$	follow fomula
3	$x_2 + h$	follow fomula
4	$x_3 + h$	follow fomula
5	$x_4 + h$	follow fomula

### Second Order Taylor Method

1. Identify  $f(x)$  and the point  $a$ .
2. Compute derivatives  $f'(x)$  and  $f''(x)$ .
3. Evaluate  $f'(a)$  and  $f''(a)$ .
4. Construct the Taylor polynomial:

$$P_2(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2$$

5. Use  $P_2(x)$  to approximate  $f(x)$  near  $a$ .

### Modified Euler's Method

1. Given  $y_n$  and  $x_n$ , compute the initial slope  $k_1 = f(x_n, y_n)$ .
2. Predict the next value  $\tilde{y}_{n+1} = y_n + k_1 h$ , where  $h$  is the step size.
3. Compute the corrected slope  $k_2 = f(x_n + h, \tilde{y}_{n+1})$ .
4. Calculate the next value  $y_{n+1} = y_n + \frac{h}{2}(k_1 + k_2)$ .
5. Update  $x_{n+1} = x_n + h$  and repeat from step 1 for further iterations.

### Runge-Kutta 4th Order (RK4)

1. Start with initial conditions  $y_n$  and  $x_n$ .
2. Calculate the first slope  $k_1 = hf(x_n, y_n)$ .
3. Compute the second slope  $k_2 = hf(x_n + \frac{h}{2}, y_n + \frac{k_1}{2})$ .
4. Calculate the third slope  $k_3 = hf(x_n + \frac{h}{2}, y_n + \frac{k_2}{2})$ .
5. Compute the fourth slope  $k_4 = hf(x_n + h, y_n + k_3)$ .
6. Estimate the next value  $y_{n+1} = y_n + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)$ .
7. Increment  $x_{n+1} = x_n + h$  and use the new  $x_{n+1}$  and  $y_{n+1}$  for the next iteration.

## Polynomial Interpolation

### Lagrange Interpolating Polynomial

1. sub abscissas into given function to get y values.
2. find the Polynomial for the amount of abscissas. Ex:  
 $(x_0, y_0), (x_1, y_1), (x_2, y_2)$   
 $\frac{(x-x_1)(x-x_2)}{(x_0-x_1)(x_0-x_2)} \cdot y_0 + \text{same thing for } y_1 \text{ and } y_2$

### Newton's Divided Difference

$$a_0 = y_0, a_1 = \frac{y_1 - y_0}{x_1 - x_0}, a_2 = \frac{\frac{y_2 - y_1}{x_2 - x_1} - \frac{y_1 - y_0}{x_1 - x_0}}{x_2 - x_0}$$

$x_i$	zeroth	first	second	third
$x_0$	$f[x_0]$	$f[x_0, x_1]$	$f[x_0, x_1, x_2]$	$f[x_0, x_1, x_2, x_3]$
$x_1$	$f[x_1]$	$f[x_1, x_2]$	$f[x_1, x_2, x_3]$	
$x_2$	$f[x_2]$	$f[x_2, x_3]$		
$x_3$	$f[x_3]$			

$$P_3(x) = f[x_0] + f[x_0, x_1](x - x_0) + f[x_0, x_1, x_2](x - x_0)(x - x_1) + f[x_0, x_1, x_2, x_3](x - x_0)(x - x_1)(x - x_2)$$

1. Sub given x and y values into first and second columns of the table.
2. Compute first divided difference using above  $a_1$  formula.
3. Compute second divided difference using above  $a_2$  formula
4. Sub into above formula.

### Newton's Forward Difference

ONLY use when the abscissas are equally spaced.

$x_i$	$y_i$	$\Delta y_i$	$\Delta^2 y_i$	$\Delta^3 y_i$
$x_0$	$y_0$	$\Delta y_0$	$\Delta^2 y_0$	$\Delta^3 y_0$
$x_1$	$y_1$	$\Delta y_1$	$\Delta^2 y_1$	
$x_2$	$y_2$	$\Delta y_2$		
$x_3$	$y_3$			

1. sub given x and y values into first and second columns of table.
2. Compute remaining columns. Ex:  $\Delta y_0 = y_1 - y_0$
3. Sub into given polynomial equation.