

# SDS291\_FinalProject

## Importing Packages

```
library(moderndiver)
library(dplyr)
```

Attaching package: 'dplyr'

The following objects are masked from 'package:stats':

filter, lag

The following objects are masked from 'package:base':

intersect, setdiff, setequal, union

```
library(tidyverse)
```

```
-- Attaching packages ----- tidyverse 1.3.2
--
```

```
v ggplot2 3.4.0      v purrr   1.0.1
v tibble  3.1.8      v stringr 1.5.0
v tidyr   1.3.0      v forcats 1.0.0
v readr   2.1.4
```

```
-- Conflicts ----- tidyverse_conflicts() --
x dplyr::filter() masks stats::filter()
x dplyr::lag()    masks stats::lag()
```

```
library(ggplot2)
library(Stat2Data)
```

## Data Importing

```
lego_clean <- read.csv("lego_clean.csv")
lego_clean$fem <- factor(lego_clean$fem, labels = c("No", "Yes"))
lego_clean$masc <- factor(lego_clean$masc, labels = c("No", "Yes"))
lego_clean$neutral <- factor(lego_clean$neutral, labels = c("No", "Yes"))
```

## Exploratory Visualizations


```
price_piece_fem <- lm(Price ~ Pieces * fem, lego_clean)
ggplot(lego_clean, aes( x = Pieces, y = Price, color = fem)) +
  geom_point(alpha = 0.5) +
  geom_smooth(method = "lm") +
  labs(title = "Scatterplot of Lego Price as a Function of \n Number of Pieces and Gender
```

`geom\_smooth()` using formula = 'y ~ x'

Warning: Removed 239 rows containing non-finite values (`stat\_smooth()`).


Warning: Removed 239 rows containing missing values (`geom\_point()`).

Scatterplot of Lego Price as a Function of Number of Pieces and Gender (Feminine or not)




```
ggplot(lego_clean, aes(x = fem)) + geom_bar() + geom_text(stat = 'count', aes(label=after_
```


Counts of Feminine vs. Non-Feminine Marketed Lego Sets



```
ggplot(lego_clean, aes(x = masc)) + geom_bar() + geom_text(stat = 'count', aes(label=after
```



```
ggplot(lego_clean, aes(x = neutral)) + geom_bar() + geom_text(stat = 'count', aes(label=af
```




```
price_piece_masc <- lm(Price ~ Pieces * masc, lego_clean)
ggplot(lego_clean, aes( x = Pieces, y = Price, color = masc)) +
  geom_point(alpha = 0.5) +
  geom_smooth(method = "lm") +
  labs(title = "Scatterplot of Lego Price as a Function of \n Number of Pieces and Gender
```

``geom_smooth()`` using formula = 'y ~ x'

Warning: Removed 239 rows containing non-finite values (``stat_smooth()``).


Warning: Removed 239 rows containing missing values (``geom_point()``).

Scatterplot of Lego Price as a Function of  
Number of Pieces and Gender (Masculine or not)






```
price_piece_neutral <- lm(Price ~ Pieces * neutral, lego_clean)
ggplot(lego_clean, aes( x = Pieces, y = Price, color = neutral)) +
  geom_point(alpha = 0.5) +
  geom_smooth(method = "lm") +
  labs(title = "Scatterplot of Lego Price as a Function of \n Number of Pieces and Gender")
```

Scatterplot of Lego Price as a Function of  
Number of Pieces and Gender (Neutral or not)




```
plot(price_piece_neutral)
```














```
plot(price_piece_masc)
```











```
plot(price_piece_fem)
```










```
price_piece <- lm(Price ~ Pieces, data = lego_clean)
plot(price_piece)
```







```
get_regression_table(price_piece)
```

```
# A tibble: 2 x 7
```

	term	estimate	std_error	statistic	p_value	lower_ci	upper_ci
	<chr>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	intercept	10.9	0.986	11.0	0	8.93	12.8
2	Pieces	0.082	0.001	64.3	0	0.08	0.085

```
get_regression_table(price_piece_fem)
```

```
# A tibble: 4 x 7
```

	term	estimate	std_error	statistic	p_value	lower_ci	upper_ci
	<chr>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	intercept	11.4	1.05	10.9	0	9.37	13.5
2	Pieces	0.082	0.001	63.3	0	0.079	0.084
3	fem: Yes	-7.69	3.80	-2.02	0.043	-15.2	-0.228
4	Pieces:femYes	0.016	0.01	1.56	0.12	-0.004	0.036

```
get_regression_table(price_piece_masc)
```

```
# A tibble: 4 x 7
  term          estimate std_error statistic p_value lower_ci upper_ci
  <chr>          <dbl>    <dbl>    <dbl>   <dbl>   <dbl>   <dbl>
1 intercept      12.8      1.23     10.4     0       10.4    15.3
2 Pieces          0.072     0.002     46.4     0        0.069   0.075
3 masc: Yes      -6.62      1.90     -3.48   0.001    -10.4    -2.89
4 Pieces:mascYes  0.026     0.002     10.5     0         0.021   0.031
```

```
get_regression_table(price_piece_neutral)
```

```
# A tibble: 4 x 7
  term          estimate std_error statistic p_value lower_ci upper_ci
  <chr>          <dbl>    <dbl>    <dbl>   <dbl>   <dbl>   <dbl>
1 intercept      5.35      1.16      4.59     0        3.06    7.63
2 Pieces          0.099     0.002     58.4     0        0.096   0.102
3 neutral: Yes    10.2      1.89      5.37     0         6.45    13.9
4 Pieces:neutralYes -0.032     0.002    -13.6     0       -0.037  -0.027
```

## Model Comparison

*compare price\_piece model (consider this are nested model) to the models that look at both price and gender (this would be the full model) null hypothesis: the nested model that only looks at price as a function of pieces is enough (coefficient of sex has no effect = 0) \*alternative hypothesis: need the full model (coefficient of sex has an effect)*

```
anova(price_piece, price_piece_neutral)
```

## Analysis of Variance Table

Model 1: Price ~ Pieces

Model 2: Price ~ Pieces \* neutral

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	1063	755013				
2	1061	639726	2	115287	95.603	< 2.2e-16 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

\*whether a set is marked as gender neutral or not has an effect on price (the full model is better the fit) - Gender neutrality is a necessary component of the model



```
anova(price_piece, price_piece_masc)
```

#### Analysis of Variance Table

Model 1: Price ~ Pieces

Model 2: Price ~ Pieces \* masc

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	1063	755013				
2	1061	678766	2	76248	59.593	< 2.2e-16 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

\*whether a set is marked as masculine or not has an effect on price (the full model is a better fit) - masculinity is a necessary component of the model.

```
anova(price_piece, price_piece_fem)
```

#### Analysis of Variance Table

Model 1: Price ~ Pieces

Model 2: Price ~ Pieces \* fem

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	1063	755013				
2	1061	752110	2	2903.1	2.0477	0.1295

\*fail to reject the null hypothesis and conclude the reduced model (price\_piece) explains more variability in the dataset than a model accounting for female marketing.

## LINE Violations

\*all models violate the normality condition at the extremity points. Since we are primarily doing a model comparison, this violation carries through and a model comparison should still hold relatively well. This will impact generalizability to all lego sets (make it less generalizable).

## Comparing The Interaction Models to Each other

```
get_regression_summaries(price_piece_neutral)
```

```
# A tibble: 1 x 9
```

	r_squared	adj_r_squared	mse	rmse	sigma	statistic	p_value	df	nobs
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	0.827	0.826	601.	24.5	24.6	1688.	0	3	1065

```
get_regression_summaries(price_piece_masc)
```

```
# A tibble: 1 x 9
```

	r_squared	adj_r_squared	mse	rmse	sigma	statistic	p_value	df	nobs
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	0.816	0.816	637.	25.2	25.3	1570.	0	3	1065

\*the adjusted r squared values indicate that the neutral model is a better predictor of price when controlling for pieces. However, the difference in the adjusted r squared values is very low (0.01). Since the number lego sets marketed to a gender neutral audience is much higher than ones marked to a masculine audience, it is likely that both of these models to a similarly good job of predicting price when controlling for number of pieces.

## Interpreting Coefficients for Masculine Model:

- Intercept: For a non-masculine model with 0 pieces the predicted price is 12.84. \*Pieces: For each additional piece in a non-masculine set, the predicted price increase is 0.07.
- masc: yes: For a masculine model with 0 pieces, the predicted price is -6.62.
- Pieces: mascYes: For each additional piece in a masculine set, the predicted price increase is 0.03.