



**NANSEN ENVIRONMENTAL AND
REMOTE SENSING CENTER**

Edvard Griegsvei 3A
N-5037 Solheimsviken/Bergen
NORWAY
Phone: + 47 55 29 72 88
Fax: + 47 55 20 00 50
e-mail: Ola.Johannessen@nrsc.no


Your ref.:

Our ref.:

Bergen: **10.09.1996**

BEKREFTELSE

Vi bekrefter herved at Amir M. Hashemi, 18.02.66, er ansatt på Nansen Senteret fra 10.06.1996. Han er ansatt som Dr. stipendiat student i 60% stilling.



A Discussion on Acoustic Propagation in the Ice-covered Region of the Arctic Sea

Amir M. Hashemi

Bergen University College, Pb. 7030, 5020, Bergen, Norway, email: ahas@hib.no

The results of a numerical investigation of the acoustic propagation in the ice-covered region of Arctic Sea where the environment can be considered range-independent are presented. Mathematical equations for Green's function and normal stress for a simple three layered model (vacuum-solid-fluid-fluid halfspace) are presented and investigated numerically. Environment parameters are motivated from measurements from Greenland Sea during Marginal Ice Zone Experiment, MIZEX [4]. This discussion is an effort to understand the acoustic plane propagation in the ice-covered region of the Arctic Sea.

PACS no. 00.00.Xx .

1. Introduction

In the recent decades, the enormous progress in computer science has stimulated acousticians to develop frequency dependent solutions for wave equation. In these numerically based techniques wave equation is solved by normal mode, fast field, and parabolic equation methods [3]. The choice of propagation model depends on the dominant characteristics of the environment and the frequency range that is under consideration. Hydrological measurements in the Marginal Ice Zone (MIZEX) of the Greenland sea reveal the need of a range-independent horizontal layered model in ice covered regions. SAFARI [2] provides accurate transmission loss values for propagation in range-independent multi-layered horizontally stratified environments and was therefore an effective tool for this investigation.

The work is structured as follows. First solution of wave equation for a range-independent environment is presented and basic equations for Green's function and normal stress are given for a three layered environment (vacuum-solid-fluid-fluid halfspace) in Sec. 2 and a numerical example (case 1) is presented to study a deep ice-covered sea, three other numerical examples study more complicated environments in Sec. 3. Even if the chosen simulated environments are very simple, remarkable results are observed. A summary ends the discussion in Sec. 4.

2. Governing Equations

By employing forward Fourier and Hankel transform respectively on the linear wave equation in a range independent environment, the depth-separated wave equation can be obtained [4]:

$$\left[\frac{\partial^2}{\partial z^2} - (k^2 - k_m^2(z))\right]G(k, z) = \frac{F(z)}{2\pi}. \quad (1)$$

Where k_m is medium wavenumber, k is the horizontal wavenumber and $F(z)$ is forcing term. The total solution of the depth-separated equation can be expressed as a superposition of a homogeneous solution G_H , satisfying boundary conditions, and a particular solution G_P , the free field produced by a source in an unbounded medium. $G_P(k, z)$ term is added to the homogeneous solution in layers containing one or more sources:

$$G(k, z) = G_H(k, z) + G_P(k, z). \quad (2)$$

$G_H(k, z)$ is found by employing the boundary conditions.

To find the total field at any range r and the time response, we apply the inverse Hankel transform.

$$g(r, z) = \int_0^\infty G(k, z) k J_0 dk. \quad (3)$$

where $J_0 = J_0(kr)$ is the Bessel function. The normal stress τ_{zz} is then given by:

$$\tau_{zz} = -\rho\omega^2 g(r, z). \quad (4)$$

where ρ the fluid density, ω the radial frequency and $g(r, z)$ is the potential field.

In this paper $G(k, z)$ and τ_{zz} are studied. The graphical results for the normal stress (NS) are given in [dB/1Pa] corresponding to the transmission loss definition, see [4], p. 67. To investigate the acoustic propagation in the ice-covered region a simple three layered model is considered (vacuum, solid, fluid, fluid half-space).

The integral representation of the homogeneous solutions in the layer 1 can be expressed as:

$$g_{11}(r, z) = \int_0^\infty [A_1^-(k)e^{-\alpha_1 z} + A_1^+(k)e^{\alpha_1 z}] k J_0 dk, \quad (5)$$

$$\text{where } \alpha_1 = \begin{cases} -\sqrt{k^2 - k_1^2}, & k > k_1, \\ i\sqrt{k^2 - k_1^2}, & k \leq k_1. \end{cases}$$

$$g_{s1}(r, z) = \int_0^\infty [B_1^-(k)e^{-\beta_1 z} + B_1^+(k)e^{\beta_1 z}] k J_0 dk, \quad (6)$$

$$\text{where } \beta_1 = \begin{cases} -\sqrt{k^2 - k_s^2}, & k > k_s, \\ i\sqrt{k^2 - k_s^2}, & k \leq k_s. \end{cases}$$

which can be explained as a plane wave decomposition of the field, where $A_1^-(k)$ and $A_1^+(k)$ are the amplitudes of the upgoing and downgoing compressional (longitudinal) field for the layer 1 and $B_1^-(k)$ and $B_1^+(k)$ are the amplitudes of the upgoing and downgoing shear field in the layer 1, $g_{11}(r, z)$ and $g_{s1}(r, z)$ are the potential fields in the solid corresponding to the compressional velocity and the shear velocity respectively. By using the Sommerfeld radiation condition, the total solution in the layer 2, where the source is located, can be expressed in the form of:

$$g_2(r, z) = \int_0^\infty \{[A_2^-(k)e^{-\alpha_2(z-D)} + A_2^+(k)e^{\alpha_2(z-D)}] + G_{P2}(k, z, z_s)\} k J_0 dk, \quad (7)$$

$$\text{where } \alpha_2 = \begin{cases} -\sqrt{k^2 - k_2^2}, & k > k_2, \\ i\sqrt{k^2 - k_2^2}, & k \leq k_2. \end{cases}$$

and $A_2^-(k)$ and $A_2^+(k)$ are the amplitudes of the upgoing and downgoing fields in the layer 2,

$G_{P2}(k, z, z_s) = \frac{S_\omega e^{\alpha_2 |z - z_s|}}{4\pi\alpha_2}$, S_ω is the source strength and D the thickness of the layer 1. Similarly the integral representation of the solution in layer 3 can be introduced as:

$$g_3(r, z) = \int_0^\infty A_3^+(k) e^{\alpha_3 [z - (D+L)]} k J_0 dk,$$

$$\text{where } \alpha_3 = \begin{cases} -\sqrt{k^2 - k_3^2}, & k > k_3, \\ i\sqrt{k^2 - k_3^2}, & k \leq k_3. \end{cases} \quad (8)$$

and A_3^+ is the amplitude of the downgoing field in layer 3 and L thickness of layer 2. The boundary conditions for such a model are presented in [4], p. 17. These lead to a linear equation system which can be written as matrix form with U as coefficient matrix and V as constant matrix:

$$U \begin{bmatrix} A_1^- \\ A_1^+ \\ B_1^- \\ B_1^+ \\ A_2^- \\ A_2^+ \\ A_3^+ \end{bmatrix} = V. \quad (9)$$

If the determinant of the coefficient matrix, $\text{Det}[U(k)]$, vanishes, then the solutions of the Eq. (9) have poles for the corresponding horizontal wavenumber values. The poles (pole) correspond to additional modes (mode) in the wavenumber spectrum. This phenomena can be observed in next part, Fig. 1 (a).

3. Numerical Examples

In this section we will study four cases to get an idea about sound propagation in the ice-covered region of the Arctic Sea. The measurements are motivated from MIZEX 87 [3] for which an ice-layer is included. In these calculations, the roughness and other inhomogeneous properties of ice are neglected.

3.1. Case 1: Three Layered Model, (vacuum-ice-fluid-fluid halfspace)

The first case considers an ice-covered sea environment for which a half-space fluid is included to neglect the effect of bottom properties. This model can be related to mathematical equations presented at previous section. The environmental parameters in the Table I were used to compute Green's function and normal stress (NS[dB/1Pa]) and results are shown in Fig. 1 and 2 for receiver depths 18 og 122m.

Table I. 3 layered waveguide (vacuum-solid-fluid-fluid)

	c_l	c_s	ρ	z
0 (vacuum)	0	0	0	∞
1 (ice)	$c_1 = 3000$	1600	0.92	$D = 4$
2 (water)	$c_2 = 1441$	0	1.0273	$L = 90$
3 (water)	$c_3 = 1457$	0	1.0279	∞

Where c_l [m/s] is the compressional sound velocity, c_s [m/s] is the shear sound velocity, ρ [gr/cm³] is density and z [m] is the layer depth and D denote ice thickness.




Figure 1: Results for Table I in layer 2 (RD=18m) and layer 3 (RD=122m) (a) modulus of depth-separated Green's function and (b) transmission loss for $f=40$ Hz, $N=4096$ (N: number of sampling points), $SD=7m$, $RD=18m$ (dashed lines) and $RD=122m$ (solid lines).




Figure 2: Results for Table I (a) modulus of depth-separated Green's function and (b) transmission loss for $f=300$ Hz, $N=16384$, $SD=7m$, $RD=18m$ (dashed lines) and $RD=122m$ (solid lines).

In Fig. 1 (a) ($f = 40$ Hz) the sharp peak is a discrete pole associated with the shear sound velocity in the ice.

In Fig. 1 (a) the last peak ($k \simeq 0.37 \text{ m}^{-1}$) is an additional pole which appears in the evanescent spectrum. It is a dramatic shear effect. The pole corresponds to vanishing of the $\text{Det}[U(k)]$, where U is given by Eq.(9). It belongs to the class of *leaky waves*. The peak disappears when the receiver is moved out deeper in the water ($RD=122m$), because in Eq.(7) the first exponential term becomes very small as the receiver depth becomes large.

In Fig. 2 (a) ($f = 300$ Hz) the sharp peak associated with sound propagation in the ice (shear velocity) is a discrete pole. The peak at $k \simeq 0.6 \text{ m}^{-1}$ is a part of the continuous spectrum, indicating a leaky mode associated with the compressional sound velocity in the ice.

In Fig. 1 (b) for $z = 18m$ the transmitted field in the layer 1 from layer 2 can be both compressional and shear fields and these can be reflected from the ice-vacuum interface and can be transmitted again as compressional field in the layer 2. The interference pattern has been dominated by the evanescent part.

3.2. Case 2: Three Layered Model (Vacuum, Arctic sound velocity profile)

The second case concerns a simple Arctic waveguide without ice layer.

Table II. Arctic waveguide 1.

	c_{l1}	c_{l2}	ρ	z
0 (vacuum)	0	0	0	∞
1 (water)	1439	1441	1.0273	80
2 (water)	1441	1456	1.0278	60
3 (water)	1456	1461	1.0279	360

Where c_{l1} is the compressional sound velocity at the top of the layer and c_{l2} compressional sound velocity at the bottom of the layer which is treated by SAFARI as a fluid with $\frac{1}{c_l^2(z)}$ varying linearly with depth. ρ is density.

The modulus of the Green's function and normal stress (NS [dB/1Pa]) for the environmental data given in Table II have been calculated in the layer 1 and 2 by SAFARI and shown in figures 3 and 4 for the frequencies 40 and 300 Hz respectively.




Figure 3: Arctic waveguide 1 (Table II), (a) modulus of depth-separated Green's function. (b) transmission loss for $f=40$ Hz, $N=4096$, $SD=3m$, $RD=18m$ (dashed lines) in layer 1 and $RD=122m$ (solid lines) in layer 2.




Figure 4: Arctic waveguide 1 (Table II), (a) modulus of depth-separated Green's function. (b) transmission loss for $f=300$ Hz, $N=16384$, $SD=3m$, $RD=18m$ (dashed lines) in layer 1, and $RD=122m$ (solid lines) in layer 2.

As one can observe evanescent field is important only for low frequencies at near field (in the number of wavelength), Fig. 3 (a).

Waves with frequencies lower than cutoff frequency $f_{cutoff} = \frac{1}{4L\sqrt{\frac{1}{c_1^2} - \frac{1}{c_2^2}}}$ can not propagate along the layer without damp-

ing. The cutoff frequency in layer 1 is about $f_{cutoff} \simeq 85$ Hz and in layer 2 about $f_{cutoff} \simeq 16$ Hz. Sound propagating with frequencies smaller than the cutoff frequency can transmit to the next layer and if the source frequency is larger than the cutoff frequency, then the sound energy can be channeled in that layer. In Fig. 3 (b) the sound energy leaks from layer 1 into layer 2 and is channeled and concentrated in the layer 2. Therefore the transmission loss at the receiver depth $z=122m$ is less than that of in the layer 1 ($z=18m$). This observation is in agreement with the ambient noise recording measurements made in Barents Sea during SIZEX 92 [1].

3.3. Case 3: Four Layered Model: Vacuum, Arctic sound velocity profile and bottom

The results of a study for the environmental data given in Table III, presented in case 4 (in which D , ice thickness, is zero) are shown in Fig. 5 and are compared with the results presented in Fig. 9 in Sec. 3.4.




Figure 5: Results for Arctic waveguide 2 (Table II), for $D=0m$, (a) modulus of depth-separated Green's function in layer 3, $|G_3(k, z)|$, (b) transmission loss in layer 3 for $f=300$ Hz, $N=16384$, $SD=3m$ and $RD=122m$.

3.4. Case 4: Five Layered Model (Vacuum, ice, Arctic sound velocity profile and bottom)

Table III. Arctic waveguide 2.

	c_{l1}	c_{l2}	c_s	ρ	z (m)
0 (vacuum)	0	0	0	0	∞
1 (ice)	3000	3000	1600	0.92	D
2 (water)	1439	1441	0	1.0273	80
3 (water)	1441	1456	0	1.0278	60
4 (water)	1456	1461	0	1.0279	$360 - D$
5 (gravel*)	1800	1800	0	2.0	∞

* A compressional attenuation ($\gamma_l = 0.6$ dB/ Λ) is applied for the absorbing bottom in the simulations.

The modulus of Green's function and normal stress for the environmental data given in Table III, with an ice thickness

$D = 4\text{ m}$, have been calculated and shown in figures and 6 for frequencies 300 Hz respectively.




Figure 6: Results for Arctic waveguide 2 (Table III), for $D=4\text{m}$ (ice thickness), (a) modulus of depth-separated Green's function in layer 3, $|G_3(k, z)|$, (b) transmission loss in layer 3 for $f=300\text{ Hz}$, $N=16384$, $SD=7\text{m}$ and $RD=122\text{m}$.

Introduction of the ice has reduced the amplitude of the sharp peak. There are no significant changes in the corresponding normal stresses at the low frequencies comparing with the last model in which there was no ice at the top.

As frequency increases to $f = 300\text{ Hz}$, normal stresses shown in Fig. 6 (b), have been damped at short ranges, comparing with the case without ice layer Fig. 5 b). The reason is that the field can be transmitted in the ice for the short ranges. The transmitted field is mostly absorbed or trapped in the ice layer.

Comparing Fig. 1 with figures 5, 6 and 7 and Fig. 2 with figures 8 and 9, indicate that reflections from the sea bottom have significant effects on Green's function and normal stress.

4. Summary and Conclusions

The study is performed in a progression from simple environments to more complicated models that can simulate an ice-covered sea environment. Each model introduces greater complexity so that the importance of the various parameters can be isolated. The analytic integral expressions for the field for a simple 3 layered model are obtained by using boundary conditions. In these calculations only compressional monochromatic point sources have been used. In each experiment both the source and receivers are localized in the water below the ice interface.

The computations by SAFARI are performed for source frequencies at 40 and 300 Hz. The main observations and results from the numerical study are:

Sound will be transmitted into an ice layer as compressional and shear waves. These shear waves can be reflected from the ice-vacuum boundary and converted back to the compressional waves into water through the ice-water boundary. There is a competition between compressional and shear

waves at the ice-water boundary.

The subsequent shear waves are trapped in the ice and even a small shear attenuation eventually removes all the energy in the shear field. The trapped compressional waves in the ice layer are only important at short ranges.

A sensitivity study for a set of environmental models simulating the ice-covered region of the Arctic Sea is performed (not shown here) and the influence of bottom properties on transmission loss is studied and the results describe that the acoustic field propagating towards the bottom, will be reflected gradually and the propagation pattern is therefore less sensitive to bottom parameters specially at higher frequencies.

Acknowledgement

I would like to acknowledge Nansen Environmental and Remote Sensing Center (NERSC) for providing software equipments and CTD measurements.

References

- [1] I. Engelsen. SIZEX 92, AMBIENT NOISE MEASUREMENTS DATA REPORT. Technical report, FFI / report-93, 2006, 1993.
- [2] H.Schmidt. SAFARI, report of the Seismo-acoustic Fast field Algorithm for Range-Independent environment. Technical report, SACLANT Undersea Research Centre, 1988.
- [3] F.B. Jensen. Numerical models in underwater acoustics, edited by L. B. Felsen (Nijhoff Dordrecht), 1984.
- [4] C. Geiger S. Sandven, Z. Kovacs and O. M. Johannesson. MIZEX 87, CTD DATA REPORT FROM POLAR CIRCLE. Technical report, NERSC, Report No. 13, 1988.



Proceedings of the
Sixth European Conference on

Underwater Acoustics

ECUA 2002


24-27 June 2002

Editor

Andrzej Stepnowski

Co-editors

Roman Salamon and Andrzej Partyka



Gdańsk University of Technology

Contents

PREFACE	V
CONTENTS	VII
PLENARY SESSION	1
Parametric Arrays: Laboratory Applications in Underwater Acoustics, <i>Victor Humphrey</i>	3
Blind Ocean Acoustic Tomography: Experimental Results on the Intifante'00 Data Set, <i>Sergio Jesus</i>	9
SEAFLOOR CHARACTERISATION	19
Multi-Sensor Analysis of the Seabed in Shallow Water Areas: Overview of the MAPLE'2001 Experiment, <i>Eric Pouliquen, M. Trevorrow, Philippe Blondel, Gaetano Canepa, F. Cernich, Reginald Hollett</i>	21
Reflection of Acoustic Waves from a Two Layered Sea Floor with Oblique Interface, <i>Panagiotis Papadakis, Hara Mizeraki</i>	31
Backscattering from a Layered Seafloor with Parameters Continuously Varying with Depth, <i>Laurent Guillon, Xavier Lurton</i>	37
Frequency Dependence of Acoustic Waves in Marine Sediments, <i>Gary Robb, Angus Best, Justin Dix, Tim Leighton, A. Harris, J.S. Riggs, Jon Bull, Paul White</i>	43
Seabed Discrimination with Downward and Forward-Looking Sonars, <i>Paul van Walree, Dick Simons, Jeroen Janmaat</i>	49
Seabed Characterization Using Backscattering Discontinuities in Sonar Images, <i>Benoit Zerr, Reginald Hollett</i>	53
Using Wavelet Analysis to Classify and Segment Sonar Signals Scattered from Underwater Sea Beds, <i>Louis Atallah, Penny Probert Smith</i>	59
Seabed Characterisation Using Spectral Moments of the Echo Signal, <i>Jaroslav Tegowski, Zbigniew Lubniewski</i>	65
Comparison of Backscattered Sea Bottom Echo Modelling in the Acoustic Pressure Domain and in the Intensity Domain, <i>Zbigniew Lubniewski</i>	71
Determination of the Seafloor Properties by Acoustic Resonances in the Reflection Coefficient, <i>Vladimir N. Fokin, Margarete S. Fokina, Viktor V. Kurin, Leonid M. Kustov</i>	77
Seabed Biotope Characterisation in Deep Water – Initial Evaluation of Single and Multi-Beam Acoustics, <i>Rudy Kloser, Gordon Keith, Tim Ryan, A. Williams, John Penrose</i>	81
Validation of a Seafloor Segmentation Algorithm for Multibeam Data, <i>Gaetano Canepa, Eric Pouliquen, Nicholas G. Pace, A. Figoli, P. Franchi</i>	89
Conditioning of Acoustic Signals for Improved Sea Floor Characterisation and Imaging, <i>Jan Ove Bakke, Freddy Pohner</i>	95
Application of Splines and Wavelets along with TIN Decimation to 3D Imaging of Seafloor from Multibeam Sonar Data, <i>Jerzy Demkowicz, Marek Moszynski, Andrzej Stepnowski</i>	99

SCATTERING AND REVERBERATION	105
Sea Surface Scattering Strength: Theory Versus Experiments and Chapman-Harris Formula, <i>Marc Gensane</i>	107
Source Localisation Through Knowledge of the Scattered Acoustic Field Near an Underwater Vehicle, <i>Paul Lepper, Gerald D'Spain, William Hodgkiss</i>	113
Spherical-Shell Scattering Computations by Ordinary Exponentials, <i>Sven Ivansson</i>	119
A Shallow-Water Reverberation PE Model, <i>E. Geraldine Bouchage, Kevin LePage</i>	125
Extraction of Modal Back-Scattering Matrix from Reverberation Data in Shallow-Water Waveguide, <i>Er Chang Shang, Tian Fu Gao, Da Jung Tang</i>	131
Sound Scattering from a Cylindrical Shell Stiffened by a Transversal Plate, <i>Aleksander Klauson, Jaan Metsaveer, Lionel Haumesser, Dominique Decultot, Gerard Maze</i>	137
3-D HP-Adaptive Structural Acoustic Finite-Element Codes for Target Scattering, <i>David S. Burnett, Mario Zampolli</i>	141
Inversion for Bottom Parameters Using Low Frequency Reverberation Data, <i>Simonette H.E. Muller, M.A. Ainslie, Wilco Boek, Dick G. Simons</i>	147
Beam Tracing Techniques for High Frequency Reverberation Modelling, <i>Matthias Meyer, Gary L. Davies</i>	153
Experimental Study of Fluctuations in Coherent Backscattering, <i>Lucie Pautet, Eric Pouliquen, A. Crawford</i>	157
Low Frequency Underwater Waveguide Sound Reflection from Continental Slopes, <i>Igor Didenkulov</i>	163
High-Frequency Bistatic Scattering: Comparison of Model and Laboratory Experiments, <i>Philippe Blondel, Paul Mc Cloghrie, Nicholas G. Pace, Gary J. Heald, Richard Brothers</i>	167
Bistatic Reverberation in Range Dependent Waveguides, <i>Kevin D. LePage</i>	177
Field Measurement of Bistatic Scattering Strength of a Sandy Seabed at 118KHz, <i>Gaetano Canepa, Nicholas Pace, Eric Pouliquen</i>	183
Wave Theoretic Calculations of Bistatic Reverberation in Range Independent Waveguides, <i>Kevin D. LePage</i>	189
Bistatic Scattering from Water-Filled Cylinders Buried in the Seabed. Model-Data Comparison, <i>Alessandra Tesei, John Fawcett, Reginald Hollett</i>	195
Multistatic Reverberation and System Modelling Using SUPREMO, <i>Chris H. Harrison, Mark Prior, A. Baldacci</i>	201
Reverberation Modelling Using Acoustic Arrivals Derived from Mode Sums and Ray Tracing, <i>Mark Prior, Chris Harrison, Kevin LePage</i>	207
Sound Scattering by Object in Shallow Water Waveguide Near Boundary, <i>Valery Grigoryev, Boris Katsnelson, Aleksey Zamislov</i>	213
SIGNAL AND DATA PROCESSING	219
Set-Membership Localization and Tracking of Autonomous Underwater Vehicles, <i>Andrea Caiti, Andrea Garulli, Flavio Livide, Domenico Prattichizzo</i>	221
A Wavefront Sensing Method for Synthetic Aperture Sonar Autofocus, <i>Hayden Callow, Michael Hayes, Peter Gough</i>	231
Interaction Between Autofocusing and Synthetic Aperture Processing: a Study Based on Simulations and Experimental Rail Data, <i>Johannes Groen, J. C. Sabel</i>	235
Adaptive Thinning Algorithm, <i>Pascal A. M. de Theije</i>	241
A Wavelet Approach to Classification of Harbour Porpoise Sonar Signals, <i>Omar Farooq, Anthony Goodson, Sekharjit Datta</i>	243

Tracking the Underwater Noise Source Using a Vector Sound-Intensity Probe, <i>Ignacy Gloza</i>	247
Design Philosophy of Model-Based Sonar System, <i>Qihu Li</i>	251
Generation of Reverberation Time Series for Activated Low Frequency Sonars, <i>Luc Leviandier, Y. Doisy</i>	257
Buried Objects Classification Issue: Parametric Transmission and Linear Transmission with a Two Dimensional Synthetic Aperture Processing, <i>Stephane Guyonic</i>	263
Multispectral Decomposition of Synthetic Aperture Sonar Images for Speckle Reduction, <i>Jocelyn Chanussot, Alain Hetet, Gwendal Le Merrer, Elisa Tireau</i>	269
Indirect Evaluation of a Noisy Vessel Passive Detection Structure, <i>Kosta Ugrinovic</i>	275
Recent Results of Ocean Acoustic Time Reversal Experiments, <i>Hee-Chung Song, W.A. Kuperman, W.S. Hodgkiss, T. Akal, Seongil Kim, G. Edelmann</i>	279
Sequential Model-Based Detection in a Shallow Ocean Acoustic Environment, <i>James Candy</i>	285
Model-Based Spatial Diversity Processing with Sparse Receive Arrays. Application to Coherent Acoustic Communication in Shallow Water, <i>Jean-Pierre Hermand</i>	289
Optimising the Sounding Pulse of the Rotational Directional Transmission Sonar, <i>Roman Salamon, Jacek Marszal</i>	295
Underwater Acoustic Target Tracking with Fixed Passive Sonar System, <i>Slawomir Henclik</i>	301
Compression of Location Signal on Base of Chirp-Signals and Noise-Like Codes, <i>Włodzimierz Pogribny, Ihor Rozhankivsky, Tomasz Marciniak</i>	307

ACOUSTIC OF INHOMOGENEOUS MEDIA

Random Horizontal Refraction and Biases in Ray Travel Time and Mode Phase, <i>Oleg Godin</i>	313
Acoustic Qualities of Surface Ship Wakes, <i>David L. Bradley, R. Lee Culver, X. Di, Leif Bjorno</i>	315
Objective Grid/Radial Selection for Efficient Transmission Loss Calculations, <i>Erik Rike, Donald Delbalzo</i>	319
A Consistent Coupled-Mode Theory for Underwater Sound Propagation in a General, Stratified Acoustic Environment, <i>Gerassimos A. Athanassoulis, Konstandinos A. Belibassakis</i>	325
Inverse Modelling of Rubber Coated Steel Plates with a Genetic Algorithm, <i>Alex N. P. Cederholm, Ilkka Karasalo</i>	333
Acoustical Properties of Waterlogged Wood, <i>Stephanie Arnott, Angus Best, Justin Dix, David Gregory</i>	339
Influence of Gradients in Inhomogeneous Elastic Media on Resonance in the Plane Wave Reflection Coefficient, <i>Margarita Fokina, Vladimir Fokin</i>	345

SOUND PROPAGATION MODELLING

Scaled Models for Underwater Acoustics and Geotechnics Applications, <i>Jean-Pierre Sessarego</i>	357
Acoustic Signal Variability in Shallow Water, <i>Peter Lourcing Nielsen, Martin Siderius, Juergen Sellschopp</i>	359
Relationship between Sound Speed Structure Near Ocean Surface and Pulses Propagating in Deep Sound Channel in the Ocean, <i>Kohji Futa, Toshio Tsuchiya, Toshiaki Kikuchi</i>	369
	375

Efficient Numerical Implementation of Discrete Transparent Boundary Conditions for Parabolic Equations, <i>Matthias Ehrhardt</i>	381
Acoustic Fields in Ocean with Currents: Effective Sound Speed Approximation, Its Justification and Extension, <i>Oleg Godin</i>	387
Numerical Simulation of Broadband Signal Dispersion in a 3-D Shallow Water Waveguide, <i>Frederic Sturm</i>	391
Influence of Parameters of Shallow Water Waveguide on Space-Time Sound Fluctuations Caused by Internal Waves, <i>Boris Katsnelson, Serguey Pereselkov</i>	397
Models of Probability Density Function of Ambient Noise, <i>Wieslaw Kicinski</i>	403

ACOUSTICS IN FISHERIES

Fish Target Strength Estimation Versus Fish Backscattering Cross-section Estimation, <i>Marek Moszynski</i>	407
<i>In Situ</i> Inter-Standardization of Acoustics Data: an Integrated Data Base for Fish School Behaviour Studies, <i>Patrice Brehmer, A. Rouault, Francois Gerlotto</i>	409
Acoustic Semi-Tomography Applied to Estimate Fish Response to Environmental Variability, <i>Andrzej Orlowski</i>	417
Acoustical Estimation of Herring Stock in the Gulf of Riga: an Encouraging Case of Method Implementation in Semi-Enclosed Basin, <i>Olavi Kaljuste, Tiit Raid, Faust Shvetsov, Guntars Strods</i>	423
Acoustic Properties of Fish as Related to External Conditions, <i>Joanna Szczucka</i>	429
Active Fish Tracking Sonar (AFTS) for Assessing Fish Behavior, <i>John B. Hedgepeth, Gary E. Johnson, John R. Skalski, Janusz Burczynski</i>	435
The Effect of Fish Migration Pattern on the Acoustical Estimates of Fish Stocks in Inland Waters, <i>Malgorzata Godlewska</i>	441
Acoustical Recognition of a Copepod Layer, <i>Calanus euxinus</i> in Response to the Dissolved Oxygen in the Black Sea, <i>Erhan Mutlu</i>	447
Application of Hydroacoustic Survey Results in Studies of Eastern Baltic Sprat Distribution and Migration Pattern, <i>Faust Shvetsov, Valeri Feldman, Vladimir Severin, Alexander Zezera, Guntars Strods</i>	451

GEOACOUSTIC INVERSION

A Light Aircraft as a Source of Sound for Performing Geo-Acoustic Inversions of the Seabed, <i>Michael J. Buckingham, Eric M. Giddens, Jonathan B. Pompa, Fernando Simonet, Thomas R. Hahn</i>	463
Bottom Reflection Properties by Inversion of Ambient Noise, <i>Chris Harrison, A. Baldacci</i>	465
Estimation of Seabed Models by Inversion of Broadband Acoustic Data, <i>Ross Chapman, Julie Desert, Ashok Agarwal, Yann Stephan, Xavier Demoulin</i>	471
Inversion of Broadband Acoustic Data on a Single Hydrophone by Use of an Equivalent Geoacoustic Model, <i>Jean-Claude Le Gac, Mark Asch, Yann Stephan, Xavier Demoulin</i>	477
Frame-Fluid Boundary in the Acoustics of Water-Saturated Sand, <i>Nicholas Chotiros, C. Pollet, J.P. Sessarego, M. Brussieux</i>	483

OCEAN AND ACOUSTIC TOMOGRAPHY

Passive Ocean Acoustic Tomography: Theory and Experiment, <i>Enrico de Marinis, Otello Gasparini, Paola Picco, Sergio Jesus, Alessandro Crise, Stefano Salon</i>	489
---	-----

Numerical Analysis of Underwater Acoustic Lens Using Alternating-Direction Implicit Method, <i>T. Tsuchiya, T. Anada, N. Endoh, T. Aoki, I. Kaihou</i>	503
Underwater Transitory Signal Characterization Using Time-Frequency Warping Operators, <i>Cornel Ioana, Andre Quinquis</i>	509
Range-Dependent Regularization of Travel-Time Tomography Based on Theoretical Modes, <i>Orlando Rodriguez, Sergio Jesus</i>	515
ACOUSTIC IMAGING	521
Detection and Imaging of Buried Objects with an Experimental Sediment Sonar, <i>Dieter Brecht, Harald Peine, Broder Fedders</i>	523
Imaging and Motion Estimation for Synthetic Aperture Sonar Based on Fast Factorised Back-Projection, <i>Simon Banks, Hugh H. Griffiths</i>	529
On Ray-Tracing and Interpolation in Seafloor Front-Scan Imaging, <i>Andrea Trucco, Maria Palmese, Stefania Repetto</i>	535
UNDERWATER COMMUNICATION	541
Acoustic Communication Experiments in the Westerschelde Shipping Lane Using Direct-Sequence Spread Spectrum Signals, <i>Paul Van Walree, Martin Van Gijzen, M. Pattipeilohy</i>	543
Blind Decision-Feedback Equalization of Shallow Water Acoustic Channels, <i>Rolf Weber, Florian Schulz, Andreas Waldhorst, Johann Friedrich Boehme</i>	549
Initializing Blind Adaptive Equalizers Using A-Priori Environmental Knowledge, <i>Andreas Waldhorst, Rolf Weber, Florian Schulz, Johann Friedrich Boehme</i>	555
NON-LINEAR ACOUSTICS	561
Cavitation Nuclei and Acoustic Cavitation Thresholds of Ocean Water, <i>Victor Akulichev</i>	563
Some Aspects of Finite Amplitude Wave Generation, <i>Grazyna Grelowska</i>	565
40 Years of Nonlinear Underwater Acoustics, <i>Leif Bjorno</i>	573
Influence of Absorption in Low Salinity Sea Water on Nonlinear Wave Propagation, <i>Grazyna Grelowska, Eugeniusz Kozaczka</i>	581
Investigation of The Difference Frequency Scattering from a Cavitation Jet, <i>Igor Didenkulov, Leonid Kustov, Alexander Martynov, Nikolay Pronchatov-Rubtsov, Pave Vyugin</i>	585
Theoretical Analysis of the Finite Amplitude Waves Interaction Problem, <i>Anna Baranowska</i>	589
ACOUSTIC IN ENVIRONMENTAL MONITORING	593
Recent Results of the ACOUS (Arctic Climat Observations Using Underwater Sound) Program, <i>Alexander Gavrilov, P.N. Mikhalevsky</i>	595
The Role of Gas Bubbles and Surface-Active Substances on the Noise Generation by Rain Drop, <i>Zygmunt Klusek</i>	605
Seismo-Acoustic Monitoring in the Pacific Ocean, <i>Pierre Piserchia, Olivier Hyvernaud, Dominique Reymond, Jacques Talandier</i>	611
A Discussion on Acoustic Propagation in the Ice-Covered Region of the Arctic Sea, <i>Amir M. Hashemi</i>	613
Numerical Study of the Compressional Reflection Coefficient in the Ice-Covered Arctic Sea, <i>Amir M. Hashemi</i>	619
An Acoustic Shadowgraph Trial to Detect Convection in the Arctic, <i>Barry J. Uscinski, Arthur Kaletsky, Clay S. Stanek, Daniel Rouseff</i>	623

TRANSDUCERS AND CALIBRATION	629
A New Type of Low Frequency High Power Terfenol-D Underwater Acoustical Transducer, <i>Xibing Mo, Houqing Zhu</i>	631
Determination of the Variation in Free-Field Hydrophone Response with Temperature and Depth, <i>Graham Beamiss, Stephen P. Robinson, Gary Hayman, T. J. Esward</i>	635
Wide-Band Circular Ultrasonic Transducers, <i>Grazyna Lypacewicz, Andrzej Nowicki, Roman Dynowski, R. Tymkiewicz</i>	641
Wideband Signatures of Acoustic Airgun Arrays, <i>Rodney Coates, John Goold</i>	645
The Effects of Mechanical Coupling on the Beam Pattern of a Multi-Element Array, <i>Andrzej Elminowicz, Waldemar Lis, Zbigniew Wojan</i>	649
 INSTRUMENTATION AND MEASUREMENTS	653
Scapa-Map Project. The Maritime Archaeology in Scapa Flow. A Detailed Measurement of the Seabed and Wrecks with a High Resolution Multi-beam Sonar, <i>Thomas Meurling</i>	655
Testing Acoustic Rain Gauges in Inland and Coastal Waters, <i>Graham D. Quartly, Katy M. Shannon, Trevor H. Guymer, Keith G. Birch, Jon M. Campbell</i>	659
Acoustic Characterisation of Panel Materials in Simulated Ocean Conditions Using the NPL Acoustic Pressure Vessel, <i>Trevor Esward, Victor Humphrey, Laurence Evans, Graham Beamiss, Gary Hayman</i>	665
Digital Scanning Sonar for Fish Feeding Monitoring in Aquaculture, <i>Tim Acker, Janusz Burczynski, John Hedgepeth, Assad Ebrahim</i>	671
 POSTER SESSION	677
Backscattering from Seagrasses: an Experience with a High Frequency Sector Scanning Sonar, <i>Roberto Bozzano, Riccardo Mantovani, Antonio Siccardi</i>	679
Wave Transmission Through a Submerged Layer with Symmetrical Density, Sound Velocity And Attenuation Gradients, <i>Rafael Carbo, Adriana C. Molero</i>	685
Acoustic Cues and Trans-Ocean Migration of Marine Megafauna, <i>Rodney Coates</i>	689
Method of Visualisation of Ultrasonic Agglomeration Gas Bubbles in Liquid, <i>Henryka Czyz, Tadeusz Gudra, Krzysztof Opielinski</i>	695
Decision Tree Fuzzy Logic Classifier of Seabed, <i>Tran Van Dung, Andrzej Stepnowski, M. Kazmierczak</i>	699
The Distribution of Acoustic Field Intensity in a Medium with Sound Velocity Gradient Using a Modified Sound Ray Density Method, <i>Jacek Gawrysiak, Roman Salamon</i>	705
Fusion Algorithms for Detection in Presence of Reverberation, <i>Guillaume Ginolhac, Jocelyn Chanussot, Cedric Pichat</i>	711
Digital Acoustic System for Ecosystem Monitoring and Mapping: Assessment of Fish, Plankton, Submerged Aquatic Vegetation and Bottom Substrata Classification, <i>Joel Hoffman, Janusz Burczynski, Bruce Sabol, M. Heilman</i>	717
Prediction of Vibroacoustical Parameters on a Cruise Ship, <i>Marek Iwaniec, Jerzy Wiciak</i>	723
Wdzydze Lake Hydroacoustic Laboratory, <i>Lech Kilian, Jacek Marszał, Jacek Gawrysiak, Krzysztof Zachariasz</i>	729
Nonlinear Multielement Radiating Antenna, <i>Eugeniusz Kozaczka, Grazyna Grelowska</i>	733
Underwater Noise Characteristics of a Small Ships, <i>Stefan Malinowski, Ignacy Gloza</i>	737

Development of Tursiops Truncatus Sonar System During First Months of Life in Captivity: Acoustic Interactions with Other Community Members, <i>Sarine Manoukian, Massimo Azzali, Cristina Farchi, Raffaella Tizzi</i>	741
Time-varied-gain with Increased Accuracy for Digital Echounders, <i>Marek Moszynski, Andrzej Stepnowski</i>	747
Technique of Information Processing in Unorthogonal Function, <i>Valery A. Pakhotin, Svetlana V. Ivanova, Vladimir A. Bessonov, Irina V. Marchenko</i>	751
Observation of Acoustical Signal Fluctuations by Time-Frequency Analysis Methods, <i>Maria Palmese, Alessio Bozzo, Sergio Jesus, Jose Onofre, Paola Picco, Andrea Trucco</i>	755
Correlation Analysis Applied to the Size Determination of Immersed Targets, <i>Patrick Schweitzer, Etienne Tisserand, Jerome Mathieu</i>	761
Ultrasonic Sizing of Immersed Targets: Separation of Superimposed Echoes, <i>Patrick Schweitzer, Etienne Tisserand, Jerome Mathieu</i>	765
Method of Underwater Object Speed Estimation by One Detector, <i>Przemyslaw Soszynski, Ignacy Gloza</i>	769
Fluctuations of Acoustic Travel Time at the Havaii-Kamchatka Path, <i>A. Stromkov, I.Didenkulov, Ya.Karlik, A.Kazarova, L.Lyubavin, E.Pelinovsky</i>	773
The Reproduction of Spatiality in Listening Rooms, <i>Lamberto Tronchin, Alessandro Giusto, V. Tarabusi, Daniele Brini</i>	777
Computer Simulator of Images in Sonar Systems, <i>Mariusz Waz, Dariusz Szulc</i>	781
AUTHOR INDEX	787