


# Push-forward and pull-back

## Pull-back of a function



Let us consider two manifolds  $M$  and  $N$ , respectively  $m$ -dimensional and  $n$ -dimensional.

Let us then consider a mapping between the two manifolds  $\phi : M \rightarrow N$  and a function  $f : N \rightarrow \mathbb{R}$ .

Let us now consider a vector  $V \in T_p M$  where  $p \in M$ . The vector  $V$  is "a derivation" operator acting on smooth functions on  $M$ . Given a set of coordinate functions  $x = (x^1, \dots, x^m)$  on  $M$ , a coordinate representation of  $V$  is  $V = V^\mu \frac{\partial}{\partial x^\mu}$ .

By definition the differential operator  $V$  can act only on functions on  $M$ . Thus it is possible to make it act on  $f$  by *pulling  $f$  back* to  $M$  through  $\phi$ .

Indeed the application  $f \circ \phi$  maps a point of  $M$  in  $\mathbb{R}$ , hence such a function can be differentiated by  $V$ . The operation of *pulling  $f$  back* to  $M$  is called *pull-back* of  $f$ . More precisely, the pull-back of any smooth function on  $N$  is a function

$$\phi^* : \mathcal{S}(N) \rightarrow \mathcal{S}(M)$$

where  $\mathcal{S}(A)$  indicates the set of smooth functions acting on  $A$ , and its action is defined as

$$V(\phi^* f)_p = V(f \circ \phi)$$

where the subscript  $p$  means that it acts on  $M$  (from now on this will be omitted considering it implicit). In coordinates  $f(\phi(p)) = f(\phi(x^{-1}(\mathbf{x}))) \Rightarrow f = f(\mathbf{x})$  where  $\mathbf{x} \in \mathbb{R}^m$ . Hence

$$V(\phi^* f) = V^\mu \frac{\partial f(\mathbf{x})}{\partial x^\mu}$$

## Push-forward of a vector

The same result above can also be seen as the result of the action of another operator, namely the *push-forward*, which we denote as  $\phi_*$ , whose role is to *push  $V$  forward* to  $N$ , so that  $\phi_* V \in T_{\phi(p)} N$  and its action on a function  $f : N \rightarrow \mathbb{R}$  is defined by requesting that

$$(\phi_* V)_{\phi(p)} f = V(\phi^* f)_p = V(f \circ \phi) \quad (1)$$

where, again, the subscript  $\phi(p)$  indicates that  $\phi_* V$  is a vector in  $T_{\phi(p)} N$  (and again from later on the subscript will be dropped and considered implicit).

What we did previously was extending  $f$  to  $M$  and working in the manifold  $M$ . Here, instead, we are extending  $V$  to  $N$  and working on  $N$ .

Now let us introduce the charts  $x = (x^1, \dots, x^m)$  in  $M$  and  $y = (y^1, \dots, y^n)$  in  $N$ . A coordinate representation of  $V$  is  $V = V^\mu \frac{\partial}{\partial x^\mu}$ .

Since  $\phi_* V$  is a vector in  $T_{\phi(p)} N$  a coordinate representation is  $\phi_* V = (\phi_* V)^\mu \frac{\partial}{\partial y^\mu}$ .

To get each component  $(\phi_* V)^\alpha$  we must make  $\phi_* V$  act on each coordinate  $y^\alpha$  so that

$$(\phi_* V) y^\alpha = \left( (\phi_* V)^\mu \frac{\partial}{\partial y^\mu} \right) y^\alpha = (\phi_* V)^\mu = (\phi_* V)^\mu \frac{\partial y^\alpha}{\partial y^\mu} = (\phi_* V)^\mu \delta_\mu^\alpha = (\phi_* V)^\alpha$$

In other words we extracted the components  $\alpha$  of the vector  $\phi_* V \in T_{\phi(p)}N$ .  
By using the definition given by 1 and the coordinate representation of  $V$

$$\begin{aligned} (\phi_* V)^\alpha &= (\phi_* V) y^\alpha = V(y^\alpha \circ \phi) = V^\mu \frac{\partial}{\partial x^\mu} (y^\alpha \circ \phi) = \\ &= V^\mu \frac{\partial}{\partial x^\mu} (y^\alpha(\phi(p))) = V^\mu \frac{\partial}{\partial x^\mu} (y^\alpha(\phi(x^{-1}(\mathbf{x})))) \end{aligned} \quad (2)$$

where  $\mathbf{x} = x(p) = (x^1, \dots, x^m) \in \mathbf{R}^m$ .

The last member of equation 1 tells us that we can consider  $y^\alpha$  as a function of  $\mathbf{x}$ .

Hence  $y^\alpha = y^\alpha(\mathbf{x})$ . Finally, equation 2 reduces to

$$(\phi_* V)^\alpha = V^\mu \frac{\partial y^\alpha}{\partial x^\mu}$$

Sounds familiar? The function  $\phi_*$  acts on  $V$  as a jacobian matrix <sup>1</sup>.

Since  $\phi_* V \in T_{\phi(p)}N$  we can make it act on a differential form  $\omega \in T_{\phi(p)}^*N$ .

$$\begin{aligned} \phi_* V &= (\phi_* V)^\mu \frac{\partial}{\partial y^\mu} \\ \omega &= \omega^\mu dy^\mu \\ (\phi_* V)(\omega) &= \omega(\phi_* V) = \langle \phi_* V, \omega \rangle = V^\mu \omega^\nu \left\langle \frac{\partial}{\partial y^\mu}, dy^\nu \right\rangle = V^\mu \omega^\mu \end{aligned}$$

### Pull-back of a differential form

The result we have just obtained can be seen in another way. Indeed, instead of *pushing*  $V$  *forward* to  $N$  (working on  $N$ ), we can *pull*  $\omega$  *back* to  $V$  (working on  $N$ ). More precisely we can introduce a new differential form  $\phi^* \omega \in T_p^*M$  by requesting that

$$\langle \phi_* V, \omega \rangle = \langle V, \phi^* \omega \rangle$$

The map  $\phi^* : T_{\phi(p)}^*N \rightarrow T_p^*M$  is called *pull-back* of  $\omega$ .

This nicely show that the pull-back  $\phi^*$  and the push forward  $\phi_*$  are dual maps one to the other.

---

<sup>1</sup>Note that since we are working on a manifold, this holds only locally! And this is why the push-forward is the generalization of the jacobian matrix. The two things coincide when we work in  $\mathbb{R}^n$ .