


Push-forward and pull-back

Pull-back of a function



Let us consider two manifolds M and N , respectively m -dimensional and n -dimensional.

Let us then consider a mapping between the two manifolds $\phi : M \rightarrow N$ and a function $f : N \rightarrow \mathbb{R}$.

Let us now consider a vector $V \in T_p M$ where $p \in M$. The vector V is "a derivation" operator acting on smooth functions on M . Given a set of coordinate functions $x = (x^1, \dots, x^m)$ on M , a coordinate representation of V is $V = V^\mu \frac{\partial}{\partial x^\mu}$.

By definition the differential operator V can act only on functions on M . Thus it is possible to make it act on f by *pulling f back* to M through ϕ .

Indeed the application $f \circ \phi$ maps a point of M in \mathbb{R} , hence such a function can be differentiated by V . The operation of *pulling f back* to M is called *pull-back* of f . More precisely, the pull-back of any smooth function on N is a function

$$\phi^* : \mathcal{S}(N) \rightarrow \mathcal{S}(M)$$

where $\mathcal{S}(A)$ indicates the set of smooth functions acting on A , and its action is defined as

$$V(\phi^* f)_p = V(f \circ \phi)$$

where the subscript p means that it acts on M (from now on this will be omitted considering it implicit). In coordinates $f(\phi(p)) = f(\phi(x^{-1}(\mathbf{x}))) \Rightarrow f = f(\mathbf{x})$ where $\mathbf{x} \in \mathbb{R}^m$. Hence

$$V(\phi_* f) = V^\mu \frac{\partial f(\mathbf{x})}{\partial x^\mu}$$

Push-forward of a vector

The same result above can also be seen as the result of the action of another operator, namely the *push-forward*, which we denote as ϕ_* , whose role is to *push V forward* to N , so that $\phi_* V \in T_{\phi(p)} N$ and its action on a function $f : N \rightarrow \mathbb{R}$ is defined by requesting that

$$(\phi_* V)_{\phi(p)} f = V(\phi^* f)_p = V(f \circ \phi) \quad (1)$$

where, again, the subscript $\phi(p)$ indicates that $\phi_* V$ is a vector in $T_{\phi(p)} N$ (and again from later on the subscript will be dropped and considered implicit).

What we did previously was extending f to M and working in the manifold M . Here, instead, we are extending V to N and working on N .

Now let us introduce the charts $x = (x^1, \dots, x^m)$ in M and $y = (y^1, \dots, y^n)$ in N . A coordinate representation of V is $V = V^\mu \frac{\partial}{\partial x^\mu}$.

Since $\phi_* V$ is a vector in $T_{\phi(p)} N$ a coordinate representation is $\phi_* V = (\phi_* V)^\mu \frac{\partial}{\partial y^\mu}$.

To get each component $(\phi_* V)^\alpha$ we must make $\phi_* V$ act on each coordinate y^α so that

$$(\phi_* V) y^\alpha = \left((\phi_* V)^\mu \frac{\partial}{\partial y^\mu} \right) y^\alpha = (\phi_* V)^\mu = (\phi_* V)^\mu \frac{\partial y^\alpha}{\partial y^\mu} = (\phi_* V)^\mu \delta_\mu^\alpha = (\phi_* V)^\alpha$$

In other words we extracted the components α of the vector $\phi_* V \in T_{\phi(p)}N$.
By using the definition given by 1 and the coordinate representation of V

$$\begin{aligned} (\phi_* V)^\alpha &= (\phi_* V) y^\alpha = V(y^\alpha \circ \phi) = V^\mu \frac{\partial}{\partial x^\mu} (y^\alpha \circ \phi) = \\ &= V^\mu \frac{\partial}{\partial x^\mu} (y^\alpha(\phi(p))) = V^\mu \frac{\partial}{\partial x^\mu} (y^\alpha(\phi(x^{-1}(\mathbf{x})))) \end{aligned} \quad (2)$$

where $\mathbf{x} = x(p) = (x^1, \dots, x^m) \in \mathbf{R}^m$.

The last member of equation 1 tells us that we can consider y^α as a function of \mathbf{x} .
Hence $y^\alpha = y^\alpha(\mathbf{x})$. Finally, equation 2 reduces to

$$(\phi_* V)^\alpha = V^\mu \frac{\partial y^\alpha}{\partial x^\mu}$$

Sounds familiar? The function ϕ_* acts on V as a jacobian matrix ¹.
Since $\phi_* V \in T_{\phi(p)}N$ we can make it act on a differential form $\omega \in T_{\phi(p)}^*N$.

$$\begin{aligned} \phi_* V &= (\phi_* V)^\mu \frac{\partial}{\partial y^\mu} \\ \omega &= \omega^\mu dy^\mu \\ (\phi_* V)(\omega) &= \omega(\phi_* V) = \langle \phi_* V, \omega \rangle = V^\mu \omega^\nu \left\langle \frac{\partial}{\partial y^\mu}, dy^\nu \right\rangle = V^\mu \omega^\mu \end{aligned}$$

Pull-back of a differential form

The result we have just obtained can be seen in another way. Indeed, instead of *pushing* V *forward* to N (working on N), we can *pull* ω *back* to V (working on N). More precisely we can introduce a new differential form $\phi^* \omega \in T_p^*M$ by requesting that

$$\langle \phi_* V, \omega \rangle = \langle V, \phi^* \omega \rangle$$

The map $\phi^* : T_{\phi(p)}^*N \rightarrow T_p^*M$ is called *pull-back* of ω .

This nicely show that the pull-back ϕ^* and the push-forward ϕ_* are dual functions one to the other.

¹Note that since we are working on a manifold, this holds only locally! And this is why the push-forward is the generalization of the jacobian matrix. The two things coincide when we work in \mathbb{R}^n .