

Deep Learning for Medical Imaging



Olivier Colliot, PhD
Research Director at CNRS
ARAMIS Lab – www.aramislab.fr
PRAIRIE – Paris Artificial Intelligence Research Institute



Maria Vakalopoulou, PhD
Assistant Professor at
CentraleSupélec
Center for Visual Computing

<https://piazza.com/centralesupelec/spring2024/3md3070>
Access code: 3md3070

Please sign up to Piazza now

<https://piazza.com/centralesupelec/spring2024/3md3070>

Access code: 3md3070

Evaluation


Part 1 – Introduction

1.1 Introduction

1.1.1 What is medical imaging?

What is medical imaging?

Techniques that allow to **see (*study*) inside the human body** in vivo and **non-invasively**



First X-ray radiography (1895) - Wilhelm Röntgen (Nobel Prize in Physics, 1901)

Medical imaging today

Multiple machines

X-ray radiography



MRI



PET / CT



Ultrasound (echography)



Image sources: <http://radiologie-abbeville.fr/files/Echo%20graphie%20g%C3%A9n%C3%A9rale.jpg>, CENIR (ICM),
https://fr.wikipedia.org/wiki/Tomographie_par_%C3%A9mission_de_positons, <https://en.wikipedia.org/wiki/Radiography>


Medical imaging today

Multiple types of images

X-ray radiography



MRI



Ultrasound



PET





Image sources: <https://fr.wikipedia.org/wiki/Radiographie>,
https://en.wikipedia.org/wiki/Obstetric_ultrasonography, Burgos et al, 2017 <https://hal.inria.fr/hal-01567343>


Medical imaging today

**Multiple types of images
with the same machine (MRI)**


Anatomical
MRI




Angiography



Functional
MRI



Diffusion
MRI





What is medical imaging?

Not only **see inside the human body**

But also **study multiple phenomena** which are inaccessible to the naked eye

- Metabolism
- Oxygenation of the blood
- Molecular structure



Part 1 – Introduction

1.1 Introduction

1.1.2 What is medical imaging used for?

What is medical imaging used for?


Medical care

- Detect lesions / anomalies
 - Find the cause of a set of symptoms

65-year old female

Presenting with
left-side weakness
and headache

MRI



Glioblastoma (brain tumour)

Source: <https://radiopaedia.org/cases/glioblastoma-nos-13?lang=us>

What is medical imaging used for?


Medical care

- Screening

56-year old female

Routine screening
Asymptomatic

Mammography



Potential breast tumour
(referred for biopsy)


Source: <https://radiopaedia.org/cases/small-breast-cancer?lang=us>

What is medical imaging used for?


- Follow up a pathology

Patient with
neurodegenerative
disease

(followed over 18
months)




M0



Medical care



M18



What is medical imaging used for?

- Quantify (extract biomarkers)
 - For instance, segmentation for volumetric analysis
 - Reduced volume of specific brain structures: biomarker of neurodegenerative diseases
 - Enlarged volume of the left ventricle of the heart

Medical care





Image source: Zheng et al, The Volume of Hippocampal Subfields in Relation to Decline of Memory Recall Across the Adult Lifespan, 2018

What is medical imaging used for?

Research

- Clinical research
 - Understand the alterations produced by a given disease

Brain alterations associated with Alzheimer's disease



Brain areas with significantly reduced metabolism in patients with Alzheimer's disease compared to healthy controls, as studied with PET imaging

What is medical imaging used for?



- Clinical research
 - Evaluate the effect of a treatment

Studying the effect of a treatment (donepezil) on the volume of the hippocampus in Alzheimer's disease

- Double-blind, randomized, placebo-controlled

Hippocampal atrophy was reduced by 45% in one year with donepezil compared to placebo

Research




What is medical imaging used for?

- Basic research
 - Cognitive neuroscience

Research

Studying which parts of the brain are used for reading




Source: Cohen et al, The visual word form area: Spatial and temporal characterization of an initial stage of reading in normal subjects and posterior split-brain patients, Brain, 2000

What is medical imaging used for?


- Basic research
 - Genetics and neuroscience

Research

Heritability of the size of brain structures



Source: Toro et al, Genomic architecture of human neuroanatomical diversity, Molecular Psychiatry, 2015



ICV: intracranial volume

BV: brain volume

VIQ: verbal IQ

PIQ: performance IQ

Part 1 – Introduction

1.1 Introduction

1.1.3 Medical image computing

Medical image computing


Scientific field dedicated to the computational analysis of medical images

At the cross-road of

- Computer vision / image processing
- Machine learning
- Statistics
- Geometry

Medical image computing – what for?

- Enhance medical images
 - Denoising




Denoising of
brain MRI

Provide higher quality images to the radiologist

Medical image computing – what for?

- Assist medical evaluation
 - Detection
 - Diagnostic classification



Automatic detection of breast cancer in mammography images


Reduce reading time for the radiologist

In the future: screen automatically and ask the radiologist only for difficult cases?

Source: McKinney et al International evaluation of an AI system for breast cancer screening, Nature, 2020

Medical image computing – what for?

- Quantify
 - Segmentation



Automatic segmentation of the hippocampus in Alzheimer's disease

Enrich visual evaluation with quantitative measures

	Alzheimer vs controls
Volume reduction	-32%
Sensitivity	84%
Specificity	84%

Source: Colliot et al, 2008

Medical image computing – what for?

- Predict
 - Predict the future state of the patient


Predict the future occurrence of
Alzheimer's disease (within 3 years)

Classifier - Features	Bal. acc.	AUC	Acc.	Sens.	Spec.
SVM - T1w MRI	0.670	0.736	0.698	0.586	0.754
SVM (trained on CN $A\beta$ - vs AD $A\beta$ +) - T1w MRI	0.679	0.764	0.708	0.547	0.811
SVM - FDG PET	0.708	0.777	0.732	0.633	0.782
SVM (trained on CN $A\beta$ - vs AD $A\beta$ +) - FDG PET	0.761	0.818	0.788	0.666	0.856
RF - Clinical _{base} + Score T ₁	0.717	0.792	0.732	0.671	0.763
RF - Clinical _{base} + Score FDG	0.760	0.831	0.791	0.669	0.852
RF - Clinical _{base} + Scores T ₁ ,FDG	0.769	0.855	0.796	0.685	0.852
RF - Clinical _{base} + RAVLT + Scores T ₁ ,FDG	0.791	0.881	0.809	0.735	0.846
RF - Clinical _{base} + ADAS + Scores T ₁ ,FDG	0.790	0.873	0.810	0.729	0.851
RF - Clinical _{base} + RAVLT + ADAS + Scores T ₁ ,FDG	0.792	0.888	0.811	0.736	0.849

Clinical_{base}: gender, education level, MMSE score, sum of boxes of CDR test

Medical image computing – what for?


- Predict
 - Predict the future observations



Predict the future MRI data in patients with Alzheimer's disease

Medical image computing – what for?

- Synthesis of medical imaging data



Source: Wei et al, Medical Image Analysis, 2019

Part 1 – Introduction

1.1 Introduction

1.1.4 Specificities of medical image computing


Specificities of medical image computing

Why it is not only computer vision
applied to medical images


Specificities of medical image computing

- 3D
 - Most medical images are 3D


Axial slices



Coronal slices




Sagittal slices




Specificities of medical image computing

- 3D
 - Most medical images are 3D




Voxel=volume element



Specificities of medical image computing

- 3D
 - Most medical images are 3D





I: Image

$I(x,y,z)$ denotes intensity value at pixel location x,y,z

Note also that whatever you see on the left is right part of the body!


Specificities of medical image computing

- 3D
 - Voxels have a physical size
 - Most often, this size needs to be taken into account in the processing
 - Quite often, the voxels are anistropic




Specificities of medical image computing

- Temporal phenomena
 - Across multiple time scales



Second

Functional brain images,
cardiac images...



M0

Year


Longitudinal studies



M18

Specificities of medical image computing


- Multiple phenomena and image types



Anatomical vs functional

Specificities of medical image computing


- Multiple phenomena and image types



Time-series

Specificities of medical image computing

- Multiple phenomena and image types



Tensor images

Specificities of medical image computing

- Validation
 - Critical
 - Can be difficult to perform
- Clinical relevance
 - Need close collaboration with physicians to identify relevant problems
- Multidisciplinary collaboration
 - Find a common language

Medical image computing with deep learning

- **Has achieved impressive results, as in other fields (CV, NLP...)**

- Even though there are some inherent difficulties, in particular the small sample size

- **Is not a universal solution**

- Other methods may remain competitive for some applications

Part 1 – Introduction

1.1 Introduction

1.1.5 About this course

Course content

■ Content

- theoretical and practical aspects of deep learning for medical imaging
- covers the main tasks involved in medical image analysis (classification, segmentation, registration, generative models...)
- deep learning techniques are presented, **alongside some more traditional image processing and machine learning approaches**
- Emphasis on validation which is critical for medical applications

■ Learning objectives

- have knowledge of state-of-the-art deep learning techniques for medical imaging
- have a deeper understanding of deep learning methods, applicable not only to medical images but also other types of data
- know how to build and validate deep learning models for medical images

Course content

- Note: today's course will be very different from the others
 - Presentation of medical image acquisition techniques
 - Closer to physics than machine learning

Part 1 – Introduction

1.2 Main medical imaging modalities



Two main types of medical images

Structural images

Also called morphological or anatomical

Visualize the structures


- Differentiate between tissues (e.g. skin, bone, muscle, fat...)
- Between organs (heart, brain...)
- Between anatomical structures within an organ (e.g. in the heart: ventricles, atria...)




Functional images

Provide information about a functional process

- Metabolism (e.g. glucose consumption)
- Blood flow
- ...






Anatomical image



Functional image (showing malignant lung tumour) superimposed on anatomical image

Two main types of medical images

Structural images



Brain MRI

Two main types of medical images

Structural images




Whole Body CT

<https://www.siemens-healthineers.com/en-sg/computed-tomography/news/mso-imaging-a-tale-of-two-cities.html>

Two main types of medical images

Functional images




Whole Body PET

<https://www.nature.com/articles/d41586-019-01833-z>

Two main types of medical images

Functional images



Functional MRI: right hand activation

Two main types of medical images

Structural images



Functional images



<https://media-cdn.tripadvisor.com/media/photos/o6/bd/47/14/shibuya-crossing-almost.jpg>


<https://www.videezy.com/transportation/39282-ho-chi-minh-city-traffic-at-intersection-vietnam>

Part 1 – Introduction

1.2.1 Structural imaging


1.2.1.1 X-ray radiography

X-Ray radiography




First X-ray radiography (1895) - Wilhelm Röntgen (Nobel Prize in Physics, 1901)

X-Ray radiography




X-Ray radiography




X-Ray radiography

Projectional radiography



X-Ray radiography



Source: Lynn N. McKinnis: *Fundamentals of Musculoskeletal Imaging*, 4th Edition:
www.FADavisPTCollection.com
Copyright © F. A. Davis Company. All rights reserved.

X-rays are absorbed by heavy atoms and not by lighter atoms

Bones: high absorbtion

Soft tissues: lower absorbtion

Air: no absorbtion

X-Ray radiography




Image source:
https://en.wikipedia.org/wiki/Dental_radiography
https://en.wikipedia.org/wiki/Chest_radiograph

X-rays are
absorbed by
heavy atoms and
not by lighter
atoms

Bones: high
absorbtion

Soft tissues: lower
absorbtion

Air: no absorbtion

Part 1 – Introduction



1.2.1 Structural imaging

1.2.1.2 CT

CT

CT: Computed Tomography


Principle : 3D reconstruction from multiple radiographic projections



Good visualization of bone
High resolution

Poor visualization of soft tissues (in particular in the brain)


CT



Multiple CT views, each made up of parallel rays incident from a different angle. Theta defines the angle of incidence.


CT

The sinogram draws the signal for each distance r and each angle θ



Sinogram formation from individual projection angles


CT



Sinogram of an example object


Source: [http://199.116.233.101/index.php/Image_Reconstruction_\(CT\)](http://199.116.233.101/index.php/Image_Reconstruction_(CT))

CT



Backprojection. Each view (intensity vs. radius) is stretched backwards in the same direction as the ray and superimposed. Here, three views are used to illustrate the technique (a), and the final result is shown (b).


CT



High pass
filter

Filtered Backprojection. Each view (intensity vs. radius) is filtered and then stretched backwards in the same direction as the ray. Here, three views are used to illustrate the technique (a), and the final result is shown (b).

CT



I: Image

$I(x,y,z)$ denotes intensity value at pixel location x,y,z

Note also that whatever you see on the left is right part of the body!

Part 1 – Introduction

1.2.1 Structural imaging

1.2.1.3 Structural MRI



MRI – Magnetic resonance imaging


High spatial resolution (millimeter), 3D

Acceptable temporal resolution (second) – for functional imaging

Very versatile modality (multiple types of images, multiple phenomena)

Not radiation





MRI Siemens Magnetom Trio/TIM of the CENIR
Centre for neuroimaging research, Paris




The signal

- It depends on the physico-chemical organization of water within tissues (mobility, chemical liaisons)
- It is produced through excitation at a specific frequency (resonance) and recorded at the return at equilibrium (relaxation)
- It is located in space through additional magnetic fields (field gradients)

The signal

- It is directly related to
 - The magnetic moment of hydrogen nuclei (protons) that are within tissues (in particular, water)
 - To their properties within an external magnetic field B_0




From (Kastler and Vetter)

- The magnetic moment μ of each proton placed into a static magnetic field B_0
 - orients itself (parallel/anti-parallel)
 - describe a movement of precession at the following angular speed (Larmor relationship)


$$\omega_0 = \gamma \cdot B_0$$

Where B_0 is the external (static) magnetic field and gamma is the gyromagnetic ratio (specific to each nucleus, here protons)





From (Kastler and Vetter)



The sum of magnetic moments μ of all protons is oriented along B_0




Excitation/relaxation

- Excitation by a coil which creates a magnetic field B_1 orthogonal to B_0
 - B_1 is a rotating magnetic field with frequency ω_0 (in order to create resonance)
- Rotation of M_0 around the direction B_1 (resonance). Creation of a component M_{xy}
- Return to equilibrium: relaxation -> production of the MR signal






Décroissance de M_{xy}




Repousse de M_z

Relaxation T1



Relaxation T2



Obtaining an image

- Spatial coding (x, y, z) relies on the use of magnetic field gradients (G_x, G_y, G_z), with the same direction as B_0 but which amplitude varies linearly along directions x, y et z
- It allows to “control” the frequencies of precession at each point, as a function of the magnetic field in this point






Fig. 6-3. Le gradient de champ magnétique se superpose au champ magnétique principal B_0 , les modifications de champ produites se rajoutant ou se retranchant de B_0 .

Le champ résultant B croît linéairement dans la direction du gradient (centré par rapport à B_0). Par conséquent, la fréquence de Larmor croît aussi de manière linéaire, proportionnellement au champ résultant B : on a maintenant $\omega = \gamma \cdot B$. Schéma : J.-P. Dillenseger.

Axial slices



Coronal slices




Sagittal slices



Anatomical MRI

T1-weighted MRI


Resolution $\sim 1\text{mm} \times 1\text{mm} \times 1\text{mm}$




Diffusion MRI

- Brownian motion
- Allow to study the structure of tissues at the microscopic level (cell membranes, fibers...)
- In the brain: allows studying the white matter connections
- Other applications: heart, muscle...


Diffusion MRI



Diffusion MRI




Neurons and axons




BOSTON



NEW YORK




White matter




Diffusion MRI

	Isotropic, unrestricted diffusion	Isotropic, restricted diffusion	Anisotropic, restricted diffusion
Diffusion Trajectory			
(free water)	(random barriers present)	(coherent axonal bundle)	
Diffusion Ellipsoid			
Diffusion Tensor	$\begin{bmatrix} D & 0 & 0 \\ 0 & D & 0 \\ 0 & 0 & D \end{bmatrix}$	$\begin{bmatrix} D_{eff} & 0 & 0 \\ 0 & D_{eff} & 0 \\ 0 & 0 & D_{eff} \end{bmatrix}$ $D_{eff} < D$	$\begin{bmatrix} D_{xx} & D_{xy} & D_{xz} \\ D_{yx} & D_{yy} & D_{yz} \\ D_{zx} & D_{zy} & D_{zz} \end{bmatrix}$

Diffusion MRI



Gradient direction



DWI z


DWI x

DWI y


(From Cauchi and Yamamoto)

Diffusion MRI


12 directions




42 directions



162 directions




642 directions



(From Le Bihan et al, 2002)

Diffusion MRI


$$\begin{bmatrix} D_{xx} & D_{xy} & D_{xz} \\ D_{yx} & D_{yy} & D_{yz} \\ D_{zx} & D_{zy} & D_{zz} \end{bmatrix} \xrightarrow{\text{blue arrow}} \Lambda = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix} = R \cdot D \cdot R^T.$$




(From Mukerjee et al, 2008)

Diffusion MRI


$$\Lambda = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix} = R \cdot D \cdot R^T.$$



Mean diffusivity



Fractional anisotropy



Fiber Orientation




$$D_{av} = \frac{\lambda_1 + \lambda_2 + \lambda_3}{3} = \text{trace}(D)/3$$

$$FA = \frac{\sqrt{(\lambda_1 - \lambda_2)^2 + (\lambda_2 - \lambda_3)^2 + (\lambda_3 - \lambda_1)^2}}{\sqrt{2} \sqrt{\lambda_1^2 + \lambda_2^2 + \lambda_3^2}}$$

(From Mukerjee et al, 2008)


Tractography

- Macroscopic estimate of the trajectories of white matter fiber bundles



(From Johansen-Berg et al, 2009)

Diffusion MRI



(From Tournier et al, 2011)

Part 1 – Introduction

1.2.2 Functional imaging

1.2.2.1 PET

PET – Positron Emission Tomography

- 1970's
- Uses radioactive compounds which emit positrons (produced by a **cyclotron**)
- Short radioactive half-life
- **Many types of compounds** → multiple functions can be explored
- **Spatial resolution** ≈ 5 mm
- **Quantitative measure**
- **Expensive**



- Intra-venous injection of a **vector**, a molecule chosen for its interest regarding the phenomenon one aims to study
- Marked by a **radioactive** atom

VECTOR




ISOTOPE

^{18}F : $T_{1/2} = 109 \text{ mn}$

^{11}C : $T_{1/2} = 20 \text{ mn}$

^{15}O : $T_{1/2} = 2 \text{ mn}$

- Intra-venous injection of a **vector**, a molecule chosen for its interest regarding the phenomenon one aims to study
- Marked by a **radioactive atom**




^{18}F : $T_{1/2} = 109 \text{ mn}$


^{11}C : $T_{1/2} = 20 \text{ mn}$

^{15}O : $T_{1/2} = 2 \text{ mn}$


- Intra-venous injection of a **vector**, a molecule chosen for its interest regarding the phenomenon one aims to study
- Marked by a **radioactive** atom





- Intra-venous injection of a **vector**, a molecule chosen for its interest regarding the phenomenon one aims to study
- Marked by a **radioactive atom**



PET






Injection of a radioactive tracer → measures the number of disintegrations ↔ concentration of the tracer




Multiple uses depending on the tracer


- Study brain metabolism
- Studying neurotransmission, inflammation, amyloid plaques...



Normal brain FDG PET



Imaging amyloid plaques in Alzheimer's disease



Part 1 – Introduction


1.2.2 Functional imaging

1.2.2.2 Functional MRI


Functional MRI

Direct measure of brain activity
BOLD (Blood Oxygen Level Dependent)

control



activation



oxy-hemoglobin HbO_2


deoxygenated hemoglobin Hb
(paramagnetic, T_2^*)

↓ [Hb / HbO_2]

↓ T_2^*

↑ signal



Functional MRI



Right hand motion

Motor activations

- **Somatotopy**



Activ 99 - CEA, PSL

Versatile technique

- Anatomical MRI: tissues and structures
- Diffusion MRI: connections, fibers
- Functional MRI: brain activity

Started a revolution in neuroscience

Application to multiple organs: brain, heart, knee...

Part 1 – Introduction

1.2.2 Functional imaging



1.2.2.3 EEG/MEG


EEG/MEG

- **Surface recording of brain activity**
 - Electrical activity
 - Electro-encephalography - EEG
 - Magnetic activity
 - Magneto-encephalography - MEG

Neuronal electrophysiology

- **Pyramidal neuron**
 - Large cell
 - Organized in parallel in macro-assemblies
 - Superimposition of electrical currents
 - The result is detectable at the surface of the scalp





Electric potential

- 1 million de neurones created a potential of around $100\mu\text{V}$

Neurons also produce magnetic fields!
Magnetoencephalography (MEG)

Instrumentation: EEG



1945



1955

Instrumentation: EEG


- More sensors
 - Up to 256
- High temporal resolution
 - < 1 millisecond



MicroMed




Electrical
Geodesics



NeuroScan

Instrumentation: MEG




Part 1 – Introduction


1.3 Main tasks in medical image computing

Classification

Assign a label to the image




Normal



Tumor

Classification – Diagnosis / Prognosis




Supervised learning

Classification

Examples:


- Differential diagnosis of neurodegenerative diseases
 - Patient presenting with cognitive deficits, what is the underlying cause?
- Predicting the future clinical state
 - Patient presenting with subtle cognitive decline at baseline, will he have lost autonomy in 5 years ?

Clustering – Discover disease subtypes




Clustering

- Different symptoms for a given biological cause
- Different biological causes for a given set of symptoms




Find homogeneous subtypes of diseases

Regression – Predict clinical scores



Regression

Example : predict age from brain MRI




	BLUP mean	SVM	6-layer CNN	ResNet	Inception V1	Ensemble prediction
MAE (SE)	5.32 (0.19)	5.31 (0.18)	4.18 (0.16)	4.02 (0.15)	3.82 (0.14)	3.46 (0.13)
$ \rho $	0.32	0.58	0.25	0.24	0.41	0.32


Classification / Regression – typical approaches

Typical approaches: 3D extensions of networks used for natural image classification

3D CNN




ResNet



Alternatives: other classification/regression approaches with pre-extracted features

- SVM
- Random forests
- LASSO
- ...

Classification




Normal



Tumor

Detection


Locate an object in the image (lesion, tumor...)



Tumor

Segmentation


Delineate “objects” in images: tissues, anatomical structures, lesions...



Source: Zheng et al, The Volume of Hippocampal Subfields in Relation to Decline of Memory Recall Across the Adult Lifespan, 2018

Registration


- Definition: put two images in spatial correspondence



[Source Tang et al]

Registration

- Problem setting




Registration

- Intra-subject, intra-modality
 - Deformation: in general rigid
 - Similarity: simple criterion (e.g. sum of square differences)


Registration

- Intra-subject, intra-modality, longitudinal
 - Deformation: non-linear


Same patient at
different time
points (e.g.
18 months
after)



M0




M18



Registration

- Intra-subject, inter-modality
 - Deformation: rigid, affine or non-linear
 - Similarity: complex criterion (e.g. mutual information)

Brain imaging




PET




MRI

Image source: (Mangin, 1995)

Thoracico-abdominal
imaging



CT



PET

Image source (Camara et al., 2007)

Registration

- Inter-subject, intra-modality
 - Deformation: non-linear
 - Account for inter-individual variability






Image source: Mangin
et al

Image reconstruction



Source: <https://de.medical.canon/wp-content/uploads/sites/17/2016/08/Metal-Artefact-Reduction-in-CT-SEMAR.pdf>


Image enhancement - Denoising



Hellier, TMI 2008

Denoising of
brain MRI

Image enhancement – Super-resolution




O. Oktay et al. IEEE TMI 2017

Source: Daniel Rueckert, Imperial College London

Image synthesis

Data augmentation



Source: Bermudez et al, Learning Implicit Brain MRI
Manifolds with Deep Learning, SPIE MI, 2018

Image translation

Synthesizing one modality from another

MRI to CT





Fig. 4: *From left to right* Input MR image, synthesized CT image, reference real CT image, and absolute error between real and synthesized CT image.

Source: Wolterink et al, Deep MR to CT synthesis using unpaired data, 2017

Tasks



Source: Daniel Rueckert, Imperial College London